Clinical characteristics of urinary tract endometriosis: A one-year national series of 232 patients from 31 endometriosis expert centers (by the FRIENDS group)

Elise Froc, Gil Dubernard, Sofiane Bendifallah, Eric Hermouet, Chrystele Rubod-Dit-Guillet, Michel Canis, Sophie Warembourg, Francois Golfier, Arnaud Fauconnier, Horace Roman, et al.

► To cite this version:

Elise Froc, Gil Dubernard, Sofiane Bendifallah, Eric Hermouet, Chrystele Rubod-Dit-Guillet, et al.. Clinical characteristics of urinary tract endometriosis: A one-year national series of 232 patients from 31 endometriosis expert centers (by the FRIENDS group). *European Journal of Obstetrics & Gynecology and Reproductive Biology*, 2021, 264, pp.155-161. 10.1016/j.ejogrb.2021.06.018 . hal-03482577

HAL Id: hal-03482577

<https://hal.sorbonne-universite.fr/hal-03482577>

Submitted on 16 Dec 2021

HAL is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers.

L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.

Clinical characteristics of urinary tract endometriosis: a one-year national series of 232 patients from 31 endometriosis expert centers (by the FRIENDS group).

Authors

Elise FROC^a, Gil DUBERNARD^a, Sofiane BENDIFALLAH^b, Eric HERMOUET^c, Chrystele RUBOD-DIT-GUILLET^d, Michel CANIS^e, Sophie WAREMBOURG^a, Francois GOLFIER^f, Arnaud FAUCONNIER^g, Horace ROMAN^h, Charles-André PHILIP^{*a}

Affiliations:

^a Department of Gynaecologic and Oncologic Surgery and Obstetrics, Hôpital de la Croix Rousse, Hospices Civils de Lyon, Université Lyon 1, France. elise.froc@gmail.com, gil.dubernard@chu-lyon.fr, sophie.warembourg@chu-lyon.fr, charles-andre.philip@chu-lyon.fr

^b Department of Gynaecology and Obstetrics, Tenon University Hospital, Assistance Publique des Hôpitaux de Paris (AP-HP), Faculté de Médecine Sorbonne Université, Institut Universitaire de Cancérologie (IUC), France, sofiane.bendifallah@aphp.fr

^c Department of Gynaecologic Surgery and Obstetrics, Clinique Sud Vendée, Fontenay le Compte, France, drerichermouet44@gmail.com

^d Gynaecology Surgery Department, CHU Lille, F-59000, Lille, France; Lille University, Faculty of Medicine, F-59000, Lille, France, chrystele.rubod@chru-lille.fr

^e CHU Clermont-Ferrand, CHU Estaing, Pôle gynécologie obstétrique et reproduction humaine - Assistance médicale à la procréation - CECOS, Clermont-Ferrand, France, mcanis@chu-clermontferrand.fr

^f Department of Gynaecologic and Oncologic Surgery and Obstetrics, Centre Hospitalier
Universitaire Lyon Sud, Hospices Civils de Lyon, Université Lyon 1, France,
francois.golfier@chu-lyon.fr

^g Department of Gyneacology and Obstetrics, CHI de Poissy St Germain en Laye, F-78303
Poissy, France Arnaud.Fauconnier@ght-yvelinesnord.fr

^h Endometriosis Center, Clinique Tivoli-Ducos, Bordeaux, France,
Horace.roman@gmail.com

Corresponding author: Charles-André PHILIP

- Address: Gynecology department, Croix-Rousse University Hospital, 103 Grande Rue de la
Croix-Rousse, 69004 Lyon, France.

- e-mail: charles-andre.philip@chu-lyon.fr

- phone number: +33472001564

37 **Highlights :**

- 38 • a series of 232 patients suffering from urinary tract endometriosis describing their
39 symptoms and the most frequent locations.
- 40 • Ureteral endometriosis was significantly associated with other deep invasive
41 endometriosis lesions
- 42 • Ureteral endometriosis was more often asymptomatic than bladder locations

43 **Key Word :** Bladder endometriosis-Diagnosis of urinary tract endometriosis-Ureteral
44 endometriosis;

45

46

47 **Word count of the text :** 3000

48 **Word count of the abstract :** 248

49 **Ethical approval :** after consulting independent ethics committee, no Institutional Review
50 Board approval was required for this study.

51 **Author Disclosure Statement:** The authors report no conflict of interest related to the
52 present study

53

ABSTRACT

(247 Words)

Objective. To review prospectively the clinical characteristics of patients suffering from urinary tract endometriosis (UTE) in France, in 2017.

Study Design. We conducted a prospective observational multicenter study including women managed surgically for UTE in 31 French endometriosis expert centers (FRIENDS group) from January 1, 2017 to December 31, 2017. We distinguished patient with isolated bladder endometriosis ("IBE") or isolated ureteral endometriosis ("IUE") and patients associating both locations (mixed locations "ML"). Surgeons belonging to FRIENDS group enrolled patients by filling a 24 items questionnaire the day of the surgery and 6 weeks later. Data on the locations of UTE, preoperative assessment, urinary symptoms and associated pelvic locations were collected in a single anonymized database.

Results. A total of 232 patients from 31 centers were included. IBE concerned 82 patients (35.3%), IUE 126 patients (54.4%) ML 24 patients (10.3%). 111 patients reported urinary symptoms (47.8%). IUE was more often asymptomatic than the rest of the locations (59.5% versus 43.3%, OR 1.92, $p = 0.017$). Associated deep infiltrating endometriosis (DIE) lesions were found in 193 patients (83.1%). IUE was significantly associated with other DIE lesions (82.5% versus 66%, OR 2.4, $p = 0.006$), particularly with rectum or sigmoid nodules (57.1% versus 36.8%, OR 2.3, $p = 0.002$) and retrocervical space nodules (31.7% versus 19.8%, OR 1.9, $p = 0.05$).

Conclusion. Our study reports the second largest series of patients operated from a UTE and shows that ureteral location seems more frequent, less symptomatic and more frequently associated to other DIE locations than bladder endometriosis.

INTRODUCTION

Deep infiltrating endometriosis (DIE) is defined as the infiltration by ectopic endometrial glands under the peritoneum for more than 5 mm. The most frequent locations are the retrocervical space, uterosacral ligaments, the ovaries, the bowel wall and the urinary tracts (1,2). Urinary tract endometriosis (UTE) is a rare location of the disease, which affects 1.2 to 6% of women (1–5). The incidence increase in populations of patients undergoing surgery for DIE (6). The bladder is usually reported as the most common location of UTE and may be present in up to 85% of cases (22, 37). The ureter is less often affected, with a prevalence ranging from 0.5 to 10% of UTE locations (7,8). Bladder endometriosis (BE) is responsible for symptoms of bladder irritation such as voiding dysfunction, polyuria, urgency, hematuria and urinary incontinence (9–15). Symptoms appear to be related to the location and size of the nodule (16).

The definition of ureteral endometriosis (UE) remains controversial. According to Nezhat *et al.*, extrinsic and intrinsic form of UE endometriosis should be differentiated. Intrinsic endometriosis, which is defined by the involvement of the ureter wall may affect up to 20% of patients with UTE (17). Conversely, extrinsic UE is **defined by an involvement of the ureter by an endometriosis nodule from adjacent structures(17)**. The incidence and characteristics of this pathology remains difficult to evaluate.

The objective of this study was to perform a prospective review of the main characteristics of patients with UTE in France during a one year period..

MATERIAL AND METHOD

Population of the study

We performed a multicentric prospective study on a population of patients managed surgically for deeply invasive UTE in France, from January 1, 2017 to December 31, 2017. All surgeons affiliated with the FRIENDS group (French coloRectal Infiltrating ENDometriosis Study group), were invited to prospectively and consecutively include all of their patients corresponding to the inclusions criteria, who underwent a surgery for UTE. The FRIEND group was founded in 2015 and consists of a network of advanced endometriosis surgeons in France. The aim was to foster shared experiences and collect routine data on surgery to improve DIE management between gynecologist surgeons interested in endometriosis (6).

In this study, patients were selected by their surgeon prior to surgery each time a surgical treatment for UTE was planned. Inclusion criteria were patients with suspected UTE who had at least one lesion of the bladder or ureter, regardless of if they were isolated or congruent with each other or congruent with another endometriotic lesion. Exclusion criteria included patients under age 18 and surgery limited to ureterolysis performed as part of the dissection for the resection of DIE or extra urinary endometriosis with absence of a bladder or ureteral lesion.

Definition of UTE locations and surgeries

We defined UTE as endometriosis involving the bladder, the ureter or both. BE (bladder endometriosis) was defined as a deep infiltration of the bladder including detrusor muscle, with or without mucosae infiltration according to Chapron *et al.* definition (18). Ureteral endometriosis (UE) was defined as any situation where there existed a compression or distortion of the ureter by ureteral endometriosis nodule, either with or without ureteral dilatation or hydronephrosis (17). Ureteral dilatation was defined as an abnormal ureteral diameter measured up to 4 mm in size (19). Hydronephrosis was defined as an abnormal renal pelvis measured up to 10mm in size (19). Ureterolysis was defined as a dissection of the ureter from the endometriosis nodule until extracting all macroscopic endometriosis tissue and total liberation of the ureter, according to Nezhat *et al* and Uccella *et al* definition (17,20). As

explained previously, ureterolysis for the protection of the ureters prior to the resection of nodules not involving the bladder and ureters (DIE, uterosacral ligaments, rectal endometriosis, etc.) were excluded in this study.

Population subgroups

In our study, we distinguished five overlapping subgroups depending on the considered variable. Some analysis included nodules confined to one urinary location either bladder (isolated bladder endometriosis “IBE”) or the ureters (isolated ureteral endometriosis “IUE”). In other part we considered all bladder lesions either associated or not with ureteral involvement (bladder endometriosis “BE”) and all ureteral lesions either associated or not with a bladder involvement (ureteral endometriosis “UE”). A final subgroup including a combination of the mixed lesions (involving both ureter and bladder) was designated "mixed locations" (ML).

Standardized questionnaire and data collection

A 24-items questionnaire was sent to surgeons to be filled right after surgery. Patient demographic data were recorded (age at endometriosis diagnosis and surgery, parity, gravidity, history of surgery), as well as symptoms (urologic and non-urologic), pre-operative management (medical treatment, imaging examinations), location of urinary tract nodule or other locations of DIE, surgical procedure performed, and operative route. A specific part of the questionnaire concerned complications, which were defined as any adverse event in relation to the surgery, up to 6 weeks post operatively, according to the modified Clavien-Dindo classification (21). The questionnaires were anonymized and returned by surgeons at the end of the study period. Data were collected by a unique investigator (E.F.) in an anonymous database. Due to word constraints, data concerning surgical management and complication will be reported in a specific publication.

Ethic consideration

The National Data Protection Authority (Commission Nationale de l'Informatique et des Libertés) approved the study. All data were fully de-identified before analysis. Our work complied with French statutes and regulations, which authorize epidemiologic surveys without advance approval of an ethics committee. Our survey involved no intervention and is thus excluded from the French statute on biomedical research (Huriet-Serusclat law, dated December 20, 1998). We complied with all French statutes concerning data about the subjects, confidentiality, and restrictions.

Statistical analysis

Data were collected in a Microsoft Excel for Windows files (Microsoft®, Redmond, Washington, USA). Median, mean, standard deviations were calculated directly in Microsoft Excel files. Odds ratios, relative risks, 95% confidence intervals and all statistical analysis were performed using “R: A Language and Environment for Statistical Computing” (R Core Team® 2008 by the R Foundation for Statistical Computing Vienna, Austria). Univariate analysis: Wilcoxon rank sum test was used to look for association with continuous data. Fisher exact test was used to assess correlation and risk associated with binomial data. Data were considered significant if p-value was below 5%. Each odds ratio and relative risk was calculated comparing the proportion of “positive” and “negative” value of each considered variable using the rest of our population of UTE patients as the reference population and not a whole endometriosis or healthy control population.

RESULTS

Population and patient's characteristics

A total of 232 patients with UTE were included in our study from January 1, 2017 to December 31, 2017. Among the members of the FRIENDS group, 31 centers participated in the study, 23 (74.2%) were from University Hospitals and 8 (25.8%) from private structures. The median number of inclusions by department was 5 and varied between 1 to 27 patients.

Surgical history of endometriosis was reported in 76 patients (32.7%) which involved a urological location in 23 patients (9.9%). The patients with IBE has a mean age of 32.6 years (Standard Deviation (SD) = 5.56) against 34.4 years (SD = 7.05) for IUE ($p=0.08$).

There was no significant difference for the mean age of first symptoms of urinary symptoms between IBE and IUE (30.7; SD 5.31 versus 31.4 SD 8.42 years, $p=0.46$). The mean time between apparition of the first urinary symptoms and the diagnosis of UTE was 0.7 years (SD 1.46) for IBE and 1.1 years for IUE (SD = 5.84, $p= 0.63$). There was no significant difference between IBE and IUE for the mean time between diagnosis of endometriosis and surgery (1.06 versus 1.68 years, respectively, $p=0.33$), neither for the mean time between the first urinary symptoms and surgery (1.75 versus 2.33 years respectively, $p = 0.85$). The existence of a preoperative ureteral dilatation or specific urinary symptoms did not influence the mean time between diagnosis and surgical management. Patients' characteristics were summarized in table 1.

Location of urinary lesions

Among the 232 patients with UTE, 126 presented an IUE, 82 an IBE and 24 a ML. In total, 150 patients presented a ureteral involvement associated or not with a bladder involvement (Ureteral Endometriosis: UE), 134 were unilateral (89.3%) and 16 were bilateral (10.6%). Similarly, 106 patients (47.7%) presented a bladder involvement associated or not with a ureteral involvement (Bladder Endometriosis: BE). Among BE lesions, 31 accounted for the bladder base (29.2%), 70 the bladder dome (66%), 8 the trigone (7.5%) including 3 patients (2.8%) which presented several bladder lesions (association of nodule of the bladder base and dome). (Table 1)

Urinary symptoms

Among the 232 patients, 111 (47.8%) presented at least one urinary symptom. Fifty-one of them had IUE (45.9%), 49 IBE (44.1%) and 11 ML (9.9%). The urinary symptoms reported were voiding dysfunction (15.9%), cystalgia (21.1%), polyuria (6.0%), urinary incontinence (0.9%), urgency (1.7%), macroscopic hematuria (5.6%), and recurrent cystitis (3.9%). The

presence of cystalgia, polyuria and hematuria were more frequent in cases of IBE than in other locations (46.3% versus 7.3%, OR = 10.9, $p < 0.001$, 12.2% versus 2.6%, OR = 5.07, $p = 0.007$, and 12.1% versus 2%, OR = 6.81, $p = 0.002$ respectively).

IUE were more often asymptomatic than other locations (59.5% versus 43.3, OR 1.92, $p = 0.017$). However, the presence of lower back pain and renal colic were more frequent in IUE cases compared to other patients (30.1% versus 6.6%, OR 6.11, $p < 0.001$ and 4.8% versus 0%, OR = $+\infty$, $p = 0.03$ respectively) (Table 2).

IBE was significantly associated with the presence of cystalgia, polyuria and hematuria as compared to other locations (OR 10.9 $p < 0.001$, OR 5.07, $p = 0.007$, and OR 6.18, $p = 0.002$ respectively). IUE was more frequent in asymptomatic patients (OR 1.92, $p = 0.017$).

Preoperative assessment

Before surgery, 172 patients (74.1%) underwent a pelvic MRI, 48 (20.7%) a pelvic ultrasound, 58 (25%) a CT scan and 50 (21.5%) a cystoscopy. Nineteen renal scintigraphy (8.1%) were performed (Table 1).

CT scan and renal scintigraphy were significantly associated to the presence of IUE (OR 2.05, $p = 0.02$ and OR $+\infty$, $p < 0.001$ respectively), whereas a cystoscopy was significantly associated with the presence of IBE (OR 3.02, $p < 0.001$). Among the patients presenting IUE, 12 underwent a preoperative cystoscopy, which was performed as part of the insertion of a Double-J stent (Table 2a, b). The existence of a hydronephrosis was significantly associated with the performance of a renal scintigraphy (OR = 13.9, $p < 0.001$).

Associated pelvic locations

One hundred and ninety three patients (83.1%) presented at least one pelvic lesion associated with urological involvement. The most frequent associated locations were the digestive tract (111, 47.8%), the uterosacral ligaments (95, 40.9%), endometriomas (60, 25.7%) and the retrocervical space (61, 26.3%).

IUE was significantly associated with the presence of other pelvic lesions, compared to the rest of the urological locations (82.5% versus 66%, OR 2.4, $p = 0.006$) and in particular with Gastrointestinal nodules (57.1% versus 36.8%, OR 2.3, $p = 0.002$) and retrocervical space nodules (31.7% versus 19.8%, OR 1.9, $p = 0.05$). IUE and ML were not significantly associated with other pelvic locations (Table 2 and 3).

High urinary tract dilatation and preoperative Double-J stent

Data regarding to ureteral dilatations and hydronephrosis were available for 164 patients (70.6%) and 144 patients (62%) respectively. Among them, a preoperative ureteral dilatation was reported in 68 patients (41.4%) whereas a hydronephrosis was present in 20 patients (13.9%). Ureteral dilatation or hydronephrosis were more commonly found in iUE cases compared to the rest of the locations (respectively 57.1% versus 18.1%, OR 6.0 and 17.8% versus 5.7%, OR = 3.6). Four isolated bladder lesions were however associated with a ureteral dilatation and affected the bladder base in one case, the trigone in one case and the bladder dome in two cases.

The insertion of a Double-J Stent before or during surgery was reported in 107 patients (46%) and concerned 66 patients with IUE (52.4%), 22 patients with IBE (26.8%) and 19 patients with ML (79.1%). It was significantly associated with the presence of a ureteral lesion whether it was isolated (52.4% for IUE versus 39.2% for BE, OR = 1.74, $p < 0.04$) or associated with bladder lesion (79.1% for UE versus 22.6% for IBE, OR = 3.57 $p < 0.001$) (Table 2 and 3).

DISCUSSION

We report the results of a prospective longitudinal study, which included 232 patients managed surgically for UTE, in 31 French endometriosis center throughout the year 2017. To our knowledge, this is the second largest documented series of patients with UTE. Indeed, Ceccaroni *et al*/ recently reported a single center experience on 160 patients with UE and 264 patients with BE over a 10 years period (21, 22).

We report 82 patients (35.3%) with IBE, 126 (54.5%) with IUE and 24 (10.3%) with ML, accounting for a total of 150 ureteral lesions (UE) and a total of 106 bladder lesions (BE). The incidence of UTE in our series varied from 8% to 27.7% depending on the center. According to Nezhat *et al.*, UTE concerns from 1% to 6% of patients suffering from endometriosis (17). In the series of Gabriel *et al.*, the prevalence of UTE was to 19.5% and up to 52.6% in the series of Knabben *et al.* (10,11).

We observed a higher rate of ureteral lesions than bladder lesions in this series (64.6% for UE versus 45.7% for BE, ratio = 1.41), while Macagnanno *et al.* reported in their review a 85% rate of bladder lesions versus 9% of ureteral lesions (7). However, our results are consistent with those of Knabben *et al.*, which reported a higher rate of ureteral lesions (94.6%) than bladder lesions (14.3%) (11). This variation in UTE repartition could be related to the existence of different definitions in UE and BE in the literature. Indeed, some authors only consider intrinsic nodule as real UE, which is defined by Nezhat *et al.* as an infiltration of ureteral wall by an endometriotic nodule. Conversely, Nezhat *et al.* also describe an extrinsic UE, which correspond to a compression or distortion of the pelvic ureter by a peritoneal nodule or an adjacent structure(17). In their study, Knabben *et al.* regrouped these two entities into UE definition (11). Similarly, we both included endometriosis nodule from the ureteral wall as well as compression of the ureter by an endometriosis nodule from an adjacent anatomical structure. However, by consensus with the whole FRIENDS group, ureterolysis performed as a prerequisite for resection of an endometrioma, a gastrointestinal nodule or utero sacral ligaments were excluded to avoid an overestimation of extrinsic ureteric involvement, which could be the case in some series (6). These distinctions raise the problem of the need for a common definition for UE, which is currently heterogenic in the literature. A new objective definition of UE could implicate pre-operative imaging.

We reported bilateral lesions in 10.7% of UTE patients, while ML accounted for 24 patients (10.3%). These results are in agreement with the study of Cavaco-Gomez *et al.* which reported

a rate of 10.6% bilateral UE and Chapron *et al.* which found a similar rate (9.3%) of mixed urinary tract locations (18,24).

We reported that UE was associated with less urinary symptoms than IBE (41.3% versus 59.8%, RR = 1.45, p = 0.008), while it was associated with “unspecific” endometriosis symptoms. Data from the literature confirm that UE remains asymptomatic in nearly 50% of the cases (17,25). Cavaco-Gomez *et al.* reported also that UE was rarely associated with urinary symptoms (34.7%) but frequently associated with unspecific symptoms (dysmenorrhea in 81.4% of cases, pelvic pain in 70.2% of cases and dyspareunia in 66.4% of cases). These findings can be justified by the high association between UE and other DIE locations, which could explain these unspecific symptoms.(24). In our series, lower back pain was much more frequent in IUE or UE than for the rest of the population (30.1%, RR = 4.57, p<0.001 and 26.7, RR = 4.35, p<0.001 respectively). History of repetitive renal colics was only found in 6 cases of IUE (4.8%, p = 0.03). This association between lower back pain as also been reported in the literature, by Bosev *et al.* (11% of patients) and by Frenna *et al.* (9.2% of patients) (26,27).

Conversely, we report that BE is significantly associated with numerous urinary symptoms. This association was also reported by Fauconnier *et al.* in 2002, confirming the high specificity of urinary symptoms for the diagnosis of BE (9). In a series assessing urinary symptoms in patients with posterior DIE isolated or associated with BE, Panel *et al.* also reported an association between the existence of cystalgia and the presence of BE (90% versus 45%, p = 0.024) (14). Villa *et al.* also confirmed that location (bladder base) and size of BE lesion were related to the presence of this symptom (16).

In our study, 83% of UTE patients also presented associated DIE lesions; in particular IUE was strongly associated with gastrointestinal endometriosis. This association had already been described in the first prospective study of the FRIENDS group carried out on 1135 patients undergoing surgery for rectal and sigmoid endometriosis, which reported 13.4% of associated ureteral lesions (6).Based on 18 studies regrouping 668 patients with UE, Cavaco Gomez *et*

304 *al.* also found 58.8% of rectal lesions associated with UTE (24). In is monocentric study
305 Ceccaroni *et al*/report associated bowel nodule in 53% of BE and in 75.6% in case of UE. (21,
306 22) We also reported that retrocervical space lesions and parametrium nodules were strongly
307 associated with IUE. These data illustrate from our point of view the importance of
308 systematically looking for UE in patients presenting a posterior DIE and especially in case of
309 rectum or sigmoid involvement.

310 CONCLUSION

311 Our study describes the main clinical characteristic of urinary tract endometriosis patient in
312 France, in 31 gynecological surgical departments over a one-year period. Schematically, we
313 distinguished two different populations of patients with their own characteristics: Patients with
314 UE are usually slightly older, often asymptomatic and present more frequently associated
315 posterior DIE lesions. Conversely, BE is associated with specific urinary symptoms (polyuria,
316 cystalgia, voiding dysfunction, hematuria) and presents less often concomitant DIE lesions.
317 The FRIENDS group, which has twice shown its ability to organize national prospective
318 studies, should continue on its path to improve the understanding of the surgical management
319 of endometriosis.

320

Acknowledgment:

The authors would like to thank the following member of the FRIENDS group, who also participated to the study : C. CHAPRON, T. HEBERT, S. DOUVIER, L. FILIPUZZI, P. DESCAMPS, A. WATRELOT, A. AGOSTINI, J. BLANCHOT, C.M. ROGER, L. FRIEDERICH, P. COLLINET, P. VON THEOBALD, A. LETOHIC, M. MIMOUNI, A. MEPHON, T. GAUTHIER, N. HABIB, A. HOST, B.COTTE, R. RAMANAH, G. TRIOPON, F. GUILIBERT, F. AFCHIN, C. HOCKE, E.LAROUZEE, B.DOUSSET, A.PIVANO, E.BEAUVILLE, and B.RESCH. We also thank Mrs. Isabelle CAIN for her help with English editing.

REFERENCES

1. Chapron C, Fauconnier A, Vieira M, Barakat H, Dousset B, Pansini V, et al. Anatomical distribution of deeply infiltrating endometriosis: surgical implications and proposition for a classification. *Hum Reprod.* 2003 Jan 1;18(1):157–61.
2. Scioscia M, Bruni F, Ceccaroni M, Steinkasserer M, Stepniewska A, Minelli L. Distribution of endometriotic lesions in endometriosis stage IV supports the menstrual reflux theory and requires specific preoperative assessment and therapy. *Acta Obstet Gynecol Scand.* 2011 Feb;90(2):136–9.
3. Berlanda N, Vercellini P, Carmignani L, Aimi G, Amicarelli F, Fedele L. Ureteral and Vesical Endometriosis: Two Different Clinical Entities Sharing the Same Pathogenesis. *Obstet Gynecol Surv.* 2009 Dec;64(12):830–42.
4. Kołodziej A, Krajewski W, Dołowy Ł, Hirnle L. Urinary Tract Endometriosis. *Urol J.* 2015 Sep 4;12(4):2213–7.
5. Nezhat C, Paka C, Gomaa M, Schipper E. Silent Loss of Kidney Secondary to Ureteral Endometriosis. *JSLS.* 2012;16(3):451–5.
6. Roman H, FRIENDS group (French coloRectal Infiltrating ENDometriosis Study group). A national snapshot of the surgical management of deep infiltrating endometriosis of the rectum and colon in France in 2015: A multicenter series of 1135 cases. *J Gynecol Obstet Hum Reprod.* 2017 Feb;46(2):159–65.
7. Maccagnano C, Pellucchi F, Rocchini L, Ghezzi M, Scattoni V, Montorsi F, et al. Diagnosis and treatment of bladder endometriosis: state of the art. *Urol Int.* 2012;89(3):249–58.
8. Yohannes P. Ureteral Endometriosis. *J Urol.* 2003 Jul;170(1):20–5.

9. Fauconnier A, Chapron C, Dubuisson J-B, Vieira M, Dousset B, Bréart G. Relation between pain symptoms and the anatomic location of deep infiltrating endometriosis. *Fertil Steril*. 2002 Oct;78(4):719–26.
10. Gabriel B, Nassif J, Trompoukis P, Barata S, Wattiez A. Prevalence and Management of Urinary Tract Endometriosis. *Urology*. 2011 Dec;78(6):1269–74.
11. Knabben L, Imboden S, Fellmann B, Nirgianakis K, Kuhn A, Mueller MD. Urinary tract endometriosis in patients with deep infiltrating endometriosis: prevalence, symptoms, management, and proposal for a new clinical classification. *Fertil Steril*. 2015 Jan;103(1):147–52.
12. Kumar S, Tiwari P, Sharma P, Goel A, Singh J, Vijay M, et al. Urinary tract endometriosis: Review of 19 cases. *Urol Ann*. 2012;4(1):6.
13. Leone Roberti Maggiore U, Ferrero S, Salvatore S. Urinary incontinence and bladder endometriosis: conservative management. *Int Urogynecology J*. 2015 Jan;26(1):159–62.
14. Panel P, Huchon C, Estrade-Huchon S, Le Tohic A, Fritel X, Fauconnier A. Bladder symptoms and urodynamic observations of patients with endometriosis confirmed by laparoscopy. *Int Urogynecology J*. 2016 Mar;27(3):445–51.
15. Vercellini P, Frontino G, Pietropaolo G, Gattei U, Daguati R, Crosignani PG. Deep endometriosis: definition, pathogenesis, and clinical management. *J Am Assoc Gynecol Laparosc*. 2004 May;11(2):153–61.
16. Villa G, Mabrouk M, Guerrini M, Mignemi G, Montanari G, Fabbri E, et al. Relationship between site and size of bladder endometriotic nodules and severity of dysuria. *J Minim Invasive Gynecol*. 2007 Sep;14(5):628–32.

17. Nezhat C, Falik R, McKinney S, King LP. Pathophysiology and management of urinary tract endometriosis. *Nat Rev Urol*. 2017 May 3;14(6):359–72.
18. Chapron C, Bourret A, Chopin N, Dousset B, Leconte M, Amsellem-Ouazana D, et al. Surgery for bladder endometriosis: long-term results and concomitant management of associated posterior deep lesions. *Hum Reprod*. 2010 Apr 1;25(4):884–9.
19. Potenta SE, D’Agostino R, Sternberg KM, Tatsumi K, Perusse K. CT Urography for Evaluation of the Ureter. *Radiogr Rev Publ Radiol Soc N Am Inc*. 2015 Jun;35(3):709–26.
20. Uccella S, Cromi A, Casarin J, Bogani G, Pinelli C, Serati M, et al. Laparoscopy for ureteral endometriosis: surgical details, long-term follow-up, and fertility outcomes. *Fertil Steril*. 2014 Jul;102(1):160-166.e2.
21. Dindo D, Demartines N, Clavien P-A. Classification of Surgical Complications: A New Proposal With Evaluation in a Cohort of 6336 Patients and Results of a Survey. *Ann Surg*. 2004 Aug;240(2):205–13.
22. Ceccaroni M, Clarizia R, Ceccarello M, De Mitri P, Roviglione G, Mautone D, et al. Total laparoscopic bladder resection in the management of deep endometriosis: “take it or leave it.” Radicality versus persistence. *Int Urogynecology J*. 2019 Sep 7;
23. Ceccaroni M, Ceccarello M, Caleffi G, Clarizia R, Scarperi S, Pastorello M, et al. Total Laparoscopic Ureteroneocystostomy for Ureteral Endometriosis: A Single-Center Experience of 160 Consecutive Patients. *J Minim Invasive Gynecol*. 2019;26(1):78–86.
24. Cavaco-Gomes J, Martinho M, Gilabert-Aguilar J, Gilabert-Estélls J. Laparoscopic management of ureteral endometriosis: A systematic review. *Eur J Obstet Gynecol Reprod Biol*. 2017 Mar;210:94–101.

25. Seracchioli R, Mabrouk M, Manuzzi L, Guerrini M, Villa G, Montanari G, et al. Importance of retroperitoneal ureteric evaluation in cases of deep infiltrating endometriosis. *J Minim Invasive Gynecol.* 2008 Aug;15(4):435–9.
26. Bosev D, Nicoll LM, Bhagan L, Lemyre M, Payne CK, Gill H, et al. Laparoscopic Management of Ureteral Endometriosis: The Stanford University Hospital Experience With 96 Consecutive Cases. *J Urol.* 2009 Dec;182(6):2748–52.
27. Frenna V, Santos L, Ohana E, Bailey C, Wattiez A. Laparoscopic management of ureteral endometriosis: Our experience. *J Minim Invasive Gynecol.* 2007 Mar;14(2):169–71.

Table 2: Univariate analysis depending on the urological location (Ureteral or bladder location)

Variable	N=	Bladder endometriosis ^a (82)			Ureteral endometriosis ^b (126)		
		YES *	OR	p-value	YES *	OR	p-value
Surgical history for endometriosis	76	16(19.5)	<u>0.36</u>	0.001	53(42.4)	2.78	0,001
<i>Urinary symptoms</i>							
Asymptomatic patients	121	33 (40.2)	<u>0.47</u>	0.008	75(59.5)	1.92	0,017
Lower back pain	45	5 (6.1)	<u>0.18</u>	<0.001	38(30.1)	6.11	<0,001
Cystalgia	49	38 (46.3)	10.9	<0.001	5(4)	<u>0.06</u>	<0,001
Dysuria	37	17 (20.7)	1.7	0.18	13(10.3)	<u>0.39</u>	0,012
Polyuria	14	10 (12.2)	5.07	0.007	4(3.2)	<u>0.31</u>	0,05
Hematuria	13	10 (12.1)	6.81	0.002	1(0.8)	<u>0.06</u>	<0,001
Recurrent cystitis	9	6 (7.3)	3.87	0.07	1(0.8)	<u>0.09</u>	0,01
Renal colics	6	0 (0)	0	0.09	6(4.8)	+∞	0,03
<i>Preoperative imaging</i>							
CT	58	9 (10.1)	<u>0.25</u>	<0.001	39(31)	2.05	0.02
MRI	172	64 (78)	1.38	0.34	95(75.4)	1.15	0.65
Pelvic or urinary ultrasound	48	16 (19.5)	0.89	0.86	29(23)	1.36	0.41
Cystoscopy	40	28 (34.1)	3.02	<0.001	12(9.5)	<u>0.19</u>	<0.001
Renal scintigraphy	19	0 (0)	<u>0</u>	<0.001	18(14.3)	+∞	<0.001
<i>High urinary tract dilatation</i>							
Ureteral dilatation	68	4 (8.2)	<u>0.07</u>	<0.001	56(57.1)	6.0	<0.001
Hydronephrosis	19	0 (0)	<u>0</u>	0.001	16(17.8)	3.60	0.04
<i>Associated lesions</i>							
Digestive tract	111	26 (31.7)	<u>0.35</u>	<0.001	72(57.1)	2.29	0,002
Adenomyosis	19	4 (4.9)	0.46	0.21	14(0.11)	2.52	0,08
Parametrium	27	4 (4.9)	<u>0.28</u>	0.018	23(18.2)	5.69	<0,001
USL	95	31 (37.8)	0.82	0.48	56(44.4)	1.37	0,28
Endometrioma	60	14 (17.1)	<u>0.46</u>	0.02	39(31)	1.81	0,07
Retrocervical space	61	15 (18.3)	<u>0.51</u>	0.04	40(31.7)	1.88	0,05
<i>Operative route</i>							
Laparoscopy	169	62(75.6)	1.24	0.54	93(73.8)	1.11	0,76
Laparotomy (total)	25	6(7.6)	0.54	0.27	12(9.5)	0.75	0.53
Laparoconversion	8	3(3.7)	1.1	1	1(0.8)	<u>0.11</u>	0.02
<i>Urological surgery</i>							
Double-J Stent	107	22(26.8)	<u>0.28</u>	<0.001	66(52.4)	1.74	0.04
Collaboration with urologist	85	24 (29.3)	0.60	0.08	46(36.8)	1	1
Ureteral resection	28	0(0)	<u>0</u>	<0.001	24(19)	6.00	<.001
Ureteroneocystostomy	23	0(0)	<u>0</u>	<0.001	20(15.9)	6.48	<0.001
Ureteroureterostomy	5	0(0)	0	0,16	4(3.2)	3.44	0.38
Ureterolysis	128	14(17.1)	<u>0.07</u>	<0.001	95(75,4)	6.78	<0.001
Partial cystectomy	100	77(93.9)	<u>85.0</u>	<0.001	0(0)	0	<0.001
<i>Associated surgical procedure</i>							
Retrocervical space	64	15(18.3)	<u>0.46</u>	0.02	41(32.5)	1.74	0,07
Digestive tract	109	25(30.1)	<u>0.34</u>	<0.001	72(57.1)	2.48	<0,001
Unilateral ovariectomy	21	2(2.4)	<u>0.17</u>	0.008	16(12,7)	2.94	0,039
Bilateral ovariectomy	14	1(1.2)	<u>0.13</u>	0.02	9(7.1)	1.55	0.44
Hysterectomy	40	6(7.3)	<u>0.27</u>	0.003	30(23,8)	3.00	0,004
Parametrium nodules	24	3(3.7)	<u>0.23</u>	0.01	21(16.7)	6.87	<0,001
Uterine adenomyosis	3	3(3.7)	+∞	0.04	0(0)	<u>0</u>	0,09
<i>Complications</i>							
Total	61	17(20.7)	0.63	0.16	35(27.8)	1.18	0,65
Urological	34	11(13.4)	0.77	0.57	18(14.3)	0.81	0,58
Second surgery	14	4(4.9)	0.72	0.77	9(7.1)	1.55	0,58

Notes: OR = odds ratio / USL: Uterosacral ligaments * n (%): ^a: comparison of isolated bladder endometriosis (YES) with the rest of urological locations (NO) / ^b: comparison of isolated ureteral endometriosis (YES) with the rest of urological locations (NO)

Table 3: Characteristics of the population depending on the urological location (Mixed location)				
		Mixed Urinary Endometriosis ^c (24)		
Variable	n	YES *	OR	p-value
Surgical history for endometriosis	76	7(29.1)	0.82	0.84
<i>Urinary symptoms</i>				
Symptomatic patients	111	11(45.8)	0.91	1
Lower back pain	45	2(8.3)	0.34	0.18
Cystalgia	49	6(25)	1.27	0.60
Dysuria	37	7(29.2)	2.44	0.07
Polyuria	14	0(0)	-	0.37
Hematuria	13	2(8.3)	1.62	0.54
Recurrent cystitis	9	2(8.3)		0.23
Renal colic	6	0(0)	-	1
<i>Preoperative imaging</i>				
CT scan	58	10(41.7)	2.38	0.07
MRI	172	13(54.2)	0.36	0.02
Pelvic or urinary ultrasound	48	3(12.5)	0.51	0.43
Cystoscopy	40	10(41.7)	3.0	0.01
Renal scintigraphy	19	1(4.2)	0.46	0.70
<i>High urinary tract dilatation</i>				
Ureteral dilatation	68	8(47)	1.29	0.62
Hydronephrosis	19	3(23.1)	2.14	0.38
<i>Associated lesions</i>				
Digestive tract	111	13(54.1)	1.33	0.53
Adenomyosis	19	1(4.2)	0.45	0.70
Parametrium	27	0(0)	0	0.09
USL	95	8(33.3)	0.69	0.51
Endometrioma	60	7(29.2)	1.20	0.80
Retrocervical space	61	6(25)	0.92	1
<i>Operative route</i>				
Laparoscopy	169	14(58.3)	0.47	0.14
Laparotomy (total)	25	7(29.2)	4.34	0.007
Laparoconversion	8	4(16.7)	10.2	0.005
<i>Urological surgery</i>				
Double-J stent	107	19(79.1)	5.18	<0.001
Association with urologist	85	15(62.5)	3.26	<0.001
Ureteral resection	28	4(16.7)	1.53	0.50
Ureteroneocystostomy	23	3(12.5)	1.34	0.71
Ureteroureterostomy	5	1(4.2)	2.21	0.42
Ureterolysis	128	19(79.2)	3.45	0.02
Partial cystectomy	100	23(95.8)	39.1	<0.001
<i>Associated surgical procedure</i>				
Retrocervical space	64	8(34.8)	1.36	0.48
Digestive	109	12(50)	1.14	0.83
Unilateral ovariectomy	21	3(12.5)	1.5	0.46
Bilateral ovariectomy	14	4(16.7)	3.96	0.04
Hysterectomy	40	4(16.7)	0.96	1
Parametrium nodules	24	0(0)	0	0.15
Uterine adenomyosis	3	0(0)	0	1
<i>Complications</i>				
Total	61	9(37.5)	1.8	0.22
Urological	34	7(29.1)	2.54	0.07
Second surgery	14	1(4.2)	0.65	0.20
Notes: RR = relative risk of presenting the variable according to the presence (YES) or not (NO) of a mixed location (ML). USL: Uterosacral ligaments * n (%): comparison between mixed locations (ML) (+) and the rest of urological locations (-)				