



HAL
open science

Letter to the editor from Behar-Cohen, et al: “The Cortisol Response of Male and Female Choroidal Endothelial Cells: Implications for Central Serous Chorioretinopathy”

Francine Behar-Cohen, Frédéric Jaisser, Min Zhao

► **To cite this version:**

Francine Behar-Cohen, Frédéric Jaisser, Min Zhao. Letter to the editor from Behar-Cohen, et al: “The Cortisol Response of Male and Female Choroidal Endothelial Cells: Implications for Central Serous Chorioretinopathy”. *Journal of Clinical Endocrinology and Metabolism*, 2021, 10.1210/clinem/dgab908 . hal-03498919

HAL Id: hal-03498919

<https://hal.sorbonne-universite.fr/hal-03498919v1>

Submitted on 21 Dec 2021

HAL is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers.

L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.

Letter to the editor from Behar-Cohen, et al: “The Cortisol Response of Male and Female Choroidal Endothelial Cells: Implications for Central Serous Chorioretinopathy”

Francine Behar-Cohen^{1,2}, Frédéric Jaisser¹, Min Zhao¹

¹Centre de Recherche des Cordeliers, Sorbonne Université, Université de Paris, Inserm, 75006 Paris, France.

²Assistance Publique-Hôpitaux de Paris, Hôpital Cochin Ophthalmopole, 75014 Paris, France

Corresponding author:

Francine Behar-Cohen (MD, PhD)

Francine.behar-cohen@uparis.fr, francine.behar@gmail.com

Disclosure statement: FBC, MZ and FJ have been cited as inventors on 2 patents belonging to public institutions, on the use of mineralocorticoid receptor antagonists for the treatment of macular edema and choroidal neovascularization.

Dear Doctor Brinks,

We have read with interest the paper of Joost Brinks et al 2021 Sep 21;dgab670. doi: 10.1210/clinem/dgab670. Online ahead of print, entitled The Cortisol Response of Male and Female Choroidal Endothelial Cells: Implications for Central Serous Chorioretinopathy (CSCR).

In this paper, the authors failed to identify the mineralocorticoid receptor (MR) in endothelial cells from the choroid of human eyes collected around 24 hrs after death.

The authors missed the important information, published 10 years ago by the team who developed the antibody they have used, that the mineralocorticoid receptor (MR) is unstable. Indeed, the team of Gomez-Sanchez explains that to use the antibody and detect MR, “prompt handling and inactivation of enzymes of degradation is required....Processing the sample immediately after harvest in a buffer containing protease and phosphatase inhibitors preserves the MR intact...” (1). It was thus expected that MR would be degraded when eyes are enucleated from donors after 24 hrs of death.

The MR immunohistochemistry images shown in this paper supports that MR might have been degraded since the MR signal in the retina is extremely low, without any signal in ganglion cells, in retinal endothelial cells and very scarce signal in few cells the inner retina, which is very different with what has been found by several other authors (2–4). There is no MR signal in the nuclei of any cells in the choroid and no MR signal in the nuclei of retinal pigment epithelial cells either, which is very different from previous results obtained on fresh tissues and cells(5,6). More importantly, the positive MR control do not show any signal in the nuclei, which could indicate methodological issues.

The authors claim that there is no animal model that recapitulate human pachychoroid phenotype but they omitted to cite the P1hMR mouse model (mouse that overexpresses human MR) (6) and the NAS (nephrectomy aldosterone salt) models (7) that show all features of human eyes with pachychoroid.

Due to the lack of similitude between the cellular model used in this study and the human physiologic conditions regarding steroid receptors, this cellular model is useful to decipher the transcriptional regulations of cortisol upon binding to the glucocorticoid receptor (GR) but cannot bring any information regarding human physiological or pathological conditions. The

conclusions on the predominance of GR pathway activation by cortisol in human choroidal endothelial eyes with CSCR is thus misleading.

In addition, it is very unlikely that the venous vasodilation observed in patients with CSCR could result from a direct activation of corticoid receptors in choroidal endothelial cells because vasodilation in the choroid is under neural control (8). Indeed, many other cells expressing both GR and MR could influence the vascular dynamics such as pericytes, smooth muscle cells, microglia and more importantly neural cells from the autonomous system, that controls the choroidal blood flow regulation.

The analysis of the cortisol response of endothelial cells alone does not allow to draw any conclusion on the involvement of MR or GR pathway in CSCR.

Reference

1. **Gomez-Sanchez CE, Warden M, Gomez-Sanchez MT, Hou X, Gomez-Sanchez EP.** Diverse immunostaining patterns of mineralocorticoid receptor monoclonal antibodies. *Steroids* 2011;76(14):1541–1545.
2. **Nitta E, Hirooka K, Tenkumo K, Fujita T, Nishiyama A, Nakamura T, Itano T, Shiraga F.** Aldosterone: a mediator of retinal ganglion cell death and the potential role in the pathogenesis in normal-tension glaucoma. *Cell Death Dis* 2013;4:e711.
3. **Wilkinson-Berka JL, Tan G, Jaworski K, Miller AG.** Identification of a retinal aldosterone system and the protective effects of mineralocorticoid receptor antagonism on retinal vascular pathology. *Circ. Res.* 2009;104(1):124–133.
4. **Bousquet E, Zhao M, Ly A, Leroux Les Jardins G, Goldenberg B, Naud M-C, Jonet L, Besson-Lescure B, Jaisser F, Farman N, De Kozak Y, Behar-Cohen F.** The aldosterone-mineralocorticoid receptor pathway exerts anti-inflammatory effects in endotoxin-induced uveitis. *PLoS ONE* 2012;7(11):e49036.
5. **Daruich A, Matet A, Dirani A, Bousquet E, Zhao M, Farman N, Jaisser F, Behar-Cohen F.** Central serous chorioretinopathy: Recent findings and new physiopathology hypothesis. *Prog Retin Eye Res* 2015;48:82–118.
6. **Canonica J, Zhao M, Favez T, Gelizé E, Jonet L, Kowalczyk L, Guegan J, Le Menuet D, Viengchareun S, Lombès M, Pussard E, Arsenijevic Y, Behar-Cohen F.** Pathogenic Effects of Mineralocorticoid Pathway Activation in Retinal Pigment Epithelium. *Int J Mol Sci* 2021;22(17):9618.
7. **Canonica J, Mehanna C, Bonnard B, Jonet L, Gelize E, Jais J-P, Jaisser F, Zhao M, Behar-Cohen F.** Effect of acute and chronic aldosterone exposure on the retinal pigment epithelium-choroid complex in rodents. *Exp. Eye Res.* 2019;187:107747.
8. **Reiner A, Fitzgerald MEC, Del Mar N, Li C.** Neural control of choroidal blood flow. *Prog Retin Eye Res* 2018;64:96–130.