



HAL
open science

Computing roadmaps in unbounded smooth real algebraic sets I: connectivity results

Rémi Prébet, Mohab Safey El Din, Éric Schost

► **To cite this version:**

Rémi Prébet, Mohab Safey El Din, Éric Schost. Computing roadmaps in unbounded smooth real algebraic sets I: connectivity results. 2022. hal-03508000v2

HAL Id: hal-03508000

<https://hal.sorbonne-universite.fr/hal-03508000v2>

Preprint submitted on 7 Mar 2022 (v2), last revised 7 Jun 2023 (v3)

HAL is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers.

L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.

Computing roadmaps in unbounded smooth real algebraic sets I: connectivity results

Rémi PRÉBET¹, Mohab SAFEY EL DIN¹, and Éric SCHOST²

¹Sorbonne Université, LIP6 CNRS UMR 7606, Paris, France

²University of Waterloo, David Cheriton School of Computer Science, Waterloo ON,
Canada

March 7, 2022

Abstract

Answering connectivity queries in real algebraic sets is a fundamental problem in effective real algebraic geometry that finds many applications in e.g. robotics where motion planning issues are topical. This computational problem is tackled through the computation of so-called *roadmaps* which are real algebraic subsets of the set V under study, of dimension at most one, and which have a connected intersection with all semi-algebraically connected components of V . Algorithms for computing roadmaps rely on statements establishing connectivity properties of some well-chosen subsets of V , assuming that V is bounded.

In this paper, we extend such connectivity statements by dropping the boundedness assumption on V . This exploits properties of so-called *generalized polar varieties*, which are critical loci of V for some well-chosen polynomial maps.

1 Introduction

Let \mathbf{Q} be a real field of real closure \mathbf{R} and let \mathbf{C} be its algebraic closure (one can think about \mathbb{Q} , \mathbb{R} and \mathbb{C} instead, for the sake of understanding) and let $n \geq 0$ be an integer. An algebraic set of $V \subset \mathbf{C}^n$ defined over \mathbf{Q} is the solution set in \mathbf{C}^n to a system of polynomial equations with coefficients in \mathbf{Q} . A real algebraic set defined over \mathbf{Q} is the set of solutions in \mathbf{R}^n to a system of polynomial equations with coefficients in \mathbf{Q} . It is also the real trace $V \cap \mathbf{R}^n$ of an algebraic set $V \subset \mathbf{C}^n$. Real algebraic sets have finitely many connected components [7, Theorem 2.4.4.]. Counting these connected components [14, 19] or answering connectivity queries over $V \cap \mathbf{R}^n$ [17] finds many applications in e.g. robotics [8, 12].

Following [8, 10], such computational issues are tackled by computing a real algebraic subset of $V \cap \mathbf{R}^n$, defined over \mathbf{Q} , which has dimension at most one and a connected intersection with all connected components of V and contains the input query points. Such a subset has been called by Canny in [8] a *roadmap* of V .

The effective construction of roadmaps, given a defining system for V , relies on connectivity statements which allow one to define real algebraic subsets of $V \cap \mathbf{R}^n$, of smaller dimension than the one of $V \cap \mathbf{R}^n$ and that have a connected intersection with the connected components of $V \cap \mathbf{R}^n$. Such existing statements in the literature make the assumption that V has finitely many singular points and $V \cap \mathbf{R}^n$ is bounded. In this paper, we focus on the problem of obtaining similar statements by dropping the boundedness assumption. We prove a new connectivity statement which generalizes the one of [16] to the unbounded case and will be used in a separate paper to obtain asymptotically faster algorithms for computing roadmaps. We start by recalling the state-of-the-art connectivity statement which allows us to introduce some material we need to state our main result.

State-of-the-art overview We start by introducing some terminology. Recall that an *algebraic set* $V \subset \mathbf{C}^n$ is the set of solutions of a finite system of polynomial equations. It can be uniquely decomposed into finitely many *irreducible components*. When all these components have the same dimension d , we say

that V is d -*equidimensional*. Those points $\mathbf{y} \in V$ at which the Jacobian matrix of a finite set of generators of its associated ideal have rank $n - d$ are called *regular* points and the set of those points is denoted by $\text{reg}(V)$. The others are called *singular* points; the set of singular points of V (its singular locus) is denoted by $\text{sing}(V)$ and is an algebraic subset of V . We refer to [18] for definitions and propositions about algebraic sets.

A *semi-algebraic set* $S \subset \mathbf{R}^n$ is the set of solutions of a finite system of polynomial equations and inequalities. We say that S is *semi-algebraically connected* if for any $\mathbf{y}, \mathbf{y}' \in S$, \mathbf{y} and \mathbf{y}' can be connected by a *semi-algebraic path* in S , that is a continuous semi-algebraic function $\gamma: [0, 1] \rightarrow S$ such that $\gamma(0) = \mathbf{y}$ and $\gamma(1) = \mathbf{y}'$. A semi-algebraic set S can be decomposed into finitely many *semi-algebraically connected components* which are semi-algebraically connected semi-algebraic sets that are both closed and open in S . Finally, for a semi-algebraic set $S \subset \mathbf{R}^n$, we denote by \bar{S} its closure for the Euclidean topology on \mathbf{R}^n . We refer to [4] and [7] for definitions and propositions about semi-algebraic sets and functions.

Let $0 \leq d \leq n$ and $V \subset \mathbf{C}^n$ be a d -equidimensional algebraic set such that $\text{sing}(V)$ is finite. For $1 \leq i \leq n$, let π_i be the canonical projection:

$$\pi_i: (\mathbf{y}_1, \dots, \mathbf{y}_n) \mapsto (\mathbf{y}_1, \dots, \mathbf{y}_i)$$

For a polynomial map $\varphi: \mathbf{C}^n \rightarrow \mathbf{C}^m$ a point $\mathbf{y} \in V$ is a *critical point* of φ if $\mathbf{y} \in \text{reg}(V)$ and $d_{\mathbf{y}}\varphi$ is not a submersion, that is

$$d_{\mathbf{y}}\varphi(T_{\mathbf{y}}V) \subsetneq \mathbf{C}^m,$$

where $d_{\mathbf{y}}\varphi$ is the differential of φ at \mathbf{y} . We will denote by $W^\circ(\varphi, V)$ the set of the critical points of φ on V . A *critical value* is the image of a critical point. We will note $K(\varphi, V) = W^\circ(\varphi, V) \cup \text{sing}(V)$. The points of $K(\varphi, V)$ are called the *singular points* of φ on V . Figure 1 show example of such critical loci.




Figure 1: Real trace of the critical locus on a sphere Z for: the projection on the first coordinate π_1 (left); the polynomial map associated to $x_1^2 + x_2^2 \in \mathbb{R}[x_1, x_2, x_3]$ (right).

For $1 \leq i \leq d$ we denote by $W(\pi_i, V)$ the i -th *polar variety* defined as the Zariski closure of the critical locus $W^\circ(\pi_i, V)$ of the restriction of π_i to V . Further, we extend this definition by considering $\varphi = (\varphi_1, \dots, \varphi_n) \subset \mathbf{Q}[x_1, \dots, x_n]$ and, for $1 \leq i \leq n$, the map

$$\varphi_i: \mathbf{C}^n \longrightarrow \mathbf{C}^i$$

$$\mathbf{y} \mapsto (\varphi_1(\mathbf{y}), \dots, \varphi_i(\mathbf{y})) \tag{1}$$

Following the ideas of [1, 2, 3] we denote similarly $W(\varphi_i, V)$ the i -th *generalized polar variety* defined as the Zariski closure of the critical locus $W^\circ(\varphi_i, V)$ of the restriction of φ_i to V . We recall below [15,

Theorem 14] (see also [6, Proposition 3.3] for a slight variant of it), making use of polar varieties to establish connectivity statements.

For $2 \leq i \leq d$, assume that the following holds:

- $V \cap \mathbf{R}^n$ is bounded;
- $W(\pi_i, V)$ is either empty or $(i - 1)$ -equidimensional and smooth outside $\text{sing}(V)$;
- $W(\pi_1, W(\pi_i, V))$ is finite;
- for any $\mathbf{y} \in \mathbf{C}^{i-1}$, $\pi_{i-1}^{-1}(\mathbf{y}) \cap V$ is either empty or $(d - i + 1)$ -equidimensional.

Let

$$K_i = W(\pi_1, W(\pi_i, V)) \cup \text{sing}(V) \quad \text{and} \quad F_i = \pi_{i-1}^{-1}(\pi_{i-1}(K_i)) \cap V.$$

Then, the real trace of $W(\pi_i, V) \cup F_i$ has a non-empty and semi-algebraically connected intersection with each semi-algebraically connected component of $V \cap \mathbf{R}^n$.

For the special case $i = 2$, this result has been originally proved in [8, 9]. A variant of this, again assuming $i = 2$, is given for general semi-algebraic sets in [10, 11].

In this paper, we focus on real algebraic sets. By dropping the restriction $i = 2$, the result in [15, Theorem 14] allows one more freedom in the choice of i in the design of roadmap algorithms to obtain better complexity. The rationale is as follows.

Restricting to $i = 2$, one expects (up to some linear change of variables or other technical manipulations) to retrieve a situation where $W(\pi_2, V)$ has dimension at most 1 and F_2 to have dimension $d - 1$ (see e.g. [15, Lemma 31]). To obtain a roadmap for $V \cap \mathbf{R}^n$ one is led to call recursively the roadmap algorithm, which applies [15, Theorem 14], on systems defining the F_i 's. Hence, the depth of the recursion is n . Besides, letting D be the maximum degree of input equations defining V , roughly speaking each recursive call requires $(nD)^{O(n)}$ arithmetic operations in \mathbf{Q} while the size of the input data grows by $(nD)^{O(n)}$ according to [15, Proposition 33]. Consequently, one obtains roadmap algorithms using $(nD)^{O(n^2)}$ arithmetic operations in \mathbf{Q} .

In [15], using a baby steps/giant steps strategy, the authors shown that one can take $i \simeq \sqrt{d}$ and then have a depth of the recursion $\simeq \sqrt{d}$. It is also proved that each recursive step needed to compute systems encoding K_i and F_i require at most $(nD)^{O(n)}$ arithmetic operations in \mathbf{Q} while the size of the input data grows by $(nD)^{O(n)}$. All in all, up to some technical details that we skip for the sake of conciseness, one obtains roadmap algorithms using $(nD)^{O(n\sqrt{n})}$ arithmetic operations in \mathbf{Q} . Finally in [16], it is shown how to apply [15, Theorem 14] with $i \simeq \frac{d}{2}$ so that the depth of the recursion becomes $\simeq \log_2(d)$. Hence, proceeding as in [15], an algorithm using $(nD)^{6n \log_2(d)}$ arithmetic operations in \mathbf{Q} is obtained in [16].

Dropping the boundedness assumption in this solving scheme is done in [5, 6] using infinitesimal deformation techniques. The algorithms proposed use respectively $(nD)^{O(n\sqrt{n})}$ and $(nD)^{O(n \log^2(n))}$ arithmetic operations in \mathbf{Q} . This induces a growth of intermediate data; also the obtained algorithm is not polynomial in its output size. Generalizing [16] to non-bounded situations requires a new connectivity statement dropping the boundedness assumption on $V \cap \mathbf{R}^n$. This is the main new result of this paper.

Main result We start with some notations and assumptions. Let $V \subset \mathbf{C}^n$ be an algebraic set defined over \mathbf{Q} and $d > 0$ be an integer. We say that V satisfies assumption (A) when

(A) V is d -equidimensional and its singular locus $\text{sing}(V)$ is finite.

Let Z be a subset of \mathbf{C}^n , $U \subset \mathbf{R}$ and $f \in \mathbf{R}[x_1, \dots, x_n]$. With a slight abuse of notation, we still denote by f the polynomial map $\mathbf{y} \in \mathbf{C}^n \mapsto f(\mathbf{y}) \in \mathbf{C}$. We write $Z_{|f \in U} = Z \cap f^{-1}(U) \cap \mathbf{R}^n$. In particular if $u \in \mathbf{R}$ we note

$$Z_{|f < u} = Z_{|f \in]-\infty, u[}, \quad Z_{|f \leq u} = Z_{|f \in]-\infty, u]} \quad \text{and} \quad Z_{|f = u} = Z_{|f \in \{u\}}.$$

Now, let $\varphi = (\varphi_1, \dots, \varphi_n) \subset \mathbf{Q}[x_1, \dots, x_n]$ and φ_i be the induced map defined as in (1) for $1 \leq i \leq n$. Recall that we say that a map $\psi: Y \subset \mathbf{R}^n \rightarrow Z \subset \mathbf{R}^m$ is a proper map if, for every closed (for Euclidean topology) and bounded subset $Z' \subset Z$, $\psi^{-1}(Z')$ is a closed and bounded subset of Y .

We say that φ satisfies assumption (P) when

(P) the restriction of the map φ_1 to $V \cap \mathbf{R}^n$ is proper and bounded from below.

We denote by $W_i = W(\varphi_i, V)$ the Zariski closure of the set of critical points of the restriction of φ_i to V . For $2 \leq i \leq d$, we say that (φ, i) satisfies assumption (B) when

(B₁) W_i is either empty or $(i - 1)$ -equidimensional and smooth outside $\text{sing}(V)$;

(B₂) for any $\mathbf{y} = (\mathbf{y}_1, \dots, \mathbf{y}_i) \in \mathbf{C}^i$, $V \cap \varphi_{i-1}^{-1}(\mathbf{y})$ is either empty or $(d - i + 1)$ -equidimensional.

Note that when B₁ holds, $\text{sing}(W_i)$ and critical loci of polynomial maps restricted to W_i are well-defined. Let $S \subset W(\varphi_1, W_i)$, we say that it satisfies assumption (C) when

(C₁) S is finite;

(C₂) S intersects every semi-algebraically connected component of $W(\varphi_1, W_i) \cap \mathbf{R}^n$.

Finally, using a construction similar to the one used in [15, Theorem 14], we let

$$K_i = W(\varphi_1, V) \cup S \cup \text{sing}(V) \quad \text{and} \quad F_i = \varphi_{i-1}^{-1}(\varphi_{i-1}(K_i)) \cap V.$$

Theorem 1.1. *For $2 \leq i \leq d$ and under the assumptions (A), (B), (C) and (P), the subset $W_i \cup F_i$ has a non-empty and semi-algebraically connected intersection with each semi-algebraically connected component of $V \cap \mathbf{R}^n$.*

This result is used in a forthcoming paper, to extend the algorithms of [16] for computing roadmaps to the case of unbounded inputs. Indeed, it allows one to design an algorithm for computing roadmaps on real algebraic sets whose real counterpart may not be bounded in kind of a straightforward way. Let us describe briefly how this would work. Let V be an equidimensional algebraic set; assume that $\text{sing}(V)$ is finite. Take

$$\varphi_1 = \sum_{k=1}^n (x_k - \mathbf{a}_k)^2 \quad \text{and for } 2 \leq j \leq n \quad \varphi_j = \sum_{k=1}^n \mathbf{b}_{j,k} x_k,$$

where $\mathbf{a} = (\mathbf{a}_1, \dots, \mathbf{a}_n) \in \mathbf{Q}^n$ and for $2 \leq j \leq n$, $\mathbf{b}_j = (\mathbf{b}_{j,1}, \dots, \mathbf{b}_{j,n}) \in \mathbf{Q}^n$. Then assumption (P) holds and according to [2, 3] for a wise choice of \mathbf{a} and \mathbf{b} one could hope to satisfy the dimensional properties of assumption (B).

Finally, constructing the set S by a sampling algorithm (for instance [4, Chap. 13]), one can apply Theorem 1.1 to V and φ . Hence, applying recursively this procedure and Theorem 1.1 to polynomial systems defining W_i and F_i one obtains a roadmap for $V \cap \mathbf{R}^n$. It should be noted that, since $F_i \cap \mathbf{R}^n$ is bounded (by assumption (P)), the algorithm of [16] directly computes a roadmap of it. Then it is enough to operate the recursive calls on the generalized polar varieties.

However, in regard of [2, 3] proving that assumption (B) holds for some generic choice of \mathbf{a} and \mathbf{b} needs more effort. It is the purpose of a future article, together with the full study of the algorithm described above.

Example 1. Let us apply the above process on an example. Let $V = \mathbf{V}(g) \subset \mathbf{C}^3$ where $g = x_1^3 + x_2^3 + x_3^3 - x_1 - x_2 - x_3 - 1 \in \mathbb{Q}[x_1, x_2, x_3]$. Since V is a hypersurface, it is 2-equidimensional and since $\text{sing}(V) = \emptyset$, V satisfies (A).

Let $\varphi = (x_1^2 + x_2^2 + x_3^2, x_1, x_2) \subset \mathbb{Q}[x_1, x_2, x_3]$. As the restriction of φ_1 to \mathbb{R}^n is the square of the Euclidean norm, (P) is satisfied. Since $2 \leq i \leq d$, necessarily $i = 2$. Then we will see later that one can write

$$W_2 = \mathbf{V}(f, 3x_2^2x_3 - 3x_2x_3^2 + x_2 - x_3).$$

One checks that W_2 is 1-equidimensional and has no singular point as well, so that $(\varphi, 2)$ satisfies (B₁). Let $K_2 = W^\circ(\varphi_1, W_2)$, it is a finite set of cardinal 11. Besides, for any $\alpha \in \mathbf{C}$,

$$V \cap \varphi_1^{-1}(\alpha) = \mathbf{V}(f, x_1^2 + x_2^2 + x_3^2 - \alpha)$$

is either empty or an equidimensional algebraic set of dimension 1. Therefore, $(\varphi, 2)$ satisfies (B). Finally, since $W^\circ(\varphi_1, W_2) \cap \mathbb{R}^3$ is a finite set, assumption (C) holds vacuously. Indeed the points stand here for the connected components.

In conclusion, by Theorem 1.1, $W_2 \cup F_2$ is a 1-roadmap of (V, \emptyset) . Figure 2 illustrates this example.




Figure 2: An illustration of Example 1. The real trace $V \cap \mathbb{R}^3$ is plotted twice as a grid. On the left, $W_2 \cap \mathbb{R}^3$ is represented as red lines, and the crosses represent all the real points of K_2 . Then, on the right, we replaced the points of K_2 by the fibers of $F_2 \cap \mathbb{R}^3$ (black lines), to repair the connectivity failures of $W_2 \cap \mathbb{R}^3$. In particular, $F_2 \cap \mathbb{R}^3$ connects the semi-algebraically connected components of $W_2 \cap \mathbb{R}^3$ that lie in the same semi-algebraically connected component of $V \cap \mathbb{R}^3$.

2 Preliminaries

Basic properties of algebraic sets Recall that given a finite set of polynomials $\mathbf{g} \subset \mathbf{C}[x_1, \dots, x_n]$ we denote by $V(\mathbf{g}) \subset \mathbf{C}^n$ the algebraic set defined by the vanishing locus of \mathbf{g} . For $\mathbf{y} \in \mathbf{C}^n$, we denote by $\text{Jac}_{\mathbf{y}}(\mathbf{g})$ the Jacobian matrix of \mathbf{g} evaluated at \mathbf{y} . Conversely, given an algebraic set $V \subset \mathbf{C}^n$, we denote by $\mathbf{I}(V)$ the *ideal of V* , that is the ideal of $\mathbf{C}[x_1, \dots, x_n]$ of polynomials vanishing on V . Such an ideal is finitely generated by Hilbert basis theorem.

Let $X \subset \mathbf{C}^n$ and $Y \subset \mathbf{C}^m$ be algebraic sets and $\mathbf{K} \subset \mathbf{C}$ be a subfield. A map $\alpha: X \rightarrow Y$ is a *regular map* defined over \mathbf{K} if there exists $(f_1, \dots, f_m) \in \mathbf{K}[x_1, \dots, x_n]$ such that $\alpha(\mathbf{y}) = (f_1(\mathbf{y}), \dots, f_m(\mathbf{y}))$ for all $\mathbf{y} \in X$. A regular map $\alpha: X \rightarrow Y$ is an *isomorphism* defined over \mathbf{K} if there exists a regular map $\beta: Y \rightarrow X$, defined over \mathbf{K} , such that $\alpha \circ \beta = \text{id}_Y$ and $\beta \circ \alpha = \text{id}_X$, where $\text{id}_Z: Z \rightarrow Z$ is the identity map on Z . We refer to [18] for further details on these notions.

The following result is straightforward.

Lemma 2.1. *Let $Y \subset \mathbf{C}^n$ and $Z \subset \mathbf{C}^m$ be two algebraic sets. Let $\alpha: Y \rightarrow Z$ be an isomorphism of algebraic sets defined over \mathbf{R} . Then the semi-algebraically connected subsets of $Y \cap \mathbf{R}^n$ and $Z \cap \mathbf{R}^m$ are in correspondence through α .*

Critical points of a polynomial map The following lemma from [16, Lemma A.2] provides an algebraic characterization of critical points.

Lemma 2.2 (Rank characterization). *Let $Z \subset \mathbf{C}^n$ be a d -equidimensional algebraic set and $\mathbf{g} = (g_1, \dots, g_p)$ be generators of $\mathbf{I}(Z)$. Let $\varphi: Z \rightarrow \mathbf{C}^m$ be a polynomial map, then the following holds.*

$$W^\circ(\varphi, Z) = \left\{ \mathbf{y} \in Z \mid \begin{array}{l} \text{rank}(\text{Jac}_{\mathbf{y}}(\mathbf{g})) = n - d \\ \text{and } \text{rank}(\text{Jac}_{\mathbf{y}}([\mathbf{g}, \varphi])) < n - d + m \end{array} \right\};$$

$$K(\varphi, Z) = \{ \mathbf{y} \in Z \mid \text{rank}(\text{Jac}_{\mathbf{y}}([\mathbf{g}, \varphi])) < n - d + m \}.$$

Let us present a direct consequence of this result, which gives a more effective criterion for the singular points of a polynomial map.

Let $\varphi = (\varphi_1, \dots, \varphi_n) \in \mathbf{C}[x_1, \dots, x_n]$ and φ_i be the deduced map defined as in (1) for $1 \leq i \leq n$.

Lemma 2.3. *Let $Z \subset \mathbf{C}^n$ be a d -equidimensional variety and \mathbf{g} be a finite set of generators of $\mathbf{I}(Z)$.*

Then for $1 \leq i \leq n$, $K(\varphi_i, Z)$ is the algebraic subset of Z defined by the vanishing of \mathbf{g} and the $(p+i)$ -minors of $\text{Jac}([\mathbf{g}, \varphi_i])$, where $p = n - d$.

Proof. One directly deduces from Lemma 2.2 that $K(\varphi_i, Z)$ is exactly the intersection of Z , the zero-set of \mathbf{g} , with the set of points $\mathbf{y} \in \mathbf{C}^n$ where $\text{rank}(\text{Jac}_{\mathbf{y}}([\mathbf{g}, \varphi_i])) < p + i$. But, by elementary linear algebra, the latter set is the zero-set of the $(p+i)$ -minors of $\text{Jac}([\mathbf{g}, \varphi_i])$, which are polynomial of $\mathbf{C}[x_1, \dots, x_n]$. \square

Definition 2.4 (Polar variety). Let $Z \subset \mathbf{C}^n$ be a d -equidimensional algebraic set, and let $1 \leq i \leq n$. We denote by $W(\varphi_i, Z)$ the Zariski closure of $W^\circ(\varphi_i, Z)$. It is called a *generalized polar variety* of Z . Remark that

$$W^\circ(\varphi_i, Z) \subset W(\varphi_i, Z) \subset K(\varphi_i, Z) \subset Z$$

by minimality of the Zariski closure. Hence $K(\varphi_i, Z) = W(\varphi_i, Z) \cup \text{sing}(Z)$ but the union is not necessarily disjoint.

3 Connectivity and critical values

In this section we consider for $n \geq 1$ an equidimensional algebraic set $Z \subset \mathbf{C}^n$ of dimension $d > 0$. We are going to prove two main connectivity results on the semi-algebraically connected components of Z through some polynomial map. These results, along with the idea of Morse theory, will be essential in the proof of Theorem 1.1. Most of the results presented here are generalizations of those present in [15, Section 3.] in the unbounded case, replacing projections by suitable polynomial maps.

3.1 Connectivity changes at critical values

The main result of this paragraph is to prove the following proposition which deals with the connectivity changes of semi-algebraically connected components when restricting close to singular values of a polynomial map.

Proposition 3.1. *Let $\varphi: \mathbf{C}^n \rightarrow \mathbf{C}$ a regular map defined over \mathbf{R} . Let $A \subset \mathbf{R}^k$ be a semi-algebraically connected semi-algebraic set, and $u \in \mathbf{R}$ and*

$$\gamma: A \rightarrow Z_{|\varphi \leq u} - (Z_{|\varphi = u} \cap K(\varphi, Z))$$

be a continuous semi-algebraic map. Then there exists a unique semi-algebraically connected component B of $Z_{|\varphi < u}$ such that $\gamma(A) \subset \overline{B}$.

Notation. In this paragraph we fix $\varphi: \mathbf{C}^n \rightarrow \mathbf{C}$ a regular (polynomial) map defined over \mathbf{R} . With a slight abuse of notation, a polynomial of $\mathbf{C}[x_1, \dots, x_n]$ associated to φ will be denoted equally.

We start by proving an extended version of [15, Lemma 6]. This can be seen as the foundation stone of all the connectivity results presented in this paper. Recall that an open Euclidean neighborhood of a point $\mathbf{y} \in \mathbf{R}^n$ is any subset of \mathbf{R}^n , that contains \mathbf{y} and is open for the Euclidean topology on \mathbf{R}^n .

Lemma 3.2. *Let $\mathbf{y} = (\mathbf{y}_1, \dots, \mathbf{y}_n)$ be in $Z \cap \mathbf{R}^n - K(\varphi, Z)$. Then, there exists a map $\alpha: Z \rightarrow \mathbf{C}^{n+1}$, such that the following holds :*

- a) *there exist open Euclidean neighborhoods $N' \subset \mathbf{R}^d$ of $\pi_d(\alpha(\mathbf{y}))$ and $N \subset \mathbf{R}^{n+1}$ of $\alpha(\mathbf{y})$, and there exists a continuous semi-algebraic map $\mathbf{f}: N' \rightarrow \mathbf{R}^{n+1-d}$ such that:*

$$\alpha(Z) \cap N = \{(z', \mathbf{f}(z')) \mid z' \in N'\};$$

- b) *$\alpha: Z \rightarrow \alpha(Z)$ is an isomorphism of algebraic sets defined over \mathbf{R} ;*

- c) *$\varphi \circ \alpha^{-1} = \pi_1$ on $\alpha(Z)$.*

Proof. Let $\mathcal{O}_{\mathbf{y}} \subset \mathbf{R}^n$ be an open Euclidean neighborhood of \mathbf{y} such that there exists $\mathbf{g} = (g_1, \dots, g_{n-d})$ in $\mathbf{C}[x_1, \dots, x_n]$, such that $Z \cap \mathcal{O}_{\mathbf{y}} = \mathbf{V}(\mathbf{g}) \cap \mathcal{O}_{\mathbf{y}}$. Such a $\mathcal{O}_{\mathbf{y}}$ is given by [7, Proposition 3.3.10] since $\mathbf{y} \in \text{reg}(Z)$. Moreover $\text{Jac}_{\mathbf{y}}(\mathbf{g})$ has full rank $n - d$ and since $\mathbf{y} \notin W(\varphi, Z)$ there exists a non-zero $(n - d + 1)$ -minor of $\text{Jac}_{\mathbf{y}}([\mathbf{g}, \varphi])$ by Lemma 2.3. Therefore, there exists a permutation σ of $\{1, \dots, n\}$ such that the matrix

$$\begin{bmatrix} \frac{\partial \mathbf{g}}{\partial x_{\sigma(i)}}(\mathbf{y}) \\ \frac{\partial \varphi}{\partial x_{\sigma(i)}}(\mathbf{y}) \end{bmatrix}_{d \leq j \leq n}$$

is invertible. Let x_0 be a new variable and define \mathbf{h} as the following finite subset of polynomials of $\mathbf{R}[x_0, x_1, \dots, x_n]$,

$$\mathbf{h} = (\tilde{\mathbf{g}}, \tilde{\varphi}) = (\mathbf{g}(\sigma^{-1} \cdot (x_1, \dots, x_n)), \varphi(\sigma^{-1} \cdot (x_1, \dots, x_n)) - x_0)$$

where $\tau \cdot (x_1, \dots, x_n) = (x_{\tau(1)}, \dots, x_{\tau(n)})$ for any permutation τ of $\{1, \dots, n\}$. Hence,

$$\mathbf{V}(\mathbf{h}) \cap (\mathbf{R} \times \mathcal{O}_{\mathbf{y}}) = \{(\varphi(\mathbf{y}), \sigma \cdot \mathbf{y}) \mid \mathbf{y} \in Z \cap \mathcal{O}_{\mathbf{y}}\} \subset \mathbf{R}^{n+1}.$$

According to the chain rule, for any $1 \leq j \leq n$ and $\mathbf{z} \in \mathbf{R}^n$,

$$\frac{\partial \tilde{\mathbf{g}}}{\partial x_j}(\varphi(\mathbf{z}), \mathbf{z}) = \frac{\partial \mathbf{g}}{\partial x_{\sigma(j)}}(\sigma^{-1} \cdot \mathbf{z}) \quad \text{and} \quad \frac{\partial \tilde{\varphi}}{\partial x_j}(\varphi(\mathbf{z}), \mathbf{z}) = \frac{\partial \varphi}{\partial x_{\sigma(j)}}(\sigma^{-1} \cdot \mathbf{z}).$$

Hence, for $\text{Jac}(\mathbf{f}, i)$ the Jacobian matrix of \mathbf{f} with respect to (x_{i+1}, \dots, x_n) , and $\tilde{\mathbf{y}} = (\varphi(\mathbf{y}), \sigma \cdot \mathbf{y})$,

$$\text{Jac}_{\tilde{\mathbf{y}}}(\mathbf{h}, d - 1) = \begin{bmatrix} \text{Jac}_{\tilde{\mathbf{y}}}(\tilde{\mathbf{g}}, d - 1) \\ \text{Jac}_{\tilde{\mathbf{y}}}(\tilde{\varphi}, d - 1) \end{bmatrix} = \begin{bmatrix} \frac{\partial \mathbf{g}}{\partial x_{\sigma(i)}}(\mathbf{y}) \\ \frac{\partial \varphi}{\partial x_{\sigma(i)}}(\mathbf{y}) \end{bmatrix}_{d \leq j \leq n},$$

which is invertible by assumption on σ .

Therefore, applying the semi-algebraic implicit function theorem [4, Th 3.30] to \mathbf{h} , there is an open Euclidean neighborhood $N' \subset \mathbf{R}^d$ of $(\varphi(\mathbf{y}), \mathbf{y}')$ where $\mathbf{y}' = (\mathbf{y}_{\sigma(\ell)}, 1 \leq \ell \leq d - 1)$, an open Euclidean neighborhood $N'' \subset \mathbf{R}^{n-d+1}$ of $\mathbf{y}'' = (\mathbf{y}_{\sigma(\ell)}, d \leq \ell \leq n)$ and a map $\mathbf{f} = (f_1, \dots, f_{n-d+1}) \in \mathcal{S}^\infty(N', N'')$ (since φ and the g_i 's are polynomials) such that:

$$\forall \mathbf{z} = (\mathbf{z}', \mathbf{z}'') \in N' \times N'', \quad [\mathbf{h}(\mathbf{z}) = 0 \iff \mathbf{z}'' = \mathbf{f}(\mathbf{z}')]]$$

Then, let $N = (N' \times N'') \cap (\mathbf{R} \times \sigma \cdot \mathcal{O}_{\mathbf{y}}) \subset \mathbf{R}^{n+1}$, the previous assertion becomes:

$$\{(\varphi(\mathbf{z}), \sigma \cdot \mathbf{z}) \mid \mathbf{z} \in Z\} \cap N = \{(\mathbf{z}', \mathbf{f}(\mathbf{z}')) \mid \mathbf{z}' \in N'\} \quad (2)$$

Finally, we claim that taking $\alpha: \mathbf{z} \in Z \mapsto (\varphi(\mathbf{z}), \sigma \cdot \mathbf{z})$ ends the proof. Indeed, by equation (2), assertion a) immediately holds since N' and N are Euclidean open neighborhood of $\pi_d(\alpha(\mathbf{y}))$ and $\alpha(\mathbf{y})$ respectively. Further, one checks that α is a Zariski isomorphism, of inverse σ^{-1} after projecting on the last n coordinates, which proves b). Finally, one sees that $\pi_1 \times \alpha = \varphi$ so that c) holds as well. \square

Remark. The previous lemma shows in particular that $Z \cap \mathbf{R}^n - K(\varphi, Z)$ is a Nash manifold (see [4, Section 3.4]) of dimension d i.e. which is locally \mathcal{S}^∞ -diffeomorphic to \mathbf{R}^d .

Lemma 3.3. *Let \mathbf{y} be in $Z \cap \mathbf{R}^n - K(\varphi, Z)$ and $u = \varphi(\mathbf{y})$. Then there exists an open Euclidean neighborhood $N(\mathbf{y})$ of \mathbf{y} such that the following holds:*

- a) $N(\mathbf{y})$ is semi-algebraically connected;
- b) $(Z \cap N(\mathbf{y}))|_{\varphi < u}$ is non-empty and semi-algebraically connected;
- c) $(Z \cap N(\mathbf{y}))|_{\varphi = u}$ is contained in $\overline{(Z \cap N(\mathbf{y}))|_{\varphi < u}}$.

This result is illustrated by Figure 3.




Figure 3: Illustration of Lemma 3.3 where $\varphi = \pi_1$, $u = \mathbf{y}_1$ and Z is isomorphic to $\mathbf{V}(x_1^2 + x_2^2 - 1) \times \mathbf{V}(x_2 + x_1^2)$. On the left, \mathbf{y} is not critical and one sees that it satisfies all the statements. On the right \mathbf{y} is critical, and $(Z \cap N(\mathbf{y}))_{|\pi_1 < \mathbf{y}_1}$ is disconnected. Note that in both cases, \mathbf{y}_1 is a critical value.

Proof. Let α, N', N and \mathbf{f} be obtained by applying Lemma 3.2. Let $\mathbf{F}: \mathbf{z}' \in N' \mapsto (\mathbf{z}', \mathbf{f}(\mathbf{z}')) \in N$. Let $\varepsilon > 0$ be such that

$$\mathcal{B} = \mathcal{B}(\pi_d(\alpha(\mathbf{y})), \varepsilon) \subset N' \subset \mathbf{R}^d$$

where $\mathcal{B}(\pi_d(\alpha(\mathbf{y})), \varepsilon)$ is the open ball of \mathbf{R}^d with radius ε and center $\pi_d(\alpha(\mathbf{y}))$. We claim that taking $N(\mathbf{y}) = \alpha^{-1}(\mathbf{F}(\mathcal{B}))$ is enough to prove the result.

First $\mathbf{F}(\mathcal{B})$ is open, semi-algebraic and semi-algebraically connected since \mathbf{F} is an open continuous map on \mathcal{B} . Then, by assumptions on α , together with Lemma 2.1, $\alpha^{-1}(\mathbf{F}(\mathcal{B}))$ is a semi-algebraically connected open neighborhood of \mathbf{y} . Hence $N(\mathbf{y})$ satisfies statement a).

Besides, remark that $\mathbf{F}(\mathcal{B}) \subset \alpha(Z)$, so that

$$(\alpha(Z) \cap \mathbf{F}(\mathcal{B}))_{|\pi_1 < u} = \mathbf{F}(\mathcal{B})_{|\pi_1 < u} = \mathbf{F}(\mathcal{B}_{|\pi_1 < u})$$

as $\pi_1(\mathbf{F}(\mathbf{z}')) = \pi_1(\mathbf{z}')$ for $\mathbf{z}' \in N'$. Since $\pi_1(\alpha(\mathbf{y})) = \varphi(\mathbf{y}) = u$, the semi-algebraic set $\mathcal{B}_{|\pi_1 < u}$ is non-empty and semi-algebraically connected (since \mathcal{B} is convex), and so is its image through \mathbf{F} by [4, Section 3.2]. But remark that for all $X \subset \mathbf{R}$,

$$(Z \cap N(\mathbf{y}))_{|\varphi \in X} = \alpha^{-1}((\alpha(Z) \cap \mathbf{F}(\mathcal{B}))_{|\pi_1 \in X}) = \alpha^{-1} \circ \mathbf{F}(\mathcal{B}_{|\pi_1 \in X}), \quad (3)$$

since $\varphi \circ \alpha^{-1} = \pi_1$. Therefore, by Lemma 2.1, $(Z \cap N(\mathbf{y}))_{|\varphi < u}$ is non-empty and semi-algebraically connected as claimed in statement b).

To prove assertion c), remark that $\mathcal{B}_{|\pi_1 = u}$ is contained in $\overline{\mathcal{B}_{|\pi_1 < u}}$, so that $\alpha^{-1} \circ \mathbf{F}(\mathcal{B}_{|\pi_1 = u})$ is contained in $\alpha^{-1} \circ \mathbf{F}(\overline{\mathcal{B}_{|\pi_1 < u}})$. Since \mathbf{F} and α^{-1} are continuous,

$$\alpha^{-1} \circ \mathbf{F}(\overline{\mathcal{B}_{|\pi_1 < u}}) \subset \overline{\alpha^{-1} \circ \mathbf{F}(\mathcal{B}_{|\pi_1 < u})}.$$

Finally by (3)

$$(Z \cap N(\mathbf{y}))_{|\varphi = u} \subset \overline{(Z \cap N(\mathbf{y}))_{|\varphi < u}}.$$

□




Figure 4: Illustration of Lemma 3.4 where $\varphi = \pi_1$, $u = \mathbf{y}_1$ and Z is isomorphic to $\mathbf{V}(x_1^2 + x_2^2 - 1) \times \mathbf{V}(x_2 + x_1^2)$. On the left \mathbf{y} is not critical and one sees that $\mathbf{y} \in \overline{B_{\mathbf{y}}}$ and $(Z \cap N(\mathbf{y}))|_{\pi_1^{-1}(\mathbf{y}_1)} \subset B_{\mathbf{y}}$. However on the right, \mathbf{y} is critical, and one observes that \mathbf{y} belongs to both $\overline{B_{\mathbf{y}}}$ and $\overline{B'_{\mathbf{y}}}$, and, in addition, that $(Z \cap N(\mathbf{y}))|_{\pi_1^{-1}(\mathbf{y}_1)}$ is not contained in any of these components. Note that in both cases, \mathbf{y}_1 is a critical value.

Lemma 3.4. *Let \mathbf{y} be in $Z \cap \mathbf{R}^n - K(\varphi, Z)$, let $u = \varphi(\mathbf{y})$ and let $N(\mathbf{y})$ as in Lemma 3.3. Then, there exists a unique semi-algebraically connected component $B_{\mathbf{y}}$ of $Z|_{\varphi < u}$ such that $\mathbf{y} \in \overline{B_{\mathbf{y}}}$. Moreover,*

$$(Z \cap N(\mathbf{y}))|_{\varphi < u} \subset B_{\mathbf{y}}.$$

This lemma is illustrated in Figure 4.

Proof. By the second item of Lemma 3.3, $(Z \cap N(\mathbf{y}))|_{\varphi < u}$ is non-empty and semi-algebraically connected. Then it is contained in a semi-algebraically connected component $B_{\mathbf{y}}$ of $Z|_{\varphi < u}$. Since the semi-algebraically connected components of $Z|_{\varphi < u}$ are pairwise disjoint, $B_{\mathbf{y}}$ is well defined and unique. Moreover by Lemma 3.3,

$$\mathbf{y} \in \overline{(Z \cap N(\mathbf{y}))|_{\varphi < u}} \subset \overline{B_{\mathbf{y}}}.$$

Finally, suppose that there exists another connected component B' of $Z|_{\varphi < u}$ such that $\mathbf{y} \in \overline{B'}$. Then \mathbf{y} belongs to the closure of B' , so that $N(\mathbf{y}) \cap B' \neq \emptyset$, since $N(\mathbf{y})$ is a neighborhood of \mathbf{y} . Thus $B' \cap B_{\mathbf{y}} \neq \emptyset$ and since they are both semi-algebraically connected component of the same set, $B' = B_{\mathbf{y}}$. \square

Let us see a geometric consequence of this result. The following lemma shows that if u is the least element of \mathbf{R} such that the hypersurface $\varphi^{-1}(\{u\})$ intersects a semi-algebraically connected component C of $Z \cap \mathbf{R}^n$, then this intersection consists entirely of singular points of φ on Z . It is illustrated by Figure 5.

Lemma 3.5. *Let $\mathbf{y} \in Z \cap \mathbf{R}^n$ with $u = \varphi(\mathbf{y})$ and let C be the semi-algebraically connected component of $Z|_{\varphi \leq u}$ containing \mathbf{y} . If $C|_{\varphi < u} = \emptyset$ then $C = C|_{\varphi = u} \subset K(\varphi, Z)$. In particular, $\mathbf{y} \in K(\varphi, Z)$.*




Figure 5: Illustration of Lemma 3.5 in two cases. On the left, $\varphi = \pi_1$ and $Z \cap \mathbb{R}^3$ is a torus. One sees that when the plane is tangent to the algebraic set, its intersection with the axis of x is reduced to a point. On the right, φ is the square of the Euclidean norm, and Z is a cylinder. Here the differential of φ at the critical point $(-r, 0, 0)$ is collinear to π_1 . The same observations as in the first case can be done.

Proof. If $C_{|\varphi < u} = \emptyset$, since $C \subset Z_{|\varphi \leq u}$ then $C = C_{|\varphi = u}$ holds. Let us prove the contrapositive of the rest of the lemma. Suppose that $C_{|\varphi = u} \not\subset K(\varphi, Z)$, and let

$$z \in C_{|\varphi = u} - K(\varphi, Z).$$

Let B_z be the semi-algebraically connected component of $Z_{|\varphi < u}$ obtained by applying Lemma 3.4. Since $\overline{B_z}$ contains z and is a semi-algebraically connected set of $Z_{|\varphi \leq u}$, then $\overline{B_z} \subset C$. Hence $C_{|\varphi < u}$ contains $(\overline{B_z})_{|\varphi < u} = B_z$, which is then not empty. \square

We prove now an important consequence of the previous lemma. It is a fundamental property of generalized polar varieties and motivates their introduction among the ingredients of a roadmap.

Proposition 3.6. *Let $u \in \mathbf{R}$ and let B be a bounded semi-algebraically connected component of $Z_{|\varphi < u}$. Then $B \cap K(\varphi, Z) \neq \emptyset$.*

Proof. Since φ is a semi-algebraic continuous map and B is semi-algebraic, then $\varphi(\overline{B})$ is a closed and bounded semi-algebraic set by [4, Theorem 3.23]. In particular, φ reaches its minimum $\varphi(z)$ on \overline{B} and since $\emptyset \neq B \subset Z_{|\varphi < u}$, then $\varphi(z) < u$. But B is a semi-algebraically connected component of $Z_{|\varphi < u}$, in particular, it is closed in $Z_{|\varphi < u}$, so that

$$\overline{B} - B \subset Z_{|\varphi = u}.$$

Therefore $z \in B$ and as $B_{|\varphi < \varphi(z)}$ is empty (z is a minimizer), $B_{|\varphi = \varphi(z)}$ and z is in $K(\varphi, Z)$ by Lemma 3.5. Finally $z \in B \cap K(\varphi, Z)$, and the latter is non-empty. \square

We are now able to prove a weaker version of Proposition 3.1, which is illustrated in Figure 6. It deals with the particular case when the map has value in some fiber $Z_{|\varphi = u}$, where $u \in \mathbf{R}$.

Lemma 3.7. *Let $u \in \mathbf{R}$ and $A \subset \mathbf{R}^k$ be a semi-algebraically connected set. Let*

$$\gamma: A \longrightarrow Z_{|\varphi = u} - K(\varphi, Z)$$

be a continuous semi-algebraic map. Then there exists a unique semi-algebraically connected component B of $Z_{|\varphi < u}$ such that $\gamma(A) \subset \overline{B}$.




Figure 6: Illustration of the proof of Proposition 3.1 where $\varphi = \pi_1$ and Z is isomorphic to $\mathbf{V}(x_1^2 + x_2^2 - 1) \times \mathbf{V}(x_2 + x_1^2)$ in two cases. On the left the $\gamma(A) \cap (Z_{|\pi_1|=u} \cap K(\pi_1, Z)) = \emptyset$ and on the right, this intersection is non-empty.

Proof. Let $\mathbf{a} \in A$ and $\mathbf{y} = \gamma(\mathbf{a})$, by assumption, $\mathbf{y} \in Z_{|\varphi|=u} - K(\varphi, Z)$. Then by Lemmas 3.3 and 3.4, there exist an open neighborhood $N(\mathbf{y})$ of \mathbf{y} and a semi-algebraically connected component $B_{\mathbf{y}}$ of $Z_{|\varphi|<u}$ such that

$$(Z \cap N(\mathbf{y}))_{|\varphi|=u} \subset \overline{(Z \cap N(\mathbf{y}))_{|\varphi|<u}} \subset \overline{B_{\mathbf{y}}}.$$

Hence for every $\mathbf{z} \in (Z \cap N(\mathbf{y}))_{|\varphi|=u} - K(\varphi, Z)$, $\mathbf{z} \in \overline{B_{\mathbf{y}}}$ so that $B_{\mathbf{z}} = B_{\mathbf{y}}$ by application of Lemma 3.4. Since γ is a continuous semi-algebraic map, there exists an open semi-algebraic neighborhood $N'(\mathbf{a})$ of \mathbf{a} such that

$$\gamma(N'(\mathbf{a})) \subset (Z \cap N(\mathbf{y}))_{|\varphi|=u} - K(\varphi, Z).$$

Hence the map $\mathbf{a} \mapsto B_{\gamma(\mathbf{a})}$ is constant on $N(\mathbf{a})$.

Since A is semi-algebraically connected, this map is actually globally constant on A and we note B the constant value that it takes on this set. Thus, by Lemma 3.4, for all $\mathbf{a} \in A$, $\gamma(\mathbf{a}) \in \overline{B_{\gamma(\mathbf{a})}} = \overline{B}$, that is $\gamma(A) \subset \overline{B}$. Besides, if B' is another semi-algebraically connected component of $Z_{|\varphi|<u}$ such that $\gamma(A) \subset \overline{B'}$, then for all $\mathbf{a} \in A$,

$$\gamma(\mathbf{a}) \in \overline{B} \cap \overline{B'} \cap Z_{|\varphi|=u} - K(\varphi, Z),$$

so that $B = B'$ by uniqueness in Lemma 3.4. \square

We can now prove the main proposition by sticking together all the pieces. The points of the map that belong to the fiber $Z_{|\varphi|=u}$ are managed by Lemma 3.7, while the remaining ones, in $Z_{|\varphi|<u}$, are more convenient to deal with. This proof is illustrated by Figure 7.

Proof of Proposition 3.1. Since γ is semi-algebraic and continuous, $\gamma(A)$ is semi-algebraically connected. Hence, if $\gamma(A) \subset Z_{|\varphi|<u}$, it is contained in a unique semi-algebraically connected component B of $Z_{|\varphi|<u}$ and we are done.

We assume now that $\gamma(A) \not\subset Z_{|\varphi|<u}$. Let $G = \gamma^{-1}(Z_{|\varphi|=u})$. It is a closed subset of A since $Z_{|\varphi|=u}$ is closed in $Z_{|\varphi|\leq u}$ and γ is continuous. Then, let G_1, \dots, G_N be the semi-algebraically connected components of G , they are closed in A since they are closed in G , which is closed in A . Besides, let H_1, \dots, H_M be the semi-algebraically connected components of $A - G$. They are open in A since they are open in $A - G$, which is open in A .

Let $\mathfrak{B}: A \rightarrow \mathcal{P}(Z_{|\varphi|<u})$ be a map, where $\mathcal{P}(Z_{|\varphi|<u})$ is the power set of $Z_{|\varphi|<u}$. The family formed by both G_1, \dots, G_N and H_1, \dots, H_M is a partition of A . Then one can define \mathfrak{B} by defining it on this partition.

H_i : Since $H_i \subset A - G$, $\gamma(H_i) \subset Z_{|\varphi < u}$ and $\gamma(H_i)$ is semi-algebraically connected as γ is continuous. Then, there exists a unique semi-algebraically connected component B_i of $Z_{|\varphi < u}$ such that $\gamma(H_i) \subset B_i \subset \overline{B_i}$.

G_i : Since G_i is semi-algebraically connected and $\gamma(G_i) \subset Z_{|\varphi = u} - K(\varphi, Z)$, Lemma 3.7 with $A = G_i$ states that there is a unique semi-algebraically connected component B'_i of $Z_{|\varphi < u}$ such that $\gamma(G_i) \subset \overline{B'_i}$.

Therefore, for all $\mathbf{a} \in A$, let \mathfrak{B} such that

$$\mathfrak{B}(\mathbf{a}) = \begin{cases} B_i & \text{if } \mathbf{a} \in H_i \\ B'_i & \text{if } \mathbf{a} \in G_i \end{cases} \text{ so that } \gamma(\mathbf{a}) \in \overline{\mathfrak{B}(\mathbf{a})}.$$

Let us show that \mathfrak{B} is locally constant, that is, for every $\mathbf{a} \in A$, there exists an open Euclidean neighborhood $N(\mathbf{a}) \subset A$ of \mathbf{a} , such that for all $\mathbf{a}' \in N(\mathbf{a})$, $\mathfrak{B}(\mathbf{a}') = \mathfrak{B}(\mathbf{a})$. Then, we will conclude by connectedness. Let $\mathbf{a} \in A$ and $1 \leq i \leq \max(M, N)$.

- If $\mathbf{a} \in H_i$, since H_i is open in A , there exists an open Euclidean neighborhood $N(\mathbf{a})$ of \mathbf{a} contained in H_i . By construction, for all $\mathbf{a}' \in N(\mathbf{a})$, $\mathfrak{B}(\mathbf{a}') = \mathfrak{B}(\mathbf{a})$. Moreover, since H_i is semi-algebraically connected, this also proves that \mathfrak{B} is actually constant on H_i , let $\mathfrak{B}(H_i)$ be this constant value.
- Else $\mathbf{a} \in G_i$, since the G_j 's are closed in A , then \mathbf{a} does not belong to the closure of any other G_j , $j \neq i$. However, the set

$$J = \{1 \leq j \leq M \mid \mathbf{a} \in \overline{H_j}\}$$

is not empty. By construction, $\gamma(\mathbf{a}) \in \overline{\mathfrak{B}(\mathbf{a})}$ and by definition of J , for every $j \in J$, $\gamma(\mathbf{a}) \in \overline{\mathfrak{B}(H_j)}$. But, by Lemma 3.4 applied with $\mathbf{y} = \gamma(\mathbf{a})$, such a semi-algebraically connected component is unique. Hence for all $j \in J$, $\mathfrak{B}(H_j) = \mathfrak{B}(\mathbf{a})$. One can then take $N(\mathbf{a}) = \mathcal{B}(\mathbf{a}, r)$ with $r > 0$ such that this open ball intersects either the H_j 's for $j \in J$ or G_i , and only them.

Finally, we proved that \mathfrak{B} is locally constant. Since A is semi-algebraically connected, \mathfrak{B} is globally constant on A . Denoting by B this constant value, we have $\gamma(A) \subset \overline{B}$ as claimed. Besides if B' is another semi-algebraically connected component of $Z_{|\varphi < u}$ such that $\gamma(A) \subset \overline{B'}$, then in particular $\overline{B} \cap \overline{B'}$ contains $\gamma(G_1) \subset Z_{|\varphi = u} - K(\varphi, Z)$, so that $B = B'$ by Lemma 3.7. \square




Figure 7: Illustration of the proof of Proposition 3.1 with $\varphi = \pi_1$ and Z is isomorphic to $\mathbf{V}(x_1^2 + x_2^2 - 1) \times \mathbf{V}(x_2 + x_1^2)$ in two cases. The intersection $\gamma(A) \cap (Z_{|\pi_1 = u} \cap K(\pi_1, Z))$ is empty on the left while, on the right, it is not.

We then deduce the following consequence on the semi-algebraically connected components of Z with respect to φ . This result is illustrated in Figure 8.

Corollary 3.8. *Let $\varphi: \mathbf{C}^n \rightarrow \mathbf{C}$ be a regular map defined over \mathbf{R} and $Z \subset \mathbf{C}^n$ be an equidimensional algebraic set of positive dimension. Let $u \in \mathbf{R}$ such that $Z|_{\varphi=u} \cap K(\varphi, Z) = \emptyset$ and let C be a semi-algebraically connected component of $Z|_{\varphi \leq u}$. Then, $C|_{\varphi < u}$ is a semi-algebraically connected component of $Z|_{\varphi < u}$.*

Proof. Let γ be the inclusion map $\gamma: C \hookrightarrow Z|_{\varphi \leq u}$. Since $Z|_{\varphi=u} \cap K(\varphi, Z) = \emptyset$, γ satisfies the assumptions of Proposition 3.1 with $A = C$. Then there exists a unique semi-algebraically connected component B of $Z|_{\varphi < u}$ such that $C \subset \overline{B}$, so that $C|_{\varphi < u} \subset \overline{B}|_{\varphi < u} = B$.

First, since $Z|_{\varphi=u} \cap K(\varphi, Z) = \emptyset$ by assumption, in particular $C|_{\varphi=u} \not\subset K(\varphi, Z)$. Then by contrapositive of Lemma 3.5, $C|_{\varphi < u}$ is not empty. Hence, since B is a semi-algebraically connected set of $Z|_{\varphi \leq u}$, containing $C|_{\varphi < u}$, then B is contained in the semi-algebraically connected component C of $Z|_{\varphi \leq u}$. Finally $B \subset Z|_{\varphi < u} \cap C = C|_{\varphi < u}$ and $C|_{\varphi < u} = B$, which is a semi-algebraically connected component of $Z|_{\varphi < u}$. \square




Figure 8: Illustration of Corollary 3.8 where $\varphi = \pi_1$ and Z is isomorphic to $\mathbf{V}(x_1^2 + x_2^2 - 1) \times \mathbf{V}(x_2 + x_1^2)$. On the left $Z|_{\pi_1=u} \cap K(\pi_1, Z) = \emptyset$ and one sees that $C|_{\pi_1 < u}$ is still a semi-algebraically connected component of $Z|_{\pi_1 < u}$. On the right $Z|_{\pi_1=u} \cap K(\pi_1, Z) \neq \emptyset$ and one sees that $C|_{\pi_1 < u}$ is disconnected.

3.2 Fibration and critical values

As in [15, Section 3.2] we are going to use a Nash version of Thom's isotopy lemma, stated in [13]. We refer to [4, Section 3.5] for the definitions of Nash diffeomorphisms, manifolds and submersions together with their properties.

Proposition 3.9. *Let $\varphi: \mathbf{C}^n \rightarrow \mathbf{C}$ be a regular map defined over \mathbf{R} and $A \subset \varphi^{-1}((-\infty, w)) \cap \mathbf{R}^n$ be a semi-algebraically connected semi-algebraic set. Let $v < w$ such that $A|_{\varphi \in (v, w)}$ is a non-empty Nash manifold, bounded, closed in $\varphi^{-1}((v, w)) \cap \mathbf{R}^n$ and such that φ is a submersion on $A|_{\varphi \in (v, w)}$. Then for all $u \in [v, w)$, $A|_{\varphi \leq u}$ is non-empty and semi-algebraically connected.*

Proof. We first prove that $\varphi: A|_{\varphi \in (v, w)} \rightarrow (v, w)$ is a proper surjective submersion. Since $A|_{\varphi \in (v, w)}$ is bounded and φ is semi-algebraic and continuous, $\varphi: A|_{\varphi \in (v, w)} \rightarrow (v, w)$ is a proper map. Let us prove that φ is also surjective on $A|_{\varphi \in (v, w)}$ that is

$$\varphi(A|_{\varphi \in (v, w)}) = (v, w).$$

By assumption, φ is a submersion from $A_{|\varphi \in (v,w)}$ to (v,w) . Then by the semi-algebraic inverse function theorem [4, Proposition 3.29], φ is an open map. Besides, as $A_{|\varphi \in (v,w)}$ is closed and bounded, there exists a closed and bounded semi-algebraic set $X \subset \mathbf{R}^n$ such that $A_{|\varphi \in (v,w)} = X \cap \varphi^{-1}((v,w)) = X_{|\varphi \in (v,w)}$. Then

$$\varphi(A_{|\varphi \in (v,w)}) = \varphi(X_{|\varphi \in (v,w)}) = \varphi(X) \cap (v,w).$$

Since X is bounded and closed, $\varphi(X)$ is closed and bounded by [4, Theorem 3.23]. Hence, $\varphi(A_{|\varphi \in (v,w)})$ is both open and closed in (v,w) . Since (v,w) is semi-algebraically connected, $\varphi(A_{|\varphi \in (v,w)}) = (v,w)$.

By the Nash version of Thom's isotopy lemma [13, Theorem 2.4], since the map $\varphi: A_{|\varphi \in (v,w)} \rightarrow (v,w)$ is a proper surjective submersion, it is a globally trivial fibration. Hence, for $\zeta \in (v,w)$, there exists a Nash diffeomorphism Ψ of the form

$$\Psi: \begin{array}{ccc} A_{|\varphi \in (v,w)} & \longrightarrow & (v,w) \times A_{|\varphi = \zeta} \\ \mathbf{y} & \longmapsto & (\varphi(\mathbf{y}), \psi(\mathbf{y})) \end{array}.$$

We now proceed to prove the main statement of the proposition. There are, at first sight, two different situations to consider: whether $u > v$ or $u = v$ (see Figure 9). Thanks to Puiseux series, we actually prove them simultaneously.

Take $u \in [v,w)$; we prove that $A_{|\varphi \leq u}$ is non-empty and semi-algebraically connected. To prove that $A_{|\varphi = u}$ is non-empty, we consider $\mathbf{z} \in A_{|\varphi = \zeta}$ and the map

$$\gamma: \begin{array}{ccc} [0,1) & \rightarrow & A_{|\varphi \in (v,w)} \\ t & \mapsto & \Psi^{-1}(tu + (1-t)\zeta, \mathbf{z}). \end{array}$$

This map is well defined and continuous, since Ψ is a Nash diffeomorphism from $A_{|\varphi \in (v,w)}$ to $(v,w) \times A_{|\varphi = \zeta}$, and satisfies $\varphi(\gamma(t)) = tu + (1-t)\zeta$ for every $t \in [0,1)$. Moreover γ is a bounded map as $A_{|\varphi \in (v,w)}$ is bounded by assumption. Then, by [4, Proposition 3.21], γ can be continuously extended to $[0,1]$. Then $\varphi(\gamma(t)) = tu + (1-t)\zeta$ is continuous on $[0,1]$, and $\varphi(\gamma(1)) = u$. Finally $\gamma(1) \in A_{|\varphi \leq u}$ and $A_{|\varphi \leq u}$ is non empty.

We prove now that $A_{|\varphi \leq u}$ is semi-algebraically connected. Consider two points \mathbf{y} and \mathbf{y}' in $A_{|\varphi \leq u}$. Since A is semi-algebraically connected by assumption, there exists a continuous path $\gamma: [0,1] \rightarrow A$ such that $\gamma(0) = \mathbf{y}$ and $\gamma(1) = \mathbf{y}'$. Let us construct, from γ , another path that lies in $A_{|\varphi \leq u}$.

Let ε be an infinitesimal, and let $\mathbf{R}' = \mathbf{R}\langle \varepsilon \rangle$ be the field of algebraic Puiseux series in ε (see [4, Section 2.6]). We denote by $A', (v,w)', \Psi', \psi', \varphi'$ and γ' the extensions of respectively $A, (v,w), \Psi, \psi, \varphi$ and γ to \mathbf{R}' in the sense of [4, Proposition 2.108]. According to [4, Exercise 2.110], $\Psi': A'_{|\varphi \in (v,w)'} \rightarrow (v,w)' \times A'_{|\varphi = \zeta}$ is a bijective map. Then let $g': [0,1]' \subset \mathbf{R}' \rightarrow A'$ be such that

$$\begin{array}{ll} g'(t) = \gamma'(t) & \text{if } \varphi'(\gamma'(t)) \leq u + \varepsilon, \\ g'(t) = \Psi'^{-1}(u + \varepsilon, \psi'(\gamma'(t))) & \text{if } u + \varepsilon \leq \varphi'(\gamma'(t)) < w. \end{array}$$

This map is well defined since $u + \varepsilon \in (v,w)$ and if $\varphi'(\gamma'(t)) = u + \varepsilon$, then $\Psi'^{-1}(u + \varepsilon, \psi'(\gamma'(t))) = \gamma'(t)$. Moreover g' is a continuous semi-algebraic map since by [4, Exercise 3.4], Ψ'^{-1}, ψ' and γ' are continuous semi-algebraic maps.

Finally one observes that g' is bounded over \mathbf{R} . Indeed if $\varphi'(\gamma'(t)) \leq u + \varepsilon$, then $g'(t) = \gamma(t)$, which is continuous on $[0,1]'$ and then bounded over \mathbf{R} . Else $\varphi'(\gamma'(t)) \in (v,w)$ and $g'(t) \in A'_{|\varphi \in (v,w)'}$, which is bounded over \mathbf{R} by [4, Proposition 3.19] since $A_{|\varphi \in (v,w)}$ is. Hence, its image $G' = g'([0,1]')$ is a semi-algebraically connected semi-algebraic set, bounded over \mathbf{R} and contained in $A'_{|\varphi \leq u + \varepsilon}$.

Let $G = \lim_{\varepsilon} G'$. By [4, Proposition 12.49], G is a closed and bounded semi-algebraic set. Then, since φ is a continuous semi-algebraic map defined over G , by [4, Lemma 3.24] for all $\mathbf{z}' \in G'$,

$$\varphi(\lim_{\varepsilon} \mathbf{z}') = \lim_{\varepsilon} \varphi(\mathbf{z}') \leq \lim_{\varepsilon} (u + \varepsilon) = u$$

So that G is contained in $A_{|\varphi \leq u}$. In addition, since G' is semi-algebraically connected and bounded over \mathbf{R} , then by [4, Proposition 12.49], G is semi-algebraically connected and contains $\mathbf{y} = \lim_{\varepsilon} g(0)$ and $\mathbf{y}' = \lim_{\varepsilon} g(1)$. We deduce that there exists, inside G , a semi-algebraic path connecting \mathbf{y} to \mathbf{y}' in $A_{|\varphi \leq u}$, which ends the proof. \square




Figure 9: Illustration of the two cases covered by the proof of Proposition 3.9 where $\varphi = \pi_1$ and Z is isomorphic to $\mathbf{V}(x_1^2 + x_2^2 - 1) \times \mathbf{V}(x_2 + x_1^2)$. The two cases are quite similar according to these figures. One sees that Ψ connects all the slices $A_{|\pi_1=u}$ for $u \in (v, w)'$. This diffeomorphism allows to transform the problematic parts (not in $A_{|\pi_1 \leq u}$) of the initial path γ (in green), into another path g (in red), that lies in $A_{|\pi_1=u} \subset A_{|\pi_1 \leq u}$.

The following result is a consequence of Proposition 3.9 as it deals with a particular case. An illustration of this statement can be found in Figure 10.

Corollary 3.10. *Let $Z \subset \mathbf{C}^n$ be an equidimensional algebraic set of positive dimension and let $\varphi: \mathbf{C}^n \rightarrow \mathbf{C}$ be a regular map defined over \mathbf{R} and proper on $Z \cap \mathbf{R}^n$. Let $v < w$ be in \mathbf{R} such that $Z_{|\varphi \in (v, w)} \cap K(\varphi, Z) = \emptyset$, and let C be a semi-algebraically connected component of $Z_{|\varphi \leq w}$. Then, $C_{|\varphi \leq v}$ is a semi-algebraically connected component of $Z_{|\varphi \leq v}$.*

Proof. As $C_{|\varphi < w} = C \cap \varphi^{-1}((-\infty, w)) \cap \mathbf{R}^n$, we are going to use Proposition 3.1 with $A = C_{|\varphi < w}$.

First we need to prove that $C_{|\varphi < w}$ is a non-empty semi-algebraically connected semi-algebraic set. Since $Z_{|\varphi=w} \cap K(\varphi, Z) = \emptyset$, by Corollary 3.8 $C_{|\varphi < w}$ is a semi-algebraically connected component of $Z_{|\varphi < w}$. Hence it is non-empty and semi-algebraically connected.

Then, we need to prove that $C_{|\varphi \in (v, w)}$ is a non-empty Nash manifold, bounded and closed in $\varphi^{-1}((v, w)) \cap \mathbf{R}^n$. Suppose first that $C_{|\varphi \in (v, w)} = \emptyset$. Then

$$C_{|\varphi \leq v} \cup C_{|\varphi=w} = C \quad \text{and} \quad C_{|\varphi \leq v} \cap C_{|\varphi=w} = \emptyset.$$

Since C is semi-algebraically connected, either $C_{|\varphi \leq v}$ or $C_{|\varphi=w}$ is empty (as they are both closed in C). In both cases our conclusion follows. It remains to tackle the case where $C_{|\varphi \in (v, w)}$ is not empty, which we assume to hold from now on.

We prove that $C_{|\varphi \in (v, w)}$ is bounded. Observe that $C_{|\varphi \in (v, w)} \subset C_{|\varphi \in [v, w]} = C \cap \mathbf{R}^n \cap \varphi^{-1}([v, w])$. Recall that φ is proper on $Z \cap \mathbf{R}^n$ by assumption, and thus on $C \cap \mathbf{R}^n$. Hence, $C_{|\varphi \in [v, w]}$ is bounded. Besides $C_{|\varphi \in (v, w)}$ is closed in $\varphi^{-1}((v, w)) \cap \mathbf{R}^n$ as

$$C_{|\varphi \in (v, w)} = C \cap \varphi^{-1}((v, w)) \cap \mathbf{R}^n,$$

and C is closed in \mathbf{R}^n as it is closed in the closed set $Z_{|\varphi \leq w}$. Since $C_{|\varphi \in (v, w)} \cap K(\varphi, Z) = \emptyset$ then by [7, Proposition 3.3.11], $C_{|\varphi \in (v, w)}$ is a Nash manifold of dimension $\dim(Z)$.

To apply Proposition 3.1, it remains to prove that φ is a Nash submersion on $C_{|\varphi \in (v, w)}$. Let $\mathbf{y} \in C_{|\varphi \in (v, w)}$. Since $\mathbf{y} \notin \text{sing}(Z)$, then $T_{\mathbf{y}}C_{|\varphi \in (v, w)} = T_{\mathbf{y}}Z \cap \mathbf{R}^n$ according to [7, Proposition 3.3.11]. Since $C_{|\varphi \in (v, w)} \cap K(\varphi, Z) = \emptyset$, $d_{\mathbf{y}}\varphi$ is onto on $T_{\mathbf{y}}Z$ and since $\dim Z > 0$, the image $d_{\mathbf{y}}\varphi(T_{\mathbf{y}}Z)$ is \mathbf{C} . Hence

$$d_{\mathbf{y}}\varphi(T_{\mathbf{y}}C_{|\varphi \in (v, w)}) = \mathbf{R}.$$

We just established that all the assumptions of Proposition 3.9 are satisfied. One can then apply it to $C_{|\varphi < w}$ and conclude that $C_{|\varphi \leq v}$ is non-empty and semi-algebraically connected. Finally, since C is a semi-algebraically connected component of $Z_{|\varphi \leq w}$, any semi-algebraically connected component of $Z_{|\varphi \leq v}$ contained in C is contained in $C_{|\varphi \leq v}$. Thus $C_{|\varphi \leq v}$ is a semi-algebraically connected component of $Z_{|\varphi \leq v}$. \square




Figure 10: Illustration of Corollary 3.10 where $\varphi = \pi_1$ and Z is isomorphic to $\mathbf{V}(x_1^2 + x_2^2 - 1) \times \mathbf{V}(x_2 + x_1^2)$ in two cases. On the left $Z_{|\pi_1 \in (v, w)} \cap K(\pi_1, Z) = \emptyset$ and we see that $C_{|\pi_1 \leq v}$ is still a semi-algebraically connected component of $Z_{|\pi_1 \leq v}$. On the right $Z_{|\pi_1 \in (v, w)} \cap K(\pi_1, Z)$ contains a point and we see that $C_{|\pi_1 \leq v}$ is semi-algebraically disconnected.

4 Proof of the main connectivity result

Recall that $\varphi = (\varphi_1, \dots, \varphi_n) \subset \mathbf{R}[x_1, \dots, x_n]$ and for $1 \leq i \leq n$, $\varphi_i: \mathbf{y} \mapsto (\varphi_1(\mathbf{y}), \dots, \varphi_i(\mathbf{y}))$. We denote by $W_i = W(\varphi_i, V)$ the Zariski closure of the set of critical points of the restriction of φ_i to V and recall that

$$K_i = W(\varphi_1, V) \cup S \cup \text{sing}(V) \quad \text{and} \quad F_i = \varphi_{i-1}^{-1}(\varphi_{i-1}(K_i)) \cap V,$$

where S is a subset of $W(\varphi_1, W_i)$. We suppose that the following assumptions holds:

- (A) V is d -equidimensional and its singular locus $\text{sing}(V)$ is finite;
- (P) the restriction of the map φ_1 to $V \cap \mathbf{R}^n$ is proper and bounded from below;
- (B₁) W_i is either empty or $(i - 1)$ -equidimensional and smooth outside $\text{sing}(V)$;
- (B₂) for any $\mathbf{y} = (\mathbf{y}_1, \dots, \mathbf{y}_i) \in \mathbf{C}^i$, $V \cap \varphi_{i-1}^{-1}(\mathbf{y})$ is either empty or $(d - i + 1)$ -equidimensional;
- (C₁) S is finite;
- (C₂) S has a non-empty intersection with every semi-algebraically connected component of $W(\varphi_1, W_i) \cap \mathbf{R}^n$.

Then the goal of this section is to prove that, $W_i \cup F_i$ intersects each semi-algebraically connected component of $V \cap \mathbf{R}^n$ and their intersection is semi-algebraically connected. Let $\mathcal{R} = F_i \cup W_i$.

We prove that the following so-called roadmap property holds:

RM: “For any semi-algebraically connected component C of $V \cap \mathbf{R}^n$, the set $C \cap \mathcal{R}$ is non-empty and semi-algebraically connected.”,

by proving a truncated version of RM and show that is is enough. For $u \in \mathbf{R}$ let

RM(u): “For any semi-algebraically connected component C of $V_{|\varphi_1| \leq u}$, the set $C \cap \mathcal{R}$ is non-empty and semi-algebraically connected.”.

Lemma 4.1. *If RM(u) holds for all $u \in \mathbf{R}$, then RM holds.*

Proof. Let C be a semi-algebraically connected component of $V \cap \mathbf{R}^n$. Since C is non-empty and semi-algebraically connected, there exist \mathbf{y} and \mathbf{y}' in C , and a semi-algebraic path $\gamma: [0, 1] \rightarrow C$ connecting them. Let

$$u = \max\{\varphi_1(\gamma(t)), t \in [0, 1]\} \in \mathbf{R}.$$

Such a maximum u exists by continuity of γ and φ_1 since $[0, 1]$ is closed and bounded. Then $\gamma([0, 1]) \subset V_{|\varphi_1| \leq u}$. Since $\gamma([0, 1])$ is semi-algebraically connected, there exists a (unique) semi-algebraically connected component B of $V_{|\varphi_1| \leq u}$ containing $\gamma([0, 1])$. In particular, B contains \mathbf{y} and \mathbf{y}' . Since RM(u) holds by assumption, then $B \cap \mathcal{R}$ is non-empty. But as $\mathbf{y} \in B \cap C$ and B is semi-algebraically connected, then C contains B . Finally, $C \cap \mathcal{R}$ contains $B \cap \mathcal{R}$ and the former is non-empty.

We can suppose now, in addition, that \mathbf{y} and \mathbf{y}' are in $C \cap \mathcal{R}$, and let B be defined as above. Then, \mathbf{y} and \mathbf{y}' are in $B \cap \mathcal{R}$, which is semi-algebraically connected by RM(u). Therefore \mathbf{y} and \mathbf{y}' are connected by a semi-algebraic path in $B \cap \mathcal{R}$. Since $B \subset C$, \mathbf{y} and \mathbf{y}' are semi-algebraically connected in $C \cap \mathcal{R}$. In conclusion, $C \cap \mathcal{R}$ is semi-algebraically connected and RM holds. \square

Remark. The previous lemma trivially holds in the case of [15, Theorem 14], since $V \cap \mathbf{R}^n$ is assumed to be bounded. Indeed, in this case, considering $u = \max_{\mathbf{y} \in V \cap \mathbf{R}^n} \varphi_1(\mathbf{y})$, one has $V_{|\varphi_1| \leq u} = V \cap \mathbf{R}^n$.

4.1 Restoring connectivity

Before proving RM(u) for all $u \in \mathbf{R}$, we need to prove the following result, which constitutes the core of the proof of Theorem 1.1. This proposition shows that the connectivity property of our roadmap candidate is satisfied when u is increasing towards singular points of φ_1 on V . This is ensured by the addition of the fibers F_i .

Proposition 4.2. *Let $u \in \mathbf{R}$ and C be a semi-algebraically connected component of $V_{|\varphi_1| \leq u}$ such that $C_{|\varphi_1| < u}$ is non-empty. Let B be a semi-algebraically connected component of $C_{|\varphi_1| < u}$, then:*

1. $\overline{B} \cap (F_i \cup W_i)$ is non-empty;
2. Any point $\mathbf{y} \in \overline{B} \cap (F_i \cup W_i)$ can be connected to a point $\mathbf{z} \in B \cap (F_i \cup W_i)$ by a semi-algebraic path in $\overline{B} \cap (F_i \cup W_i)$.

Let us begin with a technical lemma:

Lemma 4.3. *Let \mathbf{K} be a real closed field containing \mathbf{R} and $\overline{\mathbf{K}}$ be its algebraic closure. Let $Z \subset \overline{\mathbf{K}}^n$ be a d -equidimensional algebraic set, where $d > 0$. Assume that for any $\mathbf{z} \in \overline{\mathbf{K}}^{i-1}$,*

$$Z \cap \varphi_{i-1}^{-1}(\mathbf{z}) \text{ is either empty or } (d - i + 1)\text{-equidimensional.}$$

Let B be a bounded semi-algebraically connected component of $Z \cap \mathbf{K}^n$ and let $\mathbf{y} \in B$. Let H be the semi-algebraically connected component of $B \cap \varphi_{i-1}^{-1}(\varphi_{i-1}(\mathbf{y}))$ containing \mathbf{y} . Then, the intersection $H \cap K(\varphi_i, Z)$ is not empty.

Proof. Let $Y = Z \cap \varphi_{i-1}^{-1}(\varphi_{i-1}(\mathbf{y}))$. By assumption, Y is an equidimensional algebraic set of dimension $d - i + 1$. Besides, H is a bounded semi-algebraically connected component of $Y \cap \mathbf{K}^n$, since B is a bounded semi-algebraically connected component of $Z \cap \mathbf{K}^n$.

Recall that $\varphi = (\varphi_1, \dots, \varphi_n)$. Then $\varphi_i(H) \subset \mathbf{R}$ is a closed and bounded semi-algebraic set by [4, Theorem 3.23]. In particular, φ_i reaches its minimum on H . Let $\mathbf{z} \in H$ be such that $\varphi_i(\mathbf{z}) = \min \varphi_i(H)$, so that $H_{|\varphi_i| < \varphi_i(\mathbf{z})}$ is empty. Then, by Lemma 3.5,

$$\mathbf{z} \in H \cap K(\varphi_i, Y).$$

Let $\mathbf{g} \subset \mathbf{K}[x_1, \dots, x_n]$ be a sequence of generators of $\mathbf{I}(Z)$, so that $Y = \mathbf{V}(\mathbf{g}, \varphi_{i-1} - \varphi_{i-1}(\mathbf{y}))$. Since Y is $(d - i + 1)$ -equidimensional, Lemma 2.2 establishes that \mathbf{z} is such that

$$\text{rank} \begin{bmatrix} \text{Jac}_{\mathbf{z}}(\mathbf{g}) \\ \text{Jac}_{\mathbf{z}}(\varphi_{i-1}) \\ \text{Jac}_{\mathbf{z}}(\varphi_i) \end{bmatrix} < n - (d - (i - 1)) + 1.$$

Since $\varphi_i = (\varphi_{i-1}, \varphi_i)$, one deduces that

$$\text{rank} \begin{bmatrix} \text{Jac}_{\mathbf{z}}(\mathbf{g}) \\ \text{Jac}_{\mathbf{z}}(\varphi_i) \end{bmatrix} < n - d + i,$$

which means that $\mathbf{z} \in H \cap K(\varphi_i, Z)$. Finally, the latter set is non-empty and the statement is proved. \square

Notation. For the rest of the subsection let u , C and B as defined in Proposition 4.2.

Let us deal with one particular case of the second item of Proposition 4.2.

Lemma 4.4. *Let \mathbf{y} be in $\overline{B} \cap F_i$. Then, there exists a point $\mathbf{z} \in B \cap (F_i \cup W_i)$ and a semi-algebraic path in $\overline{B} \cap (F_i \cup W_i)$ connecting \mathbf{y} to \mathbf{z} .*

Proof. Let $\mathbf{y} \in \overline{B} \cap F_i$, we assume that $\mathbf{y} \notin B$ so that $\varphi_1(\mathbf{y}) = u$, otherwise taking $\mathbf{z} = \mathbf{y}$ would end the proof. Since $\mathbf{y} \in \overline{B}$, by the curve selection lemma [4, Th. 3.22], there exists a semi-algebraic path $\gamma: [0, 1] \rightarrow \mathbf{R}^n$ such that $\gamma(0) = \mathbf{y}$ and $\gamma(t) \in B$ for all $t \in (0, 1]$. Let ε be an infinitesimal, $\mathbf{R}' = \mathbf{R}(\varepsilon)$ be the field of algebraic Puiseux series and $\psi = (\psi_1, \dots, \psi_n)$ be the semi-algebraic germ of γ at the right of the origin (see [4, Section 3.3]). According to [4, Theorem 3.17], we can identify ψ with an element of $(\mathbf{R}')^n$. With some notation abuse we will denote them equally. Hence by [4, Proposition 3.21], $\lim_{\varepsilon} \psi = \mathbf{y}$. Let finally

$$H = \text{ext}(B, \mathbf{R}') \cap \varphi_{i-1}^{-1}(\varphi_{i-1}(\psi)) \subset (\mathbf{R}')^n$$

where $\text{ext}(B, \mathbf{R}')$ is the extension of B to \mathbf{R}' and φ_j for $1 \leq j \leq n$, with some notation abuse, still denote the extension of φ_j to \mathbf{R}' .

Since $\gamma((0, 1]) \subset B$, then, by [4, Proposition 3.19], $\psi \in \text{ext}(B, \mathbf{R}')$. Hence $\psi \in H$ and H is non-empty. Moreover B is bounded since $\varphi_1: V \cap \mathbf{R}^n \rightarrow \mathbf{R}$ is a proper map bounded below by assumption (P). Then [4, Proposition 3.19] states that $\text{ext}(B, \mathbf{R}')$ and then H are bounded over \mathbf{R} . Hence the map \lim_{ε} is well defined on H and

$$\mathbf{y} \in \lim_{\varepsilon} H = \{\lim_{\varepsilon} \mathbf{y}', \mathbf{y}' \in H\} \subset \mathbf{R}^n.$$

Finally, as φ_{i-1} is semi-algebraic and continuous, $\lim_{\varepsilon} H$ is contained in $\overline{B} \cap \varphi_{i-1}^{-1}(\varphi_{i-1}(\mathbf{y}))$ by [4, Lemma 3.24]. But $\mathbf{y} \in F_i$, so that

$$\varphi_{i-1}^{-1}(\varphi_{i-1}(\mathbf{y})) \subset \varphi_{i-1}^{-1}(\varphi_{i-1}(K_i)),$$

and finally $\lim_{\varepsilon} H$ is actually in $\overline{B} \cap F_i$.

Let H_1 be the semi-algebraically connected component of H containing ψ . By [4, Proposition 5.24], $\lim_{\varepsilon} H_1$ is the semi-algebraically connected component of $\lim_{\varepsilon} H$ containing \mathbf{y} . Actually, we just proved that every \mathbf{w} in $\lim_{\varepsilon} H_1$ can be semi-algebraically connected to \mathbf{y} into $\overline{B} \cap F_i$. We find now some $\mathbf{w} \in \lim_{\varepsilon} H_1$ that can be connected to a point $\mathbf{z} \in B \cap (F_i \cup W_i)$ to end the proof. Such a \mathbf{w} must be the origin of a germ of semi-algebraic functions that lives in $B \cap (W_i \cup F_i)$.

By assumption (A) V is d -equidimensional. By assumption (B₂), for all $\mathbf{z} \in V$, the algebraic set $V \cap \varphi_{i-1}^{-1}(\varphi_{i-1}(\mathbf{z}))$ is $(d - i + 1)$ -equidimensional. Then, if we denote by \mathbf{C}' the algebraic closure of \mathbf{R}' , it is an algebraic closed extension of \mathbf{C} , so that the algebraic sets of $(\mathbf{C}')^n$

$$Z = \{\mathbf{z} \in (\mathbf{C}')^n \mid \forall h \in \mathbf{I}(V), h(\mathbf{z}) = 0\} \quad \text{and} \quad Z \cap \varphi_{i-1}^{-1}(\varphi_{i-1}(\psi))$$

are equidimensional of dimension respectively d and $(d - i + 1)$. Since B is a semi-algebraically connected component of $V_{|\varphi_1 < u}$, then, by [4, Proposition 5.24], $\text{ext}(B, \mathbf{R}')$ is a semi-algebraically connected component of

$$\text{ext}(V_{|\varphi_1 < u}, \mathbf{R}') = \text{ext}(V \cap \mathbf{R}^n, \mathbf{R}')_{|\varphi_1 < u} = Z_{|\varphi_1 < u},$$

The first item is then proved. Let $\mathbf{y} \in \overline{B} \cap (F_i \cup W_i)$. To prove the second item, one only needs to consider the case where $\mathbf{y} \in \overline{B} \cap (W_i - F_i)$ according to Lemma 4.4. Moreover one can assume that $\mathbf{y} \notin B$ and then $\varphi_1(\mathbf{y}) = u$, otherwise, taking $\mathbf{z} = \mathbf{y}$, would end the proof.

Let D be the semi-algebraically connected component of $(W_i)_{|\varphi_1 \leq u}$ containing \mathbf{y} , one considers two disjoint cases.

1. If $D \not\subset \overline{B}$, there exists $\mathbf{y}' \in D$ such that $\mathbf{y}' \notin \overline{B}$. Then let $\gamma: [0, 1] \rightarrow D$ such that $\gamma(0) = \mathbf{y}$ and $\gamma(1) = \mathbf{y}'$. Hence, if

$$t_1 = \max\{t \in [0, 1] \mid \gamma(t) \in \overline{B}\},$$

then $\gamma(t_1) \in K(\varphi_1, V)$ by contrapositive of statement *c*) of Lemma 3.3. Since $K(\varphi_1, V) \subset F_i$, we can apply Lemma 4.4 to $\gamma(t_1)$ and find $\mathbf{z} \in B \cap (F_i \cup W_i)$ that is connected to $\gamma(t_1)$ and then to \mathbf{y} by a semi-algebraic path in $\overline{B} \cap (F_i \cup W_i)$.

2. If $D \subset \overline{B}$, we claim that there exists some $\mathbf{z} \in D \cap F_i$. Indeed since D is a semi-algebraically connected component of $(W_i)_{|\varphi_1 \leq u}$ and φ_1 is a proper map, D is bounded. Then by Proposition 3.6 there exists $\mathbf{y}' \in D \cap K(\varphi_1, W_i)$. If $\mathbf{y}' \in \text{sing}(W_i)$ then $\mathbf{y}' \in \text{sing}(V)$ by assumption B_1 and taking $\mathbf{z} = \mathbf{y}' \in F_i$ one concludes as in the first item.

Else $\mathbf{y}' \in W(\varphi_1, W_i)$, so let E be the semi-algebraically connected component of $W(\varphi_1, W_i)$ containing \mathbf{y}' . Since $\varphi_1(W(\varphi_1, W_i))$ is finite by Sard's lemma, $\varphi_1(E) = \{\varphi_1(\mathbf{y}')\}$, so that $E \subset (W_i)_{|\varphi_1 \leq u}$. Hence, since E is semi-algebraically connected, $E \subset D$. By assumption C_2 , there exists $\mathbf{z} \in E \cap S$, so that $\mathbf{z} \in D \cap S \subset D \cap F_i$ and we are done.

Then we can connect \mathbf{y} to \mathbf{z} inside $D \subset \overline{B} \cap W_i$ and since $\mathbf{z} \in D \cap F_i$ which is contained in $\overline{B} \cap F_i$, we can connect similarly \mathbf{z} to some $\mathbf{z}' \in B \cap (F_i \cup W_i)$ inside $\overline{B} \cap F_i$ by Lemma 4.4. Putting things together, \mathbf{y} is connected to some $\mathbf{z}' \in B \cap (F_i \cup W_i)$ by a semi-algebraic path in $\overline{B} \cap F_i$. \square

Corollary 4.5. *Let $u \in \mathbf{R}$ such that for all $u' < u$, $\text{RM}(u')$ holds. Let C be a semi-algebraically connected component of $V_{|\varphi_1 \leq u}$ such that $C_{|\varphi_1 < u}$ is non-empty. If B is a semi-algebraically connected component of $C_{|\varphi_1 < u}$, then $\overline{B} \cap \mathcal{R}$ is non-empty and semi-algebraically connected.*

Proof. Let \mathbf{y} and \mathbf{y}' be in $\overline{B} \cap \mathcal{R}$. According to Proposition 4.2, they can respectively be connected to some \mathbf{z} and \mathbf{z}' in $B \cap \mathcal{R}$, by a semi-algebraic path in $\overline{B} \cap \mathcal{R}$. As B is semi-algebraically connected, there exists a semi-algebraic path $\gamma: [0, 1] \rightarrow B$ connecting \mathbf{z} to \mathbf{z}' . Let

$$u' = \max\{\varphi_1(\gamma(t)) \mid t \in [0, 1]\},$$

so that $\gamma([0, 1]) \subset V_{|\varphi_1 \leq u'}$. Such a u' exists by continuity of γ , and satisfies $u' < u$, as $[0, 1]$ is closed and bounded.

Let B' be the semi-algebraically connected component of $B_{|\varphi_1 \leq u'}$ that contains $\gamma([0, 1])$. Since B' is also a semi-algebraically connected component of $V_{|\varphi_1 \leq u'}$, property $\text{RM}(u')$ states that $B' \cap \mathcal{R}$ is non-empty and semi-algebraically connected. Then, as \mathbf{z} and \mathbf{z}' are in $B' \cap \mathcal{R}$, they can be connected by a semi-algebraic path in $B' \cap \mathcal{R}$, and then, in $B \cap \mathcal{R}$. Thus \mathbf{y} and \mathbf{y}' are connected by a semi-algebraic path in $\overline{B} \cap \mathcal{R}$ and we are done. \square

4.2 Recursive proof of the truncated roadmap property

In order to prove that $\text{RM}(u)$ holds for all $u \in \mathbf{R}$, one can consider two disjoint cases: whether u is a real singular value of φ_1 , that is $u \in \varphi_1(K_i)$, or not. The following lemma allows us to proceed by induction.

Lemma 4.6. *The set $\varphi_1(K_i)$ is non-empty and finite.*

Proof. According to the algebraic version of Sard's theorem [16, Proposition B.2.], the set of critical values of φ_1 on V is an algebraic set of \mathbf{C} of dimension 0. Then, it is either empty or non-empty but finite. Hence, $\varphi_1(K_i)$ is either empty or non-empty but finite, as S , $\text{sing}(V)$ and \mathcal{P} are, by assumption. Moreover since φ_1 is a proper map bounded from below on $V \cap \mathbf{R}^n$ by assumption (P), for any $u \in \mathbf{R}$, $Z_{|\varphi_1 < u}$ is bounded. Then, since V is not empty, by Proposition 3.6 the sets $K(\varphi_1, V)$ and then $\varphi_1(K_i)$ are not empty. \square

We denote by $v_1 < \dots < v_\ell$ the points of $\varphi_1(K_i \cap \mathbf{R}^n)$ and, in addition, let $v_{\ell+1} = +\infty$. We proceed by proving the two following steps.

Step 1: Let $u \in \mathbf{R}$, if $\text{RM}(u')$ holds for all $u' < u$, then $\text{RM}(u)$ holds.

Step 2: Let $j \in \{1, \dots, \ell\}$, if $\text{RM}(v_j)$ holds, then for all $u \in (v_j, v_{j+1})$, $\text{RM}(u)$ holds.

Remark that, by Lemma 3.5, $v_1 = \min_{V \cap \mathbf{R}^n} \varphi_1$, since $V \cap \mathbf{R}^n$ is closed. Then for $u' < v_1$, $V_{|\varphi \leq u'} = \emptyset$ and $\text{RM}(u')$ trivially holds. Hence, proving these two steps is enough to prove $\text{RM}(u)$ for all u in \mathbf{R} , by an immediate induction.

Proposition 4.7 (Step 1). *Let $u \in \mathbf{R}$. Assume that for all $u' < u$, $\text{RM}(u')$ holds. Then $\text{RM}(u)$ holds.*

The proof of this proposition is illustrated by Figure 12.

Proof. Let $u \in \mathbf{R}$ be such that for all $u' < u$, $\text{RM}(u')$ holds and let C be a semi-algebraically connected component of $V_{|\varphi_1 \leq u}$. We have to prove that $C \cap \mathcal{R}$ is non-empty and semi-algebraically connected.

If $C_{|\varphi_1 < u}$ is empty, then, by Lemma 3.5, $C \subset K(\varphi_1, V)$. But the points of $K(\varphi_1, V)$ are either in W_i or in $\text{sing}(V) \subset F_i$. Hence $K(\varphi_1, V) \subset \mathcal{R}$ and $C \cap \mathcal{R} = C$, which is non-empty and semi-algebraically connected by definition.

From now on, $C_{|\varphi_1 < u}$ is supposed to be non-empty and let B_1, \dots, B_r be its semi-algebraically connected components. According to Corollary 4.5, for all $1 \leq j \leq r$, $\overline{B_j} \cap \mathcal{R}$ is non-empty and semi-algebraically connected. Then, as $\overline{B_j} \subset C$,

$$\overline{B_j} \cap \mathcal{R} \subset C \cap \mathcal{R}$$

for every $1 \leq j \leq r$, and $C \cap \mathcal{R}$ is non-empty.

Let us now prove that $C \cap \mathcal{R}$ is semi-algebraically connected. Let \mathbf{y} and \mathbf{y}' in $C \cap \mathcal{R}$. As C is semi-algebraically connected, there exists a semi-algebraically continuous map $\gamma: [0, 1] \rightarrow C$ such that $\gamma(0) = \mathbf{y}$ and $\gamma(1) = \mathbf{y}'$. Now let

$$G = \gamma^{-1}(C_{|\varphi_1 = u} \cap K(\varphi_1, V)) \quad \text{and} \quad H = [0, 1] - G.$$

We denote by G_1, \dots, G_N the connected components of G and H_1, \dots, H_M those of H . The sets H_j for $1 \leq j \leq M$ are open intervals of $[0, 1]$, and we note $\ell_j = \inf(H_j)$ and $r_j = \sup(H_j)$. Since $\gamma(G)$ already lies in $C \cap \mathcal{R}$, let us establish that for every $1 \leq j \leq M$, $\gamma(\ell_j)$ and $\gamma(r_j)$ can be connected by another semi-algebraic path τ_j in $C \cap \mathcal{R}$.

Let $1 \leq j \leq M$, then $\gamma(H_j) \cap (C_{|\varphi_1 = u} \cap K(\varphi_1, V)) = \emptyset$ by definition. Moreover, $\gamma(H_j) \subset C$ so that

$$\gamma(H_j) \cap (V_{|\varphi_1 = u} \cap K(\varphi_1, V)) = \emptyset.$$

Hence, since H_j is connected, there exists, by Proposition 3.1, a unique semi-algebraically connected component B of $V_{|\varphi_1 < u}$ such that $\gamma(H_j) \subset \overline{B}$. But $\gamma(H_j) \subset C$, so that \overline{B} and thus B is actually contained in C . Therefore, B is actually a semi-algebraically connected component of $C_{|\varphi_1 < u}$ and there exists $1 \leq k \leq r$ such that $B = B_k$. At this step $\gamma(H_j) \subset \overline{B_k}$, so that

$$\gamma([\ell_j, r_j]) = \gamma(\overline{H_j}) \subset \overline{\gamma(H_j)} \subset \overline{B_k},$$

and both $\gamma(\ell_j)$ and $\gamma(r_j)$ are in $\overline{B_k}$. Remark that both ℓ_j and r_j are in G , so that both $\gamma(\ell_j)$ and $\gamma(r_j)$ are in $K(\varphi_1, V) \subset F_i \subset \mathcal{R}$. Thus, both $\gamma(\ell_j)$ and $\gamma(r_j)$ are in $\overline{B_k} \cap \mathcal{R}$. According to Corollary 4.5, they can be connected by a semi-algebraic path $\tau_j: [0, 1] \rightarrow \overline{B_k} \cap \mathcal{R} \subset C \cap \mathcal{R}$.

In conclusion, we have proved that for $1 \leq j \leq M$, $\gamma(\ell_j)$ and $\gamma(r_j)$ can be connected by a semi-algebraic path τ_j in $C \cap \mathcal{R}$. Therefore the semi-algebraic sub-paths $\gamma|_{H_j}$ can be replaced by the τ_j 's, which lie in $C \cap \mathcal{R}$. Moreover, for all $1 \leq j \leq N$

$$\gamma(G_j) \subset C \cap \mathcal{R}.$$

Since the H_j 's and G_j 's form a partition of $[0, 1]$, by putting together alternatively the τ_j 's and the $\gamma|_{G_j}$'s, one obtains a semi-algebraic path in $C \cap \mathcal{R}$ connecting $\mathbf{y} = \gamma(0)$ to $\mathbf{y}' = \gamma(1)$. And we are done. \square




Figure 12: Illustration of proof of Proposition 4.7 with $\varphi_1 = \pi_1$ and V is isomorphic to $V(x_1^2 + x_2^2 - 1) \times V(x_2 + x_1^2)$. Here, only \mathbf{y}' belongs to $C_{|\pi_1=u} \cap K(\pi_1, V)$. Then we replace the path $\gamma = \gamma|_{H_1}$ by a path τ_1 that lies in the intersection of the roadmap and the semi-algebraically connected component C .

Proposition 4.8 (Step 2). *Let $j \in \{1, \dots, \ell\}$, if $\text{RM}(v_j)$ holds, then for all $u \in (v_j, v_{j+1})$, $\text{RM}(u)$ holds.*

The proof of this proposition is illustrated by Figure 13.

Proof. Let $j \in \{0, \dots, \ell\}$ and $u \in (v_j, v_{j+1})$. Let C be a semi-algebraically connected component of $V_{|\varphi_1 \leq u}$; we have to prove that $C \cap \mathcal{R}$ is non-empty and semi-algebraically connected.

Let us first prove that $C_{|\varphi_1 \leq v_j} \cap \mathcal{R}$ is non-empty and semi-algebraically connected. By assumption (A), V is an equidimensional algebraic set of positive dimension, and by assumption (P), the restriction of φ_1 to $V \cap \mathbf{R}^n$ is a proper map bounded below. Moreover, as $\varphi_1(K(\varphi_1, V) \cap \mathbf{R}^n) \subset \{v_1, \dots, v_\ell\}$, then

$$V_{|\varphi_1 \in (v_j, u]} \cap K(\varphi_1, V) = \emptyset.$$

Then using Corollary 3.10, one deduces that $C_{|\varphi_1 \leq v_j}$ is a semi-algebraically connected component of $V_{|\varphi_1 \leq v_j}$. Hence, by property $\text{RM}(v_j)$, the set $C_{|\varphi_1 \leq v_j} \cap \mathcal{R}$ is non-empty and semi-algebraically connected. In particular, $C \cap \mathcal{R}$ is non-empty.

Let us now prove that $C \cap \mathcal{R}$ is semi-algebraically connected. Let \mathbf{y} be in $C \cap \mathcal{R}$. According to the previous paragraph, one just need to be able to connect \mathbf{y} to a point \mathbf{z} of $C_{|\varphi_1 \leq v_j} \cap \mathcal{R}$ by a semi-algebraic path in $C \cap \mathcal{R}$ and then apply $\text{RM}(v_j)$. First, if $\mathbf{y} \in C_{|\varphi_1 \leq v_j} \cap \mathcal{R}$, there is nothing to do. Suppose now that $\mathbf{y} \in C_{|\varphi_1 \in (v_j, u]} \cap \mathcal{R}$. We claim that actually

$$\mathbf{y} \in C \cap W_i.$$

Indeed, if $\mathbf{y} \in C \cap F_i$, then $\varphi_{i-1}(\mathbf{y}) \in \varphi_{i-1}(K_i)$ and $\varphi_1(\mathbf{y})$ would be one of the v_1, \dots, v_ℓ .

Let D be the semi-algebraically connected component of $(C \cap W_i)_{|\varphi_1 \leq u}$ containing \mathbf{y} . Remark that D is a semi-algebraically connected component of $(W_i)_{|\varphi_1 \leq u}$, as it contains \mathbf{y} and is contained in C . Since $\varphi_1(W(\varphi_1, W_i))$ is finite by Sard's lemma, then $\varphi_1(W(\varphi_1, W_i)) = \varphi_1(S)$. Hence

$$(v_j, u) \cap \varphi_1(W(\varphi_1, W_i)) = \emptyset.$$

Since W_i is equidimensional and smooth outside $\text{sing}(V)$, then by Corollary 3.10, $D_{|\varphi_1 \leq v_j}$ is a semi-algebraically connected component of $(W_i)_{|\varphi_1 \leq v_j}$. Therefore, let $\mathbf{z} \in D_{|\varphi_1 \leq v_j}$. Since D is semi-algebraically connected, there exists a semi-algebraic path, connecting $\mathbf{y} \in D \subset C \cap \mathcal{R}$ to

$$\mathbf{z} \in D_{|\varphi_1 \leq v_j} \subset C_{|\varphi_1 \leq v_j} \cap \mathcal{R}$$

in $D \subset C \cap \mathcal{R}$. We are done. \square




Figure 13: Illustration of proof of Proposition 4.8 with $\varphi_1 = \pi_1$ and V is isomorphic to $V(x_1^2 + x_2^2 - 1) \times V(x_2 + x_1^2)$. We connect the points \mathbf{y} and \mathbf{y}' in $C \cap W_i$ to respectively \mathbf{z} and \mathbf{z}' in $C_{|\pi_1 \leq v_j}$. Then we are reduced to the case of Step 1.

Acknowledgments

The first and second authors are supported by ANR grants ANR-18-CE33-0011 SESAME, and ANR-19-CE40-0018 DE RERUM NATURA, the joint ANR-FWF ANR-19-CE48-0015 ECARP project, the European Union's Horizon 2020 research and innovative training network program under the Marie Skłodowska-Curie grant agreement N° 813211 (POEMA) and the Grant FA8665-20-1-7029 of the EOARD-AFOSR. The third author was supported by an NSERC Discovery Grant.

References

- [1] B. Bank, M. Giusti, J. Heintz, and L. M. Pardo. Generalized polar varieties and an efficient real elimination. *Kybernetika*, 40(5):519–550, 2004.
- [2] B. Bank, M. Giusti, J. Heintz, and L. M. Pardo. Generalized polar varieties: Geometry and algorithms. *Journal of complexity*, 21(4):377–412, 2005.
- [3] B. Bank, M. Giusti, J. Heintz, M. Safey El Din, and E. Schost. On the geometry of polar varieties. *Applicable Algebra in Engineering, Communication and Computing*, 21(1):33–83, 2010.
- [4] S. Basu, R. Pollack, and M.-F. Roy. *Algorithms in Real Algebraic Geometry*. 10. Springer, Berlin, Heidelberg, 3 edition, 2016.
- [5] S. Basu and M.-F. Roy. Divide and conquer roadmap for algebraic sets. *Discrete & Computational Geometry*, 52(2):278–343, 2014.
- [6] S. Basu, M.-F. Roy, M. Safey El Din, and É. Schost. A baby step–giant step roadmap algorithm for general algebraic sets. *Foundations of Computational Mathematics*, 14(6):1117–1172, 2014.
- [7] J. Bochnak, M. Coste, and M.-F. Roy. *Real algebraic geometry*, volume 36. Springer Science & Business Media, 2013.
- [8] J. Canny. *The complexity of robot motion planning*. MIT press, 1988.
- [9] J. Canny. Constructing roadmaps of semi-algebraic sets i: Completeness. *Artificial Intelligence*, 37(1-3):203–222, 1988.

- [10] J. Canny. Computing roadmaps of general semi-algebraic sets. *The Computer Journal*, 36(5):504–514, 1993.
- [11] J. F. Canny. Computing roadmaps of general semi-algebraic sets. In *International Symposium on Applied Algebra, Algebraic Algorithms, and Error-Correcting Codes*, pages 94–107. Springer, 1991.
- [12] J. Capco, M. Safey El Din, and J. Schicho. Robots, computer algebra and eight connected components. In *Proceedings of the 45th International Symposium on Symbolic and Algebraic Computation*, pages 62–69, 2020.
- [13] M. Coste and M. Shiota. Nash triviality in families of nash manifolds. *Inventiones mathematicae*, 108(1):349–368, 1992.
- [14] D. Grigoryev and N. Vorobjov. Counting connected components of a semialgebraic set in subexponential time. *Computational Complexity*, 2:133–186, 06 1992.
- [15] M. Safey El Din and É. Schost. A baby steps/giant steps probabilistic algorithm for computing roadmaps in smooth bounded real hypersurface. *Discrete & Computational Geometry*, 45(1):181–220, 2011.
- [16] M. Safey El Din and É. Schost. A nearly optimal algorithm for deciding connectivity queries in smooth and bounded real algebraic sets. *Journal of the ACM (JACM)*, 63(6):1–37, 2017.
- [17] J. T. Schwartz and M. Sharir. On the “piano movers” problem. ii. general techniques for computing topological properties of real algebraic manifolds. *Advances in applied Mathematics*, 4(3):298–351, 1983.
- [18] I. R. Shafarevich. *Basic Algebraic Geometry*, volume 1. Springer, 1994.
- [19] N. Vorobjov and D. Y. Grigoriev. Determination of the number of connected components of a semi-algebraic set in subexponential time. *Dokl. Akad. Nauk SSSR*, 314(5), 1990.