



**HAL**  
open science

**The immune microenvironment in patients with mismatch-repair-proficient oligometastatic colorectal cancer exposed to chemotherapy: the randomized MIROX GERCOR cohort study**

Marine Jary, Wen-wei Liu, Dongyao Yan, Isaac Bai, Andrea Muranyi, Elise Colle, Isabelle Brocheriou, Anthony Turpin, Nina Radosevic-robin, Pierre Bourgoïn, et al.

► **To cite this version:**

Marine Jary, Wen-wei Liu, Dongyao Yan, Isaac Bai, Andrea Muranyi, et al.. The immune microenvironment in patients with mismatch-repair-proficient oligometastatic colorectal cancer exposed to chemotherapy: the randomized MIROX GERCOR cohort study. *Molecular Oncology*, 2021, 10.1002/1878-0261.13173 . hal-03510003

**HAL Id: hal-03510003**

**<https://hal.sorbonne-universite.fr/hal-03510003>**

Submitted on 4 Jan 2022

**HAL** is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers.

L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.

1

2 DR. MARINE JARY (Orcid ID : 0000-0001-7502-7708)

3

4

5 Received Date : 17-May-2021

6 Revised Date : 26-Nov-2021

7 Accepted Date : 23-Dec-2021

8 Article type : Research Article

9

10

11 **The immune microenvironment in patients with mismatch-repair-proficient oligometastatic**  
12 **colorectal cancer exposed to chemotherapy: the randomized MIROX GERCOR cohort**  
13 **study**

14

15 Marine Jary<sup>1,2</sup>, Wen-Wei Liu<sup>3</sup>, Dongyao Yan<sup>3</sup>, Isaac Bai<sup>3</sup>, Andrea Muranyi<sup>3</sup>, Elise Colle<sup>4</sup>, Isabelle  
16 Brocheriou<sup>5</sup>, Anthony Turpin<sup>1,6</sup>, Nina Radosevic-Robin<sup>7</sup>, Pierre Bourgoin<sup>8</sup>, Frédérique Penault-  
17 Llorca<sup>7</sup>, Romain Cohen<sup>1,9</sup>, Dewi Vernerey<sup>1,10,11</sup>, Thierry André<sup>1,9</sup>, Christophe Borg<sup>1,11,12</sup>, Kandavel  
18 Shanmugam<sup>3</sup> #, Magali Svrcek<sup>1,8</sup> #

19

20 #: *KS and MS contributed equally to this work*

21

22 1. Multidisciplinary Group in Oncology (GERCOR), Paris, France

23 2. Department of Surgical and Medical Oncology, University Hospital of Clermont-Ferrand,  
24 Clermont-Ferrand, France

25 3. Ventana Medical Systems Inc., Tucson, Arizona, USA

26 4. Department of Medical Oncology, University Hospital of Beaujon, Clichy, France

27 5. Sorbonne University, Department of Pathology, Assistance Publique-Hôpitaux de Paris, Pitié-  
28 Salpêtrière Hospital, Paris, France

This article has been accepted for publication and undergone full peer review but has not been through the copyediting, typesetting, pagination and proofreading process, which may lead to differences between this version and the [Version of Record](#). Please cite this article as [doi: 10.1002/1878-0261.13173](#)

Molecular Oncology (2020) © 2020 The Authors. Published by FEBS Press and John Wiley & Sons Ltd.

This is an open access article under the terms of the Creative Commons Attribution License, which permits use, distribution and reproduction in any medium, provided the original work is properly cited.

- 1 6. Department of Medical Oncology, University Hospital of Lille, Lille, France  
2 7. IMoST - Molecular Imaging & Theranostic Strategies, Clermont-Ferrand, France  
3 8. Sorbonne University, Department of Pathology, Assistance Publique-Hôpitaux de Paris, Saint-  
4 Antoine Hospital, Paris, France  
5 9. Sorbonne University, Department of Medical Oncology, Assistance Publique-Hôpitaux de  
6 Paris, Saint-Antoine Hospital, Paris, France  
7 10. Methodology and Quality of Life in Oncology Unit, Besançon University Hospital, Besançon,  
8 France  
9 11. INSERM, EFS BFC, UMR1098, RIGHT, University of Bourgogne Franche-Comté,  
10 Interactions hôte-greffon-tumeur/Ingénierie Cellulaire et Génique, Besançon, France  
11 12. Department of Medical Oncology, University Hospital of Besançon, Besançon, France  
12

13 *Running title:* Immune profile of pMMR omCRC  
14

15 **Keywords:**

16 Oligometastatic colorectal cancer, pMMR, immune profile, T lymphocytes, PD-L1  
17  
18

19 **Correspondence to:**

20 Dr Kandavel Shanmugam, Medical Innovation, Clinical Science, Roche Tissue Diagnostics, Tucson,  
21 AZ, USA; Email: kandavel.shanmugam@roche.com

22 Pr Magali SVRCEK, Sorbonne Université, Assistance Publique-Hôpitaux de Paris, Department of  
23 Pathology, Saint-Antoine Hospital, Paris, France; Tel: +33(0)149282179, Fax: +33(0)149282878,  
24 Email: magali.svrcek@aphp.fr  
25  
26

## 1 **ABBREVIATIONS**

- 2
- 3 CC1 buffer: Cell Conditioning 1 buffer
- 4 CEA: CarcinoEmbryonic Antigen
- 5 CI: Confidence Interval
- 6 CMS: Consensus Molecular Subtype
- 7 CRC: Colorectal cancer
- 8 DFS: Disease-free survival
- 9 dMMR/MSI: deficient mismatch repair /microsatellite instable
- 10 FFPE: Formalin-fixed, paraffin-embedded
- 11 FOXP3: Forkhead box protein 3
- 12 GERCOR: Groupe Coopérateur Multidisciplinaire en Oncologie
- 13 HR: Hazard ratios
- 14 ICIs: Immune checkpoint inhibitors
- 15 IE: intra-epithelial
- 16 IF: invasive front
- 17 IFN- $\gamma$ : Interferon Gamma
- 18 IHC: Immunohistochemistry
- 19 mCRC: metastatic CRC
- 20 MDSC: Myeloid derived suppressor cell
- 21 MEK: Mitogen-activated protein kinase kinase
- 22 MHC: Major histocompatibility complex
- 23 MLH1: MutL homolog 1
- 24 MSH2: MutS homolog 2
- 25 MSH6: MutS homolog 6
- 26 omCRC: oligometastatic CRC
- 27 PD-1: Programmed cell death protein 1
- 28 PD-L1: Programmed death ligand 1
- 29 pMMR/MSS: proficient mismatch repair/microsatellite stable
- 30 PMS2: PMS1 homolog 2
- 31 TILs: Tumor-infiltrating lymphocytes
- 32 TRG: Tumor regression grade

1

2

3 **ABSTRACT**

4 In the era of immune checkpoint inhibitors, understanding the metastatic microenvironment of  
5 proficient mismatch repair/microsatellite stable (pMMR/MSS) colorectal cancer (CRC) is of  
6 paramount importance to both prognostication and the development of more effective novel  
7 therapies. In this study, primary and paired metastasis tissue samples were collected from patients  
8 with resectable metastatic CRC treated with adjuvant FOLFOX or peri-operative chemotherapy in  
9 the MIROX phase III prospective study. In total, 74 cancer tissues were stained for CD3, CD8,  
10 Forkhead box protein 3 (FOXP3), Programmed cell Death protein-1(PD-1, invasive front, stromal,  
11 intra-epithelial compartments) and Programmed Death-Ligand 1 (PD-L1, tumor, immune cells).  
12 The immune profiling of primary CRC had a limited value to predict the immune context of paired  
13 metastases for all markers but CD3+. The expression of CD8 and PD-L1 was higher in metastases  
14 after neoadjuvant FOLFOX. In metastases, both CD3 T cells at the invasive front and PD-L1  
15 expressions on immune cells were predictive of better disease-free survival. These results show  
16 that the effect of FOLFOX on modifying the immune microenvironment in resected CRC  
17 metastases and measurement of PD-L1 expression and tumor-infiltrating CD8 T cells in  
18 pMMR/MSS metastatic tissue samples could improve treatment strategies of metastatic CRC  
19 patients.

20

21

22

23

## 1 1. Introduction

2 Accumulating evidence suggests that the adaptive immune system can influence cancer  
3 progression and that the quantification of tumor-infiltrating lymphocytes (TILs) may improve  
4 prognostic ability of the staging system in patients with solid tumors. In colorectal cancer (CRC),  
5 the impact of immune cell infiltration in the primary tumor on survival has been demonstrated  
6 [1,2]. Patients with metastatic CRC (mCRC) in the liver have heterogeneous clinical outcomes.  
7 Indeed, 70% of patients with curatively resected metastases will relapse and half of these will  
8 ultimately die [3,4]. Clinico-pathological prognostic factors like the tumor regression grade (TRG)  
9 have been proposed to identify patients who may be at risk for recurrence [5], but none of these  
10 markers has been sufficiently informative to correctly predict the outcome. In the era of  
11 personalized medicine, an identification of prognostic and predictive biomarkers is essential.  
12 Regarding mCRC, the immune microenvironment of the liver metastases reflects an important  
13 aspect of the overall portrait of the patient's disease, especially the heterogeneity compared to the  
14 primary tumor and its clinical impact [6–9]. The immune microenvironment of colorectal  
15 metastases has not been fully investigated, and published studies are often limited to CD4, CD8,  
16 and regulatory T cells [10]. In addition, several chemotherapy regimens such as oxaliplatin seem  
17 to have a preponderant role in anti-tumor immunologic infiltration, with a stimulating effect on the  
18 peritumoral immune response [11,12]. In particular, immunogenic cell death is provoked by  
19 FOLFOX and accompanied by tumor-targeting immune responses, release of damage-associated  
20 molecular patterns, and recruitment of antigen-presenting immune cells. Interestingly, the EORTC  
21 phase III study 40983 of 82 patients with resected colorectal liver metastases (38 in the surgery  
22 with perioperative FOLFOX chemotherapy arm and 44 in the surgery alone arm) [13] showed for  
23 the first time that chemotherapy influences immune cell profiles, independent of patient  
24 characteristics. In this latter study, abundance of CD3 T cell lymphocytes at the invasive margin of  
25 the resected metastasis specimens appeared to be prognostic. Moreover, immune infiltration of  
26 lymphocytes was associated with increased progression-free survival.

27 The efficacy of immune checkpoint inhibitors (ICIs) in mismatch repair deficient  
28 (dMMR)/microsatellite instable (MSI) mCRC is now well established [14–18]. However,  
29 questions remain regarding the role of ICIs for the treatment of MMR-proficient  
30 (pMMR)/microsatellite stable (MSS) mCRC. The combination of ICIs with other anticancer drugs  
31 is currently being evaluated in pMMR/MSS mCRC. The disappointing results of the phase III  
32 IMblaze 370 trial (atezolizumab with or without cobimetinib *versus* regorafenib) raise concerns

1 regarding the testing ICI-based strategies without decision-guiding biomarkers in pMMR/MSS  
2 mCRC. Contrarily, the NICHE study provided hypothesis-generating data for patients with  
3 localized pMMR/MSS colon cancer [19]. Thus, deeper understanding of the metastatic  
4 microenvironment of pMMR/MSS mCRC, and particularly oligometastatic CRC (omCRC) could  
5 improve selection of patients who may benefit from ICI combinations and other immunogenic  
6 drugs such as oxaliplatin.

7 Programmed Death-Ligand 1 (PD-L1) is expressed by tumor cells and certain immune cell  
8 types (dendritic cells, macrophages, B lymphocytes, Natural Killer cells). T-cells expressing  
9 Programmed cell Death protein-1 (PD-1) exhibit suppressed proliferation through PD-1/PD-L1  
10 interaction. However, PD-1 and PD-L1 expression in primary CRC are associated with favorable  
11 outcomes [20,21]. In most cancers treated with anti-PD-1 or anti-PD-L1 antibodies, the response  
12 rate is often higher in tumors expressing higher levels of PD-L1-positive immune cells. PD-L1-  
13 expressing tumor cells have been shown to regulate host immunity in the CRC microenvironment  
14 [22]. In the pMMR cohort of the NICHE study, the presence of T cells with co-expression of CD8  
15 and PD-1 was the only biomarker found to predict major or partial pathological response [19].  
16 Data regarding the expression of PD-L1 in CRC liver metastases and notably the interactions of  
17 PD-L1 with elements of the immune tumor microenvironment, as well as patient outcome, have  
18 recently been described [23]. No correlation between tumor-specific PD-L1 expression and  
19 survival was shown, confirming the results from other studies [24]. However, contradictory results  
20 are observed between the metastatic and early-stage settings [25], and few study yet assessed the  
21 expression of PD-L1 in a homogeneous cohort of matched primary and metastatic pMMR/MSS  
22 omCRC [26].

23 In this study we aimed to extensively characterize the immune microenvironment in  
24 patients with resectable mCRC treated or not with neoadjuvant FOLFOX to highlight an  
25 immunologic signature in this setting. We described potential tumor heterogeneity in chemo-naive  
26 patients with synchronous metastases and investigated if the immune infiltrate in resected  
27 colorectal metastases could be predictive of survival. The influence of FOLFOX-based  
28 chemotherapy and PD-1 and PD- L1 expression on the immune microenvironment and patient  
29 survival was further investigated.

## 1 2. MATERIALS AND METHODS

### 2 2.1. Study population

3 In total, 74 mCRC patients with available tissues from both primary tumors and paired  
4 metastases out of 284 included in the open-label prospective phase III MIROX trial were analyzed  
5 in this analysis. Patients with resectable or resected synchronous or metachronous metastases  
6 (only one site in liver, lung, ovary, or peritoneum) were treated with 6 cycles of FOLFOX4  
7 (oxaliplatin 85 mg/m<sup>2</sup>) or FOLFOX7 (oxaliplatin 130 mg/m<sup>2</sup>) before metastasis resection followed  
8 by adjuvant chemotherapy (FOLFOX or FOLFIRI). The dose of oxaliplatin was randomly  
9 assigned at the beginning of the study [27]. The primary CRC was resected before diagnosis of  
10 metastasis and neoadjuvant chemotherapy. This trial was approved by the local Ethics Committees  
11 at participating GERCOR (*Groupe Coopérateur Multidisciplinaire en Oncologie*) centers. All  
12 patients provided their written informed consent to receive treatment and participate in  
13 translational analysis.

14 Formalin-fixed, paraffin-embedded (FFPE) primary CRC tumor specimens were obtained  
15 prior to chemotherapy. Paired metastatic lesions were collected prior to adjuvant chemotherapy or  
16 after neoadjuvant chemotherapy and centralized at the Department of Pathology, Saint-Antoine  
17 Hospital (Paris, France), constituting the study cohort, BIOMIROX. Pathological response of CRC  
18 liver metastasis in patients treated with perioperative chemotherapy was estimated by the TRG  
19 pathological response score [5]. The study methodologies conformed to the standards set by the  
20 Declaration of Helsinki.

### 22 2.2. Immunohistochemistry analysis

23 Immunohistochemistry (IHC) staining was performed on serial sections from surgically  
24 resected specimens (VENTANA BenchMark ULTRA automated staining instrument at Pitié-  
25 Salpêtrière Hospital). Briefly, FFPE tissue sections were deparaffinized, pretreated with Cell  
26 Conditioning 1 for antigen retrieval, and treated to inactivate the endogenous peroxidase and then  
27 incubated with CONFIRM anti-CD3 (2GV6) rabbit monoclonal antibody, anti-CD8 (SP239)  
28 rabbit monoclonal antibody, anti-FOXP3 (SP97) rabbit monoclonal antibody, anti-PD-1  
29 (NAT105) mouse monoclonal antibody, and anti-PD-L1 (SP263) rabbit monoclonal antibody.  
30 Staining was visualized using the OptiView DAB IHC Detection Kit (Ventana Medical System,  
31 Inc.). Following detection, all slides were counterstained with hematoxylin II and bluing reagent  
32 for 4 minutes each, and coverslips were applied.



1 The density of tumor infiltrating immune CD3 and CD8 T cells was semi-quantitatively  
2 scored at the invasive front (IF; cells localized in stroma adjacent to the invasive tumor margin)  
3 and the intra-tumoral (or stromal) compartment [28] and graded as follows: 0, no positive cells; 1,  
4 scattered positive cells; 2, moderate number of positive cells; 3, abundant occurrence of positive  
5 cells. For CD3 T cells, the intra-epithelial (IE) compartment was also assessed. PD-1 and FOXP3  
6 were scored semi-quantitatively, both at the IF and stromal compartment, as follows: 0, no  
7 positive cell or scattered cells; 1, moderate number of positive cells, and 2, numerous positive  
8 cells. Positive PD-L1 expression was defined as any staining  $\geq 1\%$ , in either infiltrating  
9 inflammatory cells or membranous-site tumor cells [29]. The percentage of cells demonstrating  
10 PD-L1 staining was scored in 5% increments and high PD-L1 level was defined as  $\geq 5\%$ , based on  
11 published literature [30-34]. Tissue samples were evaluated by an experienced pathologist (MS)  
12 blinded to clinical information, treatment regimens, and outcomes (**Table S1**).

13 Tumor tissue sections were double-stained for PD-L1 and CD8 using a fully automated  
14 procedure in a Benchmark Ultra automate (Ventana/Roche). Epitope retrieval was performed in  
15 Cell Conditioning 1 (CC1 buffer) for 60 min at 95°C. PD-L1 was detected by Ventana PD-L1  
16 SP263 Assay (Roche Diagnostics, Meylan, France), per manufacturer's instructions. CD8 was  
17 detected by clone SP239 (Abcam, Cambridge, UK) at 1/100 for 60 min at room temperature. For  
18 PD-L1, the antigen-antibody reaction was revealed by OptiView DAB IHC Detection Kit, and for  
19 CD8 by UltraView Universal AP Red Detection Kit (red signal), both from Roche Diagnostics  
20 (Meylan, France). Co-staining was appreciated by a semi-quantitative method.

21 The expression of MutL homolog 1 (MLH1, dilution 1/70, clone G168-728, Pharmingen, San  
22 Diego, CA), MutS homolog 2 (MSH2, dilution 1/100, clone FE11, Calbiochem, Oncogene  
23 Research Products, Cambridge, MA), MutS homolog 6 (MSH6, dilution 1/100, clone 44, Becton  
24 Dickinson, Lexington, NC), and PMS1 homolog 2 (PMS2, clone A16-4, 1:150 dilution, BD  
25 PharMingen, Le Pont de Claix, France) was assessed. Immunostaining of MLH1, MSH2, MSH6,  
26 and PMS2 in tumor cells was evaluated as positive (mismatch repair [MMR]-proficient [pMMR])  
27 or negative (MMR-deficient [dMMR]). Tumors were considered negative when there was a  
28 complete absence of nuclear staining of neoplastic cells in the presence of an internal positive  
29 control.

### 30 31 2.3. Statistical analysis

1 All IHC markers but PD-L1 were semi-quantitatively assessed. Therefore, they were  
2 treated as categorical variables in this study. The Cox proportional-hazards model was used to  
3 estimate hazard ratios (HRs) and *P* values with disease-free survival (DFS). The association  
4 between clinical, biomarker parameters, and survival was estimated with univariate Cox  
5 proportional hazards models and HR with 95% Confidence Interval (CI) were given. Multivariate  
6 Cox models were investigated including clinico-biological parameters with *p* value <0.05 in  
7 univariate analysis. *P* value <0.05 was considered statistically significant. Spearman correlations  
8 between metastatic immune infiltrate and TRG scoring were estimated. Analyses were performed  
9 by SAS version 9.4 (SAS Institute Inc, Cary, NC).

10

11

## 12 3. RESULTS

### 13 3.1. Study cohort characteristics

14 A total of 74 patients with pMMR CRC were included in the current study. All baseline  
15 characteristics are summarized in **Table S2** and in the flowchart (**Fig. S1**). The median age of  
16 patients was 61 years (range 29-75). Forty-one patients had nearby lymph node metastases of the  
17 primary, and 82.6% were left-sided tumors (including rectum). Most patients (*n* = 65) had liver-  
18 only metastases and nine had another only one site of distant metastases (lung [*n* = 3], ovary [*n* =  
19 1], peritoneal location [*n* = 5]). The median number of metastases was 2 (range, 1-7). Twenty-four  
20 patients had metachronous metastases. The mean preoperative CarcinoEmbryonic Antigen (CEA)  
21 level was 55.9 ng/mL (*n* = 70). Out of the 74 patients analyzed, 34 received adjuvant  
22 chemotherapy after surgery of their metastases (“chemo-naïve” patients). Among patients treated  
23 with neoadjuvant chemotherapy, two did not receive oxaliplatin.

24

### 25 3.2. Immune profiling between the primary CRC and matched metastasis

26 The distribution of semi-quantitative scores of CD3, CD8, FOXP3, PD-1, and PD-L1 (**Fig.**  
27 **1, Table 1, Table S1, Fig.S2**) showed considerable heterogeneity, both in primary tumors and  
28 metastases, in chemo-naïve patients and those treated with neoadjuvant chemotherapy before  
29 surgery. Infiltration of CD3 T lymphocytes was the strongest in stroma and the IF both in primary  
30 tumors and metastases.

1 To avoid the potential confounding effect of chemotherapy on immune cell infiltration, we  
2 further examined the distribution scores by restricting the analyses to 34 chemo-naïve patients  
3 only (**Fig. S2**). The vast majority of these patients ( $n = 31$ ) demonstrated high density of CD3 T  
4 cells (IHC score 2-3) in the IF of metastatic sites (liver [ $n = 25$ ], lung [ $n = 2$ ], ovary [ $n = 1$ ],  
5 peritoneal [ $n = 3$ ]). Seventeen patients in the chemo-naïve cohort had a high expression of CD3  
6 cells (IHC 2 and 3) in the IF of the primary tumor (**Table 1**). Interestingly, this expression was  
7 strongly correlated (16 out of 17 patients, 94%) with that observed in the metastases compared but  
8 it was not the case for stromal and intraepithelial compartments CD3 cells (IHC 2 and 3) of the  
9 primary tumor and their matched metastases (**Table 1 and Table S1**).

10 PD-1 overexpression on CD8 T cells is known as an exhaustion biomarker, but it also  
11 reflects antigen-experienced lymphocytes. Conversely, PD-L1 expression on tumor or immune  
12 cells is induced by an interferon-mediated signaling. We characterized therefore CD3 and CD8  
13 expression according to PD-L1 expression in the pMMR/MSS mCRC study cohort. A high level  
14 of PD-L1 expression was observed predominantly in immune cells (20 [58.8%] vs 3 [8.8%] in  
15 tumoral cells; **Table S1**) in CRC. Among the 33 patients for whom CD8 staining was available,  
16 four had high CD8 immune infiltrate (IHC score 2-3) in the IF of the primary tumor (**Table S1**).  
17 Three of the latter patients had a PD-L1 expression  $>5\%$  in immune cells (CD8<sup>high</sup>/PD-L1<sup>high</sup>  
18 patients; **Fig. 1**). Of the 29 patients with low CD8 score, 17 showed high PD-L1 expression  
19 (CD8<sup>low</sup>/PD-L1<sup>high</sup> patients; **Table 1 and Table S1**). By comparison, twenty-six patients had low  
20 CD8 immune infiltrate in the IF of metastases, 10 of whom were classified as PD-L1<sup>low</sup> ( $<5\%$ ;  
21 CD8<sup>low</sup>/PD-L1<sup>low</sup>). Seven patients had high CD8 T cell infiltration in the IF. In all these patients, a  
22 high PD-L1 score ( $\geq 5\%$  of expression by immune cells; **Fig. 1, Table 1, and Table S1**) was  
23 observed. In order to better appreciate the co-localization of CD8 and PD-L1, their co-staining was  
24 performed in six of these seven patients. The co-localization of CD8 and PD-L1 was observed in  
25  $\geq 20\%$  of inflammatory cells. In four out of seven cases, a co-localization was detected in over  
26 50% of cells. Negative controls had no more than 10% of co-localization detected (**Fig. S3**).

27 Regulatory T cells in CRC have been described in the non-inflamed tumors because of  
28 their tolerance properties, but they also play a role for effector T cells, which is to maintain  
29 immune homeostasis, even when the anti-tumor immune response is active [35-37]. In our study,  
30 half of the CD8<sup>high</sup>/PD-L1<sup>high</sup> patients had high immune IF FOXP3 infiltration.

31 Contrary to CD3 staining, the analysis showed that CD8 and PD-L1 expressions were  
32 slightly less correlated between the primary tumors and matched metastases. Fifteen of the 20

1 patients with a high PD-L1 expression (i.e. at least 5% of positive immune or tumoral cells) in the  
2 primary tumor had also high PD-L1 expression in the matched metastasis (**Table 1 and Table S1**).  
3 Three out of four patients with high CD8 expression in the primary tumor had also more CD8 T-  
4 cell infiltration in their metastasis. Three patients showing positive PD-L1 staining in the primary  
5 tumor were negative/low for PD-L1 in the matched metastasis but showed a high CD3 immune  
6 infiltration (IHC score 2-3) in the IF of the metastatic tumor.

7 The above data suggested that the immune profiling performed on the primary tumor has a  
8 limited predictive value to estimate the immune context of metastasis. Further analysis focused on  
9 the tumor-infiltrating immune cell in metastasis.

### 10 11 **3.3. Patients' characteristics according to the expression of IF CD3, IF CD8, and immune** 12 **PD-L1 in metastases of chemo-naïve patients**

13 The clinical characteristics of 34 chemo-naïve patients according to the expression of IF  
14 CD3, IF CD8, and immune cells PD-L1 in metastatic tumors are described in **Table 2**. Most of  
15 patients with high IF CD3 expression in metastatic tumors (out of 31 [96.7%]; IHC score 2+ and  
16 3+) and all patients with and CD8+ and PD-L1 in metastases ( $n = 7$ ) had tumors less than 5 cm in  
17 diameter and 70% (21 out of 31 patients) and 85%, respectively, had only one metastatic site. The  
18 CEA level before metastases resection was low in these patients.

### 19 20 **3.4. Patient and biomarker characteristics according to neoadjuvant chemotherapy: the** 21 **impact of neoadjuvant FOLFOX on immune infiltrate in metastases**

22 Oxaliplatin-based chemotherapy can release antigens and promote immunogenic cell death  
23 leading to a specific anti-tumor immune response. Therefore, we further examined the distribution  
24 and type of the immune cell infiltration in patients with or without neoadjuvant FOLFOX  
25 chemotherapy (**Table 1, Table S3**). Patients with chemotherapy-treated metastatic sites had  
26 metastases strongly (IHC 3) positive for CD3 cells at IF (28.2% versus 14.7%), moderately (IHC  
27 2)/strongly positive for CD3 cells in stroma (47.5% versus 32.4%), and moderately/strongly  
28 positive for CD8 cells at IF (41.1% versus 21.2%) than patients whose metastases were not  
29 exposed to chemotherapy. FOXP3 reg T cell staining was relatively weak but significantly  
30 decreased in the IF after chemotherapy ( $P = 0.0010$ ). Of interest, seven chemo-naïve patients (out

1 of 33; 21%) had high CD8 and high /PD-L1 staining versus 13 who received FOLFOX-based  
2 chemotherapy (out of 38; 34%).

3 These results suggested that chemotherapy may impact the immune microenvironment of  
4 patients and increase CD8 and PD-L1 expression in metastases.

5  
6 Finally, a significant inverse correlation between CD3<sup>high</sup> T cell infiltration in stroma and  
7 TRG was observed, reflecting a better pathological response in CD3 T cells-inflamed tumors  
8 (Spearman correlation -0.33,  $P = 0.0484$ ; **Table S4**).

### 9 10 3.5. Association between the immune infiltrate and survival in metastases

11 In the univariate analysis, low IF CD3 in metastases correlated with a shorter DFS (**Table**  
12 **3**). Patients with a higher number of metastases and neoadjuvant chemotherapy had shorter DFS.  
13 The only variable identified as a prognostic factor for DFS in the multivariate analysis was IF  
14 CD3<sup>high</sup> T-cell infiltrate (HR = 0.31, 95% CI: 0.15-0.67,  $P = 0.002$ ). DFS was significantly better  
15 in patients with high IF CD3 expression (IHC 2-3) than in those with low IF CD3 expression  
16 (median DFS of 2.2 years, 95% CI: 1.2-3.9 versus 0.59 years, 95% CI: 0.13-1.13, respectively, HR  
17 = 0.36,  $P = 0.005$ , **Fig. 2A**).

18 A similar pattern of differential DFS according to the IF CD3<sup>high</sup> score in metastases was  
19 observed when the analysis was restricted to patients with synchronous metastases (**Fig. S4**).  
20 Given the relatively small size of our cohort, we did not analyze IF CD3<sup>high</sup> T cell expression  
21 impact on DFS in patients with metachronous metastases.

22 According to stratification based on PD-L1 expression by immune cells in metastases, the median  
23 DFS was significantly better in patients with high PD-L1 expression (the median DFS = 2.56  
24 years, 95% CI: 1.11-8.34 versus 1.17 years, 95% CI: 0.73-2.22, respectively, HR = 0.58, 95% CI:  
25 0.34-1.02,  $P = 0.05$ , **Fig. 2B**). The DFS was not significantly different regarding metastatic  
26 CD8<sup>high</sup> expression in stroma or at the IF.

27 A schematic graphical summary of the key points is presented in **Fig. S5**.

28

## 1 4. DISCUSSION

2

3 Assessment of tumor-immune infiltrating cells is emerging as an important prognostic tool  
4 to stratify cancer patients according to the immune microenvironment. However, there are  
5 currently limited data on such analysis between the primary and paired metastatic tumor in  
6 patients with mCRC. In this study, we observed the intra-patient heterogeneity of immune  
7 infiltrates in both chemotherapy-naïve primary tumors and in the matched metastases (mainly in  
8 the liver) in 74 patients with mCRC.

9 Many studies explored immune infiltrate across tumorigenesis, with approaches outlining  
10 the difference in the immune contexture between primary and metastatic tumors [38-39]. In the  
11 study by Angelora et al., a thorough genomic and immunological characterization of primary and  
12 the matched metastatic tumor of two patients with CRC did not show any correlation, suggesting a  
13 high level of tumor heterogeneity between all lesions [38]. This observation was further clearly  
14 confirmed by a Consensus Molecular Subtype (CMS) characterization in omCRC patients [39].  
15 The authors showed that CMS subtypes, determined in patients undergoing partial hepatectomy of  
16 resectable CRC liver metastases, were variable. Gene expression analysis showed the absence of  
17 CMS1 (1%) and CMS3 (0%) subtypes in liver metastases and their presence in in the primary  
18 tumors (14% and 13%, respectively). Therefore, the immune profiling performed on the primary  
19 CRC has a limited predictive value to estimate the immune contexture in metastases.

20 In our analysis, the immune infiltrate in the primary tumor was not associated with survival  
21 (data not shown), except for stromal FOXP3. Previously published reports showed discordant  
22 results regarding the prognostic value of FOXP3 probably reflecting the plasticity of these  
23 immune subsets. Our results provide evidence that CD3 score in the IF of metastases has an  
24 independent prognostic value for DFS, doubling the survival rate, independent of TRG, further  
25 confirming the previously published data [1,13,40]. Contrary to Mlecnik et al. [2], we did not  
26 observe higher IF CD3 in patients with lower TRG (data not published). The median DFS was  
27 also significantly better in patients with higher immune PD-L1 expression in metastases in our  
28 study. In another study, PD-L1 expression in tissue microarray of surgically excised pMMR CRC  
29 patients was associated with improved overall survival, but analyses were performed only on  
30 primary tumors [20]. Considering these observations, it appears more pertinent to explore and  
31 characterize the immune infiltrate of metastases and to correlate it with patient prognosis for  
32 stratification of patients for appropriate therapies.

1 An efficient immune response is characterized by activation of the interferon signaling  
2 pathway generating mature cytotoxic lymphocytes. PD-L1 expression on tumor or immune cells is  
3 induced by an interferon-mediated signaling and hence a subpopulation of patients with such  
4 expression seems to be of particular interest. In general, CD8<sup>high</sup>/PD-L1<sup>high</sup> cells were previously  
5 described as functional effector cells as they produce significantly higher level of Interferon  
6 Gamma (IFN- $\gamma$ ) and express more the degranulation marker CD107a than CD8<sup>low</sup>/PD-L1<sup>low</sup> cells  
7 [41]. Although we did not assessed survival for the immune PD-L1 and CD8<sup>high</sup> groups due to the  
8 small number of patients in each group, the characterization of this subgroup was of importance in  
9 pMMR CRC patients. For the first time, we showed that this entity seems to be associated with  
10 lower tumoral mass and lower CEA. Lower stroma stiffness is one of the hypotheses explaining  
11 the enhancement of the CD8<sup>high</sup>/PD-L1<sup>high</sup> population, supported by recent observation correlating  
12 desmoplastic angiogenic stroma and CD8<sup>high</sup> T cell immune infiltration in CRC liver metastasis  
13 [42]. This is in accordance with the previous observations by Wang et al. [40], showing the  
14 immune infiltrate associated with metastatic size, number of metastases, and Fong clinical Risk  
15 Score in resected CRC liver metastases.

16 The effect of chemotherapy on the CD8<sup>high</sup>/PD-L1<sup>high</sup> signature is of importance to better  
17 personalize treatment strategies. In our cohort, we observed significantly more patients harboring  
18 CD8<sup>high</sup>/PD-L1<sup>high</sup> staining in the group exposed to neoadjuvant FOLFOX-based chemotherapy.  
19 The T lymphocyte density and location in metastatic melanomas were reported to have predictive  
20 value for treatment outcome of patients receiving anti-PD-1/PD-L1 therapies [43]. Although an  
21 increase in CD8<sup>high</sup> and PD-L1 immune infiltrate by chemotherapy did not significantly correlate  
22 with survival, we hypothesize that this specific subgroup of patients as identified herein may  
23 benefit from personalized treatment including immune checkpoint inhibitors. Effect of  
24 chemotherapy on the immune infiltrate was described in non-metastatic rectal cancer, with an  
25 increased stromal CD8<sup>high</sup> and CD4 T cells after chemoradiotherapy, associated with a better  
26 prognosis [44]. Recently, the pooled analysis of post-chemotherapy resected metastases of pMMR  
27 CRC confirmed the good prognosis of patients with such an inflamed microenvironment [23].  
28 However, this analysis did not include paired tumors before and after treatment. Using a  
29 combination of chemotherapies with immune checkpoint inhibitors may improve response rate, as  
30 recent data showed up to 27% of pathological response in pMMR CRC patients after nivolumab  
31 and ipilimumab neoadjuvant therapy [19]. The tissue immune profiling could allow design of  
32 immunotherapies for the CD8<sup>high</sup>/PD-L1<sup>high</sup> subset of patients in the adjuvant setting of

1 metastasectomy. It could also help to plan strategies in the following chemo-immunotherapy lines  
2 of treatment. This is consistent with the recent results reported by Kumagai et al. [45] showing that  
3 a profound reactivation of effector PD-1<sup>+</sup>CD8<sup>+</sup> T cells is necessary for tumor regression, which  
4 paves the way for a promising predictive biomarker for PD-1 blockade therapies.

5 Our study has several limitations. Firstly, the immune infiltrate was investigated in only  
6 one metastatic lesion per patient. A major obstacle to a refined definition of the immune  
7 contexture of human CRC liver metastases resides in the heterogeneity between the different  
8 metastases in the same patient, in addition to the profound heterogeneity of tumor lesions across  
9 patients. Galon et al. demonstrated that the immune phenotype of the least-infiltrated metastasis  
10 had a stronger association with patient outcome than other metastases [6]. This effort of selection  
11 could not be performed here. Nevertheless, in our study, IHC analyses were performed with the  
12 semi-quantitative method on the entire slide within three separate compartments of the tumor.  
13 Secondly, this study does not include *RAS* mutational status. It was reported that CRC harboring  
14 *RAS* mutations or Mitogen-activated protein kinase kinase (MEK) activation had less major  
15 histocompatibility complex-I (MHC-I) expression and lower CD8 T cell activation [46].  
16 Neoadjuvant chemotherapy was previously shown to enhance CD8 and immune PD-L1 expression  
17 in metastases of *RAS* wild-type cancers [47], and this will be crucial to analyze in validation  
18 cohorts. Finally, the peripheral immune response could not be assessed in complement to the intra-  
19 tumoral approach. The initiation of cell death and subsequent activation of T cells when antigens  
20 are released can be monitored by different methods. The peripheral immune response against  
21 tumor antigens before and after administration of the FOLFOX regimen has been previously  
22 assessed in mCRC patients [48]. An epitope spreading stimulating immune response against a  
23 broad spectrum of tumor antigens hinders monitoring the effect of immune-based chemotherapy in  
24 the circulating blood of patients. Peripheral blood leucocytes phenotypic profiling is easy to  
25 perform and decreased levels of myeloid-derived suppressor cells (MDSC) and increased levels of  
26 circulating CD8<sup>+</sup> T cells lymphocytes after 5-fluorouracil-based chemotherapy have been detected  
27 by flow cytometric analyzes [49]. Doublet chemotherapies with the FOLFOX +/- bevacizumab  
28 induce a significant decrease in the number of MDSC, specifically granulocytic MDSC, which  
29 was associated with better progression-free survival in patients receiving this combination [50-51].  
30 A combined peripheral and intra-tumoral assessment of the immune response is a promising  
31 approach for the future studies.

32



## 1 **5. CONCLUSION**

2 In conclusion, our findings suggest an effect of chemotherapy on modifying the immune  
3 microenvironment in resected CRC metastases and highlight the relevance of CD8<sup>high</sup>/PD-L1<sup>high</sup> in  
4 pMMR/MSS metastases for adjuvant treatments including immunotherapy strategies. This key  
5 immune signature based on the assessment of CD8 and PD-L1 by IHC, adapted for a routine  
6 practice and a cost-effective method, paves the way for further prospective analyses in mCRC and  
7 encourages the efforts for such rare data sharing.

## 8 9 **ACKNOWLEDGMENTS**

10 We thank Linh Pham for managing the project. The authors also thank the technical staff of the  
11 Department of Pathology of Pitié-Salpêtrière and Magdalena Benetkiewicz (Sc.D.; GERCOR) for  
12 editorial assistance.

### 13 14 **Data Accessibility:**

15 The datasets used and/or analyzed during the present study are available within the supplementary  
16 figures and tables.

17  
18 **Disclosure:** The authors have declared no conflict of interest.

### 19 **Author contributions**

20 MJ: Formal analysis, Validation, Writing-original draft

21 WWL: Statistical analysis, Critically reviewal of the manuscript

22 DY: Formal analysis, Critically reviewal of the manuscript

23 IB: Statistical analysis, Critically reviewal of the manuscript

24 AM, EC, AT, RC, DV: Critically reviewal of the manuscript

25 IB: Data curation, Critically reviewal of the manuscript

26 NR, PB, FPL: Complementary investigation, Critically reviewal of the manuscript

27 TA: Conceptualization, Formal analysis, Validation, Investigation, Critically reviewal of the  
28 manuscript

29 CB, KS: Formal analysis, Validation, Critically reviewal of the manuscript

30 MS: Conceptualization, Data curation, Formal analysis, Validation, Investigation, Writing-original  
31 draft

32

1 **Financial support:** No funding was received for this article.

2

Accepted Article

## 1 REFERENCES

- 2 1. Galon J, Costes A, Sanchez-Cabo F, Kirilovsky A, Mlecnik B, Lagorce-Pagès C, Tosolini M,  
3 Camus M, Berger A, Wind P, et al (2006). Type, density, and location of immune cells within  
4 human colorectal tumors predict clinical outcome. *Science* 313,1960–4.
- 5 2. Mlecnik B, Tosolini M, Kirilovsky A, Berger A, Bindea G, Meatchi T, Bruneval P,  
6 Trajanoski Z, Fridman WH, Pagès F, et al (2011). Histopathologic-Based Prognostic Factors  
7 of Colorectal Cancers Are Associated With the State of the Local Immune Reaction. *J Clin*  
8 *Oncol. American Society of Clinical Oncology* 29,610–8.
- 9 3. Siegel RL, Miller KD, Sauer AG, Fedewa SA, Butterly LF, Anderson JC, Cercek A, Smith  
10 RA, Jemal A (2020). Colorectal cancer statistics. *CA Cancer J Clin.* 70,145–64.
- 11 4. Manfredi S, Lepage C, Hatem C, Coatmeur O, Faivre J, Bouvier A-M (2006). Epidemiology  
12 and management of liver metastases from colorectal cancer. *Ann Surg.* 244, 254–9.
- 13 5. Rubbia-Brandt L, Giostra E, Brezault C, Roth AD, Andres A, Audard V, Sartoretti P, Dousset  
14 B, Majno PE, Soubrane O, et al (2007). Importance of histological tumor response assessment  
15 in predicting the outcome in patients with colorectal liver metastases treated with neo-  
16 adjuvant chemotherapy followed by liver surgery. *Ann Oncol Off J Eur Soc Med Oncol.* 18,  
17 299–304.
- 18 6. Van den Eynde M, Mlecnik B, Bindea G, Fredriksen T, Church SE, Lafontaine L, Haicheur  
19 N, Marliot F, Angelova M, Vasaturo A, et al (2018). The Link between the Multiverse of  
20 Immune Microenvironments in Metastases and the Survival of Colorectal Cancer Patients.  
21 *Cancer Cell* 34, 1012-1026.e3.
- 22 7. Halama N, Michel S, Kloor M, Zoernig I, Benner A, Spille A, Pommerenke T, von Knebel  
23 DM, Folprecht G, Luber B, et al (2011). Localization and density of immune cells in the  
24 invasive margin of human colorectal cancer liver metastases are prognostic for response to  
25 chemotherapy. *Cancer Res.* 71, 5670–7.
- 26 8. Maker AV, Ito H, Mo Q, Weisenberg E, Qin L-X, Turcotte S, Maithel S, Shia J, Blumgart L,  
27 Fong Y, et al (2015). Genetic evidence that intratumoral T-cell proliferation and activation are  
28 associated with recurrence and survival in patients with resected colorectal liver metastases.  
29 *Cancer Immunol Res.* 3, 380–8.
- 30 9. Halama N, Spille A, Lerchl T, Brand K, Herpel E, Welte S, Keim S, Lahrmann B, Klupp F,  
31 Kahlert C, et al (2013). Hepatic metastases of colorectal cancer are rather homogeneous but

- 1 differ from primary lesions in terms of immune cell infiltration. *Oncoimmunology*. 2:e24116.
- 2 10. Zwing N, Failmezger H, Ooi C-H, Hibar DP, Cañamero M, Gomes B, Gaire F, Korski K  
3 (2020). Analysis of Spatial Organization of Suppressive Myeloid Cells and Effector T Cells in  
4 Colorectal Cancer—A Potential Tool for Discovering Prognostic Biomarkers in Clinical  
5 Research. *Front Immunol*. 11, 550250
- 6 11. Zitvogel L, Apetoh L, Ghiringhelli F, André F, Tesniere A, Kroemer G (2008). The  
7 anticancer immune response: indispensable for therapeutic success? *J Clin Invest*. 118, 1991–  
8 2001.
- 9 12. Vanmeerbeek I, Sprooten J, De Ruyscher D, Tejpar S, Vandenberghe P, Fucikova J, Spisek  
10 R, Zitvogel L, Kroemer G, Galluzzi L, et al (2020). Trial watch: chemotherapy-induced  
11 immunogenic cell death in immuno-oncology. *Oncoimmunology* 9, 1703449.
- 12 13. Tanis E, Julié C, Emile J-F, Mauer M, Nordlinger B, Aust D, Roth A, Lutz MP, Gruenberger  
13 T, Wrba F, et al (2015). Prognostic impact of immune response in resectable colorectal liver  
14 metastases treated by surgery alone or surgery with perioperative FOLFOX in the randomised  
15 EORTC study 40983. *Eur J Cancer* 51, 2708–17.
- 16 14. Le DT, Uram JN, Wang H, Bartlett BR, Kemberling H, Eyring AD, Skora AD, Luber BS,  
17 Azad NS, Laheru D, et al (2015). PD-1 Blockade in Tumors with Mismatch-Repair  
18 Deficiency. *N Engl J Med*. 372, 2509–20.
- 19 15. Le DT, Durham JN, Smith KN, Wang H, Bartlett BR, Aulakh LK, Lu S, Kemberling H, Wilt  
20 C, Luber BS, et al (2017). Mismatch-repair deficiency predicts response of solid tumors to  
21 PD-1 blockade. *Science* 357, 409–13.
- 22 16. Overman MJ, McDermott R, Leach JL, Lonardi S, Lenz H-J, Morse MA, Desai J, Hill A,  
23 Axelson M, Moss RA, et al (2017). Nivolumab in patients with metastatic DNA mismatch  
24 repair-deficient or microsatellite instability-high colorectal cancer (CheckMate 142): an open-  
25 label, multicentre, phase 2 study. *Lancet Oncol*. 18, 1182–91.
- 26 17. Overman MJ, Lonardi S, Wong KYM, Lenz H-J, Gelsomino F, Aglietta M, Morse MA, Van  
27 Cutsem E, McDermott R, Hill A, et al (2018). Durable Clinical Benefit With Nivolumab Plus  
28 Ipilimumab in DNA Mismatch Repair-Deficient/Microsatellite Instability-High Metastatic  
29 Colorectal Cancer. *J Clin Oncol Off J Am Soc Clin Oncol*. 36, 773–9.
- 30 18. André T, Shiu K-K, Kim TW, Jensen BV, Jensen LH, Punt C, Smith D, Garcia-Carbonero R,  
31 Benavides M, Gibbs P, et al (2020). Pembrolizumab in Microsatellite-Instability-High  
32 Advanced Colorectal Cancer. *N Engl J Med*. 383, 2207–18.

- 1 19. Chalabi M, Fanchi LF, Dijkstra KK, Van den Berg JG, Aalbers AG, Sikorska K, Lopez-  
2 Yurda M, Grootscholten C, Beets GL, Snaebjornsson P, et al (2020). Neoadjuvant  
3 immunotherapy leads to pathological responses in MMR-proficient and MMR-deficient early-  
4 stage colon cancers. *Nat Med.* 26, 566–76.
- 5 20. Droeser RA, Hirt C, Viehl CT, Frey DM, Nebiker C, Huber X, Zlobec I, Eppenberger-Castori  
6 S, Tzankov A, Rosso R, et al (2013). Clinical impact of programmed cell death ligand 1  
7 expression in colorectal cancer. *Eur J Cancer* 49, 2233–42.
- 8 21. Li Y, Liang L, Dai W, Cai G, Xu Y, Li X, Li Q, Cai S (2016). Prognostic impact of  
9 programed cell death-1 (PD-1) and PD-ligand 1 (PD-L1) expression in cancer cells and tumor  
10 infiltrating lymphocytes in colorectal cancer. *Mol Cancer* 15:55.
- 11 22. Masugi Y, Nishihara R, Yang J, Mima K, da Silva A, Shi Y, Inamura K, Cao Y, Song M,  
12 Nowak JA, et al (2017). Tumour CD274 (PD-L1) expression and T cells in colorectal cancer.  
13 *Gut.* 66, 1463–73.
- 14 23. Moretto R, Corallo S, Belfiore A, Rossini D, Boccaccino A, Lonardi S, Centonze G, Morano  
15 F, Germani MM, Loupakis F, et al (2020). Prognostic impact of immune-microenvironment  
16 in colorectal liver metastases resected after triplets plus a biologic agent: A pooled analysis of  
17 five prospective trials. *Eur J Cancer Oxf Engl* 1990. 135, 78–88.
- 18 24. Berntsson J, Eberhard J, Nodin B, Leandersson K, Larsson AH, Jirström K (2018).  
19 Expression of programmed cell death protein 1 (PD-1) and its ligand PD-L1 in colorectal  
20 cancer: Relationship with sidedness and prognosis. *Oncoimmunology* 7:e1465165.
- 21 25. Ni X, Sun X, Wang D, Chen Y, Zhang Y, Li W, Wang L, Suo J (2019). The  
22 clinicopathological and prognostic value of programmed death-ligand 1 in colorectal cancer: a  
23 meta-analysis. *Clin Transl Oncol.* 21(5), 674-686.
- 24 26. Ahtiainen M, Elomaa H, Väyrynen JP, Wirta EV, Kuopio T, Helminen O, Seppälä TT,  
25 Kellokumpu I, Mecklin JP (2021). Immune Contexture of MMR-Proficient Primary  
26 Colorectal Cancer and Matched Liver and Lung Metastases. *Cancers (Basel)* 13(7), 1530.
- 27 27. Hebbbar M, Chibaudel B, André T, Mineur L, Smith D, Louvet C, Dutel JL, Ychou M, Legoux  
28 JL, Mabro M, et al (2015). FOLFOX4 versus sequential dose-dense FOLFOX7 followed by  
29 FOLFIRI in patients with resectable metastatic colorectal cancer (MIROX): a pragmatic  
30 approach to chemotherapy timing with perioperative or postoperative chemotherapy from an  
31 open-label, randomized phase III trial. *Ann Oncol.* 26, 340–7.
- 32 28. Dahlin AM, Henriksson ML, Van Guelpen B, Stenling R, Oberg A, Rutegård J, Palmqvist R

- 1 (2011). Colorectal cancer prognosis depends on T-cell infiltration and molecular  
2 characteristics of the tumor. *Mod Pathol Off J U S Can Acad Pathol Inc.* 24, 671–82.
- 3 29. Ma C, Patel K, Singhi AD, Ren B, Zhu B, Shaikh F, Sun W (2016). Programmed Death-  
4 Ligand 1 Expression Is Common in Gastric Cancer Associated With Epstein-Barr Virus or  
5 Microsatellite Instability. *Am J Surg Pathol.* 40, 1496–506.
- 6 30. Hellmann MD, Kim T-W, Lee CB, Goh B-C, Miller WH, Oh D-Y, et al (2019). Phase Ib study  
7 of atezolizumab combined with cobimetinib in patients with solid tumors. *Ann Oncol Off J*  
8 *Eur Soc Med Oncol.* 30, 1134–42
- 9 31. Kong P, Wang J, Song Z, Liu S, He W, Jiang C, et al (2019). Circulating Lymphocytes, PD-L1  
10 Expression on Tumor-infiltrating Lymphocytes, and Survival of Colorectal Cancer Patients  
11 with Different Mismatch Repair Gene Status. *J Cancer* 10, 1745–54.
- 12 32. Taube JM, Klein A, Brahmer JR, Xu H, Pan X, Kim JH, et al (2014). Association of PD-1,  
13 PD-1 ligands, and other features of the tumor immune microenvironment with response to  
14 anti-PD-1 therapy. *Clin Cancer Res Off J Am Assoc Cancer Res.* 20, 5064–74.
- 15 33. Kim JH, Park HE, Cho N-Y, Lee HS, Kang GH (2016). Characterisation of PD-L1-positive  
16 subsets of microsatellite-unstable colorectal cancers. *Br J Cancer* 115, 490–6.
- 17 34. Lee KS, Kim BH, Oh H-K, Kim D-W, Kang S-B, Kim H, et al (2018). Programmed cell death  
18 ligand-1 protein expression and CD274/PD-L1 gene amplification in colorectal cancer:  
19 Implications for prognosis. *Cancer Sci.* 109, 2957–69.
- 20 35. Chen DS, Mellman I (2017). Elements of cancer immunity and the cancer-immune set point.  
21 *Nature* 541, 321–30.
- 22 36. Müller P, Kreuzaler M, Khan T, Thommen DS, Martin K, Glatz K, Savic S, Harbeck N, Nitz  
23 U, Gluz O, et al (2015). Trastuzumab emtansine (T-DM1) renders HER2+ breast cancer  
24 highly susceptible to CTLA-4/PD-1 blockade. *Sci Transl Med.* 7, 315ra188.
- 25 37. Yamaguchi T, Teraguchi S, Furusawa C, Machiyama H, Watanabe TM, Fujita H, Sakaguchi S,  
26 Yanagida T (2019). Theoretical modeling reveals that regulatory T cells increase T-cell  
27 interaction with antigen-presenting cells for stable immune tolerance. *Int Immunol.* 31, 743–  
28 53.
- 29 38. Angelova M, Mlecnik B, Vasaturo A, Bindea G, Fredriksen T, Lafontaine L, Buttard B,  
30 Morgand E, Bruni D, Jouret-Mourin A, et al (2018). Evolution of Metastases in Space and  
31 Time under Immune Selection. *Cell.* 175, 751-765.e16.
- 32 39. Pitroda SP, Khodarev NN, Huang L, Uppal A, Wightman SC, Ganai S, Joseph N, Pitt J,

- 1 Brown M, Forde M, et al (2018). Integrated molecular subtyping defines a curable  
2 oligometastatic state in colorectal liver metastasis. *Nat Commun.* 9, 1793.
- 3 40. Wang Y, Lin H, Huang M, Shao Q, Wang Z, Wang F, Yuan Y, Li B, Wang D, Ding P, et al  
4 (2018). The Immunoscore system predicts prognosis after liver metastasectomy in colorectal  
5 cancer liver metastases. *Cancer Immunol Immunother.* 67, 435–44.
- 6 41. Liu X, Wu X, Cao S, Harrington SM, Yin P, Mansfield AS, Dong H (2016). B7-H1 antibodies  
7 lose antitumor activity due to activation of p38 MAPK that leads to apoptosis of tumor-  
8 reactive CD8<sup>+</sup> T cells. *Sci Rep.*, 6:36722.
- 9 42. Höppener DJ, Nierop PMH, Hof J, Sideras K, Zhou G, Visser L, Gouw ASH, de Jong KP,  
10 Sprengers D, Kwekkeboom J, et al (2020). Enrichment of the tumour immune  
11 microenvironment in patients with desmoplastic colorectal liver metastasis. *Br J Cancer.* 123,  
12 196–206.
- 13 43. Tumeh PC, Harview CL, Yearley JH, Shintaku IP, Taylor EJM, Robert L, Chmielowski B,  
14 Spasic M, Henry G, Ciobanu V, et al (2014). PD-1 blockade induces responses by inhibiting  
15 adaptive immune resistance. *Nature* 515, 568–71.
- 16 44. Teng F, Mu D, Meng X, Kong L, Zhu H, Liu S, Zhang J, Yu J (2015). Tumor infiltrating  
17 lymphocytes (TILs) before and after neoadjuvant chemoradiotherapy and its clinical utility for  
18 rectal cancer. *Am J Cancer Res.* 5, 2064–74.
- 19 45. Kumagai S, Togashi Y, Kamada T, Sugiyama E, Nishinakamura H, Takeuchi Y, Vitaly K,  
20 Itahashi K, Maeda Y, Matsui S, et al (2020). The PD-1 expression balance between effector  
21 and regulatory T cells predicts the clinical efficacy of PD-1 blockade therapies. *Nat Immunol.*  
22 21, 1346–58.
- 23 46. Ebert PJR, Cheung J, Yang Y, McNamara E, Hong R, Moskalenko M, Gould SE, Maecker H,  
24 Irving BA, Kim JM, et al (2016). MAP Kinase Inhibition Promotes T Cell and Anti-tumor  
25 Activity in Combination with PD-L1 Checkpoint Blockade. *Immunity* 44, 609–21.
- 26 47. Ledys F, Klopfenstein Q, Truntzer C, Arnould L, Vincent J, Bengrine L, Remark R, Boidot R,  
27 Ladoire S, Ghiringhelli F, et al (2018). RAS status and neoadjuvant chemotherapy impact  
28 CD8<sup>+</sup> cells and tumor HLA class I expression in liver metastatic colorectal cancer. *J*  
29 *Immunother Cancer* 6:123.
- 30 48. Galaine J, Turco C, Vauchy C, Royer B, Mercier-Letondal P, Queiroz L, Loyon R, Mouget V,  
31 Boidot R, Laheurte C, et al (2019). CD4 T cells target colorectal cancer antigens upregulated  
32 by oxaliplatin. *Int J Cancer* 145, 3112–25.

- 1 49. Vincent J, Mignot G, Chalmin F, Ladoire S, Bruchard M, Chevriaux A, Martin F, Apetoh L,  
2 Rébé C, Ghiringhelli F (2010). 5-Fluorouracil selectively kills tumor-associated myeloid-  
3 derived suppressor cells resulting in enhanced T cell-dependent antitumor immunity. *Cancer*  
4 *Res.* 70, 3052–61.
- 5 50. Kanterman J, Sade-Feldman M, Biton M, Ish-Shalom E, Lasry A, Goldshtein A, et al (2014).  
6 Adverse immunoregulatory effects of 5FU and CPT11 chemotherapy on myeloid-derived  
7 suppressor cells and colorectal cancer outcomes. *Cancer Res.* 74, 6022–35.
- 8 51. Limagne E, Euvrard R, Thibaudin M, Rébé C, Derangère V, Chevriaux A, et al (2016).  
9 Accumulation of MDSC and Th17 Cells in Patients with Metastatic Colorectal Cancer  
10 Predicts the Efficacy of a FOLFOX-Bevacizumab Drug Treatment Regimen. *Cancer Res.* 76,  
11 5241–52.
- 12  
13  
14

## 15 SUPPORTING INFORMATION

16 **Table S1.** Raw data

17 **Table S2.** The cohort baseline characteristics

18 **Table S3.** Patients' and metastatic biomarkers characteristics according to neoadjuvant or adjuvant  
19 chemotherapy

20 **Table S4:** Spearman correlation between metastatic immune infiltrate and TRG scoring

21 **Figure S1:** Flow chart of the study

22 **Figure S2:** Heatmaps of biomarkers expression in the primary and metastatic tumor

23 **Figure S3:** Representative images of immunohistochemistry co-staining of CD8 and PD-L1.

24 **Figure S4:** Kaplan-Meier curve showing the association between CD3<sup>high</sup> in the invasive front  
25 and DFS in mCRC patients with synchronous metastases.

26 **Figure S5:** Graphical summary of the results

27  
28



**Table 1. Distribution of biomarkers in the primary tumor and metastatic sites**

	IHC	CD3 IF	CD3 stroma	CD3 IE	CD8 IF	CD8 stroma	PD-1 IF	PD-1 stroma	Foxp3 IF	Foxp3 stroma		PD-L1 tumoral cells	PD-L1 immune cells
<b>Primary Tumors</b>													
All (N = 74)	<b>IHC 3+</b>	11 (14.9%)	15 (20.5%)	14 (19.2%)	0	0	NA	NA	NA	NA			
	<b>IHC 2+ and 3+</b>	33 (44.6%)	42 (57.5)	26 (35.6%)	6 (8.5%)	8 (11.3%)	2 (2.7%)	1 (1.4%)	3 (4.1%)	11 (14.9%)	<b>≥5%</b>	5 (6.8%)	42 (57.5%)
	<b>IHC 0+ and 1+</b>	41 (55.4%)	31 (42.5%)	47 (64.4%)	65 (91.5%)	63 (88.7%)	71 (97.3%)	73 (98.6%)	71 (95.9%)	63 (85.1%)	<b>&lt;5%</b>	68 (93.2%)	31 (42.5%)
Associated with neo-adjuvant metastases (N = 40)	<b>IHC 3+</b>	4 (10.0%)	8 (20.0%)	7 (17.5%)	0	0	NA	NA	NA	NA			
	<b>IHC 2+ and 3+</b>	16 (40.0%)	22 (55.0%)	13 (32.5%)	2 (5.3%)	5 (13.2%)	1 (2.5%)	0	2 (5.0%)	4 (10.0%)	<b>≥5%</b>	2 (5.1%)	22 (56.4%)
	<b>IHC 0+ and 1+</b>	24 (60.0%)	18 (45.0%)	27 (67.5%)	36 (94.7%)	33 (86.8%)	39 (97.5%)	40 (100%)	38 (95.0%)	36 (90.0%)	<b>&lt;5%</b>	37 (94.9%)	17 (43.6%)
Associated with chemo-naive metastases (N = 34)	<b>IHC 3+</b>	7 (20.6%)	7 (21.2%)	7 (21.2%)	0	0	NA	NA	NA	NA			
	<b>IHC 2+ and 3+</b>	17 (50.0%)	20 (60.6%)	13 (39.4%)	4 (12.1%)	3 (9.1%)	1 (3.0%)	1 (2.9%)	1 (2.9%)	7 (20.6%)	<b>≥5%</b>	3 (8.8%)	20 (58.8%)
	<b>IHC 0+ and 1+</b>	17 (50.0%)	13 (39.4%)	20 (60.6%)	29 (87.9%)	30 (90.9%)	32 (97%)	33 (97.1%)	33 (97.1%)	27 (79.4%)	<b>&lt;5%</b>	31 (91.2%)	14 (41.2%)
<b>Metastases</b>													
All (N = 74)	<b>IHC 3+</b>	16 (21.9%)	10 (13.5%)	12 (16.4%)	2 (2.8%)	3 (4.2%)	NA	NA	NA	NA			
	<b>IHC 2+ and 3+</b>	63 (86.3%)	30 (40.5%)	20 (27.4%)	23 (31.9%)	16 (22.2%)	9 (12.7%)	4 (5.6%)	0	4 (5.6%)	<b>≥5%</b>	7 (9.7%)	44 (61.1%)
	<b>IHC 0+ and 1+</b>	10 (13.7%)	44 (59.5%)	53 (72.6%)	49 (68.1%)	56 (77.8%)	62 (87.3%)	67 (94.4%)	72 (100%)	68 (94.4%)	<b>&lt;5%</b>	65 (90.3%)	28 (38.9%)
Neoadjuvant (N = 40)	<b>IHC 3+</b>	11 (28.2%)	7 (17.5%)	5 (12.5%)	1 (2.6%)	2 (5.1%)	NA	NA	NA	NA			
	<b>IHC 2+ and 3+</b>	32 (82.1%)	19 (47.5%)	8 (20.0%)	16 (41.0%)	10 (25.6%)	6 (15.8%)	3 (7.9%)	0	0	<b>≥5%</b>	5 (13.2%)	21 (55.3%)
	<b>IHC 0+ and 1+</b>	7 (17.9%)	21 (52.5%)	32 (80.0%)	23 (59.0%)	29 (74.4%)	32 (84.2%)	35 (92.1%)	39 (100%)	39 (100%)	<b>&lt;5%</b>	33 (86.8%)	17 (44.7%)
Chemo-naive (N = 34)	<b>IHC 3+</b>	5 (14.7%)	3 (8.8%)	7 (21.2%)	1 (3.0%)	1 (3.0%)	NA	NA	NA	NA			
	<b>IHC 2+ and 3+</b>	31 (91.2%)	11 (32.4%)	12 (36.4%)	7 (21.2%)	6 (18.2%)	3 (9.1%)	1 (3.0%)	0	4 (12.1%)	<b>≥5%</b>	2 (5.9%)	23 (67.6%)
	<b>IHC 0+ and 1+</b>	3 (8.8%)	23 (37.6%)	21 (63.6%)	26 (78.8%)	27 (81.8%)	30 (90.9%)	32 (97.0%)	33 (100%)	29 (87.9%)	<b>&lt;5%</b>	32 (94.1%)	11 (32.4%)

Abbreviations: IHC, immunohistochemistry; IF, invasive front

**Table 2. Patients' characteristics according to invasive front CD3, invasive front CD8, and immune cell PD-L1 expressions in chemo-naive patients with metastases**

Parameter	Invasive front CD3 in metastases			Invasive front CD8 and immune cells PD-L1 in metastases				
		CD3 high	CD3 low	<i>P</i>	CD8high PD-L1 high	CD8 low PD-L1 high	CD8 low PD-L1 low	<i>P</i>
<b>Age (years)</b>	<i>N</i>	31	3		7	16	10	
	<b>Median (range)</b>	<b>61 (45-75)</b>	<b>68 (66-74)</b>	0.089	<b>64.2 (45-75)</b>	<b>59.6 (50-71)</b>	<b>64.6 (53-75)</b>	0.332
<b>Longest diameter of metastases (cm), <i>N</i> (%)</b>	<i>N</i>	30	3		7	16	9	
	<=5	29 (96.7)	2 (66.7)	0.176	7 (100.0)	16 (100.0)	7 (77.8)	0.115
	>5	1 (3.3)	1 (33.3)		0 (0.0)	0 (0.0)	2 (22.2)	
<b>N-stage, <i>N</i> (%)</b>	<i>N</i>	30	3		7	16	9	
	N0	17 (56.7)	1 (33.3)	0.579	4 (57.1)	11 (68.8)	3 (33.3)	0.247
	N+	13 (43.3)	2 (66.7)		3 (42.9)	5 (31.3)	6 (66.7)	
<b>Number of metastases, <i>N</i> (%)</b>	<i>N</i>	30	3		7	16	9	
	<b>Mean (SD)</b>	<b>1.4 (0.77)</b>	<b>3.3 (3.21)</b>	0.153	<b>1.1 (0.38)</b>	<b>1.6 (0.73)</b>	<b>2.1 (2.09)</b>	0.408
	<=1	21 (70.0)	1 (33.3)	0.252	6 (85.7)	9 (56.3)	6 (66.7)	0.439
	>1	9 (30.0)	2 (66.7)		1 (14.3)	7 (43.8)	3 (33.3)	
<b>Preoperative CEA level (ng/mL)</b>	<i>N</i>	28	3		7	16	8	
	<b>Mean (SD)</b>	<b>28.0 (64.22)</b>	<b>6.3 (7.33)</b>	0.640	<b>7.5 (5.17)</b>	<b>13.9 (34.10)</b>	<b>66.0 (105.31)</b>	0.052
<b>Timing of metastases, <i>N</i> (%)</b>	<i>N</i>	31	3		7	16	10	
	Metachronous	11 (35.5)	1 (33.3)	1.000	2 (28.6)	7 (43.8)	3 (30.0)	0.715
	Synchronous	20 (64.5)	2 (66.7)		5 (71.4)	9 (56.3)	7 (70.0)	
<b>Sex, <i>N</i> (%)</b>	<i>N</i>	31	3		7	16	10	
	Male	18 (58.1)	3 (100.0)	0.270	6 (85.7)	7 (43.8)	8 (80.0)	0.081
	Female	13 (41.9)	0 (0.0)		1 (14.3)	9 (56.3)	2 (20.0)	
<b>Tumor Sidedness, <i>N</i> (%)</b>	<i>N</i>	30	3		7	16	10	
	Right-Sided	6 (20.0)	1 (33.3)	0.524	1 (14.3)	3 (18.8)	3 (30.0)	0.742
	Left-Sided(with Rectum)	24 (80.0)	2 (66.7)		6 (85.7)	13 (81.3)	7 (70.0)	

**Table 3. Univariate and multivariate analyses for disease-free survival**

Parameter	Univariate analysis		Multivariate analysis	
	HR (95% CI)	P	HR (95% CI)	P
Age (years)	1 (0.970-1.029)	0.951		
Preoperative CEA level	1 (1.000-1.002)	0.130		
Chemotherapy	No	1	1	
	Neoadjuvant	1.99 (1.147-3.453)	<b>0.014</b>	1.682 (0.918-3.082) 0.093
Longest diameter of metastases (cm)	<=5	1		
	>5	1.29 (0.604-2.749)	0.513	
N-stage	N0	1		
	N+	1.38 (0.789-2.413)	0.259	
Number of metastases	<=1	1	1	
	>1	2.12 (1.214-3.688)	<b>0.008</b>	1.816 (0.987-3.344) 0.055
Timing of metastases	Metachronous	1		
	Synchronous	0.93 (0.489-1.778)	0.832	
TRG	2-3	1		
	4-5	1.02 (0.465-2.256)	0.952	
Sex	Male	1		
	Female	1.41 (0.817, 2.451)	0.216	
Tumor Sidedness	Right-Sided	1		
	Left-Sided(with Rectum)	1.71 (0.728, 4.020)	0.218	
CD3+ IF	Low	1	1	
	High	0.36 (0.175-0.755)	<b>0.007</b>	0.309 (0.145-0.657) <b>0.002</b>
CD3+ Stroma	Low	1		
	High	1.19 (0.692-2.032)	0.534	
CD3+ IE	Low	1		
	High	0.71 (0.380-1.333)	0.288	
CD8+ IF	Low	1		
	High	1.21 (0.688-2.133)	0.507	
CD8+ Stroma	Low	1		
	High	1.47 (0.776-2.787)	0.237	
PD-L1 TC	Low	1		
	High	0.80 (0.286-2.213)	0.661	
PD-L1 IC	Low	1		
	High	0.58 (0.335-1.020)	0.059	
CD8+ IF and PD-L1 IC	CD8low PD-L1low	1		
	CD8hi PD-L1hi	0.79 (0.405-1.526)	0.477	
	CD8hi PD-L1low	1.21 (0.278-5.215)	0.803	
	CD8low PD-L1hi	0.48 (0.238-0.981)	<b>0.044</b>	
FOXP3 IF	Staining 0	1		
	Staining 1	0.66 (0.313-1.412)	0.288	
FOXP3 Stroma	Staining 0	1		
	Staining 1	0.59 (0.263-1.301)	0.188	

	Staining 2	0.57 (0.138-2.356)	0.437	
<b>PD1 IF</b>	Low	1		
	High	0.49 (0.192-1.264)	0.141	
<b>PD1 Stroma</b>	Low	1		
	High	0.57 (0.138-2.351)	0.437	

**Abbreviations:** TRG, tumor regression grades; IF, invasive front; IE, intra-epithelial; IC, immune cell; CT, tumor cell

## Figure Legends













**Figure 1: Representative images of immunohistochemistry staining of CD8 and PD-L1.** (A) High CD8 expression in the invasive front of the primary tumor, (B) High PD-L1 expression in the invasive front of the primary tumor of the same patient, (C) High CD8 expression on immune cell in the invasive front of the liver metastasis of the same patient; (D) High PD-L1 expression on immune cell in the invasive front of the liver metastasis of the same patient, (E) High CD8 expression in the invasive front of a liver metastasis of a second patient, (F) High PD-L1 expression in the invasive front of a liver metastasis of the same second patient, (G) High CD8 expression in the invasive front of an ovary metastasis of a third patient, (H) High PD-L1 expression in the invasive front of an ovary metastasis of the same third patient, (I) High CD8 expression in the invasive front of a liver metastasis of a fourth patient, (J) High PD-L1 expression in the invasive front of a liver metastasis of the same fourth patient, (K) High CD8 expression in the invasive front of a liver metastasis of a fifth patient, (L) High PD-L1 expression in the invasive front of a liver metastasis of the same fifth patient.


*Scale bar for 1A, 1B, 1C, 1D = 200  $\mu$ m.*

*Scale bar for 1E to 1L = 100  $\mu$ m.*


**Figure 2: Kaplan-Meier curves showing the association between CD3<sup>high</sup> T cell and PD-L1 markers and DFS in patients with mCRC.** (A) High and low CD3 in the invasive front at the metastatic site (B) High and low PD-L1 expression on immune cells at the metastatic site

*CI, confidence interval*



**A**

High	10	8	7	6	4	1	0
Low	10	2	1	1	0	1	0

**B**

High	11	8	4	3	4	1	0
Low	10	4	2	0	1	0	0