

Portal hypertension and hepatocellular carcinoma: Navigating uncharted waters

Sabrina Sidali, Jean-charles Nault

▶ To cite this version:

Sabrina Sidali, Jean-charles Nault. Portal hypertension and hepatocellular carcinoma: Navigating uncharted waters. United European Gastroenterology Journal, 2022, $10.1002/\mathrm{ueg}2.12193$. hal-03519185

HAL Id: hal-03519185 https://hal.sorbonne-universite.fr/hal-03519185

Submitted on 10 Jan2022

HAL is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers. L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.

Medtronic

Small capsule. Big impact.

Celebrating 20 years of PillCam[™] capsule endoscopy.

PillCam

Join the celebration



EMEA-DG-2100120 © 2021 Medtronic.

EDITORIAL

uegjournal WILEY

Portal hypertension and hepatocellular carcinoma: Navigating uncharted waters

Portal hypertension (PHT) is the main determinant of liver cirrhosis decompensation, defined by the occurrence of ascites, variceal bleeding or hepatic encephalopathy.^{1,2} A portal pressure gradient \geq 10 mmHg, usually estimated by the hepatic venous pressure gradient (HVPG), defines clinically significant or evident PHT (clinically significant portal hypertension or clinically evident portal hypertension (CSPH or CEPH)), since liver decompensation might occur above this threshold.³ However, HVPG is not performed routinely in clinical practice, and the presence of esophageal/gastric varices and low platelet count associated with splenomegaly have been considered as surrogate markers of CEPH.⁴

Hepatocellular carcinoma (HCC) is another complication of cirrhosis and is closely linked to PHT. Hepatocellular carcinoma increases HVPG through the presence of arteriovenous shunting within the tumor and modifications of liver architecture whereas CEPH is predictive of HCC occurrence, independently of the severity of the underlying cirrhosis.^{5,6} The occurrence of acute variceal bleeding and/ or clinical ascites is also associated with poor prognosis in patients with HCC.^{7,8} The impact of PHT in the treatment of patients with HCC was described more than 30 years ago when the Barcelona Clinic Liver Cancer team showed that presence of CEPH was associated with a high risk of postoperative hepatic decompensation and poor long term outcomes after liver resection.⁹ Consequently, current guidelines do not recommend hepatic resection for patients with CEPH.^{10,11} Unfortunately, HCC is frequently diagnosed at intermediate and advanced stages when curative treatment cannot be applied. Trans arterial chemoembolization (TACE) is the treatment of choice in patients with intermediate-stage HCC and may also be considered in selected patients with unresectable and not ablatable early stage HCC.¹¹

For patients undergoing TACE, CEPH status has not been considered in the European Association for the Study of the Liver guidelines.¹¹ However, TACE causes a decrease in hepatic arterial blood flow, accompanied by a transient increase in portal blood flow. The clinical implications of these changes in hepatic hemodynamics are unclear, with limited data on TACE outcomes in HCC patients with CEPH. Nevertheless, two retrospective, Korean studies in 2018 reported CEPH was associated with a decrease in overall survival (OS) after chemoembolization.^{12,13}

In the current issue of the United European Gastroenterology Journal, Müller et al. assess the prevalence and the prognostic impact

of CEPH in a cohort of western patients with HCC that had undergone TACE.¹⁴ CEPH was defined by the presence of at least one of the following criteria: radiological ascites, esophageal/gastric varices, splenomegaly (>12 cm) and a low-platelet count (<100 G/L). A total of 349 patients were included in the study of which 304 (87.1%) had cirrhosis. Among patients with cirrhosis, CEPH was observed in 241 (69.1%) patients. Median OS were 10.6 and 17.1 months in patients with and without CEPH, respectively (p = 0.036). In multivariate analysis, CEPH was not a significant risk factor for death (p = 0.190), and among the CEPH defining criteria, only ascites at imaging was associated with a lower OS (p < 0.003).

Müller et al. thus described some important observations. They reported that CEPH is present in more than two thirds of the patients with HCC that underwent TACE, in line with previous studies.^{12,13} The absence of impact of CEPH on survival in multivariate analysis precludes its application to stratify use of TACE in intermediate HCC, but this finding should be interpreted with caution as OS was significantly impaired in univariate analysis. It could suggest that that the severity of CEPH could be more relevant than its presence by itself, as evidenced by the influence of ascites on outcomes. Furthermore, severe CEPH reflected by an inverse portal flow at ultrasonography is usually an exclusion criterion for TACE and it is unclear how many patients were excluded for this reason.

Overall, this study raises important questions in the field of HCC and PHT. Transjugular intrahepatic portosystemic shunt could be discussed in selected patients with small HCC and CEPH responsible of symptoms, in order to allow locoregional treatment or as a bridge to liver transplantation.¹⁵ The emergence of first-line combined treatment using antiangiogenic (anti-VEGF, bevacizumab) and anti-PD-L1 therapy (atezolizumab) as the gold standard in advanced HCC has led to new controversy on the risk of variceal bleeding due to antiangiogenic treatment. The modality of esophageal varices screening, choice of therapeutics for PHT and timing of introduction of the combined treatment in patients with PHT are still a matter of debate. Finally, more data in preclinical models and in real life are warranted to better decipher the complex interplay between PHT and outcomes after locoregional and systemic treatments in patients with HCC.

ACKNOWLEDGMENT

Jean Charles Nault received research fundings from Bayer and Ipsen.

This is an open access article under the terms of the Creative Commons Attribution License, which permits use, distribution and reproduction in any medium, provided the original work is properly cited.

© 2022 The Authors. United European Gastroenterology Journal published by Wiley Periodicals LLC. on behalf of United European Gastroenterology.

The authors have no conflict of interest to declare.

FUNDING INFORMATION

Bayer and Ipsen

Sabrina Sidali^{1,2}

Jean-Charles Nault^{2,3,4} 🕩

¹Université de Paris, Assistance-Publique Hôpitaux de Paris, Hôpital Beaujon, Service d'Hépatologie, DMU DIGEST, Clichy, France

²Centre de Recherche des Cordeliers, Sorbonne Université, Inserm, Université de Paris, Team Functional Genomics of Solid Tumors, Equipe labellisée Ligue Nationale Contre le Cancer, Labex Oncolmmunology, Paris, France

³Liver Unit, Hôpital Avicenne, Hôpitaux Universitaires Paris-Seine-Saint-Denis, Assistance-Publique Hôpitaux de Paris, Bobigny, France

⁴Unité de Formation et de Recherche Santé Médecine et Biologie Humaine, Université Paris 13, Communauté d'Universités et Etablissements Sorbonne Paris Cité, Paris, France

Correspondence

Jean-Charles Nault, Hôpital Avicenne, Service d'Hépatologie, 125 rue de Stalingrad 93000 Bobigny, France. Email: naultjc@gmail.com

Prevalence and clinical significance of clinically evident portal hypertension in patients with hepatocellular carcinoma undergoing transarterial chemoembolization.

DATA AVAILABILITY STATEMENT

No data, editorial.

ORCID

Jean-Charles Nault 🕩 https://orcid.org/0000-0002-4875-9353

REFERENCES

 Tsochatzis EA, Bosch J, Burroughs AK. Liver cirrhosis. Lancet. 2014;383(9930):1749–61.

- D'Amico G, Garcia-Tsao G, Pagliaro L. Natural history and prognostic indicators of survival in cirrhosis: a systematic review of 118 studies. J Hepatol. 2006;44(1):217-31.
- Ripoll C, Groszmann R, Garcia-Tsao G, Grace N, Burroughs A, Planas R, et al. Hepatic venous pressure gradient predicts clinical decompensation in patients with compensated cirrhosis. Gastroenterology. 2007;133(2):481–8.
- de Franchis R. Expanding consensus in portal hypertension. J Hepatol. 2015;63(3):743–52.
- Ripoll C, Groszmann RJ, Garcia-Tsao G, Bosch J, Grace N, Burroughs A, et al. Hepatic venous pressure gradient predicts development of hepatocellular carcinoma independently of severity of cirrhosis. J Hepatol. 2009;50(5):923–8.
- Allaire M, Rudler M, Thabut D. Portal hypertension and hepatocellular carcinoma: des liaisons dangereuses. Liver Int. 2021;41(8):1734–43.
- Hsu C-Y, Lee Y-H, Huang Y-H, Hsia C-Y, Su C-W, Lin H-C, et al. Ascites in patients with hepatocellular carcinoma: prevalence, associated factors, prognostic impact, and staging strategy. Hepatol Int. 2013;7(1):188–98.
- Lee YR, Park SY, Tak WY. Treatment outcomes and prognostic factors of acute variceal bleeding in patients with hepatocellular carcinoma. Gut Liver. 2020;14(4):500–8.
- Bruix J, Castells A, Bosch J, Feu F, Fuster J, Garcia-Pagan J, et al. Surgical resection of hepatocellular carcinoma in cirrhotic patients: prognostic value of preoperative portal pressure. Gastroenterology. 1996;111(4):1018–22.
- Berzigotti A, Reig M, Abraldes JG, Bosch J, Bruix J. Portal hypertension and the outcome of surgery for hepatocellular carcinoma in compensated cirrhosis: a systematic review and meta-analysis. Hepatology. 2015;61(2):526–36.
- Galle PR, Forner A, Llovet JM, Mazzaferro V, Piscaglia F, Raoul J-L, et al. EASL clinical practice guidelines: management of hepatocellular carcinoma. J Hepatol. 2018;69(1):182–236.
- Choi JW, Chung JW, Lee DH, Kim H-C, Hur S, Lee M, et al. Portal hypertension is associated with poor outcome of transarterial chemoembolization in patients with hepatocellular carcinoma. Eur Radiol. 2018;28(5):2184–93.
- Kim NH, Lee T, Cho YK, Kim BI, Kim HJ. Impact of clinically evident portal hypertension on clinical outcome of patients with hepatocellular carcinoma treated by transarterial chemoembolization. J Gastroenterol Hepatol. 2018;33(7):1397–406.
- Müller L, Hahn F, Mähringer-Kunz A, Stoehr F, Gairing SJ, Foerster F, et al. Prevalence and clinical significance of clinically evident portal hypertension in patients with hepatocellular carcinoma undergoing transarterial chemoembolization. United Eur Gastroenterol J. 2021.
- Padia SA, Chewning RH, Kogut MJ, Ingraham CR, Johnson GE, Bhattacharya R, et al. Outcomes of locoregional tumor therapy for patients with hepatocellular carcinoma and transjugular intrahepatic portosystemic shunts. Cardiovasc Intervent Radiol. 2015;38(4): 913-21.