



HAL
open science

Rapid Bursts of Magnetically Gated Accretion in the Intermediate Polar V1025 Cen

Colin Littlefield, Jean-Pierre Lasota, Jean-Marie Hameury, Scaringi Simone, Peter Garnavich, Paula Szkody, Mark Kennedy, Mckenna Leichty

► **To cite this version:**

Colin Littlefield, Jean-Pierre Lasota, Jean-Marie Hameury, Scaringi Simone, Peter Garnavich, et al.. Rapid Bursts of Magnetically Gated Accretion in the Intermediate Polar V1025 Cen. *The Astrophysical journal letters*, 2022, 10.3847/2041-8213/ac4262 . hal-03519494

HAL Id: hal-03519494

<https://hal.sorbonne-universite.fr/hal-03519494v1>







Submitted on 10 Jan 2022

HAL is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers.

L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.



Rapid Bursts of Magnetically Gated Accretion in the Intermediate Polar V1025 Cen

Colin Littlefield^{1,2,3} , Jean-Pierre Lasota^{4,5} , Jean-Marie Hameury⁶ , Simone Scaringi⁷, Peter Garnavich¹ , Paula Szkody² ,
Mark Kennedy⁸ , and McKenna Leichty¹

¹ Department of Physics, University of Notre Dame, Notre Dame, IN 46556, USA; clittlef@alumni.nd.edu

² Department of Astronomy, University of Washington, Seattle, WA 98195, USA

³ Bay Area Environmental Research Institute, Moffett Field, CA 94035, USA

⁴ Institut d'Astrophysique de Paris, CNRS et Sorbonne Université, UMR 7095, 98bis Bd Arago, F-75014, Paris, France

⁵ Nicolaus Copernicus Astronomical Center, Polish Academy of Sciences, Bartycka 18, 00-716, Warsaw, Poland

⁶ Observatoire Astronomique de Strasbourg, Université de Strasbourg, CNRS UMR 7550, F-67000, Strasbourg, France

⁷ Centre for Extragalactic Astronomy, Department of Physics, University of Durham, South Road, Durham DH1 3LE, UK

⁸ Department of Physics, University College Cork, Cork, Ireland

Received 2021 October 26; revised 2021 December 3; accepted 2021 December 7; published 2022 January 4

Abstract

Magnetically gated accretion has emerged as a proposed mechanism for producing extremely short, repetitive bursts of accretion onto magnetized white dwarfs in intermediate polars (IPs), but this phenomenon has not been detected previously in a confirmed IP. We report the 27 day TESS light curve of V1025 Cen, an IP that shows a remarkable series of 12 bursts of accretion, each lasting for less than 6 hours. The extreme brevity of the bursts and their short recurrence times ($\sim 1\text{--}3$ days) are incompatible with the dwarf-nova instability, but they are natural consequences of the magnetic gating mechanism developed by Spruit and Taam to explain the Type II bursts of the accreting neutron star known as the Rapid Burster. In this model, the accretion flow piles up at the magnetospheric boundary and presses inward until it couples with the star's magnetic field, producing an abrupt burst of accretion. After each burst, the reservoir of matter at the edge of the magnetosphere is replenished, leading to cyclical bursts of accretion. A pair of recent studies applied this instability to the suspected IPs MV Lyr and TW Pic, but the magnetic nature of these two systems has not been independently confirmed. In contrast, previous studies have unambiguously established the white dwarf in V1025 Cen to be significantly magnetized. The detection of magnetically gated bursts in a confirmed IP therefore validates the extension of the Spruit and Taam instability to magnetized white dwarfs.

Unified Astronomy Thesaurus concepts: Cataclysmic variable stars (203); DQ Herculis stars (407); White dwarf stars (1799); Stellar magnetic fields (1610); Stellar accretion disks (1579)

1. Introduction

Accreting, nonmagnetic white dwarfs (WDs) in cataclysmic variable stars (CVs) occasionally undergo dwarf-nova outbursts, which are brief episodes of greatly enhanced accretion. The mechanism that produces these outbursts, the dwarf-nova instability (DNI), is the runaway ionization of the accretion disk when it exceeds a critical effective temperature (for reviews, see Osaki 1996; Lasota 2001; Hameury 2020). As it ionizes, the disk's temperature and viscosity skyrocket, resulting in a temporary increase in the accretion rate onto the WD. A typical outburst will result in a multimagnitude brightening in optical photometry for several days.

Despite the many successes of the DNI model, it has struggled to explain the properties of outbursts observed in intermediate polars (IPs), the subset of CVs that contains WDs whose magnetic-field strengths are high enough to disrupt the accretion flow without synchronizing the WD's spin to the binary orbital period (for a review, see Patterson 1994). Although some IPs, such as CC Scl, exhibit bona fide dwarf-nova outbursts, outbursts in other IPs can last for less than one day, with long, irregular recurrence intervals (Table 1 in Hameury & Lasota 2017a); their amplitudes also tend to be lower, typically less than about $\sim 1\text{--}2$ mag. Hameury & Lasota (2017a) concluded that the DNI cannot explain the very short (\sim hours-long) outbursts observed in some

IPs, implying that a different mechanism is responsible for very short IP outbursts.

Recently, Scaringi et al. (2017, 2021) identified a repetitive series of rapid, low-amplitude bursts of accretion in the nominally nonmagnetic systems MV Lyr and TW Pic, respectively. Both studies attributed these bursts to magnetically gated accretion via the Spruit & Taam (1993) instability, which occurs in systems whose inner disk radius, defined as the magnetospheric radius, is close to the corotation radius. In such a situation, the rapidly rotating magnetosphere creates a centrifugal barrier that suppresses accretion and causes the accretion flow to pile up just outside the magnetospheric boundary. Eventually, this material compresses the magnetosphere until it is able to couple onto magnetic field lines and accrete. Once the reservoir of matter outside the magnetosphere is depleted, the cycle repeats itself, giving rise to episodic spurts of accretion. Scaringi et al. (2017, 2021) determined that this process would occur at a critical mass-transfer rate determined by the WD's magnetic-field strength and its rotational frequency. However, neither of these two systems was previously known to possess a magnetized WD, and the bursts themselves provided the main evidence for the WD's magnetism.

1.1. V1025 Cen

V1025 Cen was identified as an IP by Buckley et al. (1998) with optical photometry and spectroscopy; they proposed an unusually long spin period ($P_{\text{spin}} = 36$ minutes) relative to the binary orbital



Original content from this work may be used under the terms of the [Creative Commons Attribution 4.0 licence](https://creativecommons.org/licenses/by/4.0/). Any further distribution of this work must maintain attribution to the author(s) and the title of the work, journal citation and DOI.




Figure 1. The combined CRTS, ASAS-SN, and ATLAS light curve of V1025 Cen. The timespan of the TESS observations is indicated in red. V1025 Cen experienced a low-accretion state between 2015 and 2018, and we believe that the unblended ATLAS data offer a better measurement of its depth compared to the blended ASAS-SN data. A single outburst, with a minimum amplitude of 0.6 mag, was detected by ASAS-SN during the TESS observations. There were several additional bursts in the months before and after the TESS observation, and the CRTS data show a single burst of unknown duration in 2009.

period ($P_{\text{orb}} = 85$ minutes). Hellier et al. (1998) confirmed this classification and the periods with X-ray observations. More recently, Hellier et al. (2002) presented phase-resolved optical spectroscopy in an effort to ascertain whether V1025 Cen accretes from a truncated accretion disk or is instead diskless, but the observations did not clearly favor either scenario.

Bailer-Jones et al. (2021) estimate a geometric distance of 196 ± 3 pc based on their analysis of Gaia EDR3 (Gaia Collaboration et al. 2016, 2021).

2. Data and Analysis

TESS observed V1025 Cen at the standard 30 minute full-frame cadence between 2019 March 27 and 2019 April 22. Survey photometry from the Catalina Real-Time Sky Survey (CRTS; Drake et al. 2009), the All-Sky Automated Survey for Supernovae (ASAS-SN; Shappee et al. 2014; Kochanek et al. 2017), and the Asteroid Terrestrial-impact Last Alert System (ATLAS; Tonry et al. 2018a; Smith et al. 2020) in Figure 1 establishes that the TESS observation occurred during the final recovery from a years-long low state independently identified by A. Covington et al. (2021, in preparation). The ASAS-SN g -band and ATLAS c -band observations disagree as to the depth of this low state, which is a likely consequence of blending in the ASAS-SN data, which uses a photometric aperture radius of 16 arcsec. The ATLAS All-Sky Stellar Reference Catalog (Tonry et al. 2018b) estimates that within a radius of 15.4 arcsec of V1025 Cen, the contaminating G -band flux from other sources is equivalent to the G -band flux of V1025 Cen at $G = 17.1$; if this modeling is reasonably accurate for blending in the g band, it could easily account for the comparatively shallow depth in the ASAS-SN g data.

We used TESSCut (Basseur et al. 2019) and lightkurve (Lightkurve Collaboration et al. 2018) to retrieve the full-frame TESS observations of V1025 Cen and to extract its background-subtracted light curve with a custom photometric aperture. As a safeguard, we also checked light curves of nearby field stars and the background to ensure that no obvious systematic problems afflicted the data. The most remarkable feature of the resulting light curve (Figure 2) is a series of 12 brief bursts, each lasting no longer than ~ 6 h. The burst

durations are all comparable, and by fitting a Gaussian to each burst, we find that their average full width at half-maximum was 3.0 ± 1.0 hr (Figure 3). The full duration of each burst was $\lesssim 6$ hr, but the time between consecutive bursts was irregular, ranging from 1.1 days to 4.6 days.

Figure 2 also includes the simultaneous ASAS-SN observations. Due to sampling limitations, only one of the bursts was detected by ASAS-SN. Curiously, the agreement between the ASAS-SN and TESS light curves degrades near the end of the observation. In some instances, the ASAS-SN observations show evidence of rapid variation on minutes-long timescales; for example, in a trio of exposures obtained within a 5 minute span near $\text{BTJD} = 1594$,⁹ V1025 Cen’s initial flux density was 3.25 ± 0.25 Jy, but the next two exposures showed it to be 1.99 ± 0.23 Jy and 2.40 ± 0.31 Jy. A separate group of observations, obtained hours earlier, also showed variability in excess of the uncertainties. The rapid variations in the ASAS-SN data do not correlate with the full width at half-maximum of the stellar point-spread functions, which suggests that they are not attributable to time-varying contamination from other sources. Furthermore, when the ASAS-SN data show this increased internal variability, the TESS light curve becomes visibly jagged. We speculate, therefore, that during portions of the TESS observation, V1025 Cen showed rapid variability that could not be resolved at the 30 minute cadence of the full-frame data.

The bursts in the TESS light curve have no precedent in the observational literature for V1025 Cen. Buckley et al. (1998) is the only published study that includes photometry of V1025 Cen, and it reported a snapshot observation at $V \sim 16.1$, along with time-series light curves showing variability between $B \sim 15.5 - 16$ for several nights in 1995 April. There is no evidence of bursts in any of their data.

Unfortunately, the WD’s spin frequency ($\omega = 40.25$ cycles d^{-1}) is above the Nyquist frequency of the TESS data (24 cycles d^{-1}), and the resulting aliasing precludes us from using the power spectrum to ascertain whether the WD was accreting from an

⁹ BTJD is defined as $\text{BJD} - 2457000$.




Figure 2. TESS light curve of V1025 Cen, with simultaneous ASAS-SN observations overlaid on the secondary y-axis. Each TESS datum is a 30 minute integration, while the ASAS-SN observations are 90 s exposures.




Figure 3. An enlargement of each outburst (blue markers) from Figure 3, superimposed upon the other 11 outbursts (gray lines). The approximate midpoint of each outburst is shown in each panel. Each burst lasted for fewer than six hours, with a mean full width at half-maximum of 3.0 ± 1.0 hr.

accretion disk or directly from the ballistic accretion stream (Ferrario & Wickramasinghe 1999). Figure 4 shows the power spectrum of the light curve during the intervals between the bursts, and while both Ω and $\omega - \Omega$ are present, there is also a Nyquist-aliased signal at $7.75 \text{ cycles d}^{-1}$, equivalent to $f_s - \omega$, where f_s is the sampling frequency of 48 cycles d^{-1} . Even more striking is a cluster of signals near 5 cycles d^{-1} , corresponding to a quasiperiodic signal that can be seen directly in the light curve in Figure 2, particularly after $\text{BTJD} \sim 1590$. We were unable to attribute these signals to Nyquist aliasing of ω , $2(\omega - \Omega)$, or other super-Nyquist frequencies; moreover, Nyquist aliasing of a periodic signal should not convert it into a cluster of aliases. It is conceivable, however, that a quasiperiodic oscillation above the Nyquist frequency could

generate a cluster of low-frequency signals with no discernible relation to V1025 Cen’s periodic variations.

3. Discussion

3.1. Magnetic Gating in V1025 Cen

Hameury & Lasota (2017a) found that for realistic disk viscosities, the DN instability cannot produce the hours-long bursts observed in some IPs. They instead suggested that these bursts might be due to some instability coupling the white dwarf magnetic field with that generated by the magnetorotational instability operating in the accretion disk. In particular, they mention the Spruit & Taam (1993) instability, which was




Figure 4. Power spectrum of V1025 Cen during the intervals between the bursts. f_s refers to the sampling frequency of 48 cycles d^{-1} , while ω is the WD spin frequency and Ω is the binary orbital frequency. Because ω (40.25 cycles d^{-1}) is above the Nyquist frequency of the data (24 cycles d^{-1}), it produces an alias at a frequency of $f_s - \omega = 7.75$ cycles d^{-1} . The power near 5 cycles d^{-1} is of unknown origin.

suggested by Mhlahlo et al. (2007) to explain EX Hya’s outbursts, but as Mhlahlo et al. (2007) pointed out, this explanation struggled to account for the long recurrence times between that system’s outbursts.

This difficulty is not present with the highly repetitive bursts of V1025 Cen, which we can readily explain using the Spruit & Taam (1993) instability. D’Angelo & Spruit (2010, 2011, 2012) applied a model based on the Spruit & Taam (1993) mechanism to outbursts in millisecond pulsars and young stellar objects (YSOs), and the light curve of V1025 Cen shows strong similarities to the synthetic light curves of D’Angelo & Spruit (2010, 2011, 2012) as well as the Type II bursts of the accreting neutron star known as the Rapid Burster. These similarities make the Spruit & Taam (1993) instability an excellent candidate to be the cause of the bursts in V1025 Cen, a hypothesis that we examine quantitatively below.

The applicability of the magnetic gating model to V1025 Cen is strongly supported by the parameters of this system. Two important radii are the corotation radius, defined as

$$R_{\text{cor}} = 3.52 \times 10^{10} M_1^{1/3} P_{\text{spin}(h)}^{2/3} \text{ cm}, \quad (1)$$

and the magnetospheric radius, given by

$$R_M = 1.4 \times 10^{10} \dot{M}_{15}^{-2/7} M_1^{-1/7} \mu_{32}^{4/7} \text{ cm}. \quad (2)$$

In these equations, M_1 is the mass of the WD in solar masses, \dot{M} is the mass-transfer rate in units of 10^{15} g s^{-1} , and μ is the magnetic moment in units of 10^{32} G cm^3 . For V1025 Cen, $R_{\text{cor}} = 2.5 \times 10^{10} M_1^{1/3} \text{ cm}$, and to have $R_{\text{cor}} \approx R_M$, the magnetic moment should be

$$\mu \approx 2.9 \times 10^{32} M_{15}^{1/2} M_1^{5/6} P_{\text{spin}(36)}^{7/6} \text{ G cm}^3, \quad (3)$$

where $P_{\text{spin}(36)}$ is the spin period expressed as a fraction of the observed 36 minute spin period of V1025 Cen. V1025 Cen satisfies the necessary condition for the instability ($R_{\text{cor}} \approx R_M$) if its accretion rate is close to the expected secular value ($\dot{M}_c \sim 10^{15} \text{ g s}^{-1}$; Knigge et al. 2011) and if the WD’s magnetic moment is typical of the assumed magnetic moments in IPs.

The fact that outbursts appear when the system is at its brightest in Figure 1 suggests that in the preceding epochs, the accretion rate was less than the critical value above which the disk would become thermally unstable and subject to DNI. Alternatively, the lack of outbursts during the low state might suggest that the accretion disk dissipated (Hameury & Lasota 2017b).

Although Spruit & Taam (1993) and D’Angelo & Spruit (2010, 2011, 2012) specifically considered a truncated accretion disk, their modeling is not predicated upon a Keplerian accretion flow. Instead, what is crucial for the mechanism to work is a ring of matter accumulated near the magnetospheric/corotation radius. A Keplerian disk would fulfill this requirement, but even if V1025 Cen were diskless during the TESS observation, modeling by King & Wynn (1999) and Norton et al. (2004) shows that diskless accretion flows can take the form of a torus that surrounds the magnetosphere.

It is unclear whether a disk exists in V1025 Cen. The circularization radius of the binary is (using Gilfanov & Arefiev 2005)

$$R_{\text{circ}} = 2.6 \times 10^9 M_1^{1/3} q^{-0.48} (1 + q)^{0.57} P_{\text{orb}(h)}^{2/3} \text{ cm}, \quad (4)$$

which for V1025 Cen is $R_{\text{circ}} = 1.0 \times 10^{10} \text{ cm}$, assuming $q = 0.1$. Therefore, $R_{\text{circ}} \lesssim R_{\text{cor}}$, and a Keplerian disk is unlikely unless q is very small. Moreover, observations by Hellier et al. (2002) were inconclusive on the existence of a disk, and in any event, they would not resolve the issue of whether one was present nearly two decades later during the TESS observation. On one hand, the unambiguous existence of a Keplerian disk would simplify the application of the Spruit & Taam (1993) mechanism to V1025 Cen because it would establish the presence of the ring of matter required by that instability. On the other hand, the arguments presented by Spruit & Taam (1993) and D’Angelo & Spruit (2010, 2011, 2012) appear to be generalizable to the torus-like accretion flows expected to form when $R_{\text{cor}} \sim R_M$, a regime in which the kick provided by magnetic forces is insufficient to expel matter to large distances. However, it will be important for theoretical studies to test our proposed generalization of the Spruit & Taam (1993) instability.

As for the outburst timescales, they depend strongly not only on \dot{M}_c , but also on the two transition widths $\Delta R/R_M$ (for the torque) and $\Delta R_2/R_M$ (for the mass flux; D’Angelo & Spruit 2011), so it is not possible to use the simulations for neutron stars or young stellar objects (YSOs) to scale them to the case of IPs. For example, the recurrence time in the simulations vary from 1 to 1000 times the viscous time at the magnetospheric radius.

Figures 1 and 2 establish that the WD continues to accrete between bursts; if the accretion rate were negligible, V1025 Cen would have faded to at least the brightness level observed during its low state. D’Angelo & Spruit (2012), in a regime that they referred to as “RII,” found that the Spruit & Taam (1993) mechanism can allow for reduced—but uninterrupted—accretion when the transition region between accreting and nonaccreting states is large. Bursts in this regime have comparatively smooth profiles and short recurrence timescales compared to those in the D’Angelo & Spruit (2012) RI regime, whose bursts have distinctive initial spikes and long intervals of negligible accretion. Moreover, the RII instability from D’Angelo & Spruit (2012) occurs only for a limited range of \dot{M} , which agrees well with the

apparent sensitivity of the bursts to the accretion rate (see Section 3.3). However, the generalization of the instability to a torus-like geometry is an important step that still requires theoretical confirmation.

In the model, the outburst amplitudes are independent of the parameters and roughly a factor two of \dot{M}_c , which seems to be compatible with observations. However, it is unclear whether this independence of parameters is a generic property of the model.

3.2. Magnetic Gating in Other IPs

Two studies have applied the Spruit & Taam (1993) model to suspected magnetic systems. Scaringi et al. (2017) found that MV Lyr shows recurring bursts of accretion every ~ 2 hr, each lasting for ~ 30 minutes, at its lowest accretion rates. MV Lyr would have to possess a substantially lower magnetic-field strength (~ 0.02 – 0.13 MG; Scaringi et al. 2017) than what is normally inferred for IPs (~ 1 – 10 MG; Patterson 1994), including V1025 Cen. The short duration (~ 30 min) and recurrence time (~ 2 hr) in MV Lyr compared to V1025 Cen would be accounted for by the weakness of the magnetic field, leading to a small R_M and hence a small viscous time at the inner edge of the disk.

Likewise, with TW Pic, magnetically gated bursts occur at diminished accretion rates, typically last for ~ 30 minutes, and recur every 1.2–2.4 hr (Scaringi et al. 2021). As with MV Lyr, the short durations and recurrence intervals of TW Pic’s bursts are consistent with it possessing a lower field strength than V1025 Cen. Scaringi et al. (2021) constrained the field strength to be $\lesssim 1$ MG but found that it is degenerate with the unknown spin period of the WD.

Because V1025 Cen has already been independently confirmed to be an IP (Buckley et al. 1998; Hellier et al. 1998), the detection of magnetic gating in its TESS light curve provides important validation of the magnetic-gating interpretation by Scaringi et al. (2017, 2021) and its extension to magnetized WDs in general. Additionally, because accreting WDs are more common than accreting neutron stars, the presence of magnetic gating in at least some of these systems offers a convenient means of amassing observations against which theoretical predictions can be tested.

3.3. Difficulty of Detecting the Bursts in Survey Photometry

Ground-based photometric surveys such as ASAS-SN have proven adept at detecting the presence of days- or weeks-long outbursts in nonmagnetic systems. In contrast, the comparison of the TESS and ASAS-SN light curves in Figure 2 shows how easily V1025 Cen’s rapid bursts can elude the relatively frequent ASAS-SN observations. Even though there were a dozen bursts in just four weeks in the TESS light curve, they are inconspicuous in the ASAS-SN photometry because of the combination of the survey’s sampling and the bursts’ low amplitudes and short durations. Only one of the 12 bursts in the TESS light curve was also observed by ASAS-SN, and that particular burst is an inconspicuous feature in the long-term ASAS-SN light curve in Figure 1. Were it not for the simultaneous TESS data, it could easily be dismissed as strong flickering or even a data-quality artifact.

Such a low detection efficiency has several important implications. For example, the ASAS-SN light curve shows at least four additional short-lived, low-amplitude brightenings in the months surrounding the TESS observation during the

2018–2019 observing season for V1025 Cen. These features resemble the lone TESS burst detected by ASAS-SN. If we assume that these are the same phenomenon observed by TESS, the presence of several bursts in the ASAS-SN light curve implies that many more eluded detection during that timespan. Therefore, we infer that the bursts in the TESS light curve were a common feature in the months surrounding the TESS observation, when V1025 Cen was finishing its recovery from a years-long low state.

A simple statistical analysis of the ASAS-SN data bolsters this inference. When it observes a target, ASAS-SN usually obtains multiple photometric measurements over the course of a few minutes, so we created a new light curve in which each unique visit was represented by a single point whose abscissa was the median time of observation and whose ordinate was the median magnitude during that visit. Using this binned light curve, we analyzed each observing season separately by detrending that season’s light curve and counting the number of measurements that were $>3\sigma$ brighter than the mean magnitude. The 2018–2019 observing season contained 4 such measurements; for comparison, a Gaussian distribution with the observed $N = 241$ would be expected to show just 0.3 measurements more than 3σ below the mean. Moreover, of the nine observing seasons covered by ASAS-SN, only 2 others showed even a single bright, $>3\sigma$ outlier. It therefore appears likely that the bursts occurred preferentially at the end of the low state in 2018 and 2019. This, in turn, provides observational evidence that magnetic gating in V1025 Cen is very sensitive to the accretion rate.

4. Conclusion

The TESS light curve of V1025 Cen contains a dozen bursts of accretion that are notable for their extremely short durations ($\lesssim 6$ hr) and rapid recurrence timescale (~ 1 – 3 days). We show that the instability proposed by Spruit & Taam (1993) and expanded in D’Angelo & Spruit (2010, 2011, 2012) can explain these bursts as episodes of magnetically gated accretion. Although Scaringi et al. (2017, 2021) have identified magnetic gating during low-accretion states in MV Lyr and TW Pic, respectively, neither of these systems is a confirmed IP. Conversely, the WD in V1025 Cen was already known to be magnetic, which bolsters the application of the magnetic gating model to accreting WDs and beyond its original context of accreting neutron stars and YSOs.

The duty cycle and amplitudes of the bursts in V1025 Cen are sufficiently low that they are difficult to recognize in survey photometry, despite reasonably good coverage. It is possible that other IPs show similar bursts, and if they do, sustained time-series photometry will be necessary to detect them unambiguously, as the TESS light curve of V1025 Cen vividly demonstrates.

We thank the anonymous referee for a helpful report that improved this manuscript. P.S. and C.L. acknowledge support from NSF grant AST-1514737.

J.P.L.’s research was supported by a grant from the French Space Agency CNES.

M.R.K. acknowledges funding from the Irish Research Council in the form of a Government of Ireland Postdoctoral Fellowship (GOIPD/2021/670: Invisible Monsters).

Software: `astropy` (Astropy Collaboration et al. 2013), `lightkurve` (Lightkurve Collaboration et al. 2018), `TESS-cut` (Brasseur et al. 2019).

ORCID iDs

Colin Littlefield  <https://orcid.org/0000-0001-7746-5795>
 Jean-Pierre Lasota  <https://orcid.org/0000-0002-6171-8396>
 Jean-Marie Hameury  <https://orcid.org/0000-0002-6412-0103>
 Peter Garnavich  <https://orcid.org/0000-0003-4069-2817>
 Paula Szkody  <https://orcid.org/0000-0003-4373-7777>
 Mark Kennedy  <https://orcid.org/0000-0001-6894-6044>

References

- Astropy Collaboration, Robitaille, T. P., Tollerud, E. J., et al. 2013, *A&A*, **558**, A33
- Bailer-Jones, C. A. L., Rybizki, J., Fouesneau, M., Demleitner, M., & Andrae, R. 2021, *AJ*, **161**, 147
- Brasseur, C. E., Phillip, C., Fleming, S. W., Mullally, S. E., & White, R. L. 2019, *Astrocute: Tools for creating cutouts of TESS images*, *Astrophysics Source Code Library*, ascl:1905.007
- Buckley, D. A. H., Cropper, M., Ramsay, G., & Wickramasinghe, D. T. 1998, *MNRAS*, **299**, 83
- D'Angelo, C. R., & Spruit, H. C. 2010, *MNRAS*, **406**, 1208
- D'Angelo, C. R., & Spruit, H. C. 2011, *MNRAS*, **416**, 893
- D'Angelo, C. R., & Spruit, H. C. 2012, *MNRAS*, **420**, 416
- Drake, A. J., Djorgovski, S. G., Mahabal, A., et al. 2009, *ApJ*, **696**, 870
- Ferrario, L., & Wickramasinghe, D. T. 1999, *MNRAS*, **309**, 517
- Gaia Collaboration, Prusti, T., de Bruijne, J. H. J., et al. 2016, *A&A*, **595**, A1
- Gaia Collaboration, Brown, A. G. A., Vallenari, A., et al. 2021, *A&A*, **649**, A1
- Gilfanov, M., & Arefiev, V. 2005, arXiv:astro-ph/0501215
- Hameury, J. M. 2020, *AdSpR*, **66**, 1004
- Hameury, J. M., & Lasota, J. P. 2017a, *A&A*, **602**, A102
- Hameury, J. M., & Lasota, J. P. 2017b, *A&A*, **606**, A7
- Hellier, C., Beardmore, A. P., & Buckley, D. A. H. 1998, *MNRAS*, **299**, 851
- Hellier, C., Wynn, G. A., & Buckley, D. A. H. 2002, *MNRAS*, **333**, 84
- King, A. R., & Wynn, G. A. 1999, *MNRAS*, **310**, 203
- Knigge, C., Baraffe, I., & Patterson, J. 2011, *ApJS*, **194**, 28
- Kochanek, C. S., Shappee, B. J., Stanek, K. Z., et al. 2017, *PASP*, **129**, 104502
- Lasota, J.-P. 2001, *NewAR*, **45**, 449
- Lightkurve Collaboration, Cardoso, J. V. d. M., Hedges, C., et al. 2018, *Lightkurve: Kepler and TESS time series analysis in Python*, *Astrophysics Source Code Library*, ascl:1812.013
- Mhlahlo, N., Buckley, D. A. H., Dhillon, V. S., et al. 2007, *MNRAS*, **380**, 353
- Norton, A. J., Wynn, G. A., & Somerscales, R. V. 2004, *ApJ*, **614**, 349
- Osaki, Y. 1996, *PASP*, **108**, 39
- Patterson, J. 1994, *PASP*, **106**, 209
- Scaringi, S., Maccarone, T. J., D'Angelo, C., Knigge, C., & Groot, P. J. 2017, *Natur*, **552**, 210
- Scaringi, S., de Martino, D., Buckley, D. A. H., et al. 2021, *NatAs*, in press
- Shappee, B. J., Prieto, J. L., Grupe, D., et al. 2014, *ApJ*, **788**, 48
- Smith, K. W., Smartt, S. J., Young, D. R., et al. 2020, *PASP*, **132**, 085002
- Spruit, H. C., & Taam, R. E. 1993, *ApJ*, **402**, 593
- Tonry, J. L., Denneau, L., Heinze, A. N., et al. 2018a, *PASP*, **130**, 064505
- Tonry, J. L., Denneau, L., Flewelling, H., et al. 2018b, *ApJ*, **867**, 105