



**HAL**  
open science

# The spontaneous mutation rate of *Drosophila pseudoobscura*

Marc Krasovec

► **To cite this version:**

Marc Krasovec. The spontaneous mutation rate of *Drosophila pseudoobscura*. G3, 2022, 11 (7), 10.1093/g3journal/jkab151 . hal-03525878

**HAL Id: hal-03525878**

**<https://hal.sorbonne-universite.fr/hal-03525878>**

Submitted on 14 Jan 2022

**HAL** is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers.

L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.

# The spontaneous mutation rate of *Drosophila pseudoobscura*

Marc Krasovec \*

CNRS, Biologie Intégrative des Organismes Marins (BIOM), Observatoire Océanologique, Banyuls-sur-Mer 66650, France

\*Corresponding author: Email: krasovec@obs-banyuls.fr

## Abstract

The spontaneous mutation rate is a very variable trait that is subject to drift, selection and is sometimes highly plastic. Consequently, its variation between close species, or even between populations from the same species, can be very large. Here, I estimated the spontaneous mutation rate of *Drosophila pseudoobscura* and *Drosophila persimilis* crosses to explore the mutation rate variation within the *Drosophila* genus. All mutation rate estimations in *Drosophila* varied fourfold, probably explained by the sensitivity of the mutation rate to environmental and experimental conditions. Moreover, I found a very high mutation rate in the hybrid cross between *D. pseudoobscura* and *D. persimilis*, in agreement with known elevated mutation rate in hybrids. This mutation rate increase can be explained by heterozygosity and fitness decrease effects in hybrids.

**Keywords:** mutation rate; spontaneous mutation; *Drosophila pseudoobscura*; hybrids

## Introduction

The spontaneous mutation rate, denoted  $\mu$ , currently has been estimated in more than 60 species [Figure 1 of Krasovec et al. (2019)] from bacteria and eukaryotes (Lynch et al. 2016; Katju and Bergthorsson 2018), and recently in one archaeal species (Kucukyildirim et al. 2020). These studies pointed out a substantial variation in the mutation rate at all scales. One of the most interesting findings is the large variation of  $\mu$  between close species, or between strains or populations from a same species. One of the most cited studies is the case of the green alga *Chlamydomonas reinhardtii* where  $\mu$  varies sevenfold between several strains (Ness et al. 2015). More recently, a study based on a mutation accumulation experiment of three populations of *Daphnia magna* reported an even greater variation in  $\mu$ , from  $3.6 \times 10^{-9}$  to  $3.4 \times 10^{-8}$  (Ho et al. 2020). A putative explanation of this variation is the effect of the environment on the mutation rate, which has been reported in yeast (Liu and Zhang 2019), *Escherichia coli* (Chu et al. 2018), and *Arabidopsis thaliana* (Jiang et al. 2014). For instance, in *A. thaliana*, stress seems to be at the origin of mutation rate increase, which was twice as high in stressful than in benign conditions. It so appears that the mutation rate is a plastic trait and its estimation can vary between experimental conditions. This raises questions about the repeatability of mutation accumulation experiments in the same species, addressed in a previous study (Behringer and Hall 2016) showing that  $\mu$  estimated from independent experiments are significantly different in *Drosophila melanogaster*, *Caenorhabditis elegans*, and *Schizosaccharomyces pombe*. In *D. melanogaster*, for example, a significant approximately twofold variation of mutation rate (Welch's t-test, P-value < 0.05) has been reported by Behringer

and Hall (2016) using previous data (Schridder et al. 2013; Keightley et al. 2014).

If environment or culture conditions have a high impact on the mutation rate, then we can reasonably expect that the mutation rate estimated under laboratory conditions may be far from the real mutation rate in the wild. This is also relevant for closely related species or strains, which should have different mutation rates as a function of their ecological and environmental ranges even with a similar genomic background. On the other hand, there are several reported cases where the mutation rate estimated from close species are similar such as in the *Caenorhabditis* genus (Denver et al. 2012); or much lower than observed in *C. reinhardtii* (Ness et al. 2015) and *D. magna* (Ho et al. 2020), such as in *Mycobacterium* (Ford et al. 2011; Kucukyildirim et al. 2016) or *Vibrio* (Dillon et al. 2016; Strauss et al. 2017) genera. In this study, I took advantage of available parents-progenies genomic data from a previous study to measure the mutation rate of *Drosophila pseudoobscura* and *Drosophila persimilis* crosses (Korunes and Noor 2019) and explore the mutation rate variation within *Drosophila* genus.

## Materials and methods

Genomic data analyzed were from a previous study on *Drosophila* crosses (Korunes and Noor 2019). Briefly, Korunes and Noor did three crosses to F2 generations with three genotypes of *D. pseudoobscura* and one genotype of *D. persimilis* [see Figure 2 of Korunes and Noor (2019)]. A grandparent female *D. pseudoobscura* PP1137 (isolated in New Mexico, USA) was crossed with three males *D. pseudoobscura* PP1134 (isolated in New Mexico, USA),

Received: March 10, 2021. Accepted: April 26, 2021

© The Author(s) 2021. Published by Oxford University Press on behalf of Genetics Society of America.

This is an Open Access article distributed under the terms of the Creative Commons Attribution License (<http://creativecommons.org/licenses/by/4.0/>), which permits unrestricted reuse, distribution, and reproduction in any medium, provided the original work is properly cited.

*D. pseudoobscura* MSH177 (isolated in California, USA), and *D. persimilis* MSH1993 (isolated in California, USA), which gave the three F1 parent generations. Then, a female from this F1 generation was crossed with a *D. pseudoobscura* PP1137 male, which gave the three F2 progeny generations. All genomes (2 grandparents, 2 F1 parents, and 10 F2 progenies for each cross) were sequenced by Illumina HiSeq 100-bp paired-end with 20× coverage (SRR numbers provided in Supplementary Table S1). A summary of the crosses is provided in Supplementary Figure S1.

The mutation identification was done following the highly stringent method developed previously (Schridder et al. 2013; Keightley et al. 2014) that gave almost 100% of true positives. Raw reads were aligned against the *D. pseudoobscura* PP1137 reference genome with bwa mem (Li and Durbin 2010), then the resulting bam files were treated with samtools (Li et al. 2009), and SNP calling was done with HaplotypeCaller from GATK (McKenna et al. 2010). Mutation calling was done separately for each cross. A candidate mutation was considered true only if it fulfilled the following conditions: the site was covered by at least 10 reads in all individuals (the 2 grandparents, the 2 F1 parents, and the 10 F2); all sites with at least one individual with 50 coverage or more were discarded to limit false positives due to repetitive elements; the mutation is identified in only one F2 individual; the site was strictly homozygous in all individuals without any alternative allele, even if support by only one read; the alternative allele had a coverage of five or more and was supported by at least 20% of total site coverage; all candidate sites were verified by visual inspection of the mpileup files (alignment text files generated by samtools) and with IGV (reads with more than 2% mismatches were considered as miss mapping) (Robinson et al. 2011); and all mutation candidates inside an indel were removed with bcftools, option -snpgap 1 (Li 2011). Last, final mutation candidates were checked in other crosses, even if all individuals did not have coverage of 10 or more. The mutation rate was calculated counting one generation for the F2 because parents and grandparents were used as ancestral lines:  $\mu = \frac{N}{G*2+n*g}$  where N is the number of mutations, G\* the number of callable sites (G\*2 because the genome is diploid and  $\mu$  is calculated per site per haploid genome), n the number of F2 progenies and g the number of generations (here g = 1).

## Data availability

Raw reads data of the crosses are available under bioproject PRJNA492790. Reference *D. pseudoobscura* and *D. persimilis* raw reads are submitted in the SRA database under numbers SRR330107 and SRR363439, respectively. Supplementary files are available on the G3 figshare portal: <https://doi.org/10.25387/g3.14501343>. Supplementary Table S1 containing SRA accession numbers of raw reads used in this study; Supplementary Figure S1 summarizing the three *Drosophila* crosses; File IGV\_denovo\_screenshots containing IGV screen shots of mutated positions.

## Results and discussion

In all, I identified 14 nucleotide mutations in 28 F2 progenies. Two F2 progenies were removed because of low coverage, one from the cross PP1134 and one from the cross MSH1993. No mutations were found in the cross PP1134, 4 mutations in the cross MSH177, and 10 mutations in the cross MSH1993. On the 15 identified mutations within the cross MSH1993, 9 arose in a same individual. The mutations were GC → AT biased, with an AT bias of 3.7 and 12.2 in the crosses MSH177 and MSH1993 (assuming one hypothetical AT to GC mutation in MSH1993 cross), respectively. This GC → AT bias is very common in metazoans and even in all eukaryotes [see Table 2 of Katju and Berghorsson

(2018), Supplementary Table S11 of Krasovec et al. (2017)]. Only one reported eukaryote species has a GC → AT mutation bias, the haptophyte *Emiliania huxleyi* (Krasovec et al. 2020).

The calculated mutation rates are  $\mu < 2.7 \times 10^{-09}$  for the cross PP1134 assuming one hypothetical mutation,  $\mu = 3.5 \times 10^{-09}$  (Poisson CI:  $0.7 \times 10^{-09}$ — $10.2 \times 10^{-09}$ ) for the cross MSH177 and  $\mu = 19.3 \times 10^{-09}$  (Poisson CI:  $9.3 \times 10^{-09}$ — $35.6 \times 10^{-09}$ ) for the cross MSH1993 (Table 1). However, the *Drosophila* genome is about 165 Mb and the callable genome for the three crosses was 20, 26, and 63 Mb (Table 2). The callable genome size of the cross MSH1993 is probably reduced due to the divergence between the two species with an average nucleotide identity of 97.7% estimated with fastANI (Jain et al. 2018). Mutation candidates due to miss mapping were removed by IGV inspection (see Materials and methods) and IGV screenshots of *de novo* mutations are provided in supplementary documentation. Considering the variation of the mutation rate within a genome, we cannot exclude that the mutation rates estimated here do not reflect the average mutation rate. The mutation rate varied by approximately fourfold between the lower and the higher measures in the *Drosophila* genus (Table 1), excluding the cross MSH1993 that is a particular case of hybrid species (see below). Taking the cross MSH1993 into account, this variation reached ~11-fold, which is the same mutation rate variation across metazoan species that counts several estimations from mammals (Uchimura et al. 2015; Besenbacher et al. 2016; Koch et al. 2019), birds (Smeds et al. 2016), worms (Denver et al. 2012; Weller et al. 2014), and arthropods (Schridder et al. 2013; Keightley et al. 2014; Flynn et al. 2017; Liu et al. 2017; Oppold and Pfenninger 2017).

Mutation rate variation within the same species, or between close species, is starting to be well documented (Ness et al. 2015; Behringer and Hall 2016; Chu et al. 2018; Liu and Zhang 2019; Ho et al. 2020). The main explanation of such variation is the effect of the environment on the mutation rate. The different isolated *Drosophila* lines used in this study came from different places in United States, Spain, and Africa, corresponding to variable ecological range. Given the sensitivity of the mutation rate to the environment, it is possible that these lines evolved several different mechanisms that control mutation rate or dedicate different amounts of energy to control it. Also, the different experimental conditions that the lines were subjected to between the published studies may have directly increased or decreased the mutation rate. Specifically, experimental conditions will influence factors such as metabolic rate, generation time, oxidative stress or energy devoted to replication/fidelity, and mutagen control, all of which impact the mutation rate (Martin and Palumbi 1993; Baer et al. 2007). Oxidative stress, which is linked to the metabolic rate, has been known for years to increase the mutation rate (Dizdaroglu 1992; Cooke et al. 2003), particularly via guanine oxidation. The generation time is also a key parameter, particularly in multicellular species, because it defines the number of germline cell divisions and the time where the germinal lines are subjected to any mutagenic process. In primates, the spontaneous mutation rate decreases when the generation time is shorter (Thomas et al. 2018; Wu et al. 2020).

The case of the cross MSH1993 is a particular because it is a cross between two species, *D. pseudoobscura* and *D. persimilis*. Hybrids often have higher mutation rate than initial species, such as was reported in *Drosophila* many years ago (Belgovsky 1937; Sturtevant 1939), and more recently in plants from the *Prunus* genus (Xie et al. 2016). However, here we do not have the mutation rate of *D. persimilis*. We so cannot exclude that *D. persimilis* may have a particularly high mutation rate compared to other *Drosophila* species, increasing the mutation rate of the hybrid cross. The mutation rate of the hybrid cross is, however,

**Table 1** Spontaneous mutation rates per site per haploid genome in *Drosophila* genus

| Species  | $\mu$ ( $\times 10^{-9}$ ) | Ref                         |
|--|----------------------------|-----------------------------|
| <i>D. melanogaster</i> (Massachusetts, USA)                            | 5.5                        | Schrider et al. (2013)      |
| <i>D. melanogaster</i> (Ghana)   | 2.8                        | Keightley et al. (2014)     |
| <i>D. melanogaster</i> DGRP360 (North Carolina, USA)                   | 5.2                        | Huang et al. (2016)         |
| <i>D. melanogaster</i> (Madrid line)                                   | 3.5                        | Keightley et al. (2009)     |
| <i>D. melanogaster</i> DGRP765 (North Carolina, USA)                   | 4.9                        | Assaf et al. (2017)         |
| <i>D. melanogaster</i> (North Carolina, USA)                           | 6.0                        | Sharp and Agrawal (2016)    |
| <i>D. melanogaster</i> (Madrid, Spain)                                 | 2.7                        | Haag-Liautard et al. (2007) |
| <i>D. melanogaster</i> Florida-33 (Florida, USA)                       | 11.7                       |                             |
| <i>D. melanogaster</i> Florida-39 (Florida, USA)                       | 6.8                        |                             |
| <i>D. pseudoobscura</i> PP1137 $\times$ <i>D. pseudoobscura</i> PP1134 | <2.7 <sup>a</sup>          | This study                  |
| <i>D. pseudoobscura</i> PP1137 $\times$ <i>D. pseudoobscura</i> MSH177 | 3.5                        |                             |
| <i>D. pseudoobscura</i> PP1137 $\times$ <i>D. persimilis</i> MSH1993   | 19.3                       |                             |

The mutation rates between MSH177 and MSH1993 (hybrids) crosses are significantly different (Chi-squared test, P-value = 0.003).

<sup>a</sup> Mutation rate corresponding to 1 mutation.

**Table 2** Identified mutations in the three crosses

| Crosses  | N <sub>F2</sub> | G*         | Mutations (contig; position; reference; mutation) |          |   |   |
|--|-----------------|------------|---|----------|---|---|
| <i>D. pseudoobscura</i> PP1137 $\times$ <i>D. pseudoobscura</i> PP1134 | 9               | 20,329,686 | No mutation                                       |          |   |   |
| <i>D. pseudoobscura</i> PP1137 $\times$ <i>D. pseudoobscura</i> MSH177 | 9               | 63,534,699 | NC_046680.1                                       | 17833723 | G | A |
|  |                 |            | NC_046681.1                                       | 6119653  | C | T |
|  |                 |            | NC_046681.1                                       | 1779949  | C | A |
|  |                 |            | NC_046679.1                                       | 918510   | T | C |
| <i>D. pseudoobscura</i> PP1137 $\times$ <i>D. persimilis</i> MSH1993   | 10              | 25,855,578 | NC_046680.1                                       | 21310719 | G | T |
|  |                 |            | NC_046680.1                                       | 13587576 | G | T |
|  |                 |            | NC_046681.1                                       | 19040615 | C | A |
|  |                 |            | NC_046681.1                                       | 20681630 | G | T |
|  |                 |            | NC_046679.1                                       | 2025226  | C | A |
|  |                 |            | NC_046679.1                                       | 21427501 | T | A |
|  |                 |            | NC_046679.1                                       | 23593472 | C | A |
|  |                 |            | NC_046680.1                                       | 11444083 | G | T |
|  |                 |            | NC_046681.1                                       | 6516206  | G | T |
|  |                 |            | NC_046681.1                                       | 26994424 | C | A |

N<sub>F2</sub> is the number of F2 progenies, G\* is the number of callable sites.

much higher than in any other *Drosophila* study and of the same order of higher mutation rate ever measured in the tree of life (Katju and Bergthorsson 2018). In *Arabidopsis*, it is hybrids between populations or ecotypes that induce a mutation rate change (Bashir et al. 2014). The reasons for this observed increase of mutation rate in hybrids remains unclear, although there are several proposed explanations. First, the heterozygosity associated mutation rate is the observation that heterozygosity is positively linked to the mutation rate (Amos 2010; Yang et al. 2015). In that case, it is proposed that the higher heterozygosity in hybrids compared to original species induces an elevated mutation rate. Another explanation is the possible fitness decrease of the hybrids compared to original species. Stress or fitness loss is known to be at the origin of mutation rate increase in *Arabidopsis* and *Drosophila* (Baer 2008; Sharp and Agrawal 2012; Jiang et al. 2014) and thus may similarly be the main contributor to higher mutation rate in hybrids.

## Acknowledgments

The author thanks Sheree Yau and Gwenaél Piganeau for their advices.

## Conflicts of interest

No conflict of interest to declare.

## Literature cited

- Amos W. 2010. Heterozygosity and mutation rate: evidence for an interaction and its implications: the potential for meiotic gene conversions to influence both mutation rate and distribution. *Bioessays*. 32:82–90.
- Assaf ZJ, Tilk S, Park J, Siegal ML, Petrov DA. 2017. Deep sequencing of natural and experimental populations of *Drosophila melanogaster* reveals biases in the spectrum of new mutations. *Genome Res*. 27:1988–2000.
- Baer CF. 2008. Does mutation rate depend on itself. *PLoS Biol*. 6:e52.
- Baer CF, Miyamoto MM, Denver DR. 2007. Mutation rate variation in multicellular eukaryotes: causes and consequences. *Nat Rev Genet*. 8:619–631.
- Bashir T, Sailer C, Gerber F, Loganathan N, Bhoopalan H, et al. 2014. Hybridization alters spontaneous mutation rates in a parent-of-origin-dependent fashion in *Arabidopsis*. *Plant Physiol*. 165:424–437.
- Behringer MG, Hall DW. 2016. The repeatability of genome-wide mutation rate and spectrum estimates. *Curr Genet*. 62:507–512.

- Belgovsky ML. 1937. A comparison of the frequency of induced mutations in *Drosophila simulans* and in its hybrid with *D. melanogaster*. *Genetica*. 19:370–386.
- Besenbacher S, Sulem P, Helgason A, Helgason H, Kristjansson H, et al. 2016. Multi-nucleotide de novo mutations in humans. *PLoS Genet*. 12:e1006315.
- Chu X-L, Zhang B-W, Zhang Q-G, Zhu B-R, Lin K, et al. 2018. Temperature responses of mutation rate and mutational spectrum in an *Escherichia coli* strain and the correlation with metabolic rate. *BMC Evol Biol*. 18:126.
- Cooke MS, Evans MD, Dizdaroglu M, Lunec J. 2003. Oxidative DNA damage: mechanisms, mutation, and disease. *FASEB J*. 17:1195–1214.
- Denver DR, Wilhelm LJ, Howe DK, Gafner K, Dolan PC, et al. 2012. Variation in base-substitution mutation in experimental and natural lineages of *Caenorhabditis* nematodes. *Genome Biol Evol*. 4:513–522.
- Dillon MM, Sung W, Sebra R, Lynch M, Cooper VS. 2016. Genome-wide biases in the rate and molecular spectrum of spontaneous mutations in *Vibrio cholerae* and *Vibrio fischeri*. *Mol Biol Evol*. 34:93–109.
- Dizdaroglu M. 1992. Oxidative damage to DNA in mammalian chromatin. *Mutat Res*. 275:331–342.
- Flynn JM, Chain FJJ, Schoen DJ, Cristescu ME. 2017. Spontaneous mutation accumulation in *Daphnia pulex* in selection-free vs. competitive environments. *Mol Biol Evol*. 34:160–173.
- Ford CB, Lin PL, Chase MR, Shah RR, Iartchouk O, et al. 2011. Use of whole genome sequencing to estimate the mutation rate of *Mycobacterium tuberculosis* during latent infection. *Nat Genet*. 43:482–486.
- Haag-Liautard C, Dorris M, Maside X, Macaskill S, Halligan DL, et al. 2007. Direct estimation of per nucleotide and genomic deleterious mutation rates in *Drosophila*. *Nature*. 445:82–85.
- Ho EKH, Macrae F, Latta LC, McIlroy P, Ebert D, et al. 2020. High and highly variable spontaneous mutation rates in *Daphnia*. *Mol Biol Evol*. 37:3258–3266.
- Huang W, Lyman RF, Lyman RA, Carbone MA, Harbison ST, et al. 2016. Spontaneous mutations and the origin and maintenance of quantitative genetic variation. *Elife*. 5:e14625
- Jain C, Rodriguez-R LM, Phillippy AM, Konstantinidis KT, Aluru S. 2018. High throughput ANI analysis of 90K prokaryotic genomes reveals clear species boundaries. *Nat Commun*. 9:5114.
- Jiang C, Mithani A, Belfield EJ, Mott R, Hurst LD, et al. 2014. Environmentally responsive genome-wide accumulation of de novo *Arabidopsis thaliana* mutations and epimutations. *Genome Res*. 24:1821–1829.
- Katju V, Bergthorsson U. 2018. Old trade, new tricks: insights into the spontaneous mutation process from the partnering of classical mutation accumulation experiments with high-throughput genomic approaches. *Genome Biol Evol*. 11:136–165.
- Keightley PD, Ness RW, Halligan DL, Haddrill PR. 2014. Estimation of the spontaneous mutation rate per nucleotide site in a *Drosophila melanogaster* full-sib family. *Genetics*. 196:313–320.
- Keightley PD, Trivedi U, Thomson M, Oliver F, Kumar S, et al. 2009. Analysis of the genome sequences of three *Drosophila melanogaster* spontaneous mutation accumulation lines. *Genome Res*. 19:1195–1201.
- Koch E, Schweizer RM, Schweizer TM, Stahler DR, Smith DW, et al. 2019. De novo mutation rate estimation in wolves of known pedigree. *Mol Biol Evol*. 36:2536–2547.
- Korunes KL, Noor MAF. 2019. Pervasive gene conversion in chromosomal inversion heterozygotes. *Mol Ecol*. 28:1302–1315.
- Krasovec M, Eyre-Walker A, Sanchez-Ferandin S, Piganeau G. 2017. Spontaneous mutation rate in the smallest photosynthetic eukaryotes. *Mol Biol Evol*. 34:1770–1779.
- Krasovec M, Rickaby REM, Filatov DA. 2020. Evolution of mutation rate in astronomically large phytoplankton populations. *Genome Biol Evol*. 12:1051–1059.
- Krasovec M, Sanchez-Brosseau S, Piganeau G. 2019. First estimation of the spontaneous mutation rate in diatoms. *Genome Biol Evol*. 11:1829–1837.
- Kucukyildirim S, Behringer M, Williams EM, Doak TG, Lynch M. 2020. Estimation of the genome-wide mutation rate and spectrum in the Archaeal species *Haloflex volcanii*. *Genetics*. 215:1107–1116.
- Kucukyildirim S, Long H, Sung W, Miller SF, Doak TG, et al. 2016. The rate and spectrum of spontaneous mutations in *Mycobacterium smegmatis*, a bacterium naturally devoid of the post-replicative mismatch repair pathway. *G3 (Bethesda)*. 6:2157–2163.
- Li H. 2011. A statistical framework for SNP calling, mutation discovery, association mapping and population genetical parameter estimation from sequencing data. *Bioinformatics*. 27:2987–2993.
- Li H, Durbin R. 2010. Fast and accurate long-read alignment with Burrows–Wheeler transform. *Bioinformatics*. 26:589–595.
- Li H, Handsaker B, Wysoker A, Fennell T, Ruan J, et al.; 1000 Genome Project Data Processing Subgroup. 2009. The sequence alignment/map format and SAMtools. *Bioinformatics*. 25:2078–2079.
- Liu H, Jia Y, Sun X, Tian D, Hurst LD, et al. 2017. Direct determination of the mutation rate in the bumblebee reveals evidence for weak recombination-associated mutation and an approximate rate constancy in insects. *Mol Biol Evol*. 34:119–130.
- Liu H, Zhang J. 2019. Yeast spontaneous mutation rate and spectrum vary with environment. *Curr Biol*. 29:1584–1591.e3.
- Lynch M, Ackerman MS, Gout J-F, Long H, Sung W, et al. 2016. Genetic drift, selection and the evolution of the mutation rate. *Nat Rev Genet*. 17:704–714.
- Martin AP, Palumbi SR. 1993. Body size, metabolic rate, generation time, and the molecular clock. *Proc Natl Acad Sci U S A*. 90:4087–4091.
- McKenna A, Hanna M, Banks E, Sivachenko A, Cibulskis K, et al. 2010. The genome analysis toolkit: a MapReduce framework for analyzing next-generation DNA sequencing data. *Genome Res*. 20:1297–1303.
- Ness RW, Morgan AD, Vasanthakrishnan RB, Colegrave N, Keightley PD. 2015. Extensive de novo mutation rate variation between individuals and across the genome of *Chlamydomonas reinhardtii*. *Genome Res*. 25:1739–1749.
- Oppold A-M, Pfenniger M. 2017. Direct estimation of the spontaneous mutation rate by short-term mutation accumulation lines in *Chironomus riparius*. *Evol Lett*. 1:86–92.
- Robinson JT, Thorvaldsdóttir H, Winckler W, Guttman M, Lander ES, et al. 2011. Integrative genomics viewer. *Nat Biotechnol*. 29:24–26.
- Schrider DR, Houle D, Lynch M, Hahn MW. 2013. Rates and genomic consequences of spontaneous mutational events in *Drosophila melanogaster*. *Genetics*. 194:937–954.
- Sharp NP, Agrawal AF. 2012. Evidence for elevated mutation rates in low-quality genotypes. *Proc Natl Acad Sci U S A*. 109:6142–6146.
- Sharp NP, Agrawal AF. 2016. Low genetic quality alters key dimensions of the mutational spectrum. *PLoS Biol*. 14:e1002419.
- Smeds L, Qvarnstrom A, Ellegren H. 2016. Direct estimate of the rate of germline mutation in a bird. *Genome Res*. 26:1211–1218. doi: 10.1101/gr.204669.116.

- Strauss C, Long H, Patterson CE, Te R, Lynch M. 2017. Genome-wide mutation rate response to pH change in the coral reef pathogen *Vibrio shilonii* AK1. *mBio*. 8:(4):e01021–17.
- Sturtevant AH. 1939. High mutation frequency induced by hybridization. *Proc Natl Acad Sci U S A*. 25:308–310.
- Thomas GWC, Wang RJ, Puri A, Harris RA, Raveendran M, et al. 2018. Reproductive longevity predicts mutation rates in primates. *Curr Biol*. 28:3193–3197.e5.
- Uchimura A, Higuchi M, Minakuchi Y, Ohno M, Toyoda A, et al. 2015. Germline mutation rates and the long-term phenotypic effects of mutation accumulation in wild-type laboratory mice and mutator mice. *Genome Res*. 25:1125–1134. doi:10.1101/gr.186148.114.
- Weller AM, Rödelberger C, Eberhardt G, Molnar RI, Sommer RJ. 2014. Opposing forces of A/T-biased mutations and G/C-biased gene conversions shape the genome of the nematode *Pristionchus pacificus*. *Genetics*. 196:1145–1152.
- Wu FL, Strand AI, Cox LA, Ober C, Wall JD, et al. 2020. A comparison of humans and baboons suggests germline mutation rates do not track cell divisions. *PLoS Biol*. 18:e3000838.
- Xie Z, Wang L, Wang L, Wang Z, Lu Z, et al. 2016. Mutation rate analysis via parent–progeny sequencing of the perennial peach. I. A low rate in woody perennials and a higher mutagenicity in hybrids. *Proc Biol Sci*. 283:(1841) 20161016.
- Yang S, Wang L, Huang J, Zhang X, Yuan Y, et al. 2015. Parent-progeny sequencing indicates higher mutation rates in heterozygotes. *Nature*. 523:463–467.

Communicating editor: R. Anholt