



HAL
open science

Mitochondrial DNA copy number as a prognostic marker is age-dependent in adult glioblastoma

Baptiste Sourty, Laure-Marie Dardaud, Céline Bris, Valérie Desquiret-Dumas, Blandine Boisselier, Laëtitia Basset, Dominique Figarella-Branger, Alain Morel, Marc Sanson, Vincent Procaccio, et al.

► To cite this version:

Baptiste Sourty, Laure-Marie Dardaud, Céline Bris, Valérie Desquiret-Dumas, Blandine Boisselier, et al.. Mitochondrial DNA copy number as a prognostic marker is age-dependent in adult glioblastoma. *Neuro-Oncology Advances*, 2022, 4 (1), pp.vdab191. 10.1093/noajnl/vdab191 . hal-03560209

HAL Id: hal-03560209

<https://hal.sorbonne-universite.fr/hal-03560209>

Submitted on 7 Feb 2022

HAL is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers.

L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.

Mitochondrial DNA copy number as a prognostic marker is age-dependent in adult glioblastoma

Baptiste Sourty, Laure-Marie Dardaud, Céline Bris, Valérie Desquiret-Dumas, Blandine Boisselier, Laëtitia Basset, Dominique Figarella-Branger[®], Alain Morel, Marc Sanson, Vincent Procaccio[†], and Audrey Rousseau[†]

Department of Pathology, University Hospital of Angers, Angers, France (B.S., L.M.D., B.B., L.B., A.R.); Department of Genetics, University Hospital of Angers and Angers University, INSERM1083, CNRS6015, MITOVASC, Angers, France (C.B., V.D.-D., V.P.); Center for Research in Cancerology and Immunology Nantes/Angers, INSERM, University of Nantes, University of Angers, Angers, France (B.B., L.B., A.R.); Aix-Marseille Univ, APHM, CNRS, INP, Inst Neurophysiopathol, CHU Timone, Service d'Anatomie Pathologique et de Neuropathologie, Marseille, France (D.F.-B.); Institut de Cancérologie de l'Ouest – Paul Papin, Angers, France (A.M.); Sorbonne University UPMC Univ Paris 06, INSERM CNRS, U1127, UMR 7225, ICM, F-75013, Groupe Hospitalier Pitié-Salpêtrière, Neurology Department 2, Paris, France (M.S.)

Corresponding Author: Audrey Rousseau, MD, PhD, Department of Pathology, Angers University Hospital, 4 rue Larrey, 49933 Angers, France (aurousseau@chu-angers.fr).

[†]These authors contributed equally to this work.

Abstract

Background. Glioblastoma (GBM) is the most common and aggressive form of glioma. GBM frequently displays chromosome (chr) 7 gain, chr 10 loss and/or *EGFR* amplification (chr7+/chr10-/*EGFR*amp). Overall survival (OS) is 15 months after treatment. In young adults, *IDH1/2* mutations are associated with longer survival. In children, *histone H3* mutations portend a dismal prognosis. Novel reliable prognostic markers are needed in GBM. We assessed the prognostic value of mitochondrial DNA (mtDNA) copy number in adult GBM.

Methods. mtDNA copy number was assessed using real-time quantitative PCR in 232 primary GBM. Methylation of *POLG* and *TFAM* genes, involved in mtDNA replication, was assessed by bisulfite-pyrosequencing in 44 and 51 cases, respectively.

Results. Median age at diagnosis was 56.6 years-old and median OS, 13.3 months. 153/232 GBM (66 %) displayed chr7+/chr10-/*EGFR*amp, 23 (9.9 %) *IDH1/2* mutation, 3 (1.3 %) *H3* mutation and 53 (22.8 %) no key genetic alterations. GBM were divided into two groups, “Low” ($n = 116$) and “High” ($n = 116$), according to the median mtDNA/nuclear DNA ratio (237.7). There was no significant difference in OS between the two groups. By dividing the whole cohort according to the median age at diagnosis, OS was longer in the “High” vs “Low” subgroup (27.3 vs 15 months, $P = .0203$) in young adult GBM ($n = 117$) and longer in the “Low” vs “High” subgroup (14.5 vs 10.2 months, $P = .0116$) in older adult GBM ($n = 115$). *POLG* was highly methylated, whereas *TFAM* remained unmethylated.

Conclusion. mtDNA copy number may be a novel prognostic biomarker in GBM, its impact depending on age.

Key Points

- High mtDNA copy number correlates with better prognosis in young adult glioblastoma.
- High mtDNA copy number correlates with worse prognosis in older adult glioblastoma.
- *POLG* and *TFAM* genes are highly methylated and unmethylated, respectively, in glioblastoma.

Importance of the Study

The prognostic value of mtDNA copy number in cancer differs according to tumor type and has rarely been investigated in gliomas. We previously demonstrated that high mtDNA copy number was associated with longer overall survival in young adult glioblastoma (<40 years). In the present study, we show that the prognostic value of mtDNA copy number may depend on age, with a longer overall survival in young adults (<56 years) and inversely, a shorter survival in older adults (≥56 years). In

young adult glioblastoma, high mtDNA content may drive oxidative metabolism, associated with cell differentiation and decreased proliferation and invasiveness. Conversely, high mtDNA copy number and mitochondria-related metabolism may not be as advantageous for tumors arising in older patients compared to young ones. Methylation levels of *POLG* and *TFAM* genes, involved in mtDNA replication, do not appear to regulate mtDNA copy number in glioblastoma.

Glioblastoma (GBM) is the most frequent primary brain tumor. This is the most aggressive form of diffuse glioma (WHO grade IV out of IV).¹ Diffuse gliomas are classified according to their *IDH1/2* (isocitrate dehydrogenase 1/2) gene status. Most GBM develop in patients older than 55 years and do not display *IDH* mutation. Those *IDH* wild-type (WT) tumors are characterized by gain of chromosome (chr) 7 (chr7+), loss of chr 10 (chr10-) and/or *EGFR* gene amplification (*EGFR*amp).¹ The median overall survival is 15 months after standard of care treatments including maximum safe resection and concomitant radiochemotherapy.² Some GBM (<10%) derive from lower-grade (II or III) *IDH*-mutant diffuse gliomas in young adults (secondary GBM). Occasionally, *IDH*-mutant GBM develop without past history of a lower-grade tumor (primary GBM). *IDH* mutation is associated with a longer survival (> 2-3 years) compared to that of *IDH*-WT diffuse gliomas. Inversely, most pediatric GBM harbor histone *H3* (*H3F3A* or *HIST1H3B/C*) gene mutations associated with a short survival (< 1 year). However, within each molecular subgroup (*IDH*-WT, *IDH*-mutant, *H3*-mutant), overall survival varies from patient to patient despite similar histopathological and genetic features. Novel reliable prognostic markers are sorely needed in GBM.

Mitochondria are cytoplasmic organelles providing energy to the cell through production of ATP. Two major pathways, glycolytic and oxidative, are involved in cell metabolism, the oxidative pathway being driven by mitochondria and requiring oxygen. The “Warburg effect” corresponds to the shift from an oxidative to a glycolytic metabolism in tumor cells despite aerobic conditions. As a result, tumor cells transform glucose into lactate through fermentation.³ Lactates have a profound impact on tumor microenvironment and display an immunosuppressive effect.⁴ Inversely, oxidative metabolism decreases tumor invasiveness and supports cell differentiation.

Mitochondrial DNA (mtDNA) is a 16 569 bp, intronless, circular, and double-stranded molecule that is maternally inherited. It encodes, among others, 13 polypeptide subunits of the electron transport chain (ETC) implicated in the oxidative phosphorylation (OXPHOS).⁵ One cell may contain hundreds to thousands of mtDNA copies.⁶ mtDNA pathogenic mutations have been described in various cancers, including colorectal and ovarian carcinomas, acute

myeloid leukemias, and GBM.⁷ mtDNA copy number has been correlated with variable prognosis in cancers.⁸⁻¹⁰ We have previously shown that high mtDNA levels were significantly associated with longer overall survival (31.8 months vs 12.9 months; $P = .013$) in young adult GBM patients (≥18 years-old and < 40 years-old; $n = 67$) strongly suggesting that mtDNA copy number may represent a novel prognostic marker in GBM.¹¹ We assessed the correlation between mtDNA levels and prognosis in a larger cohort of GBM patients with a wider age range, older patients being the most affected by the disease.

mtDNA copy number is strictly regulated by several nuclear-encoded transcription and replication factors. Among them, mitochondrial transcription factor A (TFAM) plays a key role in activating the transcription of mtDNA and initiating its replication.^{12,13} mtDNA replication is driven by mitochondria-specific DNA Polymerase Gamma (POLG) that consists of one catalytic subunit (POLGA), encoded by *POLG* gene, and a dimeric form of its accessory subunit (POLGB), encoded by *POLG2* gene. POLGA acts as a DNA polymerase with exonuclease activity and needs an RNA-DNA hybrid primer transcript encoded by *TFAM* and recognized by POLGB to replicate mtDNA.¹⁴ Thus, *TFAM* and *POLG* are essential nuclear-encoded genes in mtDNA copy number regulation.

Gene expression is strictly regulated by epigenetic changes affecting either histones or DNA. DNA methylation is one of the main epigenetic mechanisms that lead to gene silencing.¹⁵ Epigenetic regulation of nuclear DNA (nDNA) has been linked to mtDNA replication through *POLG* exon 2 methylation.¹⁶ Whether mtDNA copy number is directly regulated by methylation of CpG islands within *TFAM* and *POLG* genes has yet to be assessed in GBM.

In the present study, we investigated the correlation between mtDNA copy number and overall survival in a large cohort of GBM patients ≥ 18 years. We demonstrate that high mtDNA copy number is associated with longer survival in young adult GBM and with shorter survival in older patients. We assessed whether tumor mtDNA content was epigenetically regulated through the methylation of two nuclear-encoded genes, *TFAM* and *POLG*, by pyrosequencing. We showed *POLG* exon 2 to be highly methylated in all GBM samples, while *TFAM* promoter remained unmethylated.

Materials and Methods

Glioblastoma Tissue Samples

All primary GBM from patients over 18 years-old operated at Angers University Hospital between 2000 and 2018 were reviewed. Only cases with available clinical, histopathological and molecular data were included in the study ($n = 192$). The histopathological diagnosis was made according to the 2016 WHO classification.¹ Molecular characterization had been previously performed for all cases using pangenomic SNP (Single Nucleotide Polymorphism) arrays (CytoSNP-850K®, Illumina, San Diego, CA, USA) and Sanger sequencing (*IDH1/2* and *H3F3A/HIST1H3B* genes). All patients had provided written informed consent for genetic analyses with the approval of the research ethics committee of Angers University Hospital (Comité de Protection des Personnes, n° CP CB 2015/08). Additionally, 11 GBM samples were provided by the Pitié-Salpêtrière University Hospital (AP-HP, OncoNeuroTek, Paris, France), 21 by the French Glioblastoma Biobank, and 8 were obtained from the Timone University Hospital (AP-HM, AP-HMTumor Bank, <http://doi.org/10.5334/ojb.63>, Marseille, France). Of the 232 cases included, 67 have already been reported.¹¹ The whole cohort is presented in [Supplementary Material S1](#).

Tumor DNA Extraction

Tumor DNA was extracted from fresh-frozen GBM tissue samples using NucleoSpin® Tissue kit (Macherey-Nagel, Hoerdt, France) according to the manufacturer's recommendations. DNA concentration was evaluated using Spectrophotometer Nanodrop ND 1000 (Thermo Fisher Scientific, Courtaboeuf, France).

Quantitative Polymerase Chain Reaction of Tumor mtDNA

Tumor mtDNA was quantified by quantitative polymerase chain reaction (qPCR) using Chromo4 System (Biorad, Hercules, CA, USA) as previously described.⁸ Two mitochondrial-encoded genes, *MT-CO1* et *MTND4*, and two nuclear-encoded genes, β 2-microglobulin (*B2M*) and glyceraldehyde 3-phosphate dehydrogenase (*GAPDH*), were analyzed using specific primers as described in [Supplementary Material S2](#). Their respective copy numbers were evaluated using a calibration range. mtDNA amount was assessed according to the ratio "mtDNA copy number: nDNA copy number". Statistical adjustment was performed for chr 12p (harboring *GAPDH* gene) and 15q (*B2M* gene) copy number alterations and ploidy status using SNP array data.

High-Throughput Sequencing of Tumor mtDNA

High-throughput mtDNA sequencing (IonTorrent, Thermo Fisher Scientific, Waltham, MA, USA) was performed in 28 GBM as previously described.⁶ An in-house bioinformatic pipeline was used to detect mtDNA point mutations, combined with eKLIPse software, designed to detect and quantify mtDNA rearrangements as described elsewhere.^{6,17}

Tumor DNA Bisulfite Conversion, Bisulfite-Specific Polymerase Chain Reaction, and Pyrosequencing

We assessed the DNA methylation status of six CpG sites located in *POLG* exon 2 and in *TFAM* promoter in 44 tumor tissue samples (26 young and 18 older patients (see below)) and 51 tumor tissue samples (31 young and 20 older patients (see below)), respectively. Tumor DNA (1 μ g) was sodium bisulfite converted using the EZ DNA Methylation-Lightning Kit (Zymo Research, Irvine, CA, USA) according to the manufacturer's recommendations.

CpG islands of *TFAM* promoter and *POLG* exon 2 were located using the website *DataBase of Transcriptional Start Sites (DBTSS)*.¹⁸ Their genomic sequences were retrieved from Alamut Visual version 2.15 (Interactive Biosoftware, Rouen, France), with the following ID number for *TFAM* transcript: NM_003201.2 (region 58385067-58385237) and for *POLG* exon 2: NM_002693.2 (region 89333509-89333745). Bisulfite-specific primers (BSP) were designed using the open access MethPrimer 2.0 program¹⁹ ([Supplementary Material S3](#)), with *POLG* reverse primer modified manually. Reverse primers were labeled with biotin at the 5' end and purified using Reverse-Phase High-Performance Liquid Chromatography (RP-HPLC). PCR products of *TFAM* and *POLG* were 171 and 237 base pairs long, respectively. Details of the PCR are provided in the [Supplementary Material S4](#).

Two different pyrosequencing primers were designed for *TFAM* and one for *POLG*, using PyroMark® ID Q96 Software (Qiagen, Venlo, Netherlands) ([Supplementary Material S5](#)). Pyrosequencing reaction was carried out using Pyromark Gold Q96 Reagents (Qiagen, Hilden, Germany), as described elsewhere,²⁰ and performed in the Pyrosequencer ID96 (Qiagen, Venlo, Netherlands). Only samples with pyrograms displaying peak height > 7 RLU (Relative Light Unit) on at least the first 6 CpG sites were taken into account.

Statistical Analyses

Data have been expressed as medians and averages for continuous variables and as absolute values and frequencies for categorical variables. Comparisons were performed using Chi-Square test or Fisher's exact test depending on the number of samples to compare categorical variables. *P* values less than .05 were considered significant. The Kaplan–Meier method was used to compare overall survival rates. Statistical analyses were performed with GraphPad Prism 9.

Results

Clinical and Histomolecular Features of the Cohort

The cohort comprised 232 GBM diagnosed according to the 2016 WHO classification. All cases were primary GBM, with no past history of lower-grade diffuse glioma. The median age at diagnosis was 56.6 years and the sex ratio (M:F) 1.6:1. The median overall survival was

13.3 months. The cohort was subdivided into four molecular groups: 1) 153 GBM (66 %) were *IDH*-WT and characterized by chr 7 gain/chr 10 loss and/or *EGFR* amplification, 2) 23 (9.9 %) were *IDH*-mutant, 3) 3 (1.3 %) displayed *H3F3A/HIST1H3B* mutations (*H3*-mutant), and 4) 53 (22.8 %) had none of the aforementioned alterations (referred to as NEC for Not Elsewhere Classified). The median age at diagnosis was 58.5 years for the *IDH*-WT group, 34 years for the *IDH*-mutant group, 39 years for the *H3*-mutant group, and 55.2 years for the NEC group. The median overall survival was 15.5 months for the *IDH*-WT group, 34.5 months for the *IDH*-mutant group, 11.3 months for the *H3*-mutant group, and 14.1 months for the NEC group.

Quantification of mtDNA copy number

The mtDNA level was assessed using the mtDNA/nDNA ratio that ranged from 28 to 3,882 (median ratio 237.7). Patients were subdivided into two groups according to the median ratio: 1) Low mtDNA level (116 patients) where the mtDNA/nDNA ratio \leq the median ratio and 2) High mtDNA level (116 patients) where the mtDNA/nDNA ratio $>$ the median ratio. The overall survival of the Low vs High group was not significantly different (14.5 months vs 17 months, respectively; $P = .75$) (Figure 1).

Since we had shown in a previous work that high mtDNA levels were significantly associated with longer overall survival in young adult GBM patients, we assessed this correlation in a larger cohort of GBM patients with a wider age range. Patients were subdivided into two groups according to the median age at diagnosis (56.6 years) in order to distinguish a young adult group (age $<$ median age, $n = 117$ patients) and an older adult group (age \geq median age, $n = 115$ patients). A correlation between mtDNA level (High or Low) and overall survival was assessed in both (young and older) age groups. Of note, mtDNA amount was not associated with patient age ($P = .44$, Kruskal-Wallis test).




Figure 1. Overall survival in the young adult group (High vs Low GBM). Kaplan–Meier survival curves in the young adult group stratified according to mtDNA content.

High mtDNA Copy Number is Associated with Longer Overall Survival in Young Adult GBM

In the young adult group, the overall survival was significantly longer in the High vs Low subgroup (27.3 months vs 15 months; $P = .0203$) (Figure 2). In this age group ($n = 117$ patients), 65 GBM (55 %) were *IDH*-WT, 22 (19 %) *IDH*-mutant, 3 (3 %) *H3*-mutant and 27 (23 %) NEC. There was no significant association between the High status and the *IDH*-mutant status ($P = .17$, Chi-Square test).

In the NEC group, the overall survival was significantly longer in the High subgroup ($n = 13$) compared to the Low subgroup ($n = 14$) (27.8 months vs 12.9 months, respectively; $P = .0214$). In the *IDH*-mutant group, there was a trend toward a longer overall survival in the High subgroup ($n = 14$) compared to the Low subgroup ($n = 8$) but that did not reach statistical significance (37.6 months vs 15 months, respectively; $P = .89$). In *IDH*-WT GBM, no significant difference in survival was found in the High ($n = 31$) vs Low group ($n = 34$) (17.6 months vs 17.1 months, respectively; $P = .69$). In *H3*-mutant GBM, overall survival was shorter in the High vs Low group (5 months vs 13.7 months, respectively; $P = .15$) but the very small number of patients in this group ($n = 3$) does not allow to draw any definitive conclusion (Table 1).

High mtDNA Copy Number is Associated with Shorter Overall Survival in Older Adult GBM

In the older age group, the overall survival was significantly shorter in the High vs Low group (10.2 months vs 14.5 months, respectively; $P = .0116$) in contrast to what was observed in the young adult group (Figure 3). In the older age group ($n = 115$ patients), 88 GBM (76 %) were *IDH*-WT, one (1 %) was *IDH*-mutant, none was *H3*-mutant, and 26 (23 %) were NEC.

For the *IDH*-WT cases, the overall survival was significantly shorter in the High vs Low group (11.4 months vs 17.1 months, respectively, $P = .0228$; 43 and 45 patients in




Figure 2. Overall survival in the young adult group (High vs Low GBM). Kaplan–Meier survival curves in the young adult group stratified according to mtDNA content.

Table 1. Median Overall Survival in Months According to mtDNA Content and Molecular Subgroup in Young Adult GBM

	High group (n = 59)	Low group (n = 58)	P value
Young adult group (n = 117)	27.3	15	.0203
IDH-WT (n = 65)	17.6	17.1	.69
NEC (n = 27)	27.8	12.9	.0214
IDH-mutant (n = 22)	37.6	15	.89
H3-mutant (n = 3)	5	13.7	.15

The High group comprised 31 *IDH*-WT GBM (53%), 14 *IDH*-mutant GBM (24%), one *H3*-mutant GBM (1%), and 13 NEC GBM (22%). The Low group comprised 34 *IDH*-WT GBM (59%), 8 *IDH*-mutant GBM (14%), two *H3*-mutant GBM (3%), and 14 NEC GBM (24%). Molecular subgroups were equally distributed between both groups (High vs Low). In the NEC subgroup, High GBM displayed a significantly longer overall survival compared to Low GBM ($P = .0214$). In the *IDH*-mutant subgroup, there was a trend towards longer overall survival in the High vs Low GBM but the difference did not reach statistical significance.


each group, respectively). For the NEC cases, there was a trend towards a shorter survival in the High vs Low group but that did not reach statistical significance (7.1 months vs 12.3 months, respectively, $P = .19$; 13 patients in each group) (Table 2).

High-Throughput Sequencing of Tumor mtDNA Reveals No Alteration

mtDNA sequencing was performed in 28 GBM (12%), from 18 young adults (10 High and 8 Low) and 10 older patients (6 High and 4 Low). Of these cases, 14 were *IDH*-WT, 7 *IDH*-mutant, and 7 NEC. Using the eKLIpse software,¹⁷ no significant accumulation of mtDNA rearrangements was detected (notably no deletion). No somatic variant was found in young adult GBM and only very few somatic variants were detected in older adult cases (average: 1 variant per tumor).

POLG Exon 2 is Highly Methylated While TFAM Promoter Remains Unmethylated in all GBM Samples

To highlight an underlying epigenetic mechanism involved in mtDNA levels in tumor cells, we focused on two genes implicated in mtDNA copy number regulation. According to the above-mentioned criteria (see Materials and Methods), 20 GBM samples (13 young and 7 older patients) yielded valid results for *POLG* exon 2. We found very high methylation levels of *POLG* exon 2 both in the Low (12 patients) and High (8 patients) groups, with similar median methylation levels of 86.3% and 82.3%, respectively (Figure 4A). With regard to *TFAM* promoter, 34 GBM samples (19 young and 15 older patients) yielded interpretable results. No CpG site was methylated (methylation level of 0%) in any GBM, whether from the Low (21 cases) or High (13 cases) group (Figure 4B).

**Figure 3.** Overall survival in the older adult group (High vs Low GBM). Kaplan–Meier survival curves in the older adult group stratified according to mtDNA content.

Discussion

We studied 232 primary GBM from patients aged 18 and over. Four molecular subgroups were defined according to the underlying genetic alterations. The aim of the study was to evaluate the prognostic impact of mitochondrial genetics in the whole cohort and within each molecular subgroup—since genetic alterations with known prognostic value may be confounding factors. mtDNA copy number was assessed in all 232 cases and mtDNA was sequenced by NGS in a series of 28 cases (28/232, 12%). Methylation status of two nuclear-encoded genes implicated in mtDNA copy number regulation (i.e. *TFAM* and *POLG*) was assessed.

We distinguished two groups of GBM, High and Low, according to the median mtDNA/nDNA ratio. No significant correlation was found between mtDNA copy number and overall survival in the whole cohort ($P = .75$). These results differ from those of a previous work where a high mtDNA

Table 2. Median Overall Survival in Months According to mtDNA Content and Molecular Subgroup in Older Adult GBM

	High group (n = 57)	Low group (n = 58)	P value
Older adult group (n = 115)	10.2	14.5	.0116
IDH-WT (n = 88)	11.4	17.1	.0228
NEC (n = 26)	7.1	12.3	.19
IDH-mutant (n = 1)	-	2.7	-

The High group comprised 43 *IDH*-WT GBM (75%), one *IDH*-mutant GBM (2%), and 13 NEC GBM (23%). The Low group comprised 45 *IDH*-WT GBM (78%), and 13 NEC GBM (22%). Molecular subgroups were equally distributed between both groups (High vs Low). In the *IDH*-WT subgroup, Low GBM displayed a significantly longer overall survival compared to High GBM. In the NEC subgroup, there was a trend towards longer overall survival in the Low vs High GBM but that did not reach statistical significance.

copy number was correlated with a longer overall survival in a cohort of 67 young adult GBM patients (18–40 years) (31.8 vs 12.9 months; $P = .013$).¹¹ However, by segregating patients according to the median age at diagnosis (56.6 years), we again demonstrated that a high mtDNA level was associated with a better overall survival in the young adult group (27.3 months vs 15 months; $P = .0203$). Hence, high mtDNA copy number appears as a positive prognostic factor in the young adult population. Interestingly, the median age of our cohort (hence the cutoff we chose to distinguish young vs older adults) was very close to the age proposed by the 2016 WHO classification (i.e. 55 years-old) to distinguish primary *IDH*-WT GBM occurring in older patients from presumably secondary *IDH*-mutant GBM arising in young adults.¹ In the young adults from the NEC group, a high mtDNA copy number was associated with a longer overall survival compared to a low mtDNA copy number (27.8 months vs 12.9 months; $P = .0214$) and a similar trend towards longer survival was observed in the High vs Low subgroup within the *IDH*-mutant group. The lack of association between High status and *IDH*-mutant status (test du Chi,² $P = .17$) indicates that *IDH* mutation is not a confounding factor for longer survival. Hence, the prognostic value of mtDNA copy number in GBM may be independent of the underlying genetic alterations.

Mitochondria play a key role in various cellular functions, including energy production, formation of reactive oxygen species, and initiation of apoptosis.^{21,22} Mitochondria are involved in aging²³ and cancers.²⁴ Somatic mtDNA variants have been reported in different cancer types and may play a role in tumorigenesis and tumor progression.^{25,26} mtDNA levels have been associated with longer or shorter survival depending on cancer types. Our results in young GBM patients concur with those of previously published studies showing high mtDNA levels to be correlated with better prognosis and low levels with shorter overall survival. One meta-analysis on 18 studies found that high mtDNA copy number was associated with a better outcome in various cancers, including digestive tract cancers, breast cancers, and gliomas.⁸ Two studies demonstrated that a low mtDNA copy number was associated with tumor progression and lymph node metastases in colorectal cancers.^{9,10} The authors linked this result to the ability of tumor cells with low mtDNA content to survive in hypoxic conditions by shifting their metabolism from mitochondrial OXPHOS to anaerobic glycolysis. Moreover, these tumors may display increased chemoresistance. By contrast, tumors with high mtDNA copy number were described as sensitive to hypoxia, slow-growing, and less invasive.⁹ Other studies showed that high mtDNA levels correlated with a poor prognosis in some tumor types (e.g. colorectal²⁷ and prostate cancers²⁸). Thus, the prognostic value of mtDNA copy number markedly differs across tumor types.

Very few studies have focused on mtDNA copy number in GBM. In diffuse gliomas, decreased mtDNA copy number was correlated with increased malignancy

according to WHO grade, with lowest mtDNA amounts in GBM (WHO grade IV) suggesting a likely role of mtDNA depletion in tumor progression.²⁹ mtDNA copy number was significantly reduced in all tumors as compared to non-neoplastic tissue. In a similar way and in accordance with our results in young adults, low mtDNA copy number was recently associated with poor outcome and treatment resistance in GBM.³⁰ GBM cancer cells maintain low copy number leading to decreased mitochondrial function and promotion of anaerobic glycolysis and cell proliferation.³¹ A very recent study identified four distinct GBM subtypes based on RNA sequencing data and key biological functions.³² Mitochondrial GBM relying on OXPHOS were associated with a better prognosis compared to glycolytic/plurimetabolic GBM depending on glycolysis/hypoxia-related functions.

Other works showed that varying mtDNA levels *in vitro* and *in vivo* in human GBM cell lines had an impact on tumorigenesis and cell differentiation.³³ mtDNA copy number is strictly regulated during cell differentiation, with mtDNA levels depending on cell type-specific energy needs. While embryonic stem cells, which are characterized by high proliferation rates, maintain low mtDNA levels, mtDNA copy number increases during cell differentiation to meet specific functional requirements.³⁴ Dickinson *et al.* demonstrated that during astrocytic differentiation, human neural stem cells displayed very high mtDNA levels contrary to human GBM cell lines, that are unable to increase their respiratory capability and differentiate.³³ However, GBM cell lines highly depleted in mtDNA displayed reduced cell proliferation *in vitro* and increased expression of early developmental genes (*OCT4*, *SHH*). Interestingly, the most aggressive tumors (fast growth and large volume) were observed with partial mtDNA depletions. This study underlines the importance (and complexity of the regulation) of mtDNA copy number in cell differentiation and tumor growth. Similarly, one study showed that mtDNA copy number might regulate methylation levels of nDNA in GBM cell lines *in vitro* and *in vivo* in a mouse model.³⁵ mtDNA depletion led to nDNA demethylation and subsequently to differentiation of tumor cells into mature astrocytes by de-repression of differentiation genes via nDNA demethylation. High mtDNA levels may lead to promoter methylation (inactivation) of some oncogenes and/or promoter demethylation (activation) of some tumor suppressor genes. This study highlights the bidirectional control of nuclear and mitochondrial genomes.

Gene expression is highly regulated by DNA methylation occurring at CpG islands. One study investigated the impact of nDNA methylation on mtDNA copy number in cancer cells from GBM cell lines, demonstrating that mtDNA is epigenetically regulated.³⁶ Demethylation agents (such as 5-azacytine and vitamin C) promoted a > 3-fold increase in mtDNA copy number, pointing out that DNA demethylation of key genomic regions is associated with increased mtDNA replication. Interestingly, *POLG* was

in the study is based on the first six CpG sites. A: *POLG* pyrograms. *Top*: Low GBM displaying an average methylation level of 76 % (case n°109, patient age: 40.6 years, ratio mtDNA/nDNA: 71.5). *Bottom*: High GBM displaying an average methylation level of 95 % (case n°173, patient age 68.9 years, ratio mtDNA/nDNA: 382.6). B: *TFAM* pyrograms. Here, only Thymines (T) (instead of cytosine (C)) are read prior to Guanines (G): the samples did not display methylation at any CpG site (methylation level of 0 %). *Top*: Low GBM (case n°219, patient age: 67.8 years, ratio mtDNA/nDNA: 154.5). *Bottom*: High GBM (case n°69, patient age: 23 years, ratio mtDNA/nDNA: 256.4).

found upregulated by demethylation, whereas *TFAM* was downregulated, suggesting their likely divergent roles in mtDNA replication.³⁶

In the present study, we found high methylation levels of *POLG* regardless of mtDNA content. To our knowledge, these results are the first to confirm on tumor tissue samples obtained on cancer cell lines by Lee *et al.*, highlighting that *POLG* exon 2 is highly methylated in human adult GBM.³⁷ Of note, high levels of gene body methylation have been found in active genes,³⁸ precluding any conclusion on *POLG* transcriptional activity in GBM. *POLG* methylation levels do not appear to be correlated to mtDNA amount, in opposition to our original hypothesis, suggesting that other mechanisms are implicated in the regulation of mtDNA copy number in GBM. However, bisulfite sequencing did not allow us to evaluate *de novo* DNA demethylation occurring through hydroxylation of 5-methylcytosine by ten-eleven translocation (TET) family demethylases,³⁹ that could be involved in mtDNA copy number increase in some GBM. Besides, the methylation analysis performed herein was done on a small number of tumors.

As already mentioned, *TFAM* is a key factor in mtDNA regulation, mtDNA transcription, and replication initiation. While the vast majority of CpG dinucleotides are methylated throughout the genome (around 80 %), most promoter-associated CpG islands appear to be largely unmethylated.⁴⁰ To the best of our knowledge, few studies investigated *TFAM* promoter methylation status in cancers. One study found that *in vitro* methylation of the nuclear respiratory factor-1 (NRF-1) binding site of the *TFAM* promoter could inhibit its downstream transcriptional activity.⁴¹ In the present study, we first hypothesized that mtDNA copy number might be regulated by *TFAM* promoter methylation. The *TFAM* promoter sequence analyzed in our study was not methylated at any CpG site in any tumor tested, suggesting that the CpG island located within *TFAM* promoter remains unmethylated in GBM. These results have to be interpreted carefully as we did not investigate *TFAM* gene expression in GBM. Moreover, alternative mechanisms have been implicated in *TFAM* regulation, including post-transcriptional changes involving micro-RNA, interactions with key regulating factors (such as NRF-1), and post-translational modifications (such as acetylation, phosphorylation, ubiquitination).^{42–45} Interestingly, *TFAM* has been found upregulated and overexpressed in glioma cell lines and glioma tissue specimens.⁴⁶ The role of *TFAM* in mtDNA regulation has yet to be fully understood.

RNA sequencing was performed in a series of 43 GBM samples and differential expression levels of *POLG* and *TFAM* were not statistically different between the High and Low groups (*data not shown*). We did not investigate other nuclear genes encoding key factors implicated in mtDNA replication (e.g. mtDNA helicase TWINKLE, mitochondrial single-stranded binding protein (mtSSB), or mitochondrial RNA polymerase POLRMT). Since *POLG* and *TFAM* methylation levels do not correlate with mtDNA levels in the present study, investigating the involvement of other key players in mtDNA replication is necessary and will be part of future experiments. Besides, mtDNA replication (hence mtDNA copy number) has been found to differ according to the underlying mitochondrial haplogroups.⁴⁷ mtDNA sequencing on a larger cohort is needed to assess the

potential contribution of these haplogroups to the differences in mtDNA copy number in GBM.

In the older adult group, low mtDNA copy number was associated with longer overall survival compared to high mtDNA copy number (14.5 vs 10.2 months; $P = .0116$). This result may be explained by profound tumor metabolism changes during aging. The latter has been shown to alter mitochondria and cell metabolism.^{48,49} Numerous studies reported accumulation of mtDNA deletions and mutations during the aging process, especially in tissues characterized by high energy demand, such as the brain.⁵⁰ Age-dependent mtDNA alterations are associated with decreased mitochondrial function and decreased oxidative metabolism.⁵¹ Given the differences in metabolism observed between cells from the same tissue according to age, one tumor arising at two different time points will depend on the metabolic characteristics of the tissue of origin and its microenvironment. It can thus be hypothesized that high mtDNA copy number and mitochondria-related metabolism may not be as advantageous for tumors arising in older patients compared to young ones.

High-throughput (NGS) sequencing of mtDNA in 28 (12 %) GBM did not show any large mtDNA rearrangement and showed very few somatic variants in older adult GBM. Larman *et al.* reported the presence of mtDNA pathogenic mutations in only 13 % of primary GBM.⁷ The extremely rapid tumor growth in primary GBM may not allow the accumulation of somatic mtDNA mutations in opposition to slow-growing tumors such as colorectal or prostate cancers.^{7,52} However, the limited number of GBM cases sequenced in our study may have hindered the detection of mtDNA alterations.

In conclusion, high mtDNA copy number is an independent prognostic factor associated with longer overall survival in young GBM patients. The prognostic value of mtDNA copy number may be independent from the underlying genetic alterations. Low mtDNA copy number was associated with better outcome in older patients, suggesting age-related mitochondrial/metabolic changes. These results need to be validated in larger cohorts. Comprehensive characterization of the methylation patterns of mtDNA copy number regulatory genes using high-throughput methyl-sequencing would help assess the impact of epigenetics on mtDNA content. GBM metabolism remains to be thoroughly evaluated to determine whether mtDNA copy number reflects metabolic reprogramming during tumorigenesis or rather regulates tumor metabolism.

Supplementary material

Supplementary data are available at *Neuro-Oncology Advances* online.

Keywords

glioblastoma | metabolism | methylation | mitochondrial DNA | prognosis

Acknowledgments

We thank Luc Fey for technical support. We thank the Assistance Publique—Hôpitaux de Marseille (AP-HM) Tumor Bank (authorization number: AC2018-31053; CRB BB-0033-00097), the Assistance Publique - Hôpitaux de Paris (AP-HP) OncoNeuroTek Tumor Bank and the French Glioblastoma Biobank for providing samples.

Funding

None declared.

Conflict of interest statement. The authors declare no conflicts of interest.

Authorship Statement. Baptiste Sourty (experimental work, data analysis, writing), Laure-Marie Dardauid (experimental work, data collection, writing), Céline Bris (data analysis, editing), Valérie Desquiret-Dumas (data analysis), Blandine Boisselier (data analysis), Laëticia Basset (data analysis), Dominique Figarella-Branger (providing samples, editing), Alain Morel (providing technical support, editing), Marc Sanson (providing samples, editing) Vincent Procaccio (study design, writing, editing), Audrey Rousseau (study design, writing, editing).

References

- Louis DN, Ohgaki H, Wiestler OD, Cavenee WK, eds. *WHO Classification of Tumours of the Central Nervous System*. Lyon: IARC; 2016.
- Stupp R, Mason WP, van den Bent MJ, et al. Radiotherapy plus concomitant and adjuvant temozolomide for glioblastoma. *N Engl J Med*. 2005;352(10):987–996.
- Liberti MV, Locasale JW. The Warburg effect: how does it benefit cancer cells? *Trends Biochem Sci*. 2016;41(3):211–218.
- Romero-Garcia S, Moreno-Altamirano MMB, Prado-Garcia H, et al. Lactate contribution to the tumor microenvironment: mechanisms, effects on immune cells and therapeutic relevance. *Front Immunol*. 2016;7:52.
- Yan C, Duanmu X, Zeng L, Liu B, Song Z. Mitochondrial DNA: distribution, mutations, and elimination. *Cells* 2019;8(4):379.
- Boucret L, Bris C, Seegers V, et al. Deep sequencing shows that oocytes are not prone to accumulate mtDNA heteroplasmic mutations during ovarian ageing. *Hum Reprod*. 2017;32(10):2101–2109.
- Larman TC, DePalma SR, Hadjipanayis AG, et al. Spectrum of somatic mitochondrial mutations in five cancers. *Proc Natl Acad Sci USA*. 2012;109(35):14087–14091.
- Chen N, Wen S, Sun X, et al. Elevated mitochondrial DNA copy number in peripheral blood and tissue predict the opposite outcome of cancer: a meta-analysis. *Sci Rep*. 2016;6:37404.
- Cui H, Huang P, Wang Z, et al. Association of decreased mitochondrial DNA content with the progression of colorectal cancer. *BMC Cancer* 2013;13:110.
- van Osch FHM, Voets AM, Schouten LJ, et al. Mitochondrial DNA copy number in colorectal cancer: between tissue comparisons, clinicopathological characteristics and survival. *Carcinogenesis* 2015;36(12):1502–1510.
- Dardauid L-M, Bris C, Desquiret-Dumas V, et al. High mitochondrial DNA copy number is associated with longer survival in young patients with glioblastoma. *Neuro-Oncol*. 2019;21(8):1084–1085.
- Fisher RP, Clayton DA. Purification and characterization of human mitochondrial transcription factor 1. *Mol Cell Biol*. 1988;8(8):3496–3509.
- Kang D, Kim SH, Hamasaki N. Mitochondrial transcription factor A (TFAM): roles in maintenance of mtDNA and cellular functions. *Mitochondrion* 2007;7(1-2):39–44.
- Falkenberg M. Mitochondrial DNA replication in mammalian cells: overview of the pathway. *Essays Biochem*. 2018;62(3):287–296.
- Easwaran H, Tsai H-C, Baylin SB. Cancer epigenetics: tumor heterogeneity, plasticity of stem-like states, and drug resistance. *Mol Cell*. 2014;54(5):716–727.
- Kelly RDW, Mahmud A, McKenzie M, et al. Mitochondrial DNA copy number is regulated in a tissue specific manner by DNA methylation of the nuclear-encoded DNA polymerase gamma A. *Nucleic Acids Res*. 2012;40(20):10124–10138.
- Goudenège D, Bris C, Hoffmann V, et al. eKLIpSe: a sensitive tool for the detection and quantification of mitochondrial DNA deletions from next-generation sequencing data. *Genet Med Off J Am Coll Med Genet*. 2019;21(6):1407–1416.
- DBTSS Home. *Database of Transcriptional Start Sites*. Available at: <https://dbtss.hgc.jp/>.
- Li L-C, Dahiya R. MethPrimer: designing primers for methylation PCRs. *Bioinformatics* 2002;18(11):1427–1431.
- Tost J, Gut IG. DNA methylation analysis by pyrosequencing. *Nat Protoc* 2007;2(9):2265–2275.
- Hekimi S, Wang Y, Noë A. Mitochondrial ROS and the effectors of the intrinsic apoptotic pathway in aging cells: the discerning killers!. *Front Genet*. 2016;7:161.
- Yee C, Yang W, Hekimi S. The intrinsic apoptosis pathway mediates the pro-longevity response to mitochondrial ROS in *C. elegans*. *Cell* 2014;157(4):897–909.
- Wang Y, Hekimi S. Mitochondrial dysfunction and longevity in animals: untangling the knot. *Science* 2015;350(6265):1204–1207.
- Porporato PE, Filigheddu N, Pedro JMB-S, et al. Mitochondrial metabolism and cancer. *Cell Res*. 2018;28(3):265–280.
- Kroemer G, Pouyssegur J. Tumor cell metabolism: cancer's Achilles' heel. *Cancer Cell* 2008;13(6):472–482.
- Baysal BE. Role of mitochondrial mutations in cancer. *Endocr Pathol*. 2006;17(3):203–212.
- Wang Y, He S, Zhu X, et al. High copy number of mitochondrial DNA predicts poor prognosis in patients with advanced stage colon cancer. *Int J Biol Markers*. 2016;31(4):e382–e388.
- Kalsbeek AMF, Chan EKF, Grogan J, et al. Altered mitochondrial genome content signals worse pathology and prognosis in prostate cancer. *Prostate* 2018;78(1):25–31.
- Correia RL, Oba-Shinjo SM, Uno M, et al. Mitochondrial DNA depletion and its correlation with TFAM, TFB1M, TFB2M and POLG in human diffusely infiltrating astrocytomas. *Mitochondrion* 2011;11(1):48–53.
- Sravva P, Nimbalkar VP, Kanuri NN, et al. Low mitochondrial DNA copy number is associated with poor prognosis and treatment resistance in glioblastoma. *Mitochondrion* 2020;55:154–163.
- Lee WTY, St John J. The control of mitochondrial DNA replication during development and tumorigenesis. *Ann N Y Acad Sci*. 2015;1350:95–106.

32. Garofano L, Migliozi S, Oh YT, et al. Pathway-based classification of glioblastoma uncovers a mitochondrial subtype with therapeutic vulnerabilities. *Nat Cancer* 2021;2(2):141–156.
33. Dickinson A, Yeung KY, Donoghue J, et al. The regulation of mitochondrial DNA copy number in glioblastoma cells. *Cell Death Differ*. 2013;20(12):1644–1653.
34. Facucho-Oliveira JM, Alderson J, Spikings EC, et al. Mitochondrial DNA replication during differentiation of murine embryonic stem cells. *J Cell Sci*. 2007;120(Pt 22):4025–4034.
35. Sun X, St John JC. Modulation of mitochondrial DNA copy number in a model of glioblastoma induces changes to DNA methylation and gene expression of the nuclear genome in tumours. *Epigenetics Chromatin* 2018;11(1):53.
36. Sun X, Johnson J, St John JC. Global DNA methylation synergistically regulates the nuclear and mitochondrial genomes in glioblastoma cells. *Nucleic Acids Res*. 2018;46(12):5977–5995.
37. Lee W, Johnson J, Gough DJ, et al. Mitochondrial DNA copy number is regulated by DNA methylation and demethylation of POLGA in stem and cancer cells and their differentiated progeny. *Cell Death Dis*. 2015;6:e1664.
38. Hellman A, Chess A. Gene body-specific methylation on the active X chromosome. *Science* 2007;315(5815):1141–1143.
39. Kohli RM, Zhang Y. TET enzymes, TDG and the dynamics of DNA demethylation. *Nature* 2013;502(7472):472–479.
40. Deaton AM, Bird A. CpG islands and the regulation of transcription. *Genes Dev*. 2011;25(10):1010–1022.
41. Choi YS, Kim S, Kyu Lee H, et al. In vitro methylation of nuclear respiratory factor-1 binding site suppresses the promoter activity of mitochondrial transcription factor A. *Biochem Biophys Res Commun*. 2004;314(1):118–122.
42. Wu K, Zhao Z, Xiao Y, et al. Roles of mitochondrial transcription factor A and microRNA-590-3p in the development of colon cancer. *Mol Med Rep*. 2016;14(6):5475–5480.
43. Wu K, Ma J, Zhan Y, et al. Down-regulation of microRNA-214 contributed to the enhanced mitochondrial transcription factor A and inhibited proliferation of colorectal cancer cells. *Cell Physiol Biochem*. 2018;49(2):545–554.
44. King GA, Hashemi Shabestari M, Taris K-KH, et al. Acetylation and phosphorylation of human TFAM regulate TFAM-DNA interactions via contrasting mechanisms. *Nucleic Acids Res*. 2018;46(7):3633–3642.
45. Dong Z, Pu L, Cui H. Mitoeigenetics and its emerging roles in cancer. *Front Cell Dev Biol*. 2020;8:4.
46. Lee H, Park J, Tran Q, et al. Mitochondrial transcription factor A (TFAM) is upregulated in glioma. *Mol Med Rep*. 2017;15(6):3781–3786.
47. Suissa S, Wang Z, Poole J, et al. Ancient mtDNA genetic variants modulate mtDNA transcription and replication. *PLoS Genet*. 2009;5(5):e1000474.
48. Tidwell TR, Sørreide K, Hagland HR. Aging, metabolism, and cancer development: from Peto's Paradox to the Warburg Effect. *Aging Dis* 2017;8(5):662–676.
49. Ren R, Ocampo A, Liu G-H, Izpisua Belmonte JC. Regulation of stem cell aging by metabolism and epigenetics. *Cell Metab*. 2017;26(3):460–474.
50. Kazachkova N, Ramos A, Santos C, et al. Mitochondrial DNA damage patterns and aging: revising the evidences for humans and mice. *Aging Dis* 2013;4(6):337–350.
51. Filograna R, Mennuni M, Alsina D, et al. Mitochondrial DNA copy number in human disease: the more the better? *FEBS Lett*. 2021;595(8):976–1002.
52. Schöpf B, Weissensteiner H, Schöäfer G, et al. OXPHOS remodeling in high-grade prostate cancer involves mtDNA mutations and increased succinate oxidation. *Nat Commun*. 2020;11(1):1487.