



HAL
open science

C1q+ macrophages: passengers or drivers of cancer progression

Margot Revel, Catherine Sautès-Fridman, Wolf Fridman, Lubka Roumenina

► **To cite this version:**

Margot Revel, Catherine Sautès-Fridman, Wolf Fridman, Lubka Roumenina. C1q+ macrophages: passengers or drivers of cancer progression. Trends in Cancer, In press, 10.1016/j.trecan.2022.02.006 . hal-03612203

HAL Id: hal-03612203

<https://hal.sorbonne-universite.fr/hal-03612203v1>

Submitted on 17 Mar 2022

HAL is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers.

L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.

1 **C1q+ macrophages: passengers or drivers of cancer progression**

2 Margot Revel¹, Catherine Sautès-Fridman^{1,2}, Wolf Herman Fridman^{1,2}, Lubka T. Roumenina^{1*}

3 ¹ Centre de Recherche des Cordeliers, INSERM, Sorbonne Université, Université de Paris, F-
4 75006 Paris, France.

5 ² Equipe labellisée Ligue contre le Cancer, 2021-2

6

7 *Correspondence to: Lubka T. Roumenina, Ph.D.; Cordeliers Research Center, INSERM UMRS
8 1138; 15 rue de l'Ecole de Medecine; 75006 Paris, France. Phone: 33-1-44-27-90-96/ Fax: 33-
9 1-40-51-04-20, E-mail: lubka.roumenina@sorbonne-universite.fr, @roumenina

10

11 **Abstract**

12 The omics era made possible the quest for efficient markers for cancer progression and
13 revealed that macrophage populations are much more complex than just the M1/M2
14 dichotomy. Complement C1q pops up as a marker of a tolerogenic and immunosuppressive
15 macrophage populations in both healthy and tumor tissues, but the specific role of C1q+
16 tumor associated macrophages is poorly understood. C1q is co-expressed in healthy and
17 tumor macrophages with HLA-DR, APOE and MRC1 (CD206), suggesting a resident origin of
18 this population. Tumor associated macrophages expressing C1q correlate with T cell
19 exhaustion and poor prognosis in numerous cancers. Herein, we discuss the plural roles of
20 C1q in these macrophages and how it could drive cancer progression.

21 Keywords: C1q, tumor-associated macrophages, T cells exhaustion, complement system

22

23 **Highlights**

- 24 • Complement component C1q is a marker of a particular sub-population of tissue-
25 resident macrophages and tumor associated macrophages (TAM), often expressing CD206,
26 HLA-DR, SEPP1, FOLR2, APOE but not SPP1, as revealed by scRNAseq in different normal
27 tissues and tumor types.
- 28 • In cancer presence of C1q+ TAM often correlates with poor prognosis.
- 29 • Presence of C1q+ TAM correlates with T cell exhaustion in cancer and immune
30 tolerance induction in healthy tissue.
- 31 • C1q is the recognition molecule of the classical complement pathway, binding to
32 immune complexes, pentraxins or other activators in the tumor microenvironment.
- 33 • C1q directly controls macrophage phenotype by interacting with surface receptors.
- 34 • C1q directly controls T cell phenotype by internalization, binding to mitochondria and
35 regulation of mitochondrial metabolism.
- 36 • C1q is likely not only a biomarker of a TAM subpopulations but a driver of cancer
37 progression.

38

39 **Outstanding Questions Box**

- 40 • Which mechanism does C1q regulate macrophage and T cell phenotypes in normal
41 tissue and in tumor microenvironment? Does C1q act intracellularly in the producing
42 macrophages or in an autocrine manner, at the cell surface?
- 43 • What is the exact origin of C1q+ macrophages?
- 44 • Could C1q, produced by other cells such as fibroblasts, regulate macrophages and T
45 cells phenotype?
- 46 • What is the interplay between the complement cascade-mediated functions of C1q
47 and its functions outside the cascade?
- 48 • What parallels can we draw between the break in immune tolerance and
49 autoimmunity, driven by the congenital C1q deficiency and the immunosuppressive tumor
50 microenvironment of the tumors that is rich in C1q+ macrophages?
- 51 • Can tumor-promoting C1q+ macrophages turn into an ally during immunotherapy
52 with anti-checkpoint inhibitors?

53

54 **C1q+ TAM in the light of scRNAseq**

55 Searching for effective biomarkers to predict cancer progression is a holy grail in oncology. Big
56 data is useful in this respect, but it is still challenging to find new biomarkers or fish out the
57 most robust ones. Single cell RNA-seq can characterize the transcriptional state of individual
58 cell types and allows one to define rare populations, otherwise lost in the bulk RNA-seq or
59 undetectable by flow cytometry or CyTOF due to lack of prior knowledge of their existence.
60 For a long time, macrophages had been divided into M1 (proinflammatory) or M2 (anti-
61 inflammatory) populations. Such dichotomy seems too simplistic and outdated. Macrophages
62 appear as a continuous spectrum of phenotypes between these two extreme populations.
63 Thanks to single cell RNAseq analysis, scientists started to explore the macrophage universe,
64 and two molecules have emerged: TREM2, a marker more often expressed by tumor
65 infiltrating macrophages and associated with pro-tumorigenic actions¹, and FOLR2, a marker
66 that suggests tissue residency². Numerous studies are now detailing another marker that is
67 expressed on macrophages, Complement component 1q (C1q). C1Q+ macrophage
68 populations in both healthy and tumor tissues (Figure 1) have been observed for many years,
69 but a deeper understanding of its function is still lacking.

70 Recent studies suggest that C1q could be used as a marker of poor prognosis for various
71 cancers. Transcriptomic data³ and protein staining on tumor sections showed that a high
72 presence of C1q+ macrophages^{3,4,5} is associated with higher post-surgical recurrence in clear
73 cell renal cell carcinoma (ccRCC)⁵ as well as in hepatocellular carcinoma⁶, and breast cancer⁷.
74 In osteosarcoma, the expression of C1Q, mostly by macrophages, negatively correlates with
75 patient survival⁸. In pancreatic ductal adenocarcinoma (PDAC), C1q expression in primary
76 tumors and hepatic metastasis is higher compared to normal tissue, and the presence of
77 C1Q+M2-like macrophages is associated with worse prognosis⁹.

78

79 Despite its potential role as a biomarker, recent studies also suggest that C1q could drive
80 tumor progression. Here we discuss recent evidence that suggests C1q+ macrophages play a
81 major role in tumor immunity and cancer progression.

82

83 **C1q+ TAM – a bunch of gene correlations**

84 In healthy tissue, the C1Q+ macrophage population is characterized by the expression of
85 C1QA, C1QB, C1QC, HLA-DRB1 and MRC1 (gene coding for the CD206 molecule)¹⁰. Different
86 cancer types exhibit an increased C1Q+ Tumor Associated Macrophages (TAM) population
87 expressing CD206, HLA-DR, SEPP1 and FOLR2^{11,12,4}. C1Q+ macrophages also express APOE in
88 renal cancer³, breast cancer⁷, and liver metastasis from colorectal cancer¹³. Nevertheless,
89 there are exceptions. In renal cancer the C1Q+ TAM do not express FOLR2 but TREM2³, which
90 is known to be associated with immunosuppression and poor prognosis in several cancers.
91 Similarly, C1Q+ TAMs express TREM2 in liver metastases in patients and in mice with PDAC¹⁴.
92 The C1Q+ TAM population has a strict mutually exclusive relationship with SPP1+ TAM^{15,16}
93 (Figure 2). This dichotomy is nearly perfect, although some studies have shown that
94 C1Q+/SPP1+ associates with FOLR2+ and/or TREM2+ TAM^{17,7}. In the colon, C1Q+
95 macrophages can be found in both healthy and tumor tissue, whereas SPP1+ macrophages
96 are only found in tumor tissue¹². These macrophages can be deciphered by the transcriptional
97 factors they express: C1Q+ TAM express mostly MAF/MAFB, while SPP1+ TAM express
98 FOS/JUN for and CEBPB/ZEB2¹². Although these cells likely originate from the same precursor,
99 they have a very different evolutionary path (Box 1). C1q+ TAM may be linked to sex, as
100 females with non-small cell lung cancer present with a higher number of C1Q+ TAMs, whereas
101 male have higher number of SPP1+ macrpahages¹⁶.
102 The ensemble of these gene expression correlations (positive correlation with APOE, HLA-DR,
103 MRC1, FOLR2 or TREM2, and negative correlation with SPP1) draws a portrait of the C1Q+
104 macrophages as a distinct immunosuppressive population. Indeed, in melanoma and basal
105 cell skin carcinoma, C1Q+TAM are enriched in non-responders to immune checkpoint
106 therapy¹⁷, suggesting that they play a role in regulating anti-tumor immunity. As such, single
107 cell RNA sequencing (scRNA-seq) is unveiling the interplay between C1Q+ macrophages and
108 other immune cells that might be responsible for the poor overall survival in cancer patients
109 and lack of response to immunotherapies.

110

111 **C1q correlates with T cells exhaustion**

112

113 The presence of C1Q+ macrophages correlates with exhausted T cells, forming a dysfunctional
114 immune circuit in ccRCC³⁻⁵. In colorectal cancer, C1Q+ TAM interact with T cell subsets.
115 Analysis of ligand-receptors pairs revealed a significant enrichment of CXCL10-CXCR3 axis in

116 C1Q+ TAM, suggesting that production of CXCL10 by C1Q+ TAM binds to its receptor CXCR3,
117 which is mostly present at the T cell surface. This finding highlights the potential role of C1Q+
118 TAM in the recruitment and activation of the Th1 response¹² (Figure 3 point 1). In lung cancer
119 a similar increase in CXCL-10 was described in C1Q+ TAM, in association with an enrichment
120 of the transcription factors IRF1, IRF7 and STAT1¹⁶. IRF1 correlates with STAT1, HLA-DR, PD-1
121 and LAG-3 in metastases of colorectal cancer¹⁸. Moreover, in ccRCC tumors, C1q+ cell density
122 correlated with expression of inhibitory receptors PD-1 and LAG3 at the CD8+ T cells surface⁵,
123 and these C1Q+ macrophages express additional immune checkpoint ligands, such as PD-L1
124 and PDL-2^{4,5}. In cervical cancer, patients with C1Q+ TAM also express high levels of immune
125 checkpoint inhibitors including CD40L, CTLA4, LAG3, PD-1, and TIGIT¹⁹ (Figure 3 point 2).
126 Moreover, in mouse models of cancer, C1Q+ macrophages specifically express EB13, a subunit
127 of the IL-35 cytokine, which allows their cross talk with intratumoral T cells and leads to their
128 dysfunction when combined with the p35 subunit of IL-12²⁰. This gene was already described
129 as a promotor of CD8+ T cell exhaustion, when it is expressed by Tregs²¹. Interestingly, Tregs
130 are also found at a higher proportion within tumors with C1Q+ TAM as compared with normal
131 samples³.

132 The maturation of dendritic cells can impact on the expression levels of C1q, ultimately
133 affecting T cell function. Indeed, immature dendritic cells express large amounts of C1q, but
134 during dendritic cell maturation, which is driven by CXCL4, C1q gene is hyper-methylated and
135 its expression decreases²². A murine model of subcutaneous injection of various murine tumor
136 cell lines (lung cancer (LLC), colorectal cancer (MC38) or melanoma cancer (B16-F10)), showed
137 that reduced C1q methylation (i.e. high C1q) promotes CD8+ T cell dysfunction and tumor
138 progression²⁰. In lung cancer and idiopathic pulmonary fibrosis, the methylation status of C1q
139 decreases as compared to healthy tissue, leading to an increase in tumor associated C1Q
140 expression, which is associated with poor prognosis²³.

141 In cervical cancer, tumors with a gene signature of C1Q+ TAM are more infiltrated by immune
142 cells and express more immune-checkpoint markers than tumors with a gene signature of
143 SPP1+ TAM¹⁹. However, it is unclear which cells express these immune checkpoints. Compared
144 to SPP1+ TAM, C1Q+ TAM also express higher levels of HLA-DR¹², which could help C1Q+ TAM
145 interact with immune cells.

146 Taken together, these lines of evidence show a clear correlation between C1q+ macrophages
147 and the activation status of T cells, but is C1q a driver or a passenger in this process?

148

149

150 **C1q in TAM... So what?**

151 Are these C1q+ macrophages accompanying other factors responsible for poor prognosis or
152 driving recurrence? Is C1q only a “marker” of these macrophages or does it play a direct role
153 in their pro-tumoral effect? As C1q is a major factor involved in the complement system,
154 recent studies are unveiling the role of the complement cascade in regulating tumor
155 progression. However, C1q is a versatile molecule and has functions extending beyond the
156 borders of the complement cascade. Below we describe emerging mechanisms by which C1q
157 regulates tumor progression.

158

159

160 *C1q in the complement system*

161 C1q is the initiating protein of the complement cascade²⁴. Complement is a part of the innate
162 immune system and its best-known function is to defend the host against invading pathogens.
163 C1q can trigger the classical pathway of the cascade when it binds to immune complexes,
164 pentraxins or one of its other over 100 different ligands. The plasma source of the majority of
165 complement proteins is the liver, but C1q is an exception, being secreted by tissue-resident
166 macrophages²⁴. To function within the cascade, C1q needs to associate with two serine
167 proteases – C1r and C1s, which trigger the proteolytic cascade that results in the generation
168 of proinflammatory anaphylatoxins C3a and C5a, the membrane attack complex C5b-9 and
169 the opsonization of target cells. How could this cascade function in the context of cancer and
170 what is the role of C1q in this scenario?

171 Transcriptomic analysis across different tumor types has revealed that cells of the tumor
172 microenvironment or tumor cells themselves express components of the classical and
173 alternative complement pathways, including C1q (Figure 1)²⁴. It appears that their co-
174 regulated overexpression is context-dependent, and their prognostic value is either favorable
175 or poor or of undetermined significance in particular types of cancers. ccRCC falls in the
176 “aggressive complement” group, where overexpression of these genes correlates with worse
177 survival. *In situ* staining, scRNAseq analysis and *ex vivo* experiments revealed that the ccRCC
178 tumors have a complement-rich environment, where some tumor cells produce C1r, C1s, C4

179 and C3 but need macrophage-derived C1q to activate the classical pathway on intratumoral
180 IgG-containing immune complexes^{5,9} (Figure 3 point 3). While complement is activated on
181 most tumor cells and promotes cell death, it does not result in cell killing and even promotes
182 tumor progression in some cancer types. This finding can be explained by the low intratumoral
183 expression of the components of the terminal pathway³ and the limited formation of a
184 membrane attack complex due to expression of specific inhibitors at the tumor cell
185 surface^{5,9,25}. Thus, chronic inflammation mediated by C5a favors an immunosuppressive
186 microenvironment and facilitates T cells exhaustion^{3,4} (Figure 3 point 4). C1Q+ macrophages
187 also expressed ApoE - a protein able to bind C1q and to activate the complement system²⁶.
188 Therefore, C1Q+ positive macrophages can induce tumor progression by triggering the
189 complement cascade.

190

191 *C1q and neoangiogenesis*

192 C1q can interact directly with endothelial cells (EC) to promote neoangiogenesis, via still
193 unknown cell surface receptors or heparan sulfate^{27,5,28}. C1q deposits can be found at the
194 surface of EC in the absence of other complement factors such as C3 or C4. C1q can induce
195 adhesion, spreading and expression of adhesion molecules by directly binding EC²⁹. In
196 addition, in a lesioned area, EC start to express C1q, which induces EC permeability,
197 proliferation, migration and endothelial tube formation *in vitro*²⁸. *In vivo*, C1q^{-/-} mice show a
198 disordered vascular network in subcutaneously implanted tumors. Other *in vitro* studies also
199 indicate that the interaction of C1q with melanoma and PDAC cells promotes proliferation,
200 migration and invasion of the tumor cells^{9,27} (Figure 3 point 5).

201

202 *C1q, immune tolerance, and T cell exhaustion*

203 In physiology C1q regulates human macrophage polarization via interactions with LAIR1, as a
204 switch towards inflammation resolution to avoid autoimmunity³⁰⁻³² (Figure 3 point 6). Indeed,
205 complete C1q deficiency, although very rare, is the strongest genetic predisposing factor to
206 systemic autoimmunity³³. C1q opsonizes apoptotic cells, enhances their uptake by
207 macrophages and immature dendritic cells, modulates cytokine release, and promotes
208 immune tolerance³⁴⁻³⁷ (Figure 3 point 7). In this context macrophages and immature dendritic
209 cells produce and secrete C1q, which will act in an autocrine manner³⁸. In addition, C1Q+ cells

210 have been found to associate with tolerance in the fetal-maternal interface during pregnancy,
211 which is reminiscent of that seen in cancer. An HLA-DR^{high} group of cells, characterized by high
212 expression of C1Q, APOE, various genes of lipid metabolism, EB13, IDO 1 and 2 (inducers of
213 cell tolerance) and the immune-checkpoint PD-L1, limits T-cell expansion driven by fetal
214 alloantigen and establishes an immune-tolerance to fetal allotransplant³⁹. These processes
215 are not well studied in cancer, but it is tempting to speculate that uptake of C1q-opsonized
216 dying cancer cells may be perceived in a tolerogenic manner by the TAM, and that this C1q
217 will re-orient their phenotype to hamper the immune response against tumor neo-antigens.

218

219

220 *C1q and cell metabolism*

221 Another function of C1q that is unrelated to the complement cascade is to play as a rheostat
222 of the mitochondrial metabolism of CD8 T cells⁴⁰ (Figure 3 point 8). Extracellular C1q is
223 internalized by CD8+ T cells and is found at the surface of mitochondria. Intracellularly C1q is
224 involved in the upregulation of mitochondria biogenesis genes, leading to the differentiation
225 of CD8+ T cells into memory T cells and not effector cells. By C1q its presence at the
226 mitochondrial surface C1q dampens CD8+ T cell responses to self-antigens. Congenital
227 deficiency of C1q is rare in humans, but it results in enhanced CD8+ T cell responses, becoming
228 the strongest genetic predisposing factor for autoimmunity. C1q is not produced by T cells,
229 but it is internalized from the extracellular milieu. Therefore, it is tempting to speculate that
230 C1q, secreted by the C1Q+ macrophages in ccRCC downregulate the capacity of adjacent
231 intratumoral CD8 T cells to respond to stimulation, thereby contributing to their exhausted
232 phenotype. The intracellular role of C1q within macrophages has not been studied, but again,
233 we speculate that it may affect their metabolism, and functional orientation towards an
234 immunosuppressive phenotype, thereby inducing T cell exhaustion. In the context of
235 atherosclerosis, C1q can modulate the cytokine expression of macrophages while they digest
236 lipid proteins, leading to an M2-like polarization⁴¹. The MafB transcription factor, which is
237 present in C1Q+TAM in colorectal cancer¹², was also described to promote M2 polarization in
238 atherosclerosis⁴².

239 These results raise the question whether pro-tumoral M2-like macrophages start to express
240 C1q or does C1q allow the polarization of these cells into pro-tumoral macrophages?

241

242 **C1q in CAF, another actor of this story**

243 Like macrophages, recent studies have begun to distinguish different sub-types of fibroblasts
244 and especially cancer-associated fibroblasts (CAF). They notably, showed that in breast and
245 pancreatic cancer a subpopulation of CAF MHC-II⁺ led to an immunosuppressive tumor micro-
246 environment^{43,44}. In breast cancer, mesenchymal stem cells (MSC) can produce exosomes
247 containing TGF- β and C1q⁴⁵. Fibroblasts are similar to MSC and can be considered as old
248 MSC⁴⁶. The exosomes from MSC but not from tumor cells drive the polarization of Monocytic
249 myeloid derived suppressor cells (M-MDSC) into M2-macrophages overexpressing CD206, PD-
250 L1 and MHC-II⁺ (Figure 3 point 9). In contrast, in lung cancer, MHC-II⁺ CAF can activate the
251 TCR of CD4⁺ T cells and rescue them from apoptosis via the C1q/C1qbp axis. Indeed, these
252 CAFs have the ability to produce C1q that will be released and will interact with its receptor
253 C1qbp present at the surface of CD4⁺ T cells⁴⁷ (Figure 3 point 10). These recent studies suggest
254 that C1q excreted by CAF influence the tumor immune microenvironment, by both directly
255 acting on T cells and on macrophage polarization. More studies are necessary to determine
256 the role of CAF upstream of macrophage actions.

257

258

259 **Concluding remarks and future perspectives**

260 Analyses of scRNAseq data and the growing literature indicate that C1q⁺ TAM are key players
261 in the tumor microenvironment, but many questions remain unanswered (see Outstanding
262 Questions). Although formal proof is needed, it seems that C1q⁺ TAM are drivers of cancer
263 progression with a direct pro-tumoral effect in the absence of immunotherapy. Interestingly,
264 ccRCC tumors with mature tertiary lymphoid structures have IgG deposition on tumor cells
265 and respond better to anti-checkpoint inhibitors⁴⁸. The tertiary lymphoid structure signature
266 contained C1q genes and APOE, and the macrophages likely contributed to the elimination of
267 tumor cells and the mounting of an anti-tumoral immune response. Further studies are
268 needed to determine how C1q and complement contribute to this process. It is still poorly
269 understood if C1q acts in an autocrine manner, on the cell surface of the macrophages or
270 intracellularly or as deposits on tumor cells. It is also unclear which cells C1q impacts on and
271 how it controls immune activation vs tolerance and exhaustion. Understanding the
272 mechanisms of action of C1q in the modulation of macrophages phenotype in health and

273 disease and whether C1q can be harnessed as a therapeutic target in combination with anti-
274 cancer checkpoint inhibitors are perspectives for the future.

275 Beyond C1q, other complement proteins also impact on the TAM phenotype. Factor H
276 differentiates CD14+ human monocytes into immunosuppressive macrophages in the context
277 of breast but not renal cancer^{49,50}. C5aR1 is also overexpressed on TAM, which exhibit an M2-
278 like functional profile. C3aR deficiency is associated with reduced accumulation and functional
279 skewing of TAM, increased T cell activation and response to anti-PD-1 therapy⁵¹ in mouse
280 models of sarcoma. In models of squamous carcinogenesis, C5aR1 inhibition improves
281 chemotherapy efficacy by reprogramming macrophages to recruit cytotoxic CD8+ T cells⁵².
282 Similar effects were observed in a mouse model of renal cancer⁵³. Future studies should
283 address how other complement proteins and activation fragments, such as FH or C3a and C5a
284 impact on C1q+ TAM vs TAM lacking C1q expression.

285

286

287

288

289

290 **Acknowledgements**

291 This work was supported by grants from: Les Entreprises contre le Cancer, Ligue Regionale
292 Contre le Cancer (GEFLUC) IDF; Institut National Du Cancer INCa_16096; Agence Nationale de
293 la Recherche ANR-21-CE14-0066-02; Comité de Paris de la Ligue contre le cancer RS22/75-37
294 and l'Idex Sorbonne Université (Programmes Investissements d'Avenir Émergences 2021-
295 2022) to LTR. Equipe labellisée Ligue contre le Cancer, 2021-2 to CSF and WHF. This work is
296 also supported by The Labex Immuno-Oncology Excellence Program, INSERM, Université de
297 Paris and Sorbonne Université.

298

299 The authors declare no competing interests.

300

301 **References**

302 1. Molgora, M. *et al.* TREM2 Modulation Remodels the Tumor Myeloid Landscape
303 Enhancing Anti-PD-1 Immunotherapy. *Cell* **182**, 886-900.e17 (2020).

304 2. Samaniego, R. *et al.* Folate Receptor β (FR β) Expression in Tissue-Resident and

- 305 Tumor-Associated Macrophages Associates with and Depends on the Expression of PU.1.
306 *Cells* **9**, 1445 (2020).
- 307 3. Obradovic, A. *et al.* Single-cell protein activity analysis identifies recurrence-
308 associated renal tumor macrophages. *Cell* **184**, 2988-3005.e16 (2021).
- 309 4. Braun, D. A. *et al.* Progressive immune dysfunction with advancing disease stage in
310 renal cell carcinoma. *Cancer Cell* **39**, 632-648.e8 (2021).
- 311 5. Roumenina, L. T. *et al.* Tumor Cells Hijack Macrophage-Produced Complement C1q to
312 Promote Tumor Growth. *Cancer Immunol Res* (2019) doi:10.1158/2326-6066.CIR-18-0891.
- 313 6. Zhang, Q. *et al.* Landscape and Dynamics of Single Immune Cells in Hepatocellular
314 Carcinoma. *Cell* **179**, 829-845.e20 (2019).
- 315 7. Ramos, R. N. *et al.* Tissue-resident FOLR2+ macrophages associate with tumor-
316 infiltrating CD8+ T cells and with increased survival of breast cancer patients.
317 2021.04.12.439412 <https://www.biorxiv.org/content/10.1101/2021.04.12.439412v1> (2021)
318 doi:10.1101/2021.04.12.439412.
- 319 8. Huang, H. *et al.* <p>Prognostic Implications of the Complement Protein C1Q and Its
320 Correlation with Immune Infiltrates in Osteosarcoma</p>. *OTT* **14**, 1737–1751 (2021).
- 321 9. Yang, J. *et al.* Integrated genomic and transcriptomic analysis reveals unique
322 characteristics of hepatic metastases and pro-metastatic role of complement C1q in
323 pancreatic ductal adenocarcinoma. *Genome Biology* **22**, 4 (2021).
- 324 10. Mulder, K. *et al.* Cross-tissue single-cell landscape of human monocytes and
325 macrophages in health and disease. *Immunity* **54**, 1883-1900.e5 (2021).
- 326 11. Sharma, A. *et al.* Onco-fetal Reprogramming of Endothelial Cells Drives
327 Immunosuppressive Macrophages in Hepatocellular Carcinoma. *Cell* **183**, 377-394.e21
328 (2020).
- 329 12. Zhang, L. *et al.* Single-Cell Analyses Inform Mechanisms of Myeloid-Targeted
330 Therapies in Colon Cancer. *Cell* **181**, 442-459.e29 (2020).
- 331 13. Donadon, M. *et al.* Macrophage morphology correlates with single-cell diversity and
332 prognosis in colorectal liver metastasis. *J Exp Med* **217**, e20191847 (2020).
- 333 14. Kemp, S. B. *et al.* Pancreatic cancer is marked by complement-high blood monocytes
334 and tumor-associated macrophages. *Life Sci Alliance* **4**, e202000935 (2021).
- 335 15. Cheng, S. *et al.* A pan-cancer single-cell transcriptional atlas of tumor infiltrating
336 myeloid cells. *Cell* **184**, 792-809.e23 (2021).
- 337 16. Yang, Q. *et al.* Single-Cell RNA Sequencing Reveals the Heterogeneity of Tumor-
338 Associated Macrophage in Non-Small Cell Lung Cancer and Differences Between Sexes. *Front*
339 *Immunol* **12**, 756722 (2021).
- 340 17. Xiong, D., Wang, Y. & You, M. A gene expression signature of TREM2hi macrophages

- 341 and $\gamma\delta$ T cells predicts immunotherapy response. *Nat Commun* **11**, 5084 (2020).
- 342 18. Shao, Y. *et al.* IRF1-mediated immune cell infiltration is associated with metastasis in
343 colon adenocarcinoma. *Medicine* **99**, e22170 (2020).
- 344 19. Li, X., Zhang, Q., Chen, G. & Luo, D. Multi-Omics Analysis Showed the Clinical Value of
345 Gene Signatures of C1QC+ and SPP1+ TAMs in Cervical Cancer. *Front Immunol* **12**, 694801
346 (2021).
- 347 20. Dong, L. *et al.* The loss of RNA N6-adenosine methyltransferase Mettl14 in tumor-
348 associated macrophages promotes CD8+ T cell dysfunction and tumor growth. *Cancer Cell*
349 (2021) doi:10.1016/j.ccell.2021.04.016.
- 350 21. Sawant, D. V. *et al.* Adaptive plasticity of IL-10+ and IL-35+ Treg cells cooperatively
351 promotes tumor T cell exhaustion. *Nat Immunol* **20**, 724–735 (2019).
- 352 22. Silva-Cardoso, S. C. *et al.* CXCL4 suppresses tolerogenic immune signature of
353 monocyte-derived dendritic cells. *European Journal of Immunology* **50**, 1598–1601 (2020).
- 354 23. Kou, W. *et al.* High complement protein C1q levels in pulmonary fibrosis and non-
355 small cell lung cancer associated with poor prognosis. *BMC Cancer* **22**, 110 (2022).
- 356 24. Roumenina, L. T., Daugan, M. V., Petitprez, F., Sautès-Fridman, C. & Fridman, W. H.
357 Context-dependent roles of complement in cancer. *Nat Rev Cancer* **19**, 698–715 (2019).
- 358 25. Daugan, M. V. *et al.* Complement C1s and C4d as prognostic biomarkers in renal
359 cancer: emergence of non-canonical functions of C1s. *Cancer Immunol Res* (2021)
360 doi:10.1158/2326-6066.CIR-20-0532.
- 361 26. Yin, C. *et al.* ApoE attenuates unresolvable inflammation by complex formation with
362 activated C1q. *Nat Med* **25**, 496–506 (2019).
- 363 27. Bulla, R. *et al.* C1q acts in the tumour microenvironment as a cancer-promoting factor
364 independently of complement activation. *Nat Commun* **7**, 10346 (2016).
- 365 28. Bossi, F. *et al.* C1q as a unique player in angiogenesis with therapeutic implication in
366 wound healing. *Proc. Natl. Acad. Sci. U.S.A.* **111**, 4209–4214 (2014).
- 367 29. Feng, X., Tonnesen, M. G., Peerschke, E. I. B. & Ghebrehiwet, B. Cooperation of C1q
368 receptors and integrins in C1q-mediated endothelial cell adhesion and spreading. *J Immunol*
369 **168**, 2441–2448 (2002).
- 370 30. Son, M. *et al.* C1q and HMGB1 reciprocally regulate human macrophage polarization.
371 *Blood* **128**, 2218–2228 (2016).
- 372 31. Liu, T. *et al.* HMGB1–C1q complexes regulate macrophage function by switching
373 between leukotriene and specialized proresolving mediator biosynthesis. *PNAS* **116**, 23254–
374 23263 (2019).
- 375 32. Fouët, G. *et al.* Molecular Basis of Complement C1q Collagen-Like Region Interaction
376 with the Immunoglobulin-Like Receptor LAIR-1. *Int J Mol Sci* **22**, 5125 (2021).

- 377 33. Stegert, M., Bock, M. & Trendelenburg, M. Clinical presentation of human C1q
378 deficiency: How much of a lupus? *Molecular Immunology* **67**, 3–11 (2015).
- 379 34. Bohlsón, S. S., O’Conner, S. D., Hulsebus, H. J., Ho, M.-M. & Fraser, D. A. Complement,
380 c1q, and c1q-related molecules regulate macrophage polarization. *Front Immunol* **5**, 402
381 (2014).
- 382 35. Verneret, M. *et al.* Relative contribution of c1q and apoptotic cell-surface calreticulin
383 to macrophage phagocytosis. *J Innate Immun* **6**, 426–434 (2014).
- 384 36. Benoit, M. E., Clarke, E. V., Morgado, P., Fraser, D. A. & Tenner, A. J. Complement
385 Protein C1q Directs Macrophage Polarization and Limits Inflammasome Activity during the
386 Uptake of Apoptotic Cells. *The Journal of Immunology* **188**, 5682–5693 (2012).
- 387 37. Fraser, D. A., Laust, A. K., Nelson, E. L. & Tenner, A. J. C1q differentially modulates
388 phagocytosis and cytokine responses during ingestion of apoptotic cells by human
389 monocytes, macrophages, and dendritic cells. *J Immunol* **183**, 6175–6185 (2009).
- 390 38. Fraser, D. A., Arora, M., Bohlsón, S. S., Lozano, E. & Tenner, A. J. Generation of
391 inhibitory NFkappaB complexes and phosphorylated cAMP response element-binding
392 protein correlates with the anti-inflammatory activity of complement protein C1q in human
393 monocytes. *J Biol Chem* **282**, 7360–7367 (2007).
- 394 39. Costa, M. L. *et al.* Two Distinct Myeloid Subsets at the Term Human Fetal–Maternal
395 Interface. *Frontiers in Immunology* **8**, (2017).
- 396 40. Ling, G. S. *et al.* C1q restrains autoimmunity and viral infection by regulating CD8+ T
397 cell metabolism. *Science* **360**, 558–563 (2018).
- 398 41. Spivia, W., Magno, P. S., Le, P. & Fraser, D. A. Complement protein C1q promotes
399 macrophage anti-inflammatory M2-like polarization during the clearance of atherogenic
400 lipoproteins. *Inflamm. Res.* **63**, 885–893 (2014).
- 401 42. Kim, H. The transcription factor MafB promotes anti-inflammatory M2 polarization
402 and cholesterol efflux in macrophages. *Sci Rep* **7**, 7591 (2017).
- 403 43. Sebastian, A. *et al.* Single-Cell Transcriptomic Analysis of Tumor-Derived Fibroblasts
404 and Normal Tissue-Resident Fibroblasts Reveals Fibroblast Heterogeneity in Breast Cancer.
405 *Cancers* **12**, 1307 (2020).
- 406 44. Elyada, E. *et al.* Cross-Species Single-Cell Analysis of Pancreatic Ductal
407 Adenocarcinoma Reveals Antigen-Presenting Cancer-Associated Fibroblasts. *Cancer Discov* **9**,
408 1102–1123 (2019).
- 409 45. Biswas, S. *et al.* Exosomes Produced by Mesenchymal Stem Cells Drive Differentiation
410 of Myeloid Cells into Immunosuppressive M2-Polarized Macrophages in Breast Cancer. *J*
411 *Immunol* **203**, 3447–3460 (2019).
- 412 46. Soundararajan, M. & Kannan, S. Fibroblasts and mesenchymal stem cells: Two sides
413 of the same coin? *J Cell Physiol* **233**, 9099–9109 (2018).

- 414 47. Kerdidani, D. *et al.* Lung tumor MHCII immunity depends on in situ antigen
415 presentation by fibroblasts. *Journal of Experimental Medicine* **219**, e20210815 (2022).
- 416 48. Meylan, M *et al.* Tertiary lymphoid structures generate and propagate anti-tumor
417 antibody-producing plasma cells in renal cell cancer. *Immunity*, in press.
- 418 49. Daugan, M. V. *et al.* Intracellular Factor H Drives Tumor Progression Independently of
419 the Complement Cascade. *Cancer Immunol Res* (2021) doi:10.1158/2326-6066.CIR-20-0787.
- 420 50. Smolag, K. I. *et al.* Complement inhibitor factor H expressed by breast cancer cells
421 differentiates CD14+ human monocytes into immunosuppressive macrophages.
422 *Oncoimmunology* **9**, 1731135 (2020).
- 423 51. Magrini, E. *et al.* Complement activation promoted by the lectin pathway mediates
424 C3aR-dependent sarcoma progression and immunosuppression. *Nat Cancer* **2**, 218–232
425 (2021).
- 426 52. Medler, T. R. *et al.* Complement C5a Fosters Squamous Carcinogenesis and Limits T
427 Cell Response to Chemotherapy. *Cancer Cell* **34**, 561-578.e6 (2018).
- 428 53. Reese, B. *et al.* Complement as Prognostic Biomarker and Potential Therapeutic
429 Target in Renal Cell Carcinoma. *J Immunol* **205**, 3218–3229 (2020).
- 430 54. Zimmerman, K. A. *et al.* Single-Cell RNA Sequencing Identifies Candidate Renal
431 Resident Macrophage Gene Expression Signatures across Species. *J Am Soc Nephrol* **30**, 767–
432 781 (2019).
- 433 55. Zilionis, R. *et al.* Single-Cell Transcriptomics of Human and Mouse Lung Cancers
434 Reveals Conserved Myeloid Populations across Individuals and Species. *Immunity* **50**, 1317-
435 1334.e10 (2019).
- 436 56. Kim, N. *et al.* Single-cell RNA sequencing demonstrates the molecular and cellular
437 reprogramming of metastatic lung adenocarcinoma. *Nat Commun* **11**, 2285 (2020).
- 438 57. Finlay, C. M. *et al.* *Genotype and Th2 cells control monocyte to tissue resident*
439 *macrophage differentiation during nematode infection of the pleural cavity.*
440 2021.12.17.472661 <https://www.biorxiv.org/content/10.1101/2021.12.17.472661v1> (2021)
441 doi:10.1101/2021.12.17.472661.
- 442 58. Niyonzima, N. *et al.* Mitochondrial C5aR1 activity in macrophages controls IL-1 β
443 production underlying sterile inflammation. *Sci Immunol* **6**, eabf2489 (2021).
- 444 59. Sun, D. *et al.* TISCH: a comprehensive web resource enabling interactive single-cell
445 transcriptome visualization of tumor microenvironment. *Nucleic Acids Research* **49**, D1420–
446 D1430 (2021).

447

448

449 **Box 1. C1q TAM = tissue resident macrophages?**

450 The origins of C1Q+ TAM is not well understood, but a Single Cell RNAseq study across species
451 (mouse, rat, pig and human) found a conserved gene signature of kidney resident
452 macrophages composed of CD81, CD74 (coding for HLA-DR-gamma), APOE and C1QC⁵⁴. A
453 similar study in lung found conserved genes between human and mice macrophages,
454 including APOE, MRC1, C1QA, C1QB and C1QC⁵⁵. These data agree with the ontogenic study
455 of multiple tumor types, describing a high similarity between C1Q+ and FOLR2+ macrophages.
456 FOLR2+ is associated with an embryonic origin of macrophages^{9,10,15}. In healthy lung tissue,
457 two types of macrophages can be distinguished, alveolar or interstitial. A sub-type of alveolar
458 macrophages expresses both C1Q and APOE⁵⁶. C1Q+ macrophages are larger in size and are
459 identified as resident in patients with colorectal cancer liver metastasis¹³ and in the peritoneal
460 cavity of mice⁵⁷. These large C1Q+ TAM confer poor prognosis¹³, and they have features of
461 foamy cells, overexpress genes of cholesterol metabolism, scavenger receptors, as well as
462 C1QA and C1QB.

463 C1Q+ TAM express, in a conserved way in multiple cancer types, HLA-DR, APOE and MRC1,
464 suggesting an embryonic, tissue resident origin. However, a recent study showed that C1Q
465 and APOE are commonly expressed by TAM in breast cancer⁷ and resident TAM express
466 FOLR2+, while infiltrating TAM express TREM2+, opening new questions on the origin of the
467 C1q+ TAM in particular contexts.

468 A transcriptional trajectory study using the example of colorectal cancer¹² shows that C1Q+
469 TAM and SPP1+ TAM populations could arise from a common precursor: CD14-expressing
470 monocytes, which differentiate towards FCN1+ monocyte-like cells and different macrophage
471 populations. One of them are the SPP1+, while the other overexpresses IL-1 β + and gives rise
472 to the sub-population of C1q+ TAM. In healthy tissue, C1q regulates the macrophages
473 polarization during the uptake of apoptotic cells by inhibiting NLRP3 gene expression, that
474 suppresses the IL-1 β cleavage³⁶. It is interesting to note that the FCN1 is defense collagen
475 and a close relative to C1q, called ficolin 1, activator of the lectin complement pathway²⁴.
476 Another complement protein, C5aR1, regulate the IL-1 β production un macrophages⁵⁸.
477 Whether a cross-talk exist between these complement proteins/pathways in macrophages,
478 being they TRM or TAM, is still unknown.

479 Finally, future work will show whether there is a difference in the C1q+ macrophages from
480 normal tissue (the ones that generate the plasma C1q pool,
481 <https://www.proteinatlas.org/ENSG00000173372-C1QA/celltype>) and the ones in tumors.

482

483

484 **Figure legends**

485

486

487 **Figure 1:** Heatmap representing the expression of C1QB gene in different immune or non-
488 immune cells depending on the cancer type. Data is retrieved from the Tumor Immune Single-
489 cell Hub (TISCH) scRNA-seq database⁵⁹. In light yellow are indicated positions for which data
490 is not available, in blue low gene expression of C1QB and in red high gene expression of C1QB.
491 Similar data were obtained for C1QA and C1QC.

492

493

494 **Figure 2:** Single Cell Analysis from the TISCH database. Gene expression of SPP1, C1QA, C1QB
495 and C1QC of Colorectal Carcinoma tumor (accession number GSE146771). On the left, the cell
496 characterization of the Single Cell RNAseq analysis. The upper panel is a large view of the gene
497 expression. The lower panel is a zoom in on the Monocytes/Macrophages population.

498

499

500 **Figure 3:** Proposed mechanisms by which C1q+ TAM drive cancer progression. C1q is produced
501 by sub-populations of TAM and TRM. 1) The C1Q+ TAM secrete the chemokine CXCL-10 that
502 binds to its receptor CXCR3 at the surface of CD8+ and CD4+ T cells, especially Th1 T cells. This
503 binding will activate and recruit T cells inside the tumor. 2) By its expression of immune
504 checkpoint inhibitors, C1Q+ TAM interact with T cell immune checkpoints, favoring T cell
505 exhaustion. 3) C1q impacts on tumor progression by activating the complement cascade in
506 the extracellular space, which will generate the anaphylatoxins C3a and C5a, and 4) promote
507 chronic inflammation. 5) The C1q molecule can interact directly with endothelial cells to
508 promote neo-angiogenesis needed for the tumor growth. 6) In healthy tissue, C1q is produced
509 by TRM and immature dendritic cells (imDC) to maintain homeostasis and opsonize apoptotic
510 cells. 7) The C1q molecule can also act in an autocrine way on macrophages, by interacting
511 with LAIR1 to regulate their polarization. 8) CD8+ T cells can internalize C1q, which will interact
512 with mitochondria to control the CD8 metabolism. 9) The secretion of C1q-containing
513 exosomes by fibroblasts leads to M2 polarization of macrophages. 10) MHC-II+ cancer

514 associated fibroblasts also produce C1q, which binds to its receptor C1qbp at the surface of
515 CD4+ T cells to activate and to rescue them from apoptosis. Figure generated with
516 biorender.com



