



HAL
open science

On fractal properties of Weierstrass-type functions

Claire David

► **To cite this version:**

Claire David. On fractal properties of Weierstrass-type functions. Proceedings of the International Geometry Center, 2019, 12 (2), 10.15673/tmgc.v12i2.1485 . hal-03698959

HAL Id: hal-03698959

<https://hal.sorbonne-universite.fr/hal-03698959v1>

Submitted on 17 Jun 2023

HAL is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers.

L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.

On fractal properties of Weierstrass-type functions

Claire David

Abstract. In the sequel, starting from the classical Weierstrass function defined, for any real number x , by $\mathcal{W}(x) = \sum_{n=0}^{+\infty} \lambda^n \cos(2\pi N_b^n x)$, where λ and N_b are two real numbers such that $0 < \lambda < 1$, $N_b \in \mathbb{N}$ and $\lambda N_b > 1$, we highlight intrinsic properties of curious maps which happen to constitute a new class of iterated function system. Those properties are all the more interesting, in so far as they can be directly linked to the computation of the box dimension of the curve, and to the proof of the non-differentiability of Weierstrass type functions.

Extended abstract

The aim of this paper is to generalize previous results on the classical Weierstrass function and its curve. It happens that this curve can be obtained as the limit of prefractals, i.e. a sequence of graphs, obtained by means of an iterated function system, which do not happen to be contraction maps as it is usually the case. They instead bear what can be viewed as an equivalent property, since, at each step of the iterative process that enable one to obtain the prefractals, they reduce the two-dimensional Lebesgue measures of a given sequence of rectangles covering the curve. Such maps play a part in the first step of the horseshoe map process introduced by Stephen Smale. Those maps are all the more interesting, in so far as they can be directly linked to the computation of the box dimension of the curve, and to the proof of the non-differentiability of the Weierstrass function. The very interesting point is thus that by considering such maps,

2010 Mathematics Subject Classification: 37F20-28A80-05C63

Keywords: Weierstrass function ; non-differentiability ; iterative function systems.

one can build a broader class of applications everywhere continuous, though nowhere differentiable, functions, as we will expose it in the sequel.

INTRODUCTION

In his seminal paper of 1981, J. E. Hutchinson [Hut81] introduces, for the first time, what will be later qualified of “iterated function system” (I.F.S.), as a finite set of contraction maps, each defined on a compact metric set K of the euclidean space \mathbb{R}^d , $d \in \mathbb{N}^*$:

$$\mathcal{S} = \{T_1, \dots, T_N\} \quad , \quad N \in \mathbb{N}^*$$

where \mathbb{N}^* denotes the set of strictly positive integers, such that

$$K = \bigcup_{i=1}^N T_i(K)$$

The compact set K is then said to be “invariant” with respect to the set \mathcal{S} (one often refer to this result as the “Gluing Lemma”).

A prequel occurrence of such maps, under the form of similitudes, can already be found in the Mandelbrot books of 1977 [Man77b], [Man77a].

Hutchinson’s novelty is to consider not the compact K itself, but the set \mathcal{S} , which arises naturally, in so far as the invariant compact K is fully determined by the set \mathcal{S} , and, interestingly, is, also, the limit of a sequence of pre-fractal graphs that can be built, in an iterative way, thanks to the maps that constitute the set \mathcal{S} .

Following this work, iterated function systems were taken up and even more developed by M. F. Barnsley et al. [BD85], as “a unified way of generating and classifying a broad class of fractals”. As explained by the authors, fractals were “traditionally viewed as being produced by a process of successive microscopic refinement taken to the limit”, which, of course, makes sense with the geometric representation one may have of fractal sets, since, when looking at smaller and smaller scales, one finds, again and again, the same form. Of course, at stake are specific and classical types of fractals, as Sierpiński gaskets, dragon curves, Cantor sets, Julia curves, ... For M. F. Barnsley and S. Demko, those fractals are to be seen as the attractors of iterated function systems, which, of course, joins the approach of J. E. Hutchinson.

M. F. Barnsley and S. Demko place themselves in a probabilistic approach. Given still a compact metric space K , the related Banach space $C(K)$ of real-valued functions defined on K , with respect to the norm

$$f \in C(K) \mapsto \|f\|_\infty = \max \{|f(x)|, x \in K\}$$

and a finite collection

$$w = \{w_1, \dots, w_N\} \quad , \quad N \in \mathbb{N}^*$$

of Borel measurable functions from K to K , they define the set $\{K, w\}$ as an iterated function system if and only if there exists an associated set of positive real numbers

$$\{p_1, \dots, p_N\} \quad , \quad \forall i \in \{1, \dots, N\} : p_i > 0 \quad , \quad \sum_{i=1}^N p_i = 1$$

such that the operator T on $C(K)$, given, for any f of $C(K)$, by

$$\forall x \in K : \quad T(f)(x) = \sum_{i=1}^N p_i (f \circ w_i)(x)$$

has the property:

$$T(C(K)) \subset C(K) \cdot$$

Treating w as a set-valued function, through

$$\forall x \in K : \quad w(x) = \{w_1(x), \dots, w_N(x)\}$$

they then naturally introduce, for the i.f.s. $\{K, w\}$, and a given x of K , the related attractor

$$\mathcal{A}(x) = \lim_{n \rightarrow +\infty} w^{\circ n}(x)$$

in the sense:

$$\lim_{n \rightarrow +\infty} \|w^{\circ n}(x) - \mathcal{A}(x)\|_\infty = 0 \cdot$$

Classical fractal sets as, for instance, the Sierpiński Gasket, fit this definition.

In our previous work on the Weierstrass curve [Dav18], which, as exposed, for instance, by A. S. Besicovitch and H. D. Ursell [BU37], or, a few years later, by B. Mandelbrot [Man77b], bears fractal properties, we showed that the curve could be obtained by means of a sequence a graphs $(\Gamma_{\mathcal{W}_m})_{m \in \mathbb{N}}$,

that approximate the studied one. This is done using a family of non-linear C^∞ maps from \mathbb{R}^2 to \mathbb{R}^2 , which happen not to be contractions, in the aforementioned classical sense. The nonlinearity does not enable one to resort to the probabilistic approach of M. F. Barnsley and S. Demko, since there does not exist a constant associated set of probabilities. Yet, even if they are not contractions, our maps bear what can be viewed as an equivalent property, since, at each step of the iterative process, they reduce the two-dimensional Lebesgue measures of a given sequence of rectangles covering the curve. This is due to the fact that they correspond, in a sense, to the composition of a contraction of ratio r_x in the horizontal direction, and a dilatation of factor r_y in the vertical one, with

$$r_x r_y < 1$$

Such maps are considered in the book of Robert L. Devaney [Dev03], where they play a part in the first step of the horseshoe map process introduced by Stephen Smale.

The Weierstrass curve is invariant with respect to the set of those maps, which makes it possible to dispose of an equivalent result of the Gluing Lemma. But what deserves to be enlightened, in our case, is that the intrinsic properties of those curious maps make them all the more interesting, in so far as they can be directly linked to the computation of the box dimension of the curve, and to the proof of the non-differentiability of the Weierstrass function, as shown in [Dav19]. All the more is the generalization to a broader class of applications that could, then, enable one to build everywhere continuous, though nowhere differentiable, functions, as we will expose it in the sequel.

1. THE CASE OF THE WEIERSTRASS FUNCTION

Notation. In the following, λ and b are two real numbers such that:

$$0 < \lambda < 1 \quad , \quad b = N_b \in \mathbb{N} \quad \text{and} \quad \lambda N_b > 1.$$

We deliberately made the choice to introduce the notation N_b which replaces the initial b , in so far as, to the origins, b is any real number, whereas we deal with the specific case of a natural integer that we consequently choose to denote by N_b , as an echo to the initial b .

The Weierstrass function, introduced in 1875 by K. Weierstrass [Wei75], known as one of these so-called pathological mathematical objects, continuous everywhere, while nowhere differentiable, is the sum of the uniformly convergent trigonometric series, defined, for any real number x , by:

$$\mathscr{W}(x) = \sum_{n=0}^{+\infty} \lambda^n \cos(2\pi N_b^n x) .$$

Definition 1.1. Weierstrass Curve

We will call *Weierstrass Curve* the restriction to $[0, 1[\times \mathbb{R}$, of the graph of the Weierstrass function, and denote it by $\Gamma_{\mathscr{W}}$.

1.2. Theoretical study. We place ourselves, in the following, in the euclidian plane of dimension 2, referred to a direct orthonormal frame. The usual Cartesian coordinates are (x, y) .

Property 1. Periodic properties of the Weierstrass function

For any real number x :

$$\mathscr{W}(x+1) = \sum_{n=0}^{+\infty} \lambda^n \cos(2\pi N_b^n x + 2\pi N_b^n) = \sum_{n=0}^{+\infty} \lambda^n \cos(2\pi N_b^n x) = \mathscr{W}(x) .$$

The study of the Weierstrass function can be restricted to the interval $[0, 1[$.

By following the method developed by J. Kigami [Kig89], we approximate the restriction $\Gamma_{\mathscr{W}}$ to $[0, 1[\times \mathbb{R}$, of the Weierstrass Curve, by a sequence of graphs, built through an iterative process. To this purpose, we introduce the iterated function system of the family of C^∞ maps from \mathbb{R}^2 to \mathbb{R}^2 :

$$\{T_0, \dots, T_{N_b-1}\}$$

where, for any integer i belonging to $\{0, \dots, N_b - 1\}$, and any (x, y) of \mathbb{R}^2 :

$$T_i(x, y) = \left(\frac{x+i}{N_b}, \lambda y + \cos\left(2\pi \left(\frac{x+i}{N_b}\right)\right) \right) .$$

Property 2.

$$\Gamma_{\mathscr{W}} = \bigcup_{i=0}^{N_b-1} T_i(\Gamma_{\mathscr{W}}).$$

Proof. We refer to [Dav18].

□

Definition 1.3. Word, on the graph $\Gamma_{\mathscr{W}}$

Let m be a strictly positive integer. We will call **number-letter** any integer \mathscr{M}_i of $\{0, \dots, N_b - 1\}$, and **word of length** $|\mathscr{M}| = m$, on the graph $\Gamma_{\mathscr{W}}$, any set of number-letters of the form:

$$\mathscr{M} = (\mathscr{M}_1, \dots, \mathscr{M}_m).$$

We will write:

$$T_{\mathscr{M}} = T_{\mathscr{M}_1} \circ \dots \circ T_{\mathscr{M}_m}.$$

Definition 1.4. For any integer i belonging to $\{0, \dots, N_b - 1\}$, let us denote by:

$$P_i = (x_i, y_i) = \left(\frac{i}{N_b - 1}, \frac{1}{1 - \lambda} \cos \left(\frac{2\pi i}{N_b - 1} \right) \right)$$

the fixed point of the map T_i .

We will denote by V_0 the ordered set (according to increasing abscissa), of the points:

$$\{P_0, \dots, P_{N_b-1}\}$$

since, for any i of $\{0, \dots, N_b - 2\}$:

$$x_i \leq x_{i+1}.$$

The set of points V_0 , where, for any i of $\{0, \dots, N_b - 2\}$, the point P_i is linked to the point P_{i+1} , constitutes an oriented graph (according to increasing abscissa), that we will denote by $\Gamma_{\mathscr{W}_0}$. V_0 is called the set of vertices

of the graph $\Gamma_{\mathcal{W}_0}$.

For any natural integer m , we set:

$$V_m = \bigcup_{i=0}^{N_b-1} T_i(V_{m-1}).$$

The set of points V_m , where two consecutive points are linked, is an oriented graph (according to increasing abscissa), which we will denote by $\Gamma_{\mathcal{W}_m}$. V_m is called the set of vertices of the graph $\Gamma_{\mathcal{W}_m}$. We will denote, in the following, by $\mathcal{N}_m^{\mathcal{S}}$ the number of vertices of the graph $\Gamma_{\mathcal{W}_m}$, and we will write:

$$V_m = \left\{ \mathcal{S}_0^m, \mathcal{S}_1^m, \dots, \mathcal{S}_{\mathcal{N}_m^{\mathcal{S}}-1}^m \right\}.$$

Property 3. For any natural integer m :

$$V_m \subset V_{m+1}.$$

Property 4. For any integer i belonging to $\{0, \dots, N_b - 2\}$:

$$T_i(P_{N_b-1}) = T_{i+1}(P_0).$$




FIGURE 1.1. The fixed points P_0, P_1, P_2 , and the graph $\Gamma_{\mathscr{W}_0}$, in the case where $\lambda = \frac{1}{2}$, and $N_b = 3$.

Definition 1.5. Vertices of the graph $\Gamma_{\mathscr{W}}$

Two points X and Y of $\Gamma_{\mathscr{W}}$ will be called *vertices* of the graph $\Gamma_{\mathscr{W}}$ if there exists a natural integer m such that:

$$(X, Y) \in V_m^2$$




FIGURE 1.2. The graph $\Gamma_{\mathscr{W}_1}$, in the case where $\lambda = \frac{1}{2}$, and $N_b = 3$. $T_0(P_2) = T_1(P_0)$ et $T_1(P_2) = T_2(P_1)$.

Definition 1.6. Consecutive vertices on the graph $\Gamma_{\mathscr{W}}$

Two points X and Y of $\Gamma_{\mathscr{W}}$ will be called *consecutive vertices* of the graph $\Gamma_{\mathscr{W}}$ if there exist a natural integer m , and an integer j of $\{0, \dots, N_b - 2\}$, such that:

$$\begin{cases} X = (T_{i_1} \circ \dots \circ T_{i_m})(P_j) \\ Y = (T_{i_1} \circ \dots \circ T_{i_m})(P_{j+1}) \end{cases} \quad \{i_1, \dots, i_m\} \in \{0, \dots, N_b - 1\}^m$$

or:

$$X = (T_{i_1} \circ T_{i_2} \circ \dots \circ T_{i_m})(P_{N_b-1}) \quad \text{and} \quad Y = (T_{i_1+1} \circ T_{i_2} \circ \dots \circ T_{i_m})(P_0).$$

Property 5. The set $\bigcup_{m \in \mathbb{N}} V_m$ is dense in $\Gamma_{\mathscr{W}}$.




FIGURE 1.3. The graphs $\Gamma_{\mathscr{W}_0}$ (in green), $\Gamma_{\mathscr{W}_1}$ (in red), $\Gamma_{\mathscr{W}_2}$ (in orange), $\Gamma_{\mathscr{W}}$ (in cyan), in the case where $\lambda = \frac{1}{2}$, and $N_b = 3$.

Definition 1.7. Edge relation, on the graph $\Gamma_{\mathscr{W}}$

Given a natural integer m , two points X and Y of $\Gamma_{\mathscr{W}_m}$ will be called **adjacent** if and only if X and Y are two consecutive vertices of $\Gamma_{\mathscr{W}_m}$. We will write:

$$X \underset{m}{\sim} Y$$

This edge relation ensures the existence of a word $\mathcal{M} = (\mathcal{M}_1, \dots, \mathcal{M}_m)$ of length m , such that X and Y both belong to the iterate:

$$T_{\mathcal{M}} V_0 = (T_{\mathcal{M}_1} \circ \dots \circ T_{\mathcal{M}_m}) V_0$$

Given two points X and Y of the graph $\Gamma_{\mathscr{W}}$, we will say that X and Y are **adjacent** if and only if there exists a natural integer m such that:

$$X \underset{m}{\sim} Y$$

Proposition 1.8. Addresses, on the Weierstrass Curve

Given a strictly positive integer m , and a word $\mathcal{M} = (\mathcal{M}_1, \dots, \mathcal{M}_m)$ of length $m \in \mathbb{N}^*$, on the graph $\Gamma_{\mathcal{M}_m}$, for any integer j of $\{1, \dots, N_b - 2\}$, any $X = T_{\mathcal{M}}(P_j)$ of $V_m \setminus V_0$, i.e. distinct from one of the N_b fixed point P_i , $0 \leq i \leq N_b - 1$, has exactly two adjacent vertices, given by:

$$T_{\mathcal{M}}(P_{j+1}) \quad \text{and} \quad T_{\mathcal{M}}(P_{j-1})$$

where:

$$T_{\mathcal{M}} = T_{\mathcal{M}_1} \circ \dots \circ T_{\mathcal{M}_m}$$

By convention, the adjacent vertices of $T_{\mathcal{M}}(P_0)$ are $T_{\mathcal{M}}(P_1)$ and $T_{\mathcal{M}}(P_{N_b-1})$, those of $T_{\mathcal{M}}(P_{N_b-1})$, $T_{\mathcal{M}}(P_{N_b-2})$ and $T_{\mathcal{M}}(P_0)$.


Notation.

For any integer j belonging to $\{0, \dots, N_b - 1\}$, any natural integer m , and any word \mathcal{M} of length m , we set:

$$T_{\mathcal{M}}(P_j) = (x(T_{\mathcal{M}}(P_j)), y(T_{\mathcal{M}}(P_j)))$$

$$L_m = x(T_{\mathcal{M}}(P_{j+1})) - x(T_{\mathcal{M}}(P_j)) = \frac{1}{(N_b - 1) N_b^m}$$

$$h_{j,m} = y(T_{\mathcal{M}}(P_{j+1})) - y(T_{\mathcal{M}}(P_j)).$$



Notation. We will denote by:

$$D_{\mathcal{M}} = 2 + \frac{\ln \lambda}{\ln N_b}$$

the Hausdorff dimension of $\Gamma_{\mathcal{M}}$ (see [BBR14], [Kel17]).

Theorem 1.9. *An upper bound and a lower bound, for the box-dimension of the Weierstrass Curve [Dav18]*

For any integer j belonging to $\{0, 1, \dots, N_b - 2\}$, each natural integer m , and each word \mathcal{M}_m of length m , let us consider the rectangle $\mathcal{R}_{j,m,\mathcal{M}_m}$, whose sides are parallel to the horizontal and vertical axes, of width:

$$L_m = x(T_{\mathcal{M}_m}(P_{j+1})) - x(T_{\mathcal{M}_m}(P_j)) = \frac{1}{(N_b - 1) N_b^m}$$

and height $|h_{j,m}|$, such that the points $T_{\mathcal{M}_m}(P_j)$ and $T_{\mathcal{M}_m}(P_{j+1})$ are two vertices of this rectangle.

We set:

$$\eta_{\mathcal{W}} = 2\pi^2 \left\{ \frac{(2N_b - 1)\lambda(N_b^2 - 1)}{(N_b - 1)^2(1 - \lambda)(\lambda N_b^2 - 1)} + \frac{2N_b}{(\lambda N_b^2 - 1)(\lambda N_b^3 - 1)} \right\}.$$

$$C_1(N_b) = \begin{cases} (N_b - 1)^{2-D_{\mathcal{W}}} \left\{ \frac{2}{1-\lambda} \sin\left(\frac{\pi}{N_b-1}\right) \min_{0 \leq j \leq N_b-1} \left| \sin\left(\frac{\pi(2j+1)}{N_b-1}\right) \right| - \frac{2\pi}{N_b(N_b-1)} \frac{1}{\lambda N_b-1} \right\} & \text{if } N_b \text{ is odd} \\ (N_b - 1)^{2-D_{\mathcal{W}}} \max \left\{ \frac{2}{1-\lambda} \sin\left(\frac{\pi}{N_b-1}\right) \min_{0 \leq j \leq N_b-1} \left| \sin\left(\frac{\pi(2j+1)}{N_b-1}\right) \right| - \frac{2\pi}{N_b(N_b-1)} \frac{1}{\lambda N_b-1}, \frac{4}{N_b^2} \frac{1-N_b^{-2}}{N_b^2-1} \right\} & \text{if } N_b \text{ is even} \end{cases}$$

and:


$$C_2(N_b) = \eta_{\mathcal{W}}(N_b - 1)^{2-D_{\mathcal{W}}}.$$

Then:

$$C_1(N_b) L_m^{2-D_{\mathcal{W}}} \leq |h_{j,m}| \leq C_2(N_b) L_m^{2-D_{\mathcal{W}}}.$$

Notation. Given a natural integer m , we set:

$$h_m = L_m^{2-D_{\mathcal{W}}} = \frac{N_b^{(D_{\mathcal{W}}-2)m}}{(N_b - 1)^{2-D_{\mathcal{W}}}}$$



$$h_{jm} \leq h_m$$

Corollary 1.10. (Of Theorem 1.9)

For any natural integer m , any integer j belonging to $\{0, 1, \dots, N_b - 2\}$, and each word \mathcal{M}_{m+1} of length $m+1$, the two-dimensional Lebesgue measure

$$\mu_{\mathcal{L}}(\mathcal{R}_{j,m+1,\mathcal{M}_{m+1}})$$

of the rectangle $\mathcal{R}_{j,m+1,\mathcal{M}_{m+1}}$, is such that, for any integer k belonging to $\{0, 1, \dots, N_b - 2\}$, any integer ℓ belonging to $\{0, 1, \dots, N_b - 2\}$, and each word \mathcal{M}_m of length m :

$$\mu_{\mathcal{L}}(\mathcal{R}_{j,m+1,\mathcal{M}_{m+1}}) < \mu_{\mathcal{L}}(\mathcal{R}_{\ell,m,\mathcal{M}_m})$$

Proof. Given a natural integer m , j in $\{0, 1, \dots, N_b - 2\}$, and a word \mathcal{M}_{m+1} of length $m + 1$, the two-dimensional Lebesgue measure of the rectangle $\mathcal{R}_{j,m+1,\mathcal{M}_{m+1}}$ can be obtained thanks to the values of the cartesian coordinates of the consecutive vertices $T_{\mathcal{M}_{m+1}}(P_j)$ and $T_{\mathcal{M}_{m+1}}(P_{j+1})$:

$$\begin{aligned} & \mu_{\mathcal{L}}(\mathcal{R}_{j,m+1,\mathcal{M}_{m+1}}) \\ &= \left\{ x(T_{\mathcal{M}_{m+1}}(P_{j+1})) - x(T_{\mathcal{M}_{m+1}}(P_j)) \right\} \times \left| y(T_{\mathcal{M}_{m+1}}(P_{j+1})) - y(T_{\mathcal{M}_{m+1}}(P_j)) \right|. \end{aligned}$$

One may then write:

$$T_{\mathcal{M}_{m+1}} = T_k \circ T_{\mathcal{M}_m} \quad , \quad k \in \{0, 1, \dots, N_b - 1\}$$

where \mathcal{M}_m is a word of length m .

Thus, due to:

$$\begin{cases} y(T_{\mathcal{M}_{m+1}}(P_{j+1})) &= \lambda y(T_{\mathcal{M}_m}(P_{j+1})) + \cos\left(2\pi \left(\frac{x(T_{\mathcal{M}_{m+1}}(P_{j+1})) + k}{N_b}\right)\right) \\ y(T_{\mathcal{M}_{m+1}}(P_j)) &= \lambda y(T_{\mathcal{M}_m}(P_j)) + \cos\left(2\pi \left(\frac{x(T_{\mathcal{M}_{m+1}}(P_j)) + k}{N_b}\right)\right) \end{cases}$$

and:

$$x(T_{\mathcal{M}_m}(P_{j+1})) - x(T_{\mathcal{M}_m}(P_j)) = L_m \leq |y(T_{\mathcal{M}_m}(P_{j+1})) - y(T_{\mathcal{M}_m}(P_j))|$$

one has:

$$\begin{aligned}
 |y(T_{\mathcal{M}_{m+1}}(P_{j+1})) - y(T_{\mathcal{M}_{m+1}}(P_j))| &\leq \lambda |y(T_{\mathcal{M}_m}(P_{j+1})) - y(T_{\mathcal{M}_m}(P_j))| \\
 &\quad + \frac{2\pi}{N_b} |x(T_{\mathcal{M}_m}(P_{j+1})) - x(T_{\mathcal{M}_m}(P_j))| \\
 &\leq \lambda |y(T_{\mathcal{M}_m}(P_{j+1})) - y(T_{\mathcal{M}_m}(P_j))| + \frac{2\pi}{N_b} L_m \\
 &\leq \left(\lambda + \frac{2\pi}{N_b} \right) |y(T_{\mathcal{M}_m}(P_{j+1})) - y(T_{\mathcal{M}_m}(P_j))|
 \end{aligned}$$

which yields:

$$\begin{aligned}
 \mu_{\mathcal{L}}(\mathcal{R}_{j,m+1}, \mathcal{M}_{m+1}) &= \frac{L_m}{N_b} \times |y(T_{\mathcal{M}_{m+1}}(P_{j+1})) - y(T_{\mathcal{M}_{m+1}}(P_j))| \\
 &\leq \frac{L_m}{N_b} \times \left\{ \lambda |y(T_{\mathcal{M}_m}(P_{j+1})) - y(T_{\mathcal{M}_m}(P_j))| + \frac{2\pi}{N_b} L_m \right\} \\
 &\leq \frac{L_m}{N_b} \times \left(\lambda + \frac{2\pi}{N_b} \right) |y(T_{\mathcal{M}_m}(P_{j+1})) - y(T_{\mathcal{M}_m}(P_j))|
 \end{aligned}$$

Due to the symmetric roles played by the integers j and ℓ , one has just to prove the result for $j = \ell$.

Since:

$$\mu_{\mathcal{L}}(\mathcal{R}_{j,m}, \mathcal{M}_m) = L_m \times |y(T_{\mathcal{M}_m}(P_{j+1})) - y(T_{\mathcal{M}_m}(P_j))|$$

and, due to $N_b \geq 3$:

$$\frac{1}{N_b} \left(\lambda + \frac{2\pi}{N_b} \right) - 1 = \frac{1}{N_b^2} \left\{ \underbrace{\lambda N_b}_{< 1} + 2\pi - N_b^2 \right\} < 0$$

which yield the expected result. □

2. A SPECIFIC CLASS OF I.F.S.

Weierstrass-type functions have been previously studied, but under the Hausdorff dimension point of view. One may refer, for instance, to the study by B. R. Hunt [Hun98], where the author considers functions defined, for any real number x , by:

$$W_{\Theta}(x) = \sum_{n=0}^{+\infty} a_n g(b_n x + \theta_n)$$

where $\sum a_n$ is a positive and convergent series, $(b_n)_{n \in \mathbb{N}}$ a positive and increasing sequence, $\Theta = (\theta_n)_{n \in \mathbb{N}}$ a uniformly distributed sequence of numbers each belonging to $[0, 1]$, and playing the part of arbitrary phases, g being a Lipschitz and 1-periodic function.

In the case where the following assumptions are satisfied:

i. there exist two strictly positive real numbers ρ and σ such that:

$$1 < \rho < \sigma \quad , \quad \forall n \in \mathbb{N} : \quad \rho b_n \leq b_{n+1} \leq \sigma b_n$$

ii. there exists D in $]1, 2[$ such that:

$$\lim_{n \rightarrow +\infty} \frac{\ln a_n}{\ln b_n} = D - 2$$

iii. there exist a positive integer p , a strictly positive real constant M , a constant ℓ in $]0, 1[$, such that for all δ in $\left[\frac{\ell}{\sigma^p}, \ell\right]$, and for any real number x chosen randomly according to a uniform distribution on $[0, 1]$, the density function of:

$$x \mapsto g(x + \delta) - g(x)$$

has a $L^{\frac{p}{p-1}}$ norm at most equal to M ;

B. R. Hunt [Hun98] shows that for almost every Θ in $[0, 1]^{\infty}$, the graph of W_{Θ} has Hausdorff dimension D . It happens that in the case of such functions, the Hausdorff dimension is equal to the box-dimension.

Yet, as concerns the lower bound estimate required to obtain the explicit value of the Hausdorff/box dimension, the author calls for strictly positive constants K and K' which, as in existing earlier works, are not given explicitly (see, in the Hunt study, section 3., page 798). Moreover, no relation is made with the non-differentiability of such functions.

One may also note that such functions cannot be described by means of a finite iterated function systems, which does not allow any use of the Gluing Lemma.

In addition, the fact that the author considers, very generally, Lipschitz functions g is not specifically justified. It is all the more interesting as evoked in the above since, if the functions g were contractant ones, one falls back more easily on classical configurations. In fact, one may just consider the limit case of functions satisfying a Lipschitz condition with a Lipschitz constant of value 1.

What seemed of interest to us was to generalize our results to, indeed, a class of Weierstrass-type functions, but defined through an iterated function system which would bear analogous properties of the maps T_i , $0 \leq i \leq N_b - 1$. First, the box-dimension can be obtained rather simply, without calling for theoretical background in dynamic systems theory, just by applying a similar method as in [Dav18]. Then, one can also simply prove the non-differentiability of such functions, as in [?].

Notation. In the sequel:

- i.* N is a strictly positive integer, greater than 2 ;
- ii.* T and M are strictly positive real numbers ;
- iii.* $(\alpha_i)_{0 \leq i \leq N-1} \in \{0, \dots, N-1\}^N$ and $(\beta_i)_{0 \leq i \leq N-1} \in \{0, \dots, N-1\}^N$ are ordered sets of positive integers:

$$\forall i \in \{0, \dots, N-2\} : \alpha_i \leq \alpha_{i+1} \quad , \quad \beta_i \leq \beta_{i+1}$$

- iv.* ψ is a T -periodic, bounded function from \mathbb{R} to \mathbb{R} satisfying a Lipschitz condition ;
- v.* r_y is a real number such that:

$$0 < r_y < 1 \quad , \quad r_y N > 1 .$$

- vi.* We set:

$$r_x = \frac{1}{N} .$$

- vii.* $\{\phi_0, \dots, \phi_{N-1}\}$ and $\{\varphi_0, \dots, \varphi_{N-1}\}$ are sets of affine contractive maps from \mathbb{R} to \mathbb{R} , of respective ratios r_x and r_y , defined, for any integer i of $\{0, \dots, N-1\}$, and for any real number x :

$$\phi_i(x) = r_x (x + \alpha_i) \quad , \quad \varphi_i(x) = r_y (x + \beta_i) .$$

- viii.* We denote by $\{\psi_0, \dots, \psi_{N-1}\}$ the set of maps from \mathbb{R} to \mathbb{R} such that, for any integer i of $\{0, \dots, N-1\}$:

$$\psi_i = \psi \circ \phi_i .$$

Notation. We introduce the set of maps from \mathbb{R}^2 to \mathbb{R}^2

$$\{\tilde{T}_0, \dots, \tilde{T}_{N-1}\}$$

such that, for any integer i of $\{0, \dots, N-1\}$, and any (x, y) of \mathbb{R}^2 :

$$\tilde{T}_i(x, y) = (\phi_i(x), \varphi_i(y) + \psi_i(x)) \cdot$$

Definition 2.1. \mathscr{W} -type function

We introduce the \mathscr{W} -type function, defined, for any real number x , by:

$$\tilde{\mathscr{W}}(x) = \sum_{n=0}^{+\infty} r_y^n \psi(T N^n x).$$

Property 6. for any real number x , the series:

$$\tilde{\mathscr{W}}(x) = \sum_{n=0}^{+\infty} r_y^n \psi(T N^n x).$$

is convergent

Proof. One may simply note that, for any real number x :

$$|r_y^n \psi(T N^n x)| \lesssim r_y^n \sup_{t \in \mathbb{R}} |\psi(t)|$$

which yields the expected result, since $\sum_{n=0}^{+\infty} r_y^n$ is a geometric convergent series. □

Definition 2.2. \mathscr{W} -type Curve

We will call \mathscr{W} -type Curve the restriction to $[0, T[\times \mathbb{R}$, of the graph of the \mathscr{W} -type function, and denote it by $\Gamma_{\widetilde{\mathscr{W}}}$.

2.3. Theoretical study. We place ourselves, in the following, in the euclidian plane of dimension 2, referred to a direct orthonormal frame. The usual Cartesian coordinates are (x, y) .

Property 7. For any integer i of $\{0, \dots, N - 1\}$, the map \widetilde{T}_i admits a fixed point, that we will denote by \widetilde{P}_i :

$$\widetilde{P}_i = \left(\frac{\alpha_i}{N - 1}, \frac{\beta_i}{1 - r_y} + \frac{1}{1 - r_y} \psi_i \left(\frac{\alpha_i}{N - 1} \right) \right).$$

Lemma 2.4. For any integer i belonging to $\{0, \dots, N - 1\}$, the map T_i is a bijection of the Weierstrass-type Curve on \mathbb{R} .

Proof. Let us consider $i \in \{0, \dots, N\}$, a point $(y, \mathscr{W}(y))$ of $\Gamma_{\widetilde{\mathscr{W}}}$, and let us look for a real number x such that:

$$T_i \left(x, \widetilde{\mathscr{W}}(x) \right) = \left(y, \mathscr{W}(y) \right).$$

One has:

$$y = \phi_i(x) = r_x (x + \alpha_i)$$

which yields:

$$x = r_x^{-1} y - \alpha_i.$$

This enables one to obtain:

$$\widetilde{\mathscr{W}}(x) = \mathscr{W}(r_x^{-1} y - \alpha_i) = \sum_{n=0}^{+\infty} r_y^n \psi(T N^{n+1} y - T \alpha_i N^n i) = \sum_{n=0}^{+\infty} r_y^n \psi(T N^{n+1} y)$$

due to the T -periodicity of the function ψ , which leads to:

$$\psi (T N^{n+1} y - T \alpha_i N^n i) = \psi (T N^{n+1} y)$$

since α_i , N and i are integers.

Also:

$$\begin{aligned} \widetilde{T}_i \left(x, \widetilde{\mathcal{W}}(x) \right) &= \left(\phi_i \left(r_x^{-1} y - \alpha_i \right), \varphi_i \left(\widetilde{\mathcal{W}}(x) \right) + \psi_i \left(r_x^{-1} y - \alpha_i \right) \right) \\ &= \left(y, r_y \sum_{n=0}^{+\infty} r_y^n \psi \left(T N^{n+1} y \right) + \psi \left(T y \right) \right) \\ &= \left(y, \sum_{n=0}^{+\infty} r_y^{n+1} \psi \left(T N^{n+1} y \right) + \psi \left(T y \right) \right) \\ &= \left(y, \sum_{n=0}^{+\infty} r_y^n \psi \left(T N^n y \right) \right) \\ &= \left(y, \widetilde{\mathcal{W}}(y) \right). \end{aligned}$$

Les ϕ_i doivent être de coeff $1/N$ There exists thus a unique real number x such that:

$$T_i \left(x, \widetilde{\mathcal{W}}(x) \right) = \left(y, \widetilde{\mathcal{W}}(y) \right).$$

□

Theorem 2.5. *An upper bound and a lower bound, for the box-dimension of the Weierstrass-type Curve*

For any integer j belonging to $\{0, 1, \dots, N-2\}$, each natural integer m , and each word \mathcal{M} of length m , let us consider the rectangle, whose sides are parallel to the horizontal and vertical axes, of width:

$$L_m = x \left(T_{\mathcal{M}} \left(P_{j+1} \right) \right) - x \left(T_{\mathcal{M}} \left(P_j \right) \right) = r_x^m$$

and height $|h_{j,m}|$, such that the points $T_{\mathcal{M}} \left(P_j \right)$ and $T_{\mathcal{M}} \left(P_{j+1} \right)$ are two vertices of this rectangle.

We set:

$$C_1(N) = \left\{ \frac{1}{1-r_y} \min_{0 \leq j \leq N-1} \left\{ (\beta_{i+1} - \beta_i) + \left\{ \psi_{i+1} \left(\frac{\alpha_{i+1}}{N-1} \right) - \psi_i \left(\frac{\alpha_i}{N-1} \right) \right\} \right\} - \frac{r_x}{r_y} \frac{1}{1-\frac{r_x}{r_y}} \frac{\alpha_{j+1} - \alpha_j}{N-1} \right\}$$

and:

$$C_2(N) = \frac{|\beta_{j+1} - \beta_j|}{1 - r_y} + \frac{1}{1 - r_y} \left| \frac{\alpha_{j+1}}{N-1} - \frac{\alpha_j}{N-1} \right| + \frac{|\alpha_{j+1} - \alpha_j|}{N(r_y - r_x)}.$$

If:

$$C_1(N) \geq 0$$

one has:

$$C_1(N) L_m^{2-D_{\tilde{\mathcal{W}}}} \leq |h_{j,m}| \leq C_2(N) L_m^{2-D_{\tilde{\mathcal{W}}}}.$$

where:

$$D_{\tilde{\mathcal{W}}} = 2 + \frac{\ln r_y}{\ln N}.$$

which yields the fractal character of the Weierstrass-type Curve, the box-dimension of which is then $D_{\tilde{\mathcal{W}}}$.

Proof. The proof is obtained as in [Dav18]. It is based on the fact that, given a strictly positive integer m , and two points X and Y of V_m such that:

$$X \underset{m}{\sim} Y$$

there exists a word \mathcal{M} of length $|\mathcal{M}| = m$, on the graph $\Gamma_{\tilde{\mathcal{W}}}$, and an integer j of $\{0, \dots, N-2\}^2$, such that:

$$X = \tilde{T}_{\mathcal{M}}(P_j) \quad , \quad Y = \tilde{T}_{\mathcal{M}}(P_{j+1}).$$

By writing $\tilde{T}_{\mathcal{M}}$ under the form:

$$\tilde{T}_{\mathcal{M}} = \tilde{T}_{i_m} \circ \tilde{T}_{i_{m-1}} \circ \dots \circ \tilde{T}_{i_1}$$

where $(i_1, \dots, i_m) \in \{0, \dots, N-1\}^m$, one gets:

$$x\left(\tilde{T}_{\mathcal{M}}(P_j)\right) = r_x^N x_j + \sum_{k=1}^m r_x^k \alpha_k \quad , \quad x\left(\tilde{T}_{\mathcal{M}}(P_{j+1})\right) = r_x^N x_{j+1} + \sum_{k=1}^m r_x^k \alpha_k$$

and:

$$\begin{cases} y\left(\tilde{T}_{\mathcal{M}}(P_j)\right) &= r_y^m y_j + \sum_{k=1}^m r_y^{m-k} \psi_k\left(r_x^k x_j + \sum_{\ell=0}^k r_x^\ell \alpha_{m-\ell}\right) \\ y\left(\tilde{T}_{\mathcal{M}}(P_{j+1})\right) &= r_y^m y_{j+1} + \sum_{k=1}^m r_y^{m-k} \psi_k\left(r_x^k x_{j+1} + \sum_{\ell=0}^k r_x^\ell \alpha_{m-\ell}\right) \end{cases}.$$

This leads to:

$$h_{j,m} - r_y^m (y_{j+1} - y_j) = \sum_{k=1}^m r_y^{m-k} \left\{ \psi_{i_k}\left(r_x^k x_{j+1} + \sum_{\ell=0}^k r_x^\ell \alpha_{m-\ell}\right) - \psi_{i_k}\left(r_x^k x_j + \sum_{\ell=0}^k r_x^\ell \alpha_{m-\ell}\right) \right\}$$

where:

$$r_y^m (y_{j+1} - y_j) = r_y^m \frac{\beta_{j+1} - \beta_j}{1 - r_y} + \frac{r_y^m}{1 - r_y} \left\{ \psi_{j+1}\left(\frac{\alpha_{j+1}}{N-1}\right) - \psi_j\left(\frac{\alpha_j}{N-1}\right) \right\}$$

Since the maps ψ_{i_k} , $1 \leq k \leq m$, satisfy a Lipschitz condition, with a Lipschitz constant equal to 1, one has thus:

$$\begin{aligned} |y(T_{\mathcal{M}}(P_j)) - y(T_{\mathcal{M}}(P_{j+1})) - r_y^m (y_{j+1} - y_j)| &\leq \sum_{k=1}^m r_y^{m-k} r_x^k |x_{j+1} - x_j| \\ &= \sum_{k=1}^m r_y^{m-k} r_x^k \frac{|\alpha_{j+1} - \alpha_j|}{N-1} \\ &= r_y^m \frac{r_x}{r_y} \frac{1 - \frac{r_x^m}{r_y^m}}{1 - \frac{r_x}{r_y}} \frac{|\alpha_{j+1} - \alpha_j|}{N-1} \\ &\leq r_y^m \frac{r_x}{r_y} \frac{1}{1 - \frac{r_x}{r_y}} \frac{|\alpha_{j+1} - \alpha_j|}{N-1} \end{aligned}$$

which leads to:

$$\begin{aligned} y(T_{\mathcal{M}}(P_j)) - y(T_{\mathcal{M}}(P_{j+1})) &\geq r_y^m \frac{\beta_{j+1} - \beta_j}{1 - r_y} + \frac{r_y^m}{1 - r_y} \left\{ \psi_{i+1}\left(\frac{\alpha_{i+1}}{N-1}\right) - \psi_i\left(\frac{\alpha_i}{N-1}\right) \right\} \\ &\quad - r_x \frac{r_y^m - r_x^m}{r_y - r_x} \frac{|\alpha_{j+1} - \alpha_j|}{N-1} \end{aligned}$$

If:

$$\frac{1}{1-r_y} \min_{0 \leq j \leq N-1} \left\{ (\beta_{j+1} - \beta_j) + \left\{ \psi_{j+1} \left(\frac{\alpha_{j+1}}{N-1} \right) - \psi_j \left(\frac{\alpha_j}{N-1} \right) \right\} \right\} - \frac{r_x}{r_y} \frac{1}{1-\frac{r_x}{r_y}} \frac{|\alpha_{j+1} - \alpha_j|}{N-1} \geq 0$$

due to the symmetric roles played by $T_{\mathcal{M}}(P_j)$ and $T_{\mathcal{M}}(P_{j+1})$, one may only consider the case when:

$$\begin{aligned} y(T_{\mathcal{M}}(P_j)) - y(T_{\mathcal{M}}(P_{j+1})) &\geq r_y^m \frac{\beta_{j+1} - \beta_j}{1-r_y} + \frac{r_y^m}{1-r_y} \left\{ \psi_{i+1} \left(\frac{\alpha_{i+1}}{N-1} \right) - \psi_i \left(\frac{\alpha_i}{N-1} \right) \right\} \\ &\quad - r_y^m \frac{r_x}{r_y} \frac{1}{1-\frac{r_x}{r_y}} \frac{|\alpha_{j+1} - \alpha_j|}{N-1} \\ &\geq 0 \end{aligned}$$

which yields:

$$y(T_{\mathcal{M}}(P_j)) - y(T_{\mathcal{M}}(P_{j+1})) \geq r_y^m \left\{ \frac{1}{1-r_y} \min_{0 \leq j \leq N-1} \left\{ (\beta_{i+1} - \beta_i) + \left\{ \psi_{i+1} \left(\frac{\alpha_{i+1}}{N-1} \right) - \psi_i \left(\frac{\alpha_i}{N-1} \right) \right\} \right\} \right. \\ \left. - \frac{r_x}{r_y} \frac{1}{1-\frac{r_x}{r_y}} \frac{|\alpha_{j+1} - \alpha_j|}{N-1} \right\}$$

The predominant term is thus:

$$r_y^m = e^{m(D_{\mathcal{M}} - 2) \ln N} = N^{m(D_{\mathcal{M}} - 2)} = L_m^{2-D_{\mathcal{M}}} (N-1)^{2-D_{\mathcal{M}}}$$

One also has:

$$\begin{aligned}
|h_{j,m}| &\leq r_y^m |y_{j+1} - y_j| \\
&+ \sum_{k=1}^m r_y^{m-k} \left| \psi_{i_k} \left(r_x^k x_{j+1} + \sum_{\ell=0}^k r_x^\ell \alpha_{m-\ell} \right) - \psi_{i_k} \left(r_x^k x_j + \sum_{\ell=0}^k r_x^\ell \alpha_{m-\ell} \right) \right| \\
&\leq r_y^m \frac{|\beta_{j+1} - \beta_j|}{1 - r_y} + \frac{r_y^m}{1 - r_y} \left| \psi_{j+1} \left(\frac{\alpha_{j+1}}{N-1} \right) - \psi_j \left(\frac{\alpha_j}{N-1} \right) \right| \\
&+ \sum_{k=1}^m r_y^{m-k} \left| \psi_{i_k} \left(r_x^k x_{j+1} + \sum_{\ell=0}^k r_x^\ell \alpha_{m-\ell} \right) - \psi_{i_k} \left(r_x^k x_j + \sum_{\ell=0}^k r_x^\ell \alpha_{m-\ell} \right) \right| \\
&\leq r_y^m \frac{|\beta_{j+1} - \beta_j|}{1 - r_y} + \frac{r_y^m}{1 - r_y} \left| \frac{\alpha_{j+1}}{N-1} - \frac{\alpha_j}{N-1} \right| + \sum_{k=1}^m r_y^{m-k} r_x^k |x_{j+1} - x_j| \\
&= r_y^m \frac{|\beta_{j+1} - \beta_j|}{1 - r_y} + \frac{r_y^m}{1 - r_y} \left| \frac{\alpha_{j+1}}{N-1} - \frac{\alpha_j}{N-1} \right| + |x_{j+1} - x_j| r_y^m \frac{r_x}{r_y} \frac{1 - \frac{r_x^m}{r_y^m}}{1 - \frac{r_x}{r_y}} \\
&= r_y^m \frac{|\beta_{j+1} - \beta_j|}{1 - r_y} + \frac{r_y^m}{1 - r_y} \left| \frac{\alpha_{j+1}}{N-1} - \frac{\alpha_j}{N-1} \right| + |x_{j+1} - x_j| r_x \frac{r_y^m - r_x^m}{r_y - r_x} \\
&\leq r_y^m \frac{|\beta_{j+1} - \beta_j|}{1 - r_y} + \frac{r_y^m}{1 - r_y} \left| \frac{\alpha_{j+1}}{N-1} - \frac{\alpha_j}{N-1} \right| + |x_{j+1} - x_j| r_x \frac{r_y^m}{r_y - r_x} \\
&\leq r_y^m \frac{|\beta_{j+1} - \beta_j|}{1 - r_y} + \frac{r_y^m}{1 - r_y} \left| \frac{\alpha_{j+1}}{N-1} - \frac{\alpha_j}{N-1} \right| + r_y^m \frac{|\alpha_{j+1} - \alpha_j|}{N(r_y - r_x)}
\end{aligned}$$

Since:

$$D_{\mathcal{W}} \approx 2 + \frac{\ln r_y}{\ln N}, \quad r_y = e^{(D_{\mathcal{W}} - 2) \ln N} = N^{(D_{\mathcal{W}} - 2)}$$

one has thus:

$$|h_{j,m}| \leq r_y^m \left\{ \frac{|\beta_{j+1} - \beta_j|}{1 - r_y} + \frac{1}{1 - r_y} \left| \frac{\alpha_{j+1}}{N-1} - \frac{\alpha_j}{N-1} \right| + \frac{|\alpha_{j+1} - \alpha_j|}{N(r_y - r_x)} \right\}$$

□

Corollary 2.6. (of Theorem 1.9)

The \mathcal{W} -type functions are non-differentiable.

Proof. One has simply to use the analogous density property as in 5.

Given a natural integer m , and two points $X = (x, \widetilde{\mathcal{W}}(x))$, $Y = (y, \widetilde{\mathcal{W}}(y))$ of the pre-fractal graph $\Gamma_{\widetilde{\mathcal{W}}_m} \subset \Gamma_{\widetilde{\mathcal{W}}}$ such that:

$$x \leq y \quad , \quad X \underset{m}{\sim} Y$$

one may write:

$$X = \widetilde{T}_{\mathcal{M}_{m,j}}(P_k) \quad , \quad Y = (x + L_m, \mathcal{W}(x + L_m)) = \widetilde{T}_{\mathcal{M}_{m,j}}(P_{k+1})$$

where $\mathcal{M}_{m,j}$, $0 \leq j \leq N^m - 1$ is a word of length m , while k denotes an integer of the set $\{0, \dots, N - 2\}$.

One may note that:

$$\left| x \left(\widetilde{T}_{\mathcal{M}_{m,j}}(P_k) \right) - x \left(\widetilde{T}_{\mathcal{M}_{m,j}}(P_{k+1}) \right) \right| = \frac{1}{N^m} = L_m \xrightarrow{m \rightarrow +\infty} 0$$

Thus:

$$\begin{aligned} \left| \widetilde{\mathcal{W}} \left(x \left(\widetilde{T}_{\mathcal{M}_{m,j}}(P_k) \right) \right) - \widetilde{\mathcal{W}} \left(x \left(\widetilde{T}_{\mathcal{M}_{m,j}}(P_{k+1}) \right) \right) \right| &\geq C_1(N) L_m^{2-D_{\widetilde{\mathcal{W}}}} \\ &= C_1(N) \left| x \left(\widetilde{T}_{\mathcal{M}_{m,j}}(P_k) \right) - x \left(\widetilde{T}_{\mathcal{M}_{m,j}}(P_{k+1}) \right) \right|^{2-D_{\widetilde{\mathcal{W}}}} \end{aligned}$$

which leads to:

$$\begin{aligned} \left| \frac{\widetilde{\mathcal{W}} \left(x \left(\widetilde{T}_{\mathcal{M}_{m,j}}(P_k) \right) \right) - \widetilde{\mathcal{W}} \left(x \left(\widetilde{T}_{\mathcal{M}_{m,j}}(P_{k+1}) \right) \right)}{x \left(\widetilde{T}_{\mathcal{M}_{m,j}}(P_k) \right) - x \left(\widetilde{T}_{\mathcal{M}_{m,j}}(P_{k+1}) \right)} \right| &\geq C_1(N) \left| x \left(\widetilde{T}_{\mathcal{M}_{m,j}}(P_k) \right) - x \left(\widetilde{T}_{\mathcal{M}_{m,j}}(P_{k+1}) \right) \right|^{1-D_{\widetilde{\mathcal{W}}}} \\ &= C_1(N) L_m^{1-D_{\widetilde{\mathcal{W}}}} \end{aligned}$$

where

$$1 - D_{\widetilde{\mathcal{W}}} = -1 - \frac{\ln r_y}{\ln N} = -\frac{\ln(r_y N)}{\ln N} < 0$$

By passing to the limit when the integer m tends towards infinity, one gets the non-differentiability expected result:

$$\lim_{m \rightarrow +\infty} \left| \frac{\widetilde{\mathcal{W}}(x + L_m) - \widetilde{\mathcal{W}}(x)}{L_m} \right| = +\infty.$$

where:

$$\lim_{m \rightarrow +\infty} L_m = 0$$

□

REFERENCES

- [BBR14] K. Barański, B. Bárány, and J. Romanowska. On the dimension of the graph of the classical Weierstrass function. *Advances in Math.*, 265:791–800, 2014.
- [BD85] M. F. Barnsley and S. Demko. Iterated function systems and the global construction of fractals. *The Proceedings of the Royal Society of London*, A(399):243–275, 1985.
- [BU37] A. S. Besicovitch and H. D. Ursell. Sets of Fractional dimensions (v): on dimensional numbers of some continuous curves. *J. London Math. Soc.*, 12(1):18–25, 1937.
- [Dav18] Cl. David. Bypassing dynamical systems : A simple way to get the box-counting dimension of the graph of the Weierstrass function. *Proceedings of the International Geometry Center*, 11(2):1–16, 2018.
- [Dav19] Cl. David. Wandering across the Weierstrass function, while revisiting its properties, to appear. 2019.
- [Dev03] R. Devaney. *An Introduction To Chaotic Dynamical Systems*. Westview Press, 2003.
- [Hun98] B. Hunt. The Hausdorff dimension of graphs of Weierstrass functions. *Proc. Amer. Math. Soc.*, 12(1):791–800, 1998.
- [Hut81] J. E. Hutchinson. Fractals and self similarity. *Indiana University Mathematics Journal*, 30:713–747, 1981.
- [Kel17] G. Keller. A simpler proof for the dimension of the graph of the classical Weierstrass function. *Ann. Inst. Poincaré*, 53(1):169–181, 2017.
- [Kig89] J. Kigami. A harmonic calculus on the Sierpiński spaces. *Japan J. Appl. Math.*, 8:259–290, 1989.
- [Man77a] B. B. Mandelbrot. *The fractal geometry of nature*. 1977.
- [Man77b] B. B. Mandelbrot. *Fractals: form, chance, and dimension*. San Francisco: Freeman, 1977.
- [Wei75] K. Weierstrass. Über continuirliche Funktionen eines reellen Arguments, die für keinen Werth des letzteren einen bestimmten Differentialquotienten besitzen. *Journal fr die reine und angewandte Mathematik*, 79:29–31, 1875.