



HAL
open science

Emerging RNA-Dependent RNA Polymerase Mutation in a Remdesivir-Treated B-cell Immunodeficient Patient With Protracted Coronavirus Disease 2019

Martin Martinot, Aude Jary, Samira Fafi-Kremer, Valentin Leducq, Heloise Delagreverie, Marc Garnier, Jérôme Pacanowski, Arsène Mékinian, France Pirenne, Pierre Tiberghien, et al.

► To cite this version:

Martin Martinot, Aude Jary, Samira Fafi-Kremer, Valentin Leducq, Heloise Delagreverie, et al.. Emerging RNA-Dependent RNA Polymerase Mutation in a Remdesivir-Treated B-cell Immunodeficient Patient With Protracted Coronavirus Disease 2019. *Clinical Infectious Diseases*, 2021, 73 (7), pp.e1762-e1765. 10.1093/cid/ciaa1474 . hal-03704027

HAL Id: hal-03704027

<https://hal.sorbonne-universite.fr/hal-03704027v1>

Submitted on 6 Sep 2024

HAL is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers.

L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.

Remdesivir failure with SARS-CoV-2 RNA-dependent RNA-polymerase mutation in a B-cell immunodeficient patient with protracted Covid-19

Martin Martinot¹, Aude Jary², Samira Fafi-Kremer³, Valentin Leducq², Heloise Delagreverie³, Marc Garnier⁴, Jérôme Pacanowski⁵, Arsène Mékinian⁶, France Pirenne^{7,8}, Pierre Tiberghien^{7,9}, Vincent Calvez², Catherine Humbrecht¹⁰, Anne-Geneviève Marcelin², Karine Lacombe¹¹

1 Infectious diseases Department, Hôpitaux Civils de Colmar, Colmar, France

2 Sorbonne Université, INSERM, Institut Pierre Louis d'Epidémiologie et de Santé Publique (iPLESP), AP-HP, Pitié-Salpêtrière Hospital, Department of Virology, Paris, France

3 Virology Laboratory, Strasbourg University Hospital, Strasbourg, France; Strasbourg University, INSERM, IRM UMR-S 1109, F-67000 Strasbourg, France

4 Sorbonne Université, GRC 29, APHP, DMU DREAM, Anesthesiology and Intensive Care Department, Saint-Antoine Hospital - Paris, France

5 Infectious diseases Department, Saint-Antoine Hospital, APHP – Sorbonne Université, Paris – France

6 Sorbonne Université, Internal Medicine Department, Inflammation-Immunopathology-Biotherapy Department (DMU i3D), Saint-Antoine Hospital, AP-HP, Paris - France

7 Etablissement Français du Sang, La Plaine St-Denis - France.

8 Institut Mondor de Recherche Biomédicale, Unité 955, Equipe 2: Transfusion et Maladies du Globule Rouge, INSERM, EFS, Université Paris-Est Créteil, Créteil - France.

9 UMR 1098 RIGHT Inserm Université de Franche-Comté Etablissement Français du Sang, Besançon

10 Etablissement Français du sang Grand Est, Strasbourg, France

11 Sorbonne Université, Inserm IPLESP, Infectious Diseases Department, Saint-Antoine Hospital, APHP, Paris – France

Corresponding author

Martin Martinot

Infectious Diseases Department

Hôpitaux Civils de Colmar

39 avenue de la liberté 68024 Colmar cedex, France

Phone No: +33389124904

Fax No: +33389124691

Email Address: martin.martinot@ch-colmar.fr

Abstract

SARS-CoV-2 is a new pandemic virus for which Remdesivir is the only antiviral available. We report the occurrence of a mutation in the RdRP (D484Y) following failure of remdesivir in a 76-year-old woman with a post-rituximab B-cell immunodeficiency and persistent SARS-CoV-2 viremia. Cure was reached after supplementation with convalescent plasma.

Key words: SARS-CoV-2, Remdesivir, SARS-CoV-2 RNA-dependent RNA-polymerase mutation, COVID-19, COVID-19 convalescent plasma.

Accepted Manuscript

Background

Coronavirus disease 2019 (COVID-19) has recently emerged as a new pandemic due to the severe acute respiratory syndrome-coronavirus 2 (SARS-CoV-2). Alongside the well-described acute pulmonary presentation of Covid-19, a more chronic evolution of the disease is being reported in patients with B-cell immunodeficiency, probably associated with the inability to produce anti-SARS-CoV-2 antibodies[1]. Remdesivir is an RNA-dependent RNA-polymerase (RdRP) inhibitor of SARS-CoV-2, the first antiviral to be given in the USA [2] and Europe [3], with an impact on the hospital duration stay but not on mortality [4]. SARS-CoV-2 mutation, especially in RNA-dependent RNA polymerase (RdRP), may affect remdesivir susceptibility [5]. More recently, anti-SARS-CoV-2 antibody supplementation by COVID-19 convalescent plasma (CP) has emerged as a promising therapy but the characterization of the patients most likely to benefit from CP is critical [6].

We report the case of a 76-year-old woman with B-cell immunodeficiency who presented with severe protracted COVID-19 and persistent SARS-CoV-2 viremia, in whom remdesivir treatment failure was associated with the emergence of a point mutation in the RdRP and cure was achieved only after CP therapy.

Methods

SARS-CoV-2 RT-PCR and antibodies testing

In house real-time reverse transcriptase PCR (RT-PCR) tests for SARS-CoV-2 nucleic acid were performed using nasopharyngeal swab, bronchoalveolar lavage (BAL) and plasma samples. Primer and probe sequences targeted two regions on the RdRP gene and were

specific to SARS-CoV-2. Assay sensitivity was around 10 copies/reaction (Institut Pasteur, Paris, France).

Sera were tested for anti-SARS-CoV-2 antibodies using the Elecsys Anti-SARS-CoV-2 electrochemiluminescence immunoassay for detection of IgA and IgG against the SARS-CoV-2 nucleocapsid antigen (Roche Cobas 6000®). This immunoassay was approved by the FDA regarding its excellent analytical performances (sensitivity 99.5% and specificity 99.8%). The positive cut-off value of optical density index ratio was 1.10.

RdRP sequencing

Screening for mutation was performed after reverse transcriptase with Transcriptase inverse SuperScript™ III (ThermoFisher), 4 nested PCR were performed to amplify the entire RdRP gene of 2795 nucleotides (nt), or 932 amino-acids (aa), in length (Supplementary data Table 2). Sanger sequencing was performed with BigDye Terminator chemistry (Thermo Fisher Scientific®) and the analysis of the reaction products on an ABI sequencer.

Sequences were analyzed on Geneious 11.1.4 as following: after cleaning base with low quality at 5' and 3' ends (error probability limit set at 0.05), the four fragments were mapped on the annotated SARS-CoV-2 reference sequence NC_045512 and consensus sequences were generated for each sample. Multiple alignments of nt and aa consensus sequences were performed with Mafft v7.388 implemented in Geneious software. Finally, the differences between sequences were identified visually.

CP collection and preparation

CP collection was approved by the ethical committee (Comité de Protection des Personnes d'Ile-de-France) on March 26th, 2020. The plasma was obtained as previously described [7].

RESULTS

A 76-year-old woman arrived at the emergency room on the 11th of April 2020 presenting with a cough, fever, and dyspnea that had persisted for 4 days. COVID-19 was diagnosed via SARS-CoV-2 positive RT-PCR on a nasopharyngeal swab.

Her medical history included recent chronic lymphocytic leukemia diagnosed in 2019. She was successfully treated by 4 doses of Rituximab / Bendamustine, last cure in October 2019. While the patient was considered in complete remission, she exhibited secondary B-immunodeficiency requiring immunoglobulin substitution stopped in February 2020.

Upon admission for COVID-19, laboratory tests exhibited a low Ig level (1.2 g/l) for which intravenous Ig were administered and Lymphocytes (Lc) immunophenotyping revealed 70 Lc/mm³ with 88% CD3+ T Lc, <1%CD19 B Lc and 11% CD16+CD56+ NK Lc. The patient was treated by cefotaxime without improvement as she consequently required increased oxygen need with fever and increased inflammation (Supplementary data Figure 1).

Broad spectrum antibiotherapy was administered and hydroxychloroquin was initiated for 9 days (29th April-7th May) with corticosteroids without improvement. On May 27, the patient still suffered from fever and hypoxemia, with an inflammatory syndrome (CRP at 137 mg/L) and a measurable SARS-CoV-2 viremia (3.88log₁₀ copies/ml) while testing negative for anti SARS-CoV-2 antibodies (Roche Cobas 6000®). Remdesivir was introduced for 5 days

(compassionate use GS-US-540-5821, 200mg at day 1 followed by 100 mg per day). The patient experienced an early clinical and biological improvement with a normalization of oxygen saturation in room air and a decreased CRP level to 38mg/L. However, SARS-CoV-2 viremia remained detectable ($3.94 \log_{10}$ copies/ml, 9th June). Five days following treatment discontinuation, the patient experienced a new clinical degradation with fever and required intra nasal oxygenation (9th June). The patient was then treated with CP (4 X 200 ml from the 12th to 15th June 2020) and experienced a rapid improvement of her clinical, biological and radiological parameters, without detection of SARS-CoV-2 in blood on day 7 and 28 after CP transfusion (22th June) (Supplementary data Table 1, Figure 1). The patient was discharged on the 22th of June and six weeks after CP transfusion, no SARS-CoV-2 RNA was detected in blood and in rhinopharynx.

Virology and RdRP mutation

Screening for mutations in the RdRP gene was performed on 2 expectorations (24th April and 15th May and 1 BAL from 27th April) and 1 blood sample (from 27th May) collected before remdesivir treatment and 2 blood samples collected after remdesivir treatment (on 9th and 12th June). Sequencing was successful for all samples except the last one collected on 12th June. The three respiratory samples collected before treatment as well as the blood sample collected the day of remdesivir initiation harbored strictly the same amino-acid sequences. In the blood sample collected after treatment (9th June), a substitution of one nucleotide in position 1449 (G>T) in the RdRP gene led to a non-synonymous mutation changing an acid aspartic to tyrosine in position 484 (Figure 1).

DISCUSSION

We demonstrate the occurrence of a mutation in RdRP polymerase following failure of remdesivir treatment in a 76-year-old woman with B-cell immunodeficiency and persistent SARS-CoV-2 viremia. RNA viruses are characterized by a high mutation rate and the occurrence of such point mutations (STPs) has already been described mainly in the different viral proteins of SARS-CoV-2 and reported as a risk for the emergence of resistance to treatment [5]. We report the first case of occurrence of a STP in RdRP *in vivo* following remdesivir treatment with identification of a single point mutation in RdRP gene not present before treatment. Although phenotypic assay was not performed, the non-synonymous mutation D484Y was only evidenced in the sample collected after treatment. Furthermore, it occurred in the finger subdomain (extended from aa 395 to 581)[8] which is one of the 3 subdomains constituting the catalytic domain of the RdRP. Lastly, D484Y is proximate to other mutations already described in SARS-CoV (F480L) and other coronaviruses (F476L) to induce resistance to remdesivir *in vitro* [9-11].

Remdesivir treatment is to date the only COVID-19 antiviral treatment with a reported efficacy and thus, identification of putative resistance mutations is of utmost importance. It is unclear whether the emergence of this mutation was favored by the profound B-cell depletion but our case illustrates however, the likelihood of appearance of SARS-CoV-2 resistant strains under treatment pressure and the need of new antiviral treatments aiming at different targets alongside the viral cycle, such as elbasvir or EIDD-1931 [10, 12].

We used a 5 day treatment duration in our patient, which was considered not inferior to a 10 day course for patients not requiring mechanical ventilation. However, the specific case of immunodeficiencies was not addressed in this trial, a subpopulation for which a longer

treatment duration could be considered. The patient did respond clinically to a 5-day treatment course with receding of fever, dyspnea and lifting of oxygen need. However, the persistence of an inflammatory syndrome at the end of treatment as well as a persistent viremia associated with the occurrence of the D484Y mutation in RdRP may be predictive of treatment failure even with a 10-day course of remdesivir.

The other originality of this case is the effectiveness of CP transfusion. By allowing for the provision of anti-SARS-CoV2 antibodies to patients unable to mount a humoral response because of B-cell immunodeficiency, CP treatment is emerging as a safe and efficient treatment in these cases characterized by an ongoing chronic viremia and a protracted Covid-19, as illustrated by a series of 17 cases that we recently reported [7].

In conclusion, mutations such as D484Y in the RdRP of SARS-CoV-2 may emerge *in vivo* during a 5-day course of remdesivir treatment and lead to treatment failure. Patients with a B-cell immunodeficiency experiencing a protracted form COVID-19 represent a population subgroup at risk of viral failure because of prolonged SARS-CoV-2 viremia. In this situation, SARS-CoV-2 convalescent plasma may represent an efficient therapeutic option.

Competing interests

KL, AGM and VC have received grants and honoraria from ViiV Healthcare, Gilead and MSD outside of COVID-19 and present work. KL reports expert meeting fees and travel grants from Gilead, Janssen, MSD, and Abbvie, outside the submitted work.

The other authors declare no competing interests.

Accepted Manuscript

REFERENCES

1. Tepasse PR, Hafezi W, Lutz M, et al. Persisting SARS-CoV-2 viraemia after rituximab therapy: two cases with fatal outcome and a review of the literature. *Br J Haematol* **2020**.
2. Beigel JH, Tomashek KM, Dodd LE, et al. Remdesivir for the Treatment of Covid-19 - Preliminary Report. *N Engl J Med* **2020**.
3. European Medicine Agency. First COVID-19 treatment recommended for EU authorisation <https://www.ema.europa.eu/en/news/first-covid-19-treatment-recommended-eu-authorisation>. Accessed 08/16/2020.
4. Goldman JD, Lye DCB, Hui DS, et al. Remdesivir for 5 or 10 Days in Patients with Severe Covid-19. *N Engl J Med* **2020**.
5. Vankadari N. Overwhelming mutations or SNPs of SARS-CoV-2: A point of caution. *Gene* **2020**; 752: 144792.
6. Valk SJ, Piechotta V, Chai KL, et al. Convalescent plasma or hyperimmune immunoglobulin for people with COVID-19: a rapid review. *Cochrane Database Syst Rev* **2020**; 5: CD013600.
7. Thomas Hueso, Cécile Poudroux, Anne-Lise Beaumont, et al. Efficacy and safety of convalescent plasma therapy for B-cell depleted patients with prolonged COVID-19 symptoms. *Blood* **2020**; In press.
8. Hillen HS, Kocic G, Farnung L, Dienemann C, Tegunov D, Cramer P. Structure of replicating SARS-CoV-2 polymerase. *Nature* **2020**; 584(7819): 154-6.
9. Agostini ML, Andres EL, Sims AC, et al. Coronavirus Susceptibility to the Antiviral Remdesivir (GS-5734) Is Mediated by the Viral Polymerase and the Proofreading Exoribonuclease. *mBio* **2018**; 9(2).

10. Brown AJ, Won JJ, Graham RL, et al. Broad spectrum antiviral remdesivir inhibits human endemic and zoonotic deltacoronaviruses with a highly divergent RNA dependent RNA polymerase. *Antiviral Res* **2019**; 169: 104541.
11. Sheahan TP, Sims AC, Zhou S, et al. An orally bioavailable broad-spectrum antiviral inhibits SARS-CoV-2 in human airway epithelial cell cultures and multiple coronaviruses in mice. *Sci Transl Med* **2020**; 12(541).
12. Balasubramaniam M, Reis RJS. Computational target-based drug repurposing of elbasvir, an antiviral drug predicted to bind multiple SARS-CoV-2 proteins. *ChemRxiv* **2020**.

Accepted Manuscript

Figure 1: RNA-dependent RNA-polymerase amino-acid sequences of SARS-CoV-2 before and after remdesivir treatment. The 3 respiratory samples were collected before treatment (Expecto_042420, BAL_042720 and Expecto_051520), one blood sample was collected the day of remdesivir initiation (Blood_052720) and the other blood sample was collected nine days after the end of remdesivir treatment (Blood_060920). Red cross shows the non-synonymous mutation identified only in sample collected after treatment (D484Y). Black cross show the amino-acid where mutation to remdesivir in vitro has already been described in SARS-CoV (F480L) and other coronaviruses (F476L).

Accepted Manuscript

Figure 1

1 20 30 40 50 60 70 80 90 100
Especro_042420 TADAAQIFLNIRVCGVIAAHLTTCGGTGTDDVYVRAFDIYNDKVAGFAKFLKTNCCRFQEKDIDDNLIDSYFVVKRHTFVNYOHEETIYNLKDFAVAKHDFKFF
BAL_042720 TADAAQIFLNIRVCGVIAAHLTTCGGTGTDDVYVRAFDIYNDKVAGFAKFLKTNCCRFQEKDIDDNLIDSYFVVKRHTFVNYOHEETIYNLKDFAVAKHDFKFF
Especro_051520 TADAAQIFLNIRVCGVIAAHLTTCGGTGTDDVYVRAFDIYNDKVAGFAKFLKTNCCRFQEKDIDDNLIDSYFVVKRHTFVNYOHEETIYNLKDFAVAKHDFKFF
Blood_051720 TADAAQIFLNIRVCGVIAAHLTTCGGTGTDDVYVRAFDIYNDKVAGFAKFLKTNCCRFQEKDIDDNLIDSYFVVKRHTFVNYOHEETIYNLKDFAVAKHDFKFF
Blood_069500 TADAAQIFLNIRVCGVIAAHLTTCGGTGTDDVYVRAFDIYNDKVAGFAKFLKTNCCRFQEKDIDDNLIDSYFVVKRHTFVNYOHEETIYNLKDFAVAKHDFKFF

110 120 130 140 150 160 170 180 190 200
Especro_042420 RIDCDMVIHSRRLTKYTMADLVYALRHFDGNCDTLKEILVYVYNCDDDFYFNKKDWDVFENIDLRVYANLGERVRGALKTVOFCDAHINAGIVGVLTD
BAL_042720 RIDCDMVIHSRRLTKYTMADLVYALRHFDGNCDTLKEILVYVYNCDDDFYFNKKDWDVFENIDLRVYANLGERVRGALKTVOFCDAHINAGIVGVLTD
Especro_051520 RIDCDMVIHSRRLTKYTMADLVYALRHFDGNCDTLKEILVYVYNCDDDFYFNKKDWDVFENIDLRVYANLGERVRGALKTVOFCDAHINAGIVGVLTD
Blood_051720 RIDCDMVIHSRRLTKYTMADLVYALRHFDGNCDTLKEILVYVYNCDDDFYFNKKDWDVFENIDLRVYANLGERVRGALKTVOFCDAHINAGIVGVLTD
Blood_069500 RIDCDMVIHSRRLTKYTMADLVYALRHFDGNCDTLKEILVYVYNCDDDFYFNKKDWDVFENIDLRVYANLGERVRGALKTVOFCDAHINAGIVGVLTD

210 220 230 240 250 260 270 280 290 300
Especro_042420 NDNLNGMYBFGDFIQITTPGSGVYVDYYLLMLITLIRALAESHVDTLTKYIKDILLKYDFTEERKLCBQYKYNBDTYHNCVNLDDRCILHIAN
BAL_042720 NDNLNGMYBFGDFIQITTPGSGVYVDYYLLMLITLIRALAESHVDTLTKYIKDILLKYDFTEERKLCBQYKYNBDTYHNCVNLDDRCILHIAN
Especro_051520 NDNLNGMYBFGDFIQITTPGSGVYVDYYLLMLITLIRALAESHVDTLTKYIKDILLKYDFTEERKLCBQYKYNBDTYHNCVNLDDRCILHIAN
Blood_051720 NDNLNGMYBFGDFIQITTPGSGVYVDYYLLMLITLIRALAESHVDTLTKYIKDILLKYDFTEERKLCBQYKYNBDTYHNCVNLDDRCILHIAN
Blood_069500 NDNLNGMYBFGDFIQITTPGSGVYVDYYLLMLITLIRALAESHVDTLTKYIKDILLKYDFTEERKLCBQYKYNBDTYHNCVNLDDRCILHIAN

310 320 330 340 350 360 370 380 390 400
Especro_042420 FVWLFSTVFLTSFGFLVRKIFVDGVFVVSIGYHRELGVHNDVNLHSSRLSPKELLVYAADAMHAAIGNLLDQRITTCFVAAALTNVAFQTKVGNFN
BAL_042720 FVWLFSTVFLTSFGFLVRKIFVDGVFVVSIGYHRELGVHNDVNLHSSRLSPKELLVYAADAMHAAIGNLLDQRITTCFVAAALTNVAFQTKVGNFN
Especro_051520 FVWLFSTVFLTSFGFLVRKIFVDGVFVVSIGYHRELGVHNDVNLHSSRLSPKELLVYAADAMHAAIGNLLDQRITTCFVAAALTNVAFQTKVGNFN
Blood_051720 FVWLFSTVFLTSFGFLVRKIFVDGVFVVSIGYHRELGVHNDVNLHSSRLSPKELLVYAADAMHAAIGNLLDQRITTCFVAAALTNVAFQTKVGNFN
Blood_069500 FVWLFSTVFLTSFGFLVRKIFVDGVFVVSIGYHRELGVHNDVNLHSSRLSPKELLVYAADAMHAAIGNLLDQRITTCFVAAALTNVAFQTKVGNFN

410 420 430 440 450 460 470 480 490 500
Especro_042420 KDFYDFAVIKGFFREGSSVELKHFFFAODGNAALIDYDYRYNLTITMCDIRQLLFVEVVDKYPFCYDGGCINANQVIVNNDKLAGFFNKIKGKARLYYDIMS
BAL_042720 KDFYDFAVIKGFFREGSSVELKHFFFAODGNAALIDYDYRYNLTITMCDIRQLLFVEVVDKYPFCYDGGCINANQVIVNNDKLAGFFNKIKGKARLYYDIMS
Especro_051520 KDFYDFAVIKGFFREGSSVELKHFFFAODGNAALIDYDYRYNLTITMCDIRQLLFVEVVDKYPFCYDGGCINANQVIVNNDKLAGFFNKIKGKARLYYDIMS
Blood_051720 KDFYDFAVIKGFFREGSSVELKHFFFAODGNAALIDYDYRYNLTITMCDIRQLLFVEVVDKYPFCYDGGCINANQVIVNNDKLAGFFNKIKGKARLYYDIMS
Blood_069500 KDFYDFAVIKGFFREGSSVELKHFFFAODGNAALIDYDYRYNLTITMCDIRQLLFVEVVDKYPFCYDGGCINANQVIVNNDKLAGFFNKIKGKARLYYDIMS

510 520 530 540 550 560 570 580 590 600
Especro_042420 YEDCDALFAYTKRNVITITOMNLYKIASAKNRARIVAGVETCTITNRGFHOKLLKIAATRIGATVIGTKFYGGHNMKTVYDVENHLMGNDYKCDR
BAL_042720 YEDCDALFAYTKRNVITITOMNLYKIASAKNRARIVAGVETCTITNRGFHOKLLKIAATRIGATVIGTKFYGGHNMKTVYDVENHLMGNDYKCDR
Especro_051520 YEDCDALFAYTKRNVITITOMNLYKIASAKNRARIVAGVETCTITNRGFHOKLLKIAATRIGATVIGTKFYGGHNMKTVYDVENHLMGNDYKCDR
Blood_051720 YEDCDALFAYTKRNVITITOMNLYKIASAKNRARIVAGVETCTITNRGFHOKLLKIAATRIGATVIGTKFYGGHNMKTVYDVENHLMGNDYKCDR
Blood_069500 YEDCDALFAYTKRNVITITOMNLYKIASAKNRARIVAGVETCTITNRGFHOKLLKIAATRIGATVIGTKFYGGHNMKTVYDVENHLMGNDYKCDR

610 620 630 640 650 660 670 680 690 700
Especro_042420 AMINMLRIMASLVLAIRKHTCCSLHRFYRLANECAOVLSEMMVGGSLYVKGGTSSGDATTAYANVFNICOAVTANVALLSTDGNKIDKYVRNLOHRLY
BAL_042720 AMINMLRIMASLVLAIRKHTCCSLHRFYRLANECAOVLSEMMVGGSLYVKGGTSSGDATTAYANVFNICOAVTANVALLSTDGNKIDKYVRNLOHRLY
Especro_051520 AMINMLRIMASLVLAIRKHTCCSLHRFYRLANECAOVLSEMMVGGSLYVKGGTSSGDATTAYANVFNICOAVTANVALLSTDGNKIDKYVRNLOHRLY
Blood_051720 AMINMLRIMASLVLAIRKHTCCSLHRFYRLANECAOVLSEMMVGGSLYVKGGTSSGDATTAYANVFNICOAVTANVALLSTDGNKIDKYVRNLOHRLY
Blood_069500 AMINMLRIMASLVLAIRKHTCCSLHRFYRLANECAOVLSEMMVGGSLYVKGGTSSGDATTAYANVFNICOAVTANVALLSTDGNKIDKYVRNLOHRLY

710 720 730 740 750 760 770 780 790 800
Especro_042420 ECLYRNRDVIDDFVNEFYAYLRKHSMMLSDDAVCFNSTYASGLVASIKNFKSVLYONNVFMSAKCWTETDLKGFHEFCQHMLVKRQDDVYVLYY
BAL_042720 ECLYRNRDVIDDFVNEFYAYLRKHSMMLSDDAVCFNSTYASGLVASIKNFKSVLYONNVFMSAKCWTETDLKGFHEFCQHMLVKRQDDVYVLYY
Especro_051520 ECLYRNRDVIDDFVNEFYAYLRKHSMMLSDDAVCFNSTYASGLVASIKNFKSVLYONNVFMSAKCWTETDLKGFHEFCQHMLVKRQDDVYVLYY
Blood_051720 ECLYRNRDVIDDFVNEFYAYLRKHSMMLSDDAVCFNSTYASGLVASIKNFKSVLYONNVFMSAKCWTETDLKGFHEFCQHMLVKRQDDVYVLYY
Blood_069500 ECLYRNRDVIDDFVNEFYAYLRKHSMMLSDDAVCFNSTYASGLVASIKNFKSVLYONNVFMSAKCWTETDLKGFHEFCQHMLVKRQDDVYVLYY

810 820 830 840 850 860 870 880 890 900
Especro_042420 DRSRILGAGCFVDDIVKIDGTLMIERFVLAIDAYLTKHPNOEYADVFRHLYOYIRKLDHDELGHMLDMYVWMLINDTISRWEFEFYAMTTHIVLQ
BAL_042720 DRSRILGAGCFVDDIVKIDGTLMIERFVLAIDAYLTKHPNOEYADVFRHLYOYIRKLDHDELGHMLDMYVWMLINDTISRWEFEFYAMTTHIVLQ
Especro_051520 DRSRILGAGCFVDDIVKIDGTLMIERFVLAIDAYLTKHPNOEYADVFRHLYOYIRKLDHDELGHMLDMYVWMLINDTISRWEFEFYAMTTHIVLQ
Blood_051720 DRSRILGAGCFVDDIVKIDGTLMIERFVLAIDAYLTKHPNOEYADVFRHLYOYIRKLDHDELGHMLDMYVWMLINDTISRWEFEFYAMTTHIVLQ
Blood_069500 DRSRILGAGCFVDDIVKIDGTLMIERFVLAIDAYLTKHPNOEYADVFRHLYOYIRKLDHDELGHMLDMYVWMLINDTISRWEFEFYAMTTHIVLQ