



HAL
open science

Sofosbuvir-Daclatasvir Is Suboptimal in Patients with Genotype 2 Chronic Hepatitis C Infection: Real-life Experience from the HEPATHER ANRS CO22 Cohort

Victor Lédighen, Clovis Lusivika-Nzinga, Jean-Pierre Bronowicki, Fabien Zoulim, Dominique Larrey, Sophie Metivier, Albert Tran, Patrick Marcellin, Didier Samuel, Olivier Chazouillères, et al.

► To cite this version:

Victor Lédighen, Clovis Lusivika-Nzinga, Jean-Pierre Bronowicki, Fabien Zoulim, Dominique Larrey, et al.. Sofosbuvir-Daclatasvir Is Suboptimal in Patients with Genotype 2 Chronic Hepatitis C Infection: Real-life Experience from the HEPATHER ANRS CO22 Cohort. *Journal of Viral Hepatitis*, 2020, 27 (10), pp.964–973. 10.1111/jvh.13321 . hal-03704126

HAL Id: hal-03704126

<https://hal.sorbonne-universite.fr/hal-03704126>

Submitted on 30 Aug 2022

HAL is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers.

L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.



Sofosbuvir-based regimens are suboptimal in patients with genotype 2 chronic hepatitis C infection: real-life experience from the HEPATHER ANRS CO22 cohort

Journal:	<i>Journal of Viral Hepatitis</i>
Manuscript ID	JVH-00011-2020.R1
Manuscript Type:	Original Paper
Date Submitted by the Author:	n/a
Complete List of Authors:	de Lédighen, Victor; Hopital Haut Leveque, Hepatology Lusivika Nzinga, Clovis; Sorbonne Université, INSERM, Institut Pierre Louis d'épidémiologie et de Santé Publique, Paris, France., Bronowicki, Jean-Pierre; CHU Nancy-Brabois Zoulim, Fabien; INSERM, U871; Larrey, Dominique; CHU Saint Eloi, Hepatology Department Metivier, Sophie; Hôpital Purpan, Tran, Albert Marcellin, Patrick; Assistance Publique - Hopitaux de Paris, Hepatology department, Hôpital Beaujon Samuel, Didier; Hopital Paul Brousse, Hepatobiliary Center Chazouilleres, Olivier Chevaliez, Stephane; Hopital Henri Mondor, French National Reference Center for Viral Hepatitis B, C and delta; Department of Virology Dorival, Celine; UPMC Faculte de Medecine; Institut Pierre Louis d'Epidemiologie et de Sante Publique Fontaine, Hélène Pawlotsky, Jean-Michel; Hopital Universitaire Henri-Mondor - APHP Carrat, Fabrice; Hôpital Saint-Antoine, AP-HP, Santé Publique; Sorbonne Universités, INSERM, UPMC Univ Paris 06, Institut Pierre Louis d'épidémiologie et de Santé Publique (IPLESP UMRS 1136), Pol, Stanislas; Hopital Cochin, Assistance Publique-Hôpitaux de Paris (AP-HP), HUPC, Hôpital Cochin, Unité d'Hépatologie, Paris, France; Université Paris Descartes, Paris, France; Inserm UMS20 and U-1223, Paris, France
Keywords:	hepatitis C, Genotype 2, Sofosbuvir, Daclatasvir, Hepather

1
2
3
4
5
6 **Sofosbuvir-based regimens are suboptimal in patients with genotype 2**
7 **chronic hepatitis C infection: real-life experience from the HEPATHER ANRS**
8 **CO22 cohort**
9

10
11
12
13 **Short title: Real-life sofosbuvir data in G2 HCV patients**
14

15 **V. de Lédighen¹, C Lusivika-Nzinga², J.P. Bronowicki³, F. Zoulim⁴, D. Larrey⁵, S. Metivier⁶,**
16 **A. Tran⁷, P. Marcellin⁸, D. Samuel⁹, O. Chazouillères¹⁰, S. Chevaliez¹¹, C. Dorival², H.**
17 **Fontaine¹², J.M. Pawlotsky¹¹, F. Carrat^{2,13}, S. Pol¹² on the behalf of the AFEF/ANRS study**
18 **group**
19
20
21
22

- 23 1. Hepatology Unit, University Hospital Bordeaux and INSERM U1053, Bordeaux University,
24 France
25
26 2. Sorbonne Université, INSERM, Institut Pierre Louis d'épidémiologie et de santé Publique,
27 Paris, France.
28
29 3. Hepato-gastroenterology, University Hospital Nancy and INSERM U1254, Lorraine
30 University, France
31
32 4. Hepatology Department, Hospices Civils de Lyon & INSERM U1052, Lyon University, Lyon,
33 France
34
35 5. Liver Unit, Saint Eloi Hospital, CHU Montpellier, Montpellier, France
36
37 6. Service Hépatologie Rangueil, CHU Toulouse, Toulouse, France
38
39 7. Université Côte d'Azur, Nice; CHU de Nice, Digestive Center, Nice; INSERM, U1065, C3M,
40 Team 8 « Chronic liver diseases associated with obesity and alcohol », Nice, France
41
42 8. University of Paris, Hepatology Unit Beaujon Hospital - APHP, Inserm UMR 1149, Clichy,
43 France
44
45 9. Hepatology Unit, APHP Paul Brousse, Villejuif, France
46
47 10. Assistance Publique - Hôpitaux de Paris, Sorbonne University, Hepatology Department,
48 INSERM, Saint-Antoine Research Center, Saint-Antoine Hospital, Paris, France
49
50 11. National Reference Center for Viral Hepatitis B, C, and Delta, Department of Virology,
51 Hôpital Henri Mondor, Université Paris-Est, Créteil, France & INSERM U955, Créteil, France
52
53 12. Hepatology Unit, APHP Cochin, Paris, France
54
55 13. Assistance Publique - Hôpitaux de Paris, Hôpital Saint-Antoine, Unité de Santé Publique,
56 Paris, France.
57
58
59
60

Disclosures

Dr V de Lédighen has received consulting and lecturing fees from Gilead, AbbVie, Echosens, Intercept Pharma, SuperSonic Imagine, Indivior, Spimaco, Pfizer.

Dr S Pol has received consulting and lecturing fees from Bristol-Myers Squibb, Janssen, Gilead, MSD, Abbvie and grants from Bristol-Myers Squibb, Gilead, Roche and MSD

Dr A Tran has received consulting and lecturing fees from Gilead, AbbVie, MSD, Ipsen and Echosens

Dr D Larrey has received consulting services from Gilead, Abbvie, MSD, Bristol-Myers Squibb, and grants from Gilead and Abbvie

Dr S Métivier has received congress invitation from Gilead

Dr H Fontaine has received congress invitations and lecturing fees from Abbvie, Gilead and MSD.

Dr O Chazouillères has received consulting and lecturing fees from Intercept Pharma, Genfit and Pliant Therapeutics

S Chevaliez has received grants from Gilead and is an advisor/speaker for Abbott Diagnostics, Abbvie, Hologic, and Cepheid

C Lusivika-Nzinga, C Dorival, F Zoulim, JP Bronowicki, S Pol, D Samuel: nothing to disclose

Dr. Carrat reports grants from INSERM-ANRS, during the conduct of the study; personal fees from IMAXIO, outside the submitted work.

Structured summary

Background

Sofosbuvir plus daclatasvir with or without ribavirin has demonstrated a high efficacy and an acceptable safety profile in clinical trials of patients infected with genotype 2 hepatitis C virus (HCV), however, there are currently no real-world data available for this regimen.

Aims

To evaluate the real-life safety and efficacy of sofosbuvir/daclatasvir with or without ribavirin in genotype 2 HCV patients in the French cohort ANRS CO22 HEPATHER (NCT01953458).

Methods

In this ongoing, national, multicentre, prospective, observational study, we observed patients with HCV genotype 2 infection who initiated treatment with sofosbuvir (400 mg/d) plus daclatasvir with or without ribavirin (1–1.2 g/d). Patients were divided into two treatment groups: sofosbuvir/daclatasvir with or without ribavirin (12 weeks/24 weeks). The primary endpoint was a sustained virological response at week 12 following the end of therapy.

Results

Overall, 88% and 91% of patients achieved a sustained virological response following 12- and 24 weeks of treatment with sofosbuvir/daclatasvir with or without ribavirin, respectively. The most common adverse events were asthenia (29%), headache (15%) and fatigue (20%) **and ribavirin addition was associated with a higher rate of adverse events and treatment discontinuation.**

Conclusions

Sofosbuvir/daclatasvir with or without ribavirin was associated with lower rates of sustained virological response in the real-life setting compared with the clinical setting, and demonstrated suboptimal efficacy for the treatment of patients with genotype 2 chronic HCV.

Keywords

Hepatitis C, genotype 2, direct-acting antivirals, HEPATHER, real-world, sofosbuvir, daclatasvir

For Peer Review

1
2
3
4
5
6
7
8
9
10
11
12
13
14
15
16
17
18
19
20
21
22
23
24
25
26
27
28
29
30
31
32
33
34
35
36
37
38
39
40
41
42
43
44
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60

Introduction

Direct-acting antiviral agents (DAAs) have transformed the treatment of hepatitis C virus (HCV), increasing sustained virologic response (SVR) rates with the associated reduction of HCV-related complications such as liver failure, hepatocellular carcinoma and liver-related death (1). DAA-based combinations are now recommended by the European Association for the Study of the Liver (EASL) for the treatment of chronic infection with all HCV genotypes (2). Genotype 2 accounts for approximately 8% of patients with chronic HCV in Europe, and approximately 13–15% of all HCV infections in the United States. Previous studies have shown 11% of patients with HCV in France are infected with genotype 2 (3). Genotype 2 infection has, historically, been 'easy-to-treat' due to favourable SVR rates of about 80% with the treatment of pegylated interferon (PegIFN) and ribavirin (RBV) (4). Now, due to the introduction of DAAs, SVR rates for patients with genotype 2 have increased to >95% (5). Today, several DAA combinations are available for the treatment of patients infected with genotype 2 (2) (1). Sofosbuvir (NS5B inhibitor) in combination with daclatasvir (NS5A inhibitor) was recommended in the EASL guidelines 2018 (2) and remains recommended as an alternative regimen by the American Association for the Study of Liver Diseases and Infectious Diseases Society of America (AASLD-IDSA). This combination has demonstrated a high efficacy in patients with HCV genotype 2, with previous clinical trials reporting SVR rates of up to 97%(6, 7)(1, 2) 12 weeks after treatment (SVR12), including patients co-infected with human Immunodeficiency virus (HIV) (7) (6). Previous reports have also indicated that this combination is well tolerated in patients with genotype 2 infection; most common adverse events were fatigue, nausea, and headache, and no patients discontinued treatment due to adverse events (7) (8).

1
2
3 Real-world data of DAAs in patients with genotype 2 chronic infection are limited; however,
4
5 available data for DAAs appear to show similar SVR rates in the real-world setting versus the
6
7 clinical trial setting. Sofosbuvir plus ribavirin has shown almost the same SVR12 rates in both
8
9 the clinical and real-life settings (91% versus 94%, respectively, in patients without cirrhosis)
10
11 (5). Real-world data has shown sofosbuvir/velpatasvir with or without ribavirin in genotype
12
13 2 patients matches the clinical setting, with SVR rates reaching $\geq 95\%$ (9) (10). Only a few
14
15 real-world studies have been published with glecaprevir/pibrentasvir (11) (12) and currently,
16
17 there are few published studies that have examined sofosbuvir with daclatasvir in in the
18
19 real-life setting. Since the generic sofosbuvir/daclatasvir combination remains highly
20
21 prescribed in low income countries, it seems relevant to report our real world results.
22
23
24
25
26
27

28 The French ANRS CO22 HEPATHER is a large multicentre study aiming to identify prognosis
29
30 factors, including response to treatment, in patients infected with hepatitis B virus (HBV)
31
32 and HCV in a real-life setting. The objective of the current prospective study was to examine
33
34 the real-life efficacy and safety of the combination of sofosbuvir plus daclatasvir with or
35
36 without ribavirin in HCV genotype 2-infected patients within the French ANRS CO22
37
38 HEPATHER cohort.
39
40
41
42
43
44
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60

Patients and Methods

Study design and participants

ANRS CO22 HEPATHER cohort (NCT01953458) is an ongoing, national, multicentre, prospective, observational study taking place across 32 investigator sites in France. This study aims to obtain real-world data on the efficacy and safety of hepatitis treatments in patients treated in 'real-life'. The study aimed to include a large population of approximately 25,000 patients (15,000 patients with HCV, 10,000 patients with HBV; of all genotypes), with a long-term follow-up of 7 or 8 years. Patients were recruited by their respective participating centre. Details of the cohort have been previously published (13).

All patients with HCV genotype 2 infection who initiated treatment with either sofosbuvir (SOF) plus ribavirin (RBV, 0.2 – 1.2 g/d), or a combination of SOF plus daclatasvir (DCV) with or without RBV (0.2 – 1.2 g/d) were included in this analysis. Patients were divided into four treatment groups: SOF/RBV (12 weeks), SOF/RBV (24 weeks), SOF/DCV with or without RBV (12 weeks) and SOF/DCV with or without RBV (24 weeks) but only results of the SOF/DCV combination should be detailed. Genotype and subtype were determined by means of sequencing of the non-structural (NS) 5B region of the HCV genome followed by phylogenetic analysis. Analysis of NS5A and NS5B resistance-associated substitutions (RASs) at baseline and at the time of failure was performed in patients treated with the DCV regimen by means of population sequencing (sensitivity of the Sanger method: 20–30%). Cirrhosis was defined by concordant reports of either a platelet count < 150 G/L or a prothrombin time < 70% or the results of liver biopsy, a fibrotest result > 0.75, a liver stiffness (FibroScan) result greater than 12.5 KPa, a fibrometer result >0.98 , a hepascore result >0.84, performed less than 1 year before and up to 3 months after inclusion. Obesity

1
2
3 was defined by a BMI \geq 30 kg/m². Due to the observational nature of the study, not all these
4
5 data were available.
6
7

8 *Outcomes*

9
10 In this report, we focus on the primary endpoint of SVR at 12 weeks (SVR12) defined by the
11
12 undetectability of HCV RNA 12 weeks after the last treatment intake. Secondary endpoints
13
14 include the undetectability of HCV RNA, 4 and 24 weeks after last treatment intake (SVR4
15
16 and SVR24, respectively), resistance analysis, premature treatment discontinuations, and
17
18 adverse events.
19
20
21
22

23 *Statistical analyses*

24
25 Descriptive analyses, including means, frequencies and percentages (with 95% confidence
26
27 intervals [CIs] for SVR12) were used to summarise the data. A univariable analysis was
28
29 performed to verify the association between SVR12 and treatment duration, patient
30
31 demographics/clinical characteristics and whether the regimen contained DCV. A
32
33 multivariable analysis was also performed to examine any potential relationships between
34
35 SVR12 and treatment duration, gender, obesity, and presence/absence of DCV.
36
37
38
39
40

41 Fisher's exact tests and Cochran–Mantel–Haenszel statistics (stratified by: 12 versus 24
42
43 weeks, and regimen with RBV versus regimen without RBV) were used to compare
44
45 treatment discontinuations and adverse events. A level of significance (α) of 5% was set for
46
47 all analyses. Collected data were stratified based on presence/absence of cirrhosis. Most
48
49 analyses were performed considering all patients included in the study as a single group.
50
51 The analyses were divided into SOF/DCV with or without RBV (12 weeks or 24 weeks), for
52
53 analysis of SVR type association, for description of treatment outcomes, and description of
54
55 the most common adverse events.
56
57
58
59
60

Results

Patient population

Between 6 August 2012 and 31 October 2016, 12,101 patients infected with chronic HCV were enrolled in the HEPATHER cohort, including 737 (6.1%) patients infected with HCV genotype 2. By 31 October 2016, 304 patients with genotype 2 in this cohort had started either a SOF/RBV (n=233) or a SOF/DCV (n=45) with or without RBV combination regimen. Overall, 278/304 (91.4%) patients completed treatment, with premature treatment discontinuation having occurred in 5% (14/278: 7 drug intolerance, 3 anaemia, 2 kidney failures, 1 asthma, 2 patient's request) of patients (Figure 1).

Across the four groups, the mean age of patients ranged from 61.1 – 67.1 years; 103 patients (37.1%) had cirrhosis, and 150 patients (54.0%) were treatment naïve (Table 1).

Baseline characteristics were similar between treatment groups, with the exception of the following: a larger proportion of treatment-naïve patients in the groups receiving 12 weeks (n= 34) compared with 24 weeks of therapy (n = 11), and a larger proportion of patients with cirrhosis, lower haemoglobin levels and lower platelet counts in the groups receiving 24 weeks of therapy versus 12 weeks. HCV subtype distribution is detailed in Supplementary appendix 1.

Efficacy

Overall, 88% (30/34) of patients achieved an SVR12 following 12 weeks of SOF/DCV with or without ribavirin. In those patients treated for 24 weeks, the SVR12 was similar (91%; [10/11]). **Treatment failed in five patients receiving SOF/DCV without RBV.** Known relapses occurred 12 weeks following the end of treatment in two patients treated with SOF/DCV

1
2
3 (12-week duration) without RBV. No significant difference in SVR12 was observed by
4
5 univariable analysis between the 12 and 24-week treatment durations ($p=0.9999$, Table 3).
6
7

8
9 No significant difference was observed in SVR12 rates for treatment-naïve patients versus
10
11 treatment-experienced patients across all treatment groups ($p=0.0753$). Whether or not the
12
13 regimen contained daclatasvir also showed no significant difference ($p=0.9999$, Table 3).
14
15

16
17 In patients without cirrhosis, the SVR12 in patients treated with SOF/DCV with or without
18
19 RBV was 94% (15/16) following 12 weeks of treatment; for those treated for 24 weeks, the
20
21 SVR12 was 100% (4/4). No significant difference was observed between treatment groups
22
23 ($p=0.9999$).
24
25

26
27 In patients with cirrhosis the SVR12 was 80% (12/15) when treated with SOF/DCV with or
28
29 without RBV for 12 weeks; for those patients treated for 24 weeks with this regimen, the
30
31 SVR12 rate was 86% (6/7). No significant difference was observed between treatment
32
33 groups ($p=0.9712$).
34
35

36
37 Sex and obesity significantly affected SVR12 rates across all treatment groups when
38
39 examined by multivariable analysis. Female patients achieved higher SVR12 rates across all
40
41 treatment groups than males (92% vs 83%, $p=0.0275$), and patients with obesity had lower
42
43 SVR12 rates across all treatment groups than those without (80% vs 90%, $p=0.0481$). Other
44
45 patient variables demonstrated no impact on SVR12 (detailed in Table 4).
46
47
48

49 50 *Resistance*

51
52
53 Resistance analysis at baseline was conducted for 42 patients who received a DCV regimen
54
55 and is detailed in Supplementary appendix 2. The resistance profile of the five patients in
56
57
58
59
60

1
2
3 whom treatment with the SOF/DCV with or without RBV regimen failed is detailed in
4
5
6 Supplementary appendix 3.
7

8 *Safety and tolerability*

9
10
11 The highest frequency of adverse events was observed in patients treated with RBV-
12 including regimen with significant differences noted between treatment groups for any
13
14 adverse event ($p < 0.05$). The most common adverse events ($\geq 10\%$ in any subgroup) were
15
16
17
18 asthenia (20%), and headache (13%). Adverse events \geq grade 3 occurred in 12% (4/12) of
19
20 patients treated with SOF/DCV with or without RBV for 12 weeks, and 18% (2/18) of
21
22 patients treated with this regimen for 24 weeks. **Anemia, intolerance and treatment**
23
24
25 **discontinuations were more frequent in the RBV group ($p < 0.05$).**
26
27

28
29
30
31
32
33
34
35
36
37
38
39
40
41
42
43
44
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60
Nine percent (3/34) of patients experienced a serious adverse event following treatment
with SOF/DCV with or without ribavirin for 12 weeks; similarly, 9% (1/11) patients treated
for 24 weeks experienced a serious adverse event. No significant difference was observed
between treatment groups regarding serious adverse events. No treatment-related deaths
were reported for any of the treatment groups.

42 **Discussion**

43
44
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60
In this real-world analysis, we examined the efficacy and safety of the DAA combination SOF
plus DCV with or without RBV in patients with genotype 2 HCV from the French ANRS CO22
HEPATHER cohort. SOF and DCV with or without RBV has shown a high efficacy and
acceptable safety profile in clinical trials, however few comparisons have been made
between clinical trial and real-world outcomes with this regimen (8) (6). Clinical trial
patients are selected using strict criteria and are closely monitored throughout treatment,
however data suggest that there is little difference in the performance of conventional

1
2
3 regimens in real-world clinical practice (e.g., SVR12 rates of 91% versus 94% in patients
4
5 without cirrhosis in previous studies with SOF/RBV) in patients with genotype 2 HCV (5).
6
7

8
9 In the current analysis, an SVR12 rate of around 90% was achieved with no significant
10
11 difference noticed between 12 or 24-week durations of treatment, treatment-naïve or
12
13 treatment-experienced patients, or whether the regimen contained DCV. These results
14
15 show that treatment with SOF/DCV with or without RBV in the real-life setting is suboptimal
16
17 compared with the clinical trial setting where reported SVR12 rates reach up to 97%.(6)
18
19 Similarly, the results are suboptimal compared with real-world data from the TRIO study of
20
21 SOF/velpatasvir with or without RBV for the treatment of patients with genotype 2 HCV,
22
23 which showed virologic responses matching the clinical setting, with SVR rates reaching
24
25 $\geq 95\%$.(14) However, generic oral directly acting agents are associated with high SVR rates in
26
27 patients with HCV infection in a real-life clinical scenario (15, 16).
28
29
30
31
32

33
34 The difference between efficacy rates reported from clinical trials and the SVR rates
35
36 reported from this real-world setting of HCV treatment regimens could be attributable to
37
38 several factors. With the addition of DCV demonstrating no significant difference in efficacy
39
40 across all treatment groups ($p=0.9999$), it is likely that the combination of SOF plus DCV with
41
42 or without RBV in patients is inherently insufficiently effective in genotype 2 patients.
43
44 However, other factors may contribute to these disappointing findings.
45
46
47
48

49
50 Resistance is another potential cause of the suboptimal results of this study. Analysis of
51
52 NS5A and NS5B RASs at baseline and the time of failure in patients treated with DCV
53
54 regimens are difficult to interpret due to the large number of patients with non-amplified
55
56 NS5A domain (and therefore do not explain the suboptimal results). However, the amino
57
58 acid mutation L31M was observed at baseline in 2/5 (40%) patients in whom the DCV
59
60

1
2
3 treatment regimen failed. In the baseline analysis, 28.6% (12/42) of patients had a known
4
5 L31M mutation, and a previous study reported that 23.1% of patients with genotype 2b had
6
7 this mutation (17). Since the L31M mutation has been reported to confer resistance to DCV
8
9 (18) this could have had some impact on the suboptimal SVR12 rates following treatment
10
11 with SOF/DCV with or without RBV observed in the real-life setting. Furthermore, negative
12
13 effects of atypical subtypes may also have had an effect of the SVR rates observed in this
14
15 real-life analysis. Within the genotypes, subtypes with nucleotide identities of 75–86% may
16
17 occur, individual isolates of each given subtype can typically differ by 8–10% and in each
18
19 patient these subtypes can differ further. Variation at the genotype level affects the
20
21 outcome of antiviral therapies (19). Within this analysis, patients with at least 17 HCV
22
23 subtypes were included, which could have potentially impacted the virologic responses
24
25 seen.
26
27
28
29
30
31
32

33 Known relapses occurred 12 weeks following treatment in two patients treated with
34
35 SOF/DCV with or without RBV (12-week duration. Extending treatment duration compared
36
37 with standard treatment, or adding ribavirin (cumulative ribavirin dose) have been
38
39 demonstrated to improve SVR rates in slow responders; this is considered likely due to a
40
41 lower rate of viral relapse associated with the time of being HCV negative during treatment
42
43 (20). However, in the current analysis, no such effect was seen since for both regimens no
44
45 significant difference in SVR12 was observed by univariable analysis between the 12 and 24-
46
47 week treatment durations.
48
49
50
51
52

53 In subtype analysis, higher SVR12 rates were consistently seen in females compared with
54
55 males, across treatment groups, however, previous studies have shown sex does not usually
56
57 impact SVR rates in patients with genotype 2 (21). Further, an association was seen between
58
59
60

1
2
3 obesity and treatment failure by multivariable analysis, with a lower SVR12 rate observed in
4 patients with obesity. This is consistent with other studies since obesity has been reported
5 to negatively influence virologic decline suggesting that the combination studied here is
6 failing to perform effectively in patients with negative baseline factors.
7
8
9
10
11
12

13 Both clinical trials and this real-world study of SOF with DCV with or without RBV have
14 demonstrated a low rate of discontinuation due to AES (6) (8). Asthenia and headache were
15 the most common AEs and were clearly manageable for all patients in this study; and
16 serious AEs were infrequent. Similar results are seen in patients treated with generics (16).
17
18
19
20
21
22
23

24 *Limitations*

25
26
27 Conclusions regarding that extension of treatment duration, prior treatments or addition of
28 DCV have an impact on virologic response are limited by the small sample sizes and the
29 uneven distribution of patients who were treatment naïve, and the number of patients who
30 had cirrhosis.
31
32
33
34
35
36
37
38
39
40

41 *Strengths*

42
43
44 The response rates reported in this study are typical of those seen in the clinic, and the
45 patient distribution is representative of the real-life setting leading to robust and consistent
46 results.
47
48
49
50

51 *Conclusions*

52
53
54 To our knowledge, this study is the first to show the safety and efficacy of SOF/DCV with or
55 without RBV for the treatment of patients with HCV infected with genotype 2 in a real-world
56 setting. Our results show that this combination is suboptimal and shows that the addition of
57
58
59
60

1
2
3 DCV demonstrates no significant difference to SVR rates. However, it is a first-line cheap
4
5 generic combination which remains frequently prescribed in low income countries.
6
7

8 9 **Statement of interests**

10
11 The ANRS CO22 cohort is sponsored by the French National Institute for Health and Medical
12
13 Research-French National Agency for Research on AIDS and Viral Hepatitis (INSERM-ANRS).
14

15
16 The sponsor contributed to the study design and writing of this report but had no role in
17
18 data collection, data analysis and data interpretation. Other funders of the study had no role
19
20 in study design, data collection, data analysis, data interpretation or writing of the report.
21
22
23
24
25
26
27
28
29
30
31
32
33
34
35
36
37
38
39
40
41
42
43
44
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60

References

1. Spearman CW, Dusheiko GM, Hellard M, Sonderup M. Hepatitis C. *Lancet* 2019;394:1451-1466.
2. European Association for the Study of the Liver. Electronic address eee, European Association for the Study of the L. EASL Recommendations on Treatment of Hepatitis C 2018. *J Hepatol* 2018;69:461-511.
3. Martinot-Peignoux M, Roudot-Thoraval F, Mendel I, Coste J, Izopet J, Duverlie G, Payan C, et al. Hepatitis C virus genotypes in France: relationship with epidemiology, pathogenicity and response to interferon therapy. *The GEMHEP. J Viral Hepat* 1999;6:435-443.
4. Andriulli A, Mangia A, Iacobellis A, Ippolito A, Leandro G, Zeuzem S. Meta-analysis: the outcome of anti-viral therapy in HCV genotype 2 and genotype 3 infected patients with chronic hepatitis. *Aliment Pharmacol Ther* 2008;28:397-404.
5. Welzel TM, Nelson DR, Morelli G, Di Bisceglie A, Reddy RK, Kuo A, Lim JK, et al. Effectiveness and safety of sofosbuvir plus ribavirin for the treatment of HCV genotype 2 infection: results of the real-world, clinical practice HCV-TARGET study. *Gut* 2017;66:1844-1852.
6. Wyles DL, Ruane PJ, Sulkowski MS, Dieterich D, Luetkemeyer A, Morgan TR, Sherman KE, et al. Daclatasvir plus Sofosbuvir for HCV in Patients Coinfected with HIV-1. *N Engl J Med* 2015;373:714-725.
7. Sulkowski MS, Gardiner DF, Rodriguez-Torres M, Reddy KR, Hassanein T, Jacobson I, Lawitz E, et al. Daclatasvir plus sofosbuvir for previously treated or untreated chronic HCV infection. *N Engl J Med* 2014;370:211-221.
8. Mangia A, Arleo A, Copetti M, Miscio M, Piazzolla V, Santoro R, Squillante MM. The combination of daclatasvir and sofosbuvir for curing genotype 2 patients who cannot tolerate ribavirin. *Liver Int* 2016;36:971-976.
9. Buggisch P, Wursthorn K, Stoehr A, Atanasov PK, Supiot R, Lee J, Ting J, et al. Real-world effectiveness and safety of sofosbuvir/velpatasvir and ledipasvir/sofosbuvir hepatitis C treatment in a single centre in Germany. *PLoS One* 2019;14:e0214795.
10. Chugh Y, Dhiman RK, Premkumar M, Prinja S, Singh Grover G, Bahuguna P. Real-world cost-effectiveness of pan-genotypic Sofosbuvir-Velpatasvir combination versus genotype dependent directly acting anti-viral drugs for treatment of hepatitis C patients in the universal coverage scheme of Punjab state in India. *PLoS One* 2019;14:e0221769.
11. Persico M, Aglitti A, Milella M, Coppola C, Messina V, Claar E, Gentile I, et al. Real-life glecaprevir/pibrentasvir in a large cohort of patients with hepatitis C virus infection: The MISTRAL study. *Liver Int* 2019;39:1852-1859.
12. Berg T, Naumann U, Stoehr A, Sick C, John C, Teuber G, Schifflholz W, et al. Real-world effectiveness and safety of glecaprevir/pibrentasvir for the treatment of chronic hepatitis C infection: data from the German Hepatitis C-Registry. *Aliment Pharmacol Ther* 2019;49:1052-1059.
13. Carrat F, Fontaine H, Dorival C, Simony M, Diallo A, Hezode C, De Ledinghen V, et al. Clinical outcomes in patients with chronic hepatitis C after direct-acting antiviral treatment: a prospective cohort study. *Lancet* 2019;393:1453-1464.
14. Curry M, Bacon B, Dieterich DT, Flamm SL, Kowdley KV, Milligan S, Tsai N, et al. Sofosbuvir/Velpatasvir in Genotype 2-6 HCV: Real-World Experience from the Trio Network. *Gastroenterology* 2017;152:S1061-S1062.
15. Gupta S, Rout G, Patel AH, Mahanta M, Kalra N, Sahu P, Sethia R, et al. Efficacy of generic oral directly acting agents in patients with hepatitis C virus infection. *J Viral Hepat* 2018;25:771-778.

16. Omar H, El Akel W, Elbaz T, El Kassas M, Elsaheed K, El Shazly H, Said M, et al. Generic daclatasvir plus sofosbuvir, with or without ribavirin, in treatment of chronic hepatitis C: real-world results from 18 378 patients in Egypt. *Aliment Pharmacol Ther* 2018;47:421-431.
17. Kliemann DA, Tovo CV, da Veiga AB, de Mattos AA, Wood C. Polymorphisms and resistance mutations of hepatitis C virus on sequences in the European hepatitis C virus database. *World J Gastroenterol* 2016;22:8910-8917.
18. Wang C, Sun JH, O'Boyle DR, 2nd, Nower P, Valera L, Roberts S, Fridell RA, et al. Persistence of resistant variants in hepatitis C virus-infected patients treated with the NS5A replication complex inhibitor daclatasvir. *Antimicrob Agents Chemother* 2013;57:2054-2065.
19. Aurora R, Donlin MJ, Cannon NA, Tavis JE. Genome-wide hepatitis C virus amino acid covariance networks can predict response to antiviral therapy in humans. *J Clin Invest* 2009;119:225-236.
20. Farnik H, Lange CM, Sarrazin C, Kronenberger B, Zeuzem S, Herrmann E. Meta-analysis shows extended therapy improves response of patients with chronic hepatitis C virus genotype 1 infection. *Clin Gastroenterol Hepatol* 2010;8:884-890.
21. Fried MW, Hadziyannis SJ, Shiffman ML, Messinger D, Zeuzem S. Rapid virological response is the most important predictor of sustained virological response across genotypes in patients with chronic hepatitis C virus infection. *J Hepatol* 2011;55:69-75.

Or Peer Review

Table 1 – Demographics and baseline characteristics of patient cohort

	SOF+RBV 12 weeks	SOF+RBV 24 weeks	SOF/DCV±RBV 12 weeks	SOF/DCV±RBV 24 weeks	P-value
	N=188	N=45	N=34	N=11	
Age, mean ± SD years	61.1 ± 11.7	67.1 ± 10	62.4 ± 11.5	64.2 ± 11.5	0.0284
Gender Male, n (%)	101 (54)	15 (33)	16 (47)	5 (45)	0.0968
BMI (kg/m ²), n (%)					
<18.5	4 (2)	0 (0)	1 (3)	0 (0)	0.1598
18.5–25	81 (44)	18 (40)	13 (38)	4 (36)	
25–30	65 (35)	9 (20)	11 (32)	3 (27)	
≥30	34 (18)	18 (40)	9 (26)	4 (36)	
Chronic hepatitis duration (years), mean ± SD	13 ± 9.5	11.9 ± 9.4	14.4 ± 7.9	13.5 ± 9.9	0.5524
Diabetes (%)	37 (20)	10 (22)	9 (26)	3 (27)	0.6917
Hypertension (%)	68 (36)	22 (49)	14 (41)	7 (64)	0.1577
Cirrhosis,* n (%)	56 (30)	25 (56)	15 (48)	7 (64)	0.0014
MELD score, mean ± SD	8.5 ± 4.3	10 ± 5.1	7.9 ± 2.7	11.5 ± 6.2	0.2837
MELD score (≥15), n (%)	3 (6)	3 (13)	1 (8)	1 (17)	0.4760
Child-Pugh score B or C	2 (4)	2 (8)	0 (0)	1 (14)	0.3074
Liver biopsy > 2 years	17 (30)	10 (40)	4 (27)	3 (43)	0.7443
Liver biopsy < 2 years	4 (7)	1 (4)	0 (0)	0 (0)	0.8813
Fibrosis stage (Liver biopsy)					
F0	0 (0)	0 (0)	1 (2)	0 (0)	0.9575
F1	0 (0)	0 (0)	1 (2)	0 (0)	
F1/F2	0 (0)	0 (0)	2 (4)	0 (0)	
F3	1 (7)	0 (0)	4 (7)	1 (4)	
F3/F4	3 (20)	0 (0)	4 (7)	2 (8)	
F4	11 (73)	7 (100)	44 (79)	22 (88)	
Prothrombin time (≤70%), n (%)	15 (9)	5 (12)	1 (3)	2 (20)	0.2803

AST (>5 x ULN), n (%)	7 (4)	0 (0)	1 (3)	1 (10)	0.2698
ALT (>5 x ULN), n (%)	19 (10)	3 (7)	4 (12)	0 (0)	0.7447
Albumin (<30g/L), n (%)	1 (1)	3 (8)	0 (0)	0 (0)	0.0533
Conjugated Bilirubin (>5µmol/L), n (%)	36 (42)	14 (58)	15 (60)	6 (67)	0.1771
HCV-RNA <60,000,000 IU/mL (%)	164 (90)	38 (88)	31 (91)	9 (90)	0.9801
Haemoglobin (<12g/dL in women or <13g/dL in men), n (%)	25 (13)	8 (18)	9 (27)	8 (73)	0.0001
Platelets (<100 G/L), n (%)	18 (10)	13 (30)	3 (9)	2 (20)	0.0065
Treatment naïve, n (%)	112 (60)	16 (36)	18 (53)	4 (36)	0.0187

*Including 5 patients with decompensated cirrhosis. SOF/RBV 12 weeks (n=1), SOF/RBV 24 weeks (n=2), SOF/DCV±RBV 24 weeks (n=2)

†Previous treatments in non-naïve patients were: PR (n=106), PR+BOC or TVR (n=2), DAA (n=2), Other (n=18)

SOF: Sofosbuvir - RBV: Ribavirin - DCV: Daclatasvir - SD: Standard Deviation – BMI: Body Mass Index – ULN: Upper limit of Normal – AST: Aspartate Aminotransferase – ALT: Alanine Aminotransferase

Table 2 – Virological response according to treatment with SOF/DCV with or without RBV for 12- and 24-week treatment durations

Negative HCV RNA	SOF/DCV ±RBV 12 weeks	SOF/DCV ±RBV 24 weeks	P- value
Week 4	19/31 (61)	7/10 (70)	0.3736
Week 12	32/33 (97)	8/10 (80)	0.1849
Week 24		7/7 (100)	0.9999
SVR4	30/33 (91)	10/11 (91)	0.9999
SVR12	30/34 (88)	10/11 (91)	0.9841
SVR24	24/28 (86)	9/10 (90)	0.4769
SVR12 in non-cirrhotic patients	15/16 (94)	4/4 (100)	0.9999
SVR12 in cirrhotic patients	12/15 (80)	6/7 (86)	0.9712
SVR12 in naïve patients	15/18 (83)	4/4 (100)	0.6325
SVR12 in treatment experienced patients	15/16 (94)	6/7 (86)	0.7902

SOF: Sofosbuvir - RBV : Ribavirin - DCV : Daclatasvir - HCV : Hepatitis C Virus - SVR : Sustained Virological Response

Table 3 – Association between treatment duration, whether the regimen contained daclatasvir, sex, obesity, liver function, treatment history and sustained virologic response, by univariable and multivariable analysis

Variable	n with SVR12 / Total (%)	Univariable		Multivariable	
		OR (95% CI)	P-Value	OR (95% CI)	P-value
Treatment duration [†]					
24 weeks	49/56 (88)	0.97 (0.38-2.79)	0.9999	1.02 (0.38-3.09)	0.9999
12 weeks	195/222 (88)			Ref	
DCV containing regimen [†]					
Yes	40/45 (89)	1.14 (0.4-3.99)	0.9999	1.21 (0.42-4.33)	0.9372
No	204/233 (88)			Ref	
Sex, male					
Yes	114/137 (83)	0.42 (0.18-0.94)	0.0344	0.4 (0.17-0.91)	0.0275
No	130/141 (92)			Ref	
Obese					
Yes	52/65 (80)	0.45 (0.2-1.04)	0.0637	0.41 (0.18-0.99)	0.0481
No	188/209 (90)			Ref	
ALT (>5 x ULN)					
Yes	26/26 (100)	5.82 (1.24-.)	0.0525		
No	213/247 (86)			NS	
Treatment history					
Treatment experienced	107/128 (84)	0.48 (0.21-1.07)	0.0753		
Treatment naïve	137/150 (91)			NS	

[†]Treatment duration and daclatasvir containing regimen were included in the multivariable model irrespective of the p-value in the univariable analysis. Other variables with p-value < 0.1 in the univariable analysis were included in a multivariable model and selected according to a backward selection. NS: not significant.

SOF: Sofosbuvir - RBV : Ribavirin - DCV : Daclatasvir - SVR : Sustained Virological Response – OR : Odds Ratio – CI : Confidence Interval – ULN : Upper limit of Normal – ALT : Alanine Aminotransferase

Table 4 – Association of sustained virological response and baseline demographics and clinical characteristics by univariable analysis

Variable	n with SVR12 / Total (%) (Yes VS No)	Univariable	
		OR (95% CI)	P-Value
Hypertension	93/111 (84) VS 151/167 (90)	0.55 (0.25-1.2)	0.1444
Suspicion of cirrhosis	86/103 (83) VS 153/170 (90)	0.56 (0.26-1.24)	0.1674
Diabetes	55/59 (93) VS 189/219 (86)	2.18 (0.72-8.87)	0.2163
Age at inclusion ≥ 65 years	112/123 (91) VS 132/154 (86)	1.69 (0.75-4.05)	0.2380
VL D0 < 6 000 000	214/242 (88) VS 23/28 (82)	1.66 (0.46-4.96)	0.4891
MELD ≥ 15	13/16 (81) VS 201/226 (89)	0.54 (0.13-3.16)	0.5551
AST > 5 ULN	11/11 (100) VS 226/259 (87)	2.21 (0.45-inf)	0.6071
Age at diagnosis ≥ 15 years	104/120 (87) VS 135/152 (89)	0.82 (0.37-1.82)	0.7216
Conjugated bilirubin ≥ 5 µmol/L	62/71 (87) VS 66/73 (90)	0.73 (0.22-2.36)	0.7462
Albumin < 30 g/L	3/4 (75) VS 211/237 (89)	0.37 (0.03-20.15)	0.7605
Haemoglobin < 12 or 13 g/dL	43/50 (86) VS 199/226 (88)	0.83 (0.33-2.42)	0.8421
Neutrophil < 1500 /mm ³	22/26 (85) VS 213/243 (88)	0.78 (0.24-3.31)	0.8451
PT ≤ 70%	20/23 (87) VS 209/233 (90)	0.77 (0.2-4.32)	0.8945
Platelets < 100 G/L	31/36 (86) VS 207/236 (88)	0.87 (0.3-3.09)	0.9591
Decompensated cirrhosis	4/5 (80) VS 240/273 (88)	0.55 (0.05-27.9)	0.9635
French ethnicity	141/160 (88) VS 103/118 (87)	1.08 (0.49-2.36)	0.9742
Overconsumption of alcohol at inclusion	0/0 (100) VS 244/277 (88)	---	

SVR : Sustained Virological Response – OR : Odds Ratio – CI : Confidence Interval – ULN : Upper limit of Normal – AST : Aspartate Aminotransferase – VL D0 : Viral Load at Day 0 – PT : Prothrombin Time

Table 5 – Treatment discontinuation, adverse events and serious adverse events according to treatment with SOF/RBV versus SOF/DCV with or without RBV for 12- and 24-week treatment durations

	SOF DCV ± RBV		+	Fisher P-value	CMH P-value (stratification: 12 vs 24w)	CMH P-value (stratification: RBV vs no RBV)
	12 weeks	24 weeks				
Number of patients	34	11				
All AEs not taken into account (missing start and end dates)						
All adverse events - any (maximum grade)	15 (44)	6 (55)		0.0146	0.0548	0.0102
NA				0.0683	0.1147	0.0969
Grade 1	4 (12)	3 (27)				
Grade 2	7 (21)	1 (9)				
Grade 3	4 (12)	2 (18)				
Grade 4						
Serious adverse events	3 (9)	1 (9)		0.5402	0.7490	0.3276
Adverse events (≥10% in any subgroup)						
Asthenia	9 (26)			0.2116	0.4884	0.8883
Headache	5 (15)	1 (9)		0.9858	0.9884	0.8421
Fatigue	1 (3)			0.0746	0.0430	0.0958
Dyspnoea				0.0687	0.0197	0.2452
Insomnia				0.1156	0.0234	0.4572
Pruritus				0.0065	0.0280	0.0042
Anaemia	1 (3)	2 (18)		0.2996	0.9036	0.3313
Hyperglycaemia	1 (3)			0.0290	0.2565	0.0068
Sleep disorder		2 (18)		0.0967	0.8390	0.8948
Leukopenia				0.0368	0.0884	0.0118
Treatment interruptions		3 (27)		0.0024	0.7088	0.0005
Intolerance		2 (18)		0.0054	0.7458	0.0005
Other		1 (9)		0.2251	0.8594	0.2717

CMH : Cochran–Mantel –Haenszel statistics – SOF: Sofosbuvir - RBV : Ribavirin - DCV : Daclatasvir – AE : Adverse Event – NA : Non Available

Figure legends

Figure 1 – Patient disposition and participant numbers per treatment group

Supplementary appendix table 1 – HCV subtype distribution of patient cohort


Supplementary appendix table 2 – resistance analysis at baseline of 42 patients

Supplementary appendix table 3 – resistance profile of the five patients who failed with the DCV regimen

For Peer Review

Figures

Figure 1 – Patient disposition



Peer Review

Appendices**Supplementary appendix table 1 – HCV subtype distribution of patient cohort**

Subtypes	Patients
	N=263
2a/2b/2c/2i/2k/2l, n	30/15/41/39/13/34
2d/2e/2f/2j/2m/2p/2q/2r, n	3/1/1/3/3/1/1/1
2k/1b variant, n	4
Undeterminate subtype, n	55
1a/1b, n	2/1
Non-amplified, n	15

For Peer Review

Supplementary appendix table 2 – resistance analysis at baseline (N=42)

	NS5A RAS	NS5B RAS
	N=42	N=42
Wild type, n	6	39
L31M, n	12	-
C92S, n	3	-
C92S/L31M, n	1	-
F28C, n	1	-
Non-amplified, n	19	3

For Peer Review

Supplementary appendix table 3 – resistance profile of the five patients in whom the SOF/DCV with or without ribavirin regimen failed

Subtype	NS5A	RAS	NS5B	RAS	NS5A RAS	NS5B RAS
	Baseline		Baseline		Failure	Failure
2 _{undeterminate}	Non-amplified		Wild type		Non-amplified	Not done
2a	L31M		Wild type		L31M	Wild type
2i	L31M		Wild type		L31M	Wild type
2 _{undeterminate}	Non-amplified		Wild type		Non-amplified	Wild type
2b	Non-amplified		Wild type		Not available	Not available

SOF: Sofosbuvir - DCV : Daclatasvir

For Peer Review

1
2
3
4
5
6
7
8
9
10
11
12
13
14
15
16
17
18
19
20
21
22
23
24
25
26
27
28
29
30
31
32
33
34
35
36
37
38
39
40
41
42
43
44
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60




Figure 1

159x87mm (300 x 300 DPI)