



HAL
open science

Position-dependent memory kernel in generalized Langevin equations: Theory and numerical estimation

Hadrien Vroylandt, Pierre Monmarché

► **To cite this version:**

Hadrien Vroylandt, Pierre Monmarché. Position-dependent memory kernel in generalized Langevin equations: Theory and numerical estimation. *The Journal of Chemical Physics*, 2022, 156 (24), pp.244105. 10.1063/5.0094566 . hal-03714956

HAL Id: hal-03714956

<https://hal.sorbonne-universite.fr/hal-03714956v1>

Submitted on 6 Jul 2022

HAL is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers.

L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.

Position-dependent memory kernel in generalized Langevin equations: theory and numerical estimation

Hadrien Vroylandt¹ and Pierre Monmarché^{2,3}

¹*Sorbonne Université, Institut des sciences du calcul et des données, ISCD,
F-75005 Paris, France*

²*Sorbonne Université, Laboratoire Jacques-Louis Lions, LJLL,
F-75005 Paris*

³*Sorbonne Université, Laboratoire de Chimie Théorique, LCT,
F-75005 Paris*

Generalized Langevin equations with non-linear forces and position-dependent linear friction memory kernels, such as commonly used to describe the effective dynamics of coarse-grained variables in molecular dynamics, are rigorously derived within the Mori-Zwanzig formalism. A fluctuation-dissipation theorem relating the properties of the noise to the memory kernel is shown. The derivation also yields Volterra-type equations for the kernel, which can be used for a numerical parametrization of the model from all-atom simulations.

I. INTRODUCTION

In molecular dynamics simulations, coarse-graining methods consist in replacing an all-atom system by a system with a smaller dimension, either by merging groups of atoms in single units or by describing the system through a relatively small set of collective variables. Such models are then used either to get some physical understanding of the system or, from a numerical viewpoint, to reduce the cost of sampling long trajectories. In both cases, an important issue is to describe the dynamics of the coarse-grained variables: in theory, obtaining their exact trajectory requires the simulation of the initial all-atom system, which one wants to avoid. In practice, it is approximated by so-called effective dynamics, namely closed equations which do not involve explicitly the withdrawn atomistic degrees of freedom (whose influence may be taken into account as a random noise)¹⁻⁵.

A class of models frequently used to perform this task is given by the Generalized Langevin Equations (GLE)⁶⁻¹², which usually take the following form:

$$\ddot{x}_t = F_e(x_t) - \int_0^t K(t-s)\dot{x}_s ds + \zeta_t \quad (1)$$

where $x_t \in \mathbb{R}^d$ is the position at time t of the variables, $F_e : \mathbb{R}^d \rightarrow \mathbb{R}^d$ is an effective force, $K : \mathbb{R}_+ \rightarrow \mathbb{R}^{d \times d}$ is a time-dependent linear friction kernel and ζ is a Gaussian noise. In many cases^{6,7,11-15}, $F_e = -\nabla W_{\text{pmf}}$ where W_{pmf} is the potential of mean force associated to the coarse variables. When K is a Dirac mass at 0 and ζ is a white noise this is the standard Langevin equation. The form (1) is heuristically motivated by the so-called Mori-Zwanzig formalism^{6,16-18}, but there are few cases where a rigorous derivation has been established. There is an important literature on linear (harmonic) cases¹⁷⁻²¹. The non-harmonic case, with a non-linear effective force given by the potential of mean force, is discussed in Refs. 22, 23, but it is observed in Ref. 24 that these derivations rely on some implicit and uncontrolled approximations, and it is argued that in general it is in fact not possible to get, within the Mori-Zwanzig formalism, an equation of the form (1) with a non-linear effective force but a friction term which is linear in the velocity. Our first contribution is a nuance to the statement of²⁴, namely: in general, it is possible to rigorously derive a GLE similar to (1) in a general non-linear coupled case, with a non-linear force and a friction which is linear in the velocity, if the kernel K is allowed to depend on the position, *i.e.* a GLE of the form

$$\ddot{x}_t = F_e(x_t) - \int_0^t K(t-s, x_s)\dot{x}_s ds + \zeta_t. \quad (2)$$

Moreover, in accordance with the standard assumption and in contrast to²⁴, in this case the memory kernel and the covariance of the noise are related by a form of fluctuation-dissipation relation (see sec. II E below for the definition), adapted to the position-dependent framework. To be clear, our framework is restricted to the case where the initial all-atom equation is an equilibrium Hamiltonian dynamics.

As we will see, along the formal derivation, the kernel K is proven to satisfy some integro-differential equations of Volterra type. As a natural continuation, our second contribution is the implementation of a numerical scheme based on these equations to estimate the kernel and parametrize GLE models from all-atom simulations, thus adapting the inverse Volterra method^{14,25-29} to the position-dependent kernel settings.

II. DERIVATION OF GENERALIZED LANGEVIN EQUATIONS

A. Preliminary considerations

The purpose of the Mori-Zwanzig approach is to extract the evolution in time of an observable from the dynamics of a microscopic state. More explicitly, it relies on the following elements:

- A first order differential equation for a microscopic state $\mathbf{X}(t)$, given by

$$\partial_t \phi(\mathbf{X}(t)) = \mathcal{L}\phi(\mathbf{X}(t)) \quad (3)$$

for all functions ϕ where \mathcal{L} denotes the evolution operator. For the sake of clarity, and since this is the main case of interest, we will only consider later the deterministic case where \mathcal{L} is the Liouville operator for Hamiltonian dynamics, in particular $\mathbf{X}(t) = (\mathbf{q}(t), \mathbf{p}(t)) \in \mathbb{R}^{2D}$ with D the position space dimension. Extensions to Markov operators for stochastic dynamics can be considered, following^{9,30} (although, when working at the level of the Kolmogorov equations for stochastic processes, one should be careful whether the goal is to describe the evolution of $\phi(\mathbf{X}(t))$ or of its expectation with respect to the initial random noise).

- A function $\mathcal{O} : \mathbb{R}^{2D} \rightarrow \mathbb{R}^d$, so that we are interested in writing a closed effective equation for the observable $\mathcal{O}(\mathbf{X}(t))$. For instance, \mathcal{O} can be a collective variable, or a collective variable and its velocity (in which case $d = 2n$ where n is the dimension of the collective variable, see sec. II C).
- A reference probability density ρ on \mathbb{R}^{2D} . For processes at equilibrium, it is simply chosen as the equilibrium Gibbs measure, but other measures are possible^{31,32}. Given ρ , we consider on $L^2(\rho, \mathbb{R}^d) = \{\phi : \mathbb{R}^{2D} \rightarrow \mathbb{R}^d, \int |\phi|^2 d\rho < \infty\}$ the scalar product

$$\langle \phi, \psi \rangle = \int_{\mathbb{R}^{2D}} \phi(\mathbf{X}) \cdot \psi(\mathbf{X}) \rho(\mathbf{X}) d\mathbf{X}. \quad (4)$$

- A set $\mathcal{E} = \{e_k, k \in \mathcal{I}\}$ of linearly independent functions from \mathbb{R}^d to \mathbb{R}^d , for some index set \mathcal{I} .

From the functions $e_k : \mathbb{R}^d \rightarrow \mathbb{R}^d$, it is convenient to introduce the functions $E_k = e_k \circ \mathcal{O}$ from \mathbb{R}^{2D} to \mathbb{R}^d , where \circ is the composition, i.e. $(e_k \circ \mathcal{O})(\mathbf{X}) = e_k(\mathcal{O}(\mathbf{X}))$.

Given the observable \mathcal{O} , the reference measure ρ and the family \mathcal{E} , we introduce $\mathcal{P}_{\mathcal{E}}$ the orthogonal projection (in $L^2(\rho)$) on the closure of the space spanned by the set $\{E_k, k \in \mathcal{I}\}$. For $\phi : \mathbb{R}^{2D} \rightarrow \mathbb{R}^d$, this orthogonal projection is formally given by

$$\mathcal{P}_{\mathcal{E}}\phi = \sum_{k \in \mathcal{I}} \left(\sum_{k' \in \mathcal{I}} G_{k,k'}^{-1} \langle E_{k'}, \phi \rangle \right) E_k \quad (5)$$

where G^{-1} the inverse of the Gram matrix of the basis, namely $G_{k,k'} = \langle E_k, E_{k'} \rangle$. For instance, when \mathcal{E} is reduced to a set of linear functions, this projector is also known as the *Mori* projector³³. By contrast, when \mathcal{E} is chosen as an Hilbert basis of $L^2(\mathbb{R}^d)$ (i.e. a sequence of functions whose linear combinations can approximate arbitrarily well any function in $L^2(\mathbb{R}^d)$), then this projector is known as the *Zwanzig* projector^{12,24}. Lastly for \mathcal{E} a finite family of function, $\mathcal{P}_{\mathcal{E}}$ is a *finite-rank* projector^{8,31}. We refer to App. A for details on projection operators.

B. The general Mori-Zwanzig derivation

For $\mathbf{X} \in \mathbb{R}^{2D}$, we denote by $\mathbf{X}(t)$ the flow associated to (3) with initial condition \mathbf{X} . It holds

$$\partial_t \mathcal{O}(\mathbf{X}(t)) = \mathcal{L}\mathcal{O}(\mathbf{X}(t)) = e^{t\mathcal{L}}\mathcal{L}\mathcal{O}(\mathbf{X}) = e^{t\mathcal{L}}\mathcal{P}_\varepsilon\mathcal{L}\mathcal{O}(\mathbf{X}) + e^{t\mathcal{L}}(1 - \mathcal{P}_\varepsilon)\mathcal{L}\mathcal{O}(\mathbf{X}).$$

Using the following Duhamel-Dyson formula:

$$e^{t\mathcal{L}} = e^{t(1-\mathcal{P}_\varepsilon)\mathcal{L}} + \int_0^t e^{(t-s)\mathcal{L}}\mathcal{P}_\varepsilon\mathcal{L}e^{s(1-\mathcal{P}_\varepsilon)\mathcal{L}}ds$$

and introducing the notation

$$\xi_t(\mathbf{X}) = e^{t(1-\mathcal{P}_\varepsilon)\mathcal{L}}(1 - \mathcal{P}_\varepsilon)\mathcal{L}\mathcal{O}(\mathbf{X}),$$

this first equation can be rewritten as

$$\begin{aligned} \partial_t \mathcal{O}(\mathbf{X}(t)) &= e^{t\mathcal{L}}\mathcal{P}_\varepsilon\mathcal{L}\mathcal{O}(\mathbf{X}) + \int_0^t e^{(t-s)\mathcal{L}}\mathcal{P}_\varepsilon\mathcal{L}\xi_s(\mathbf{X})ds + \xi_t(\mathbf{X}) \\ &= \mathcal{P}_\varepsilon\mathcal{L}\mathcal{O}(\mathbf{X}(t)) + \int_0^t \mathcal{P}_\varepsilon\mathcal{L}\xi_s(\mathbf{X}(t-s))ds + \xi_t(\mathbf{X}). \end{aligned} \quad (6)$$

From Eq. (5), we introduce the coefficients f_k and for $s \geq 0$ the time-dependent coefficients $g_k(s)$ such that

$$\mathcal{P}_\varepsilon\mathcal{L}\mathcal{O} = \sum_{k \in \mathcal{I}} f_k E_k, \quad \mathcal{P}_\varepsilon\mathcal{L}\xi_s = \sum_{k \in \mathcal{I}} g_k(s) E_k. \quad (7)$$

The projected evolution equation is then, writing $\mathcal{O}(t) = \mathcal{O}(\mathbf{X}(t))$,

$$\begin{aligned} \frac{\partial \mathcal{O}(t)}{\partial t} &= \sum_k f_k e_k(\mathcal{O}(t)) + \int_0^t \sum_k g_k(s) e_k(\mathcal{O}(t-s))ds + \xi_t(\mathbf{X}) \\ &:= f(\mathcal{O}(t)) + \int_0^t g(s, \mathcal{O}(t-s))ds + \xi_t(\mathbf{X}), \end{aligned} \quad (8)$$

with $f = \sum_k f_k e_k$ and $g(s, \cdot) = \sum_k g_k(s) e_k$ for $s \geq 0$. This is the generic form of a GLE where on the right hand side, the first term is known as the mean force term, the second term is related to the memory and the last term $\xi_t(\mathbf{X})$, which is interpreted as a noise, follows the orthogonal dynamics³¹

$$\partial_t \xi_t(\mathbf{X}) = (1 - \mathcal{P}_\varepsilon)\mathcal{L}\xi_t(\mathbf{X}) \quad \text{with} \quad \xi_0(\mathbf{X}) = (1 - \mathcal{P}_\varepsilon)\mathcal{L}\mathcal{O}(\mathbf{X}). \quad (9)$$

Notice that (8) is an exact equality. The approximation comes afterward when, in order to withdraw the orthogonal degrees of freedom and close the equation without solving the orthogonal dynamics, $\xi_t(\mathbf{X})$ is replaced by a random noise ζ_t independent from $\mathcal{O}(0)$.

The key property of $\xi_t(\mathbf{X})$, under which relies this approximation, is that it is by design uncorrelated to $e_k(\mathcal{O}(0))$ for all $k \in \mathcal{I}$. Indeed, assuming that \mathbf{X} is a random variable distributed according to ρ , we have for all $k \in \mathcal{I}$

$$\mathbb{E}(e_k(\mathcal{O}(0)) \cdot \xi_t(\mathbf{X})) = \langle E_k, \xi_t \rangle = 0, \quad (10)$$

and if for instance all constant functions are in the span of $\{e_k, k \in \mathcal{I}\}$, we obtain that the average of $\xi_t(\mathbf{X})$ with respect to ρ is zero.

The validity in some asymptotic regimes of the approximation obtained by replacing $\xi_t(\mathbf{X})$ by an independent noise can be shown in some cases, see e.g.³⁴⁻³⁶. Gaussian processes are then naturally motivated by the central limit theorem, although real applications don't always have particular reasons to fall in this regime³⁷⁻³⁹. We won't discuss further this point from a theoretical point of view.

Taking the scalar product of (8) with E_ℓ for $\ell \in \mathcal{I}$, the orthogonality condition (10) yields⁴⁰

$$\left\langle E_\ell(0), \frac{\partial \mathcal{O}(t)}{\partial t} \right\rangle = \sum_k f_k \langle E_\ell(0), E_k(t) \rangle + \int_0^t \sum_k g_k(s) \langle E_\ell(0), E_k(t-s) \rangle ds, \quad (11)$$

where $E_k(t) = e^{t\mathcal{L}} E_k$. Therefore, we obtain a set of Volterra integral equations of the first kind. When the set \mathcal{I} is finite, they can be numerically inverted using a time discretization of the integral as illustrated in sec. III. A set of Volterra equations of the second kind can also be derived as explained in the appendix E.

C. Hamiltonian systems

Let us now focus on Hamiltonian systems as it is the main case of interest. We consider a system of particles with position \mathbf{q} and momentum \mathbf{p} in \mathbb{R}^D . The Hamiltonian is $\mathbf{p} \cdot \mathbf{M}^{-1} \cdot \mathbf{p}/2 + V(\mathbf{q})$, with V the potential energy and \mathbf{M} the mass matrix. The evolution operator (3) is then given for any function $\phi(\mathbf{q}, \mathbf{p})$ of position and momentums by

$$\mathcal{L}\phi(\mathbf{q}, \mathbf{p}) = \mathbf{p} \cdot \mathbf{M}^{-1} \cdot \nabla_{\mathbf{q}}\phi(\mathbf{q}, \mathbf{p}) - \nabla_{\mathbf{q}}V(\mathbf{q}) \cdot \nabla_{\mathbf{p}}\phi(\mathbf{q}, \mathbf{p})$$

and the reference measure is given by the equilibrium Gibbs measure

$$\rho_{eq}(\mathbf{q}, \mathbf{p}) = e^{-\beta V(\mathbf{q}) - \beta \mathbf{p} \cdot \mathbf{M}^{-1} \cdot \mathbf{p}/2} / Z$$

with Z a normalisation factor and $\beta = 1/k_B T$ the inverse temperature of the system. Integrating by parts, it is well-known that \mathcal{L} is skew-symmetric in $L^2(\rho_{eq})$, namely

$$\langle \mathcal{L}\phi, \psi \rangle = -\langle \phi, \mathcal{L}\psi \rangle \quad (12)$$

for all functions ϕ, ψ . In particular, applying this with $\psi = 1$ so that $\mathcal{L}\psi = 0$, we get that ρ_{eq} is left invariant by the Hamiltonian dynamics.

In view of the general framework described in the previous section, let us now present the specific choices made in the present work, motivated by the specific form of GLE we want to derive. First, we work with a particular choice of observable. Decomposing the microscopic state in its position and momentum coordinates, namely $\mathbf{X} = (\mathbf{q}, \mathbf{p})$, we consider a first observable $\mathcal{O}_x(\mathbf{X}) = \varphi(\mathbf{q})$ for some collective variable $\varphi : \mathbb{R}^D \rightarrow \mathbb{R}^n$, and a second observable $\mathcal{O}_v(\mathbf{X}) = \mathcal{L}\mathcal{O}_x(\mathbf{X}) = \mathbf{p} \cdot \mathbf{M}^{-1} \cdot \nabla_{\mathbf{q}}\varphi(\mathbf{q})$. Then our observable is $\mathcal{O}(\mathbf{X}) = (\mathcal{O}_x(\mathbf{X}), \mathcal{O}_v(\mathbf{X}))$ of dimension $d = 2n$. We call \mathcal{O}_x the position observable and \mathcal{O}_v the velocity observable. Notice that, by design,

$$\partial_t \mathcal{O}_x(\mathbf{X}(t)) = \mathcal{O}_v(\mathbf{X}(t)) = \mathbf{p}(t) \cdot \mathbf{M}^{-1} \cdot \nabla_{\mathbf{q}}\varphi(\mathbf{q}(t)). \quad (13)$$

From this choice of observable, the GLE (8) is now written as

$$\partial_t \begin{pmatrix} \mathcal{O}_x(t) \\ \mathcal{O}_v(t) \end{pmatrix} = \begin{pmatrix} f^x(\mathcal{O}_x(t), \mathcal{O}_v(t)) \\ f^v(\mathcal{O}_x(t), \mathcal{O}_v(t)) \end{pmatrix} + \int_0^t \begin{pmatrix} g^x(s, \mathcal{O}_x(t-s), \mathcal{O}_v(t-s)) \\ g^v(s, \mathcal{O}_x(t-s), \mathcal{O}_v(t-s)) \end{pmatrix} ds + \begin{pmatrix} \xi_t^x(\mathbf{X}) \\ \xi_t^v(\mathbf{X}) \end{pmatrix}, \quad (14)$$

where f^x and f^v are the mean force terms for position and velocity respectively, g^x and g^v the memory parts and ξ^x and ξ^v the noise on position and velocity respectively. It remains to choose a set of functions to perform the projection. Given $\{h_k, k \in \mathcal{J}\}$ a family of functions from \mathbb{R}^n to \mathbb{R}^n , we consider the family of basis functions from \mathbb{R}^{2n} to \mathbb{R}^{2n} given by

$$\mathcal{E} = \left\{ a : \begin{pmatrix} o_1 \\ o_2 \end{pmatrix} \mapsto \begin{pmatrix} o_2 \\ 0 \end{pmatrix} \right\} \cup \left\{ b_k : \begin{pmatrix} o_1 \\ o_2 \end{pmatrix} \mapsto \begin{pmatrix} 0 \\ h_k(o_1) \end{pmatrix} \right\}_{k \in \mathcal{J}} \cup \left\{ c_k : \begin{pmatrix} o_1 \\ o_2 \end{pmatrix} \mapsto \begin{pmatrix} 0 \\ \nabla h_k(o_1) o_2 \end{pmatrix} \right\}_{k \in \mathcal{J}},$$

where ∇h_k stands for the Jacobian matrix of h_k .

Let us now comment this particular choice. Our family is divided into three sets of functions $\mathcal{E} = \{a\} \cup \{b_k, k \in \mathcal{J}\} \cup \{c_k, k \in \mathcal{J}\}$. The singleton $\{a\}$ is related to the fact $\partial_t \mathcal{O}_x(t) = \mathcal{O}_v(t)$. The second set $\{b_k, k \in \mathcal{J}\}$ is introduced to model the position dependency of the mean force term for the velocity. The last set $\{c_k, k \in \mathcal{J}\}$ is required to obtain the desired form of the memory part (i.e. linear in velocity). Notice that the functions c_k are related to the functions b_k by the fact that $\partial_t h_k(\mathcal{O}_x(t)) = \nabla h_k(\mathcal{O}_x(t)) \mathcal{O}_v(t)$ hence $\partial_t b_k(\mathcal{O}_x(t), \mathcal{O}_v(t)) = c_k(\mathcal{O}_x(t), \mathcal{O}_v(t))$.

As in the general framework, we write $A = a \circ \mathcal{O}$, $B_k = b_k \circ \mathcal{O}$ and $C_k = c_k \circ \mathcal{O}$.

The three parts of \mathcal{E} correspond to subspaces of $L^2(\rho)$ which are orthogonal one with each other. Indeed the scalar products $\langle A, B_k \rangle$ and $\langle A, C_k \rangle$ are both zero, and

$$\langle B_k, C_{k'} \rangle = \int_{\mathbb{R}^D} h_k(\varphi(\mathbf{q})) \cdot \nabla h_{k'}(\varphi(\mathbf{q})) \left(\int_{\mathbb{R}^D} \mathbf{p} \rho_{eq}(\mathbf{q}, \mathbf{p}) d\mathbf{p} \right) \cdot \nabla \varphi(\mathbf{q}) d\mathbf{q} = 0,$$

where we used (13) and that the average of the momentum under ρ_{eq} is zero. As a consequence, in the representation (5) of the orthogonal projection, the Gram matrix and its inverse have three diagonal blocks, corresponding to the three parts of \mathcal{E} . We introduce G_b (resp. G_c) the Gram matrix of the family $\{b_k, k \in \mathcal{J}\}$ (resp. $\{c_k, k \in \mathcal{J}\}$).

Finally, we also define for any function f and g of the observable, the scalar product

$$\langle f, g \rangle_\varphi = \langle f \circ \mathcal{O}_x, g \circ \mathcal{O}_x \rangle = \int_{\mathbb{R}^n} f(z)^T \cdot g(z) \rho_\varphi(z) dz$$

where $\rho_\varphi : \mathbb{R}^n \rightarrow \mathbb{R}_+$ is the marginal density of $\rho_{eq}(\mathbf{q}, \mathbf{p})$, namely is such that

$$\int_{\mathbb{R}^{2D}} g(\varphi(\mathbf{q}, \mathbf{p})) \rho_{eq}(\mathbf{q}, \mathbf{p}) d\mathbf{q} d\mathbf{p} = \int_{\mathbb{R}^n} g(z) \rho_\varphi(z) dz$$

for all function g of the observable, see also appendix B for a definition using the coarea formula. When f and g are matrix-valued function, the scalar product is defined as

$$\langle f, g \rangle_\varphi = \sum_{i,j} \int_{\mathbb{R}^n} f(z)_{i,j} g(z)_{i,j} \rho_\varphi(z) dz.$$

D. Continuing the derivation

Equipped with our choice of observables and basis, we can now make explicit the form of the mean force and the memory kernel in (8), namely the coefficients f_k and $g_k(s, \cdot)$ in (7). The projection of the position evolution equation follows trivially from Eq.(13) and the presence of the singleton $\{a\}$ in our choice of basis. Using the notations of (14), we get $f^x(o_1, o_2) = o_2$ and $g^x(s, \cdot) = \xi_s^x = 0$ for all $s \geq 0$.

In the following we focus on the second line of (14), namely the projection of the velocity evolution equation. From (7) and the explicit expression of the projector (5), the question is thus to compute the scalar products of $\mathcal{L}\mathcal{O}_v$ and of $\mathcal{L}\xi_s^v = \mathcal{L}e^{s(1-\mathcal{P}_\varepsilon)}\mathcal{L}\mathcal{O}_v$ with B_k and C_k (here and in the following, with a slight abuse of notations, we denote by B_k and C_k the second n -dimensional component of B_k and C_k).

For all $k \in \mathcal{J}$,

$$\begin{aligned} \langle C_k, \mathcal{L}\mathcal{O}_v \rangle &= \int_{\mathbb{R}^{2D}} [\nabla h_k(\varphi(\mathbf{q})) (\mathbf{p} \cdot \mathbf{M}^{-1} \cdot \nabla \varphi(\mathbf{q}))] \\ &\quad \cdot [-\nabla V(\mathbf{q}) \cdot \nabla \varphi(\mathbf{q}) + \mathbf{p} \cdot \mathbf{M}^{-1} \cdot \nabla^2 \varphi(\mathbf{q}) \mathbf{p}] \rho_{eq}(\mathbf{q}, \mathbf{p}) d\mathbf{q} d\mathbf{p} = 0, \end{aligned} \quad (15)$$

where we used that the integrand is odd in \mathbf{p} and ρ_{eq} is invariant by reflection of the momentum. The mean force term f^v in (14) is then

$$f^v(o_1, o_2) = \sum_{k \in \mathcal{J}} \left(\sum_{k' \in \mathcal{J}} (G_b^{-1})_{k,k'} \langle B_{k'}, \mathcal{L}\mathcal{O}_v \rangle \right) h_k(o_1) =: f_b(o_1). \quad (16)$$

In general, this mean force f_b is non-linear (if $\{h_k, k \in \mathcal{J}\}$ contains non-linear functions on \mathbb{R}^n). It can be build from the potential of mean force W_{pmf} and an effective mass matrix M_φ . The potential of mean force is the function $W_{\text{pmf}} = -\frac{1}{\beta} \ln \rho_\varphi$. The effective mass matrix is defined as the conditional expectation

$$\begin{aligned} M_\varphi(z)^{-1} &= \beta \mathbb{E} (\mathcal{O}_v(\mathbf{q}, \mathbf{p}) \mathcal{O}_v(\mathbf{q}, \mathbf{p})^T \mid \varphi(\mathbf{q}) = z) \\ &= \mathbb{E} (\nabla \varphi(\mathbf{q}) \cdot \mathbf{M}^{-1} \cdot \nabla \varphi(\mathbf{q})^T \mid \varphi(\mathbf{q}) = z), \end{aligned} \quad (17)$$

where the second equality follows from $\int_{\mathbb{R}^D} \mathbf{p} \mathbf{p}^T \rho_{eq}(\mathbf{q}, \mathbf{p}) d\mathbf{p} = \rho_{eq}(\mathbf{q}) \mathbf{M} / \beta$. Here, M_φ is interpreted as an effective mass in view of the equipartition theorem. It is then detailed in the appendix C that

$$\langle B_k, \mathcal{L}\mathcal{O}_v \rangle = \langle B_k, \Psi \circ \mathcal{O}_x \rangle = \langle h_k, \Psi \rangle_\varphi \quad (18)$$

with $\Psi : \mathbb{R}^n \rightarrow \mathbb{R}^n$

$$\Psi(z) = \frac{1}{\beta} (\nabla \cdot M_\varphi(z)^{-1} + M_\varphi(z)^{-1} \cdot \nabla \ln \rho_\varphi(z)). \quad (19)$$

If Ψ is in the space spanned by $\{h_k, k \in \mathcal{J}\}$, then (18) implies that $f_b = \Psi$. Otherwise, f_b is a projection of Ψ on this space (this is typically the case in practice since a finite set of basis functions is used).

When the effective mass matrix is independent of the position, we get

$$\Psi(z) = \frac{1}{\beta} M_\varphi^{-1} \cdot \nabla \ln \rho_\varphi(z) = -\frac{1}{\beta} M_\varphi^{-1} \cdot \nabla W_{\text{pmf}}(z). \quad (20)$$

For instance, this holds if $n = 1$ and the reaction coordinate is a distance between two atoms, or for linear reaction coordinates (taking $\varphi_i(\mathbf{q}) = e_i \cdot \mathbf{q}$ with e_1, \dots, e_n an orthonormal basis of the range of the reaction coordinate).

We now turn to the projection of $\mathcal{L}\xi_s^v$. Using the skew-symmetry of \mathcal{L} , for all $k \in \mathcal{J}$,

$$\langle B_k, \mathcal{L}\xi_s^v \rangle = -\langle \mathcal{L}B_k, \xi_s^v \rangle = -\langle C_k, \xi_s^v \rangle = 0$$

since the noise is orthogonal to any function in \mathcal{E} . Therefore the memory part of the equation is (using again the skew-symmetry of \mathcal{L})

$$g_s^v(o_1, o_2) = -\sum_{k \in \mathcal{I}} \left(\sum_{k' \in \mathcal{I}} (G_c^{-1})_{k,k'} \langle \mathcal{L}C_{k'}^v, \xi_s^v \rangle \right) \nabla h_k(o_1) o_2 =: -K_b(s, o_1) o_2. \quad (21)$$

The value of the scalar product $\langle \mathcal{L}C_k, \xi_s^v \rangle$ is discussed in sec. II E. To follow the usual convention, we refer to $K_b(s, o_1)$ as the *memory kernel*.

Combining together all these elements, we get

$$\begin{cases} \partial_t \mathcal{O}_x(t) &= \mathcal{O}_v(t) \\ \partial_t \mathcal{O}_v(t) &= f_b(\mathcal{O}_x(t)) - \int_0^t K_b(s, \mathcal{O}_x(t-s)) \mathcal{O}_v(t-s) ds + \xi_t^v(\mathbf{X}). \end{cases} \quad (22)$$

Closing this equation by approximating $\xi_t^v(\mathbf{X})$ by a random noise ζ_t we get, as announced, a GLE for the position $x_t = \mathcal{O}_x(t)$ of the form:

$$\ddot{x}_t = f_b(x_t) - \int_0^t K_b(s, x_{t-s}) \dot{x}_{t-s} ds + \zeta_t.$$

Up to now, we have left unspecified the set $\{h_k, k \in \mathcal{J}\}$. Several possibilities are available. First, for instance, we can choose $\{h_k, k \in \mathcal{J}\}$ as a basis of the linear and constant functions (*i.e.* the Mori projection), in which case we retrieve the linear form of the GLE

$$\ddot{x}_t = -\omega(x_t - y_0) - \int_0^t K_M(s) \dot{x}_{t-s} ds + \zeta_t, \quad (23)$$

with ω and y_0 some constants characterizing the force.

Second, an alternative is to take a basis of $L^2(\rho, \mathbb{R}^d)$ (*i.e.* the Zwanzig projection). Assuming position-independent effective mass matrix, the GLE then reads

$$\ddot{x}_t = -M_\varphi^{-1} \cdot \nabla W_{\text{pmf}}(x_t) - \int_0^t K_Z(s, x_{t-s}) \dot{x}_{t-s} ds + \zeta_t, \quad (24)$$

where the mean force term is now the derivative of the potential of mean force. More generally, the mean force is the conditional expectation Ψ .

Third, if the mean force term is known a priori (*e.g.* when the potential of mean force is computed apart), then we can choose the set $\{h_k, k \in \mathcal{J}\}$ to be reduced to the singleton $\{\nabla W_{\text{pmf}}\}$, which yields for position-independent mass matrix

$$\ddot{x}_t = -M_\varphi^{-1} \cdot \nabla W_{\text{pmf}}(x_t) - \int_0^t K_{\text{eff}}(s) \nabla^2 W_{\text{pmf}}(x_{t-s}) \dot{x}_{t-s} ds + \zeta_t. \quad (25)$$

This choice is somehow the minimal set-up to get a mean force which derives from ρ_φ . In that case, the dependency in position of the memory kernel is completely fixed via the Hessian of the potential of mean force. We recover the initial form (1) when this Hessian is approximately constant over the studied range of position (i.e. close to an harmonic case), and only in this case.

Notice, that if a gradient form of the mean force is wanted, we can take $h_k = \nabla \ell_k$ where $\{\ell_k, k \in \mathcal{J}\}$ is an Hilbert basis of the Sobolev space $\mathcal{H}^1(\mathbb{R}^n)$. In that case f_b is an Helmholtz projection of the mean force obtained with the Zwanzig projection.

In both cases (24) and (25), as required in Ref. 24, the mean force derives from ρ_φ and the memory kernel is linear in the velocity (although with a non-linear dependency in the position). Indeed, linear reaction coordinates are considered in Ref. 24, which means a position independent effective mass matrix, possibly up to a linear change of variable.

To conclude this section, let us highlight a slight subtlety: in the general derivation of sec. II B, we assumed the functions e_k to be linearly independent. In fact from a theoretical point of view this is not important since the only thing that matters in the definition of the projection is the space spanned by $\{E_k, k \in \mathcal{I}\}$. However, from a practical point of view, this is used when considering the inverse of the Gram matrix in the explicit formula (5). Then, with the specific choice of basis functions described in sec. II C, it can be convenient to take $\{h_k, k \in \mathcal{J}\}$ which contains some constant functions (in particular to have a mean-zero noise, see next section), in which case the corresponding c_k is zero (since $\nabla h_k = 0$), which means the functions $E_k, k \in \mathcal{I}$, are not linearly independent. Again, this is not important for the theoretical derivation, but it means that, in practice, one should be careful when applying (5) and possibly use the pseudo-inverse of the Gram matrix or reduce the set of functions.

E. Characteristics of the noise

Once the mean force and memory kernel coefficients have been estimated, the standard way to close the equation (8) is to replace $\xi_t(\mathbf{X})$ by a Gaussian process ζ_t with the same mean and variance. However, this approximation may in fact not be suitable in general^{37–39}. In any cases, let us compute these characteristics. In the following, we consider the specific case introduced in sec. II C and such we are only interested to the properties of the noise ξ_t^v in the second line of of (22).

1. Mean value

Since $\xi_t(\mathbf{X})$ is replaced in practice by a random process which is independent from $\mathcal{O}(0)$, a minima, it should be enforced that $\xi_t(\mathbf{X})$ is not correlated to $e_k(\mathcal{O}(0))$ for $k \in \mathcal{I}$. As we saw, this is true if the average of $\xi_t(\mathbf{X})$ is 0 under ρ .

The simplest way to enforce this is to assume that all constant functions are in the span of $\{e_k, k \in \mathcal{I}\}$, in which case the orthogonality condition (10) concludes. Since we only consider ξ_t^v , the previous condition is satisfied as soon as all constant functions from \mathbb{R}^n to \mathbb{R}^n are in the space spanned by $\{h_k, k \in \mathcal{J}\}$. In which case we obtain $\langle \xi_t^v \rangle = 0$.

2. Variance and 2-FDT

For GLEs, a Fluctuation-Dissipation Theorem of second type (2-FDT) is a relation which expresses the covariance function of the noise $\xi_t(\mathbf{X})$ in terms of the memory kernel K . It has been established in very general frameworks for *linear* cases^{6,30,39}. In fact, although (8) may be non-linear in general, it is always based on a linear projection, and we can always write a linear GLE for the vector $(e_k(\mathcal{O}(t)))_{k \in \mathcal{I}}$. Hence, a 2-FDT holds for this (possibly infinite-dimensional) equation, which implies that a 2-FDT holds for (8) if for instance the identity function is in the span of $\mathcal{E} = \{e_k, k \in \mathcal{I}\}$.

This is true if the identity function on \mathbb{R}^n is in the span of $\{h_k, k \in \mathcal{J}\}$. In fact, without loss of generality (up to a change of basis), in that case we simply assume that $h_0(o_1) = o_1$, in which case the corresponding c_0 is exactly the required function. Then, using the skew-symmetry of \mathcal{L} and the orthogonality (10) of the noise,

$$\begin{aligned} \langle C_0, \mathcal{L}\xi_s^v \rangle &= -\langle \mathcal{L}C_0, \xi_s^v \rangle = -\langle \mathcal{L}\mathcal{O}_v, \xi_s^v \rangle \\ &= -\langle (1 - \mathcal{P}_{\mathcal{E}})\mathcal{L}\mathcal{O}_v, \xi_s^v \rangle = -\langle \xi_0^v, \xi_s^v \rangle. \end{aligned}$$

The left-hand side is a coefficient involved in the decomposition (21) of K . More precisely, if the basis function has been orthonormalized, this is exactly the time-dependent coefficient of C_0

If the ambient space dimension n is larger than 1, then we need the full covariance matrix of the noise, i.e. $\langle (\xi_0^v)_i, (\xi_s^v)_j \rangle$ for $1 \leq i, j \leq n$. The previous computation is still correct if we assume now that all the linear functions on \mathbb{R}^n are in the span of $\{h_k, k \in \mathcal{J}\}$. In that case, denoting by $M_{i,j}$ the matrix with all coefficients 0 except the coefficient (i, j) equal to 1, $h_{i,j} : o_1 \mapsto M_{i,j}o_1$ and the corresponding $C_{i,j} : \mathbf{X} \mapsto M_{i,j}\mathcal{O}_v(\mathbf{X})$, we get again

$$\langle C_{i,j}, \mathcal{L}\xi_s \rangle = -\langle (\xi_0^v)_i, (\xi_s^v)_j \rangle.$$

As a conclusion, in this case, we can say that a 2-FDT holds in the sense that the covariance matrix of the noise in (22) is determined by some particular coefficients of the kernel K in (21). In practice, this means that the covariance of the noise can be computed through the Volterra equations (11).

3. Position-dependent average and 2-FDT

The fact that $\xi_t(\mathbf{X})$ is uncorrelated to $e_k(\mathcal{O}(0))$ for all $t \geq 0$ and $k \in \mathcal{I}$ gives little information on a possible dependency of the law of $\xi_t(\mathbf{X})$ with respect to $\mathcal{O}(t)$. In particular, if we use the covariance function computed in the previous section to approximate $\xi_t(\mathbf{X})$ by a Gaussian noise ζ_t with the same characteristics, then the law of ζ_t does not depend on $\mathcal{O}(t)$. This approximation may not be relevant, having in mind some alternative ways to derive an effective dynamics^{3,41}. We discuss in the following some results about the position dependence average and variance of the noise.

Let us introduce the conditional average of the noises $\bar{\xi}_t$ and $\bar{\xi}_t^z$ given by

$$\bar{\xi}_t(z) = \mathbb{E}(\xi_t^v(\mathbf{q}, \mathbf{p}) \mid \varphi(\mathbf{q}) = z), \quad (26)$$

and

$$\bar{\xi}_t^z(z) = \mathbb{E}(\xi_t^v(\mathbf{q}, \mathbf{p})\mathcal{O}_v(\mathbf{q}, \mathbf{p})^T \mid \varphi(\mathbf{q}) = z). \quad (27)$$

Eq. (10) induces on $\bar{\xi}_t$ and $\bar{\xi}_t^z$ the set of constraints:

$$\forall k \in \mathcal{J}, \quad \langle h_k, \bar{\xi}_t \rangle_\varphi = 0 \quad \text{and} \quad \langle \nabla h_k, \bar{\xi}_t^z \rangle_\varphi = 0. \quad (28)$$

Similarly consider $\sigma_\varphi(t, z)$ the conditional variance of the noise given by

$$\sigma_\varphi(t, z) = \mathbb{E} \left(\xi_0^v(\mathbf{q}, \mathbf{p}) \xi_t^v(\mathbf{q}, \mathbf{p})^T \mid \varphi(\mathbf{q}) = z \right). \quad (29)$$

As detailed in the appendix C, we can decompose the memory kernel K_b as

$$K_b(t, o_1) = K_\sigma(t, o_1) + K_{\bar{\xi}}(t, o_1), \quad (30)$$

where $K_\sigma(t, o_1)$ and $K_{\bar{\xi}}(t, o_1)$ are defined by

$$K_\sigma(t, o_1) = \sum_{k \in \mathcal{I}} \left(\sum_{k' \in \mathcal{I}} (G_c^{-1})_{k, k'} \langle \nabla h_{k'}, \sigma_\varphi(t, \cdot) \rangle_\varphi \right) \nabla h_k(o_1) \quad (31)$$

and

$$K_{\bar{\xi}}(t, o_1) = \sum_{k \in \mathcal{I}} \left(\sum_{k' \in \mathcal{I}} (G_c^{-1})_{k, k'} \langle \nabla h_{k'}, f_b \bar{\xi}_t - \nabla \cdot (\bar{\xi}_t^z \rho_\varphi) / \rho_\varphi \rangle_\varphi \right) \nabla h_k(o_1), \quad (32)$$

with f_b is the mean force of Eq.(16). To obtain a position-dependent 2-FDT, namely a pointwise relation between K_b and σ , $K_{\bar{\xi}}$ should cancel. This is not the case in general. Indeed in (32), the terms multiplying the conditional average of the noise do not necessary belong to the space spanned by our function set and, as a consequence, the conditions (28) cannot be used to prove that this term is zero. This fact is illustrated in sec. III B.

However, $K_{\bar{\xi}}(t, o_1)$ is indeed zero when the set $\{h_k, k \in \mathcal{J}\}$ is a basis of $L^2(\rho, \mathbb{R}^d)$ (*i.e.* in the case of the Zwanzig projection $K_b = K_Z$). From the conditions (28), we get

$$\forall z \in \mathbb{R}^d, \quad \bar{\xi}_t(z) = 0 \quad \text{and} \quad \bar{\xi}_t^z(z) = 0,$$

from which $K_{\bar{\xi}}(t, o_1) = 0$. We then get a position-dependent 2-FDT from the effective mass matrix and conditionnal variance of the noise,

$$K_Z(t, o_1) = \beta M_\varphi(o_1) \cdot \sigma_\varphi(t, o_1)$$

where K_Z is the memory kernel of Eq. (24).

Finally, when the Mori projection is used, we emphasize that $K_{\bar{\xi}}(t, o_1)$ is also zero and we recover the result of sec. II E 2.

F. Other forms of GLE

The particular choices of observable and of function set made in sec. II C lead to the GLE (22), but other choices are possible. In fact, a priori, although it may be appealing from a physical point of view, there is no reason for the best model of approximation of an effective dynamics to be a position/velocity GLE with a mean force term deriving from the potential of mean force and a friction kernel which is linear in the velocity (which was the motivation of these choices). Here are some other possibilities.

The form of the observable in Ref. 24 is similar to the one in sec. II C, but the function set is chosen as

$$\mathcal{E}_{GS} = \left\{ a : \begin{pmatrix} o_1 \\ o_2 \end{pmatrix} \mapsto \begin{pmatrix} o_2 \\ 0 \end{pmatrix} \right\} \cup \left\{ f_k : \begin{pmatrix} o_1 \\ o_2 \end{pmatrix} \mapsto \begin{pmatrix} 0 \\ f_k(o_1, o_2) \end{pmatrix} \right\}_{k \in \mathcal{J}} \quad (33)$$

where $(f_k)_{k \in \mathcal{J}}$ is a basis of $L^2(\mathbb{R}^{2d})$. This results in a GLE of the form

$$\ddot{x}_t = F_e(x_t) - \int_0^t K_{GS}(s, x_{t-s}, \dot{x}_{t-s}) ds + \zeta_t, \quad (34)$$

where the memory is no longer linear in the velocity.

In contrast, we could have chosen a smaller function set containing only the two first families of sec. II C, namely

$$\mathcal{E}_{NV} = \{a\} \cup \{b_k, k \in \mathcal{J}\}.$$

In that case, we obtain a GLE of the form

$$\ddot{x}_t = F_e(x_t) - \int_0^t K_{NV}(s, x_{t-s}) ds + \zeta_t, \quad (35)$$

where there is no dependence in velocity in the memory term.

A slightly larger function set is obtained by adding a function linear in velocity

$$\mathcal{E}_H = \{a\} \cup \{b_k, k \in \mathcal{J}\} \cup \left\{ \begin{pmatrix} o_1 \\ o_2 \end{pmatrix} \mapsto \begin{pmatrix} 0 \\ o_2 \end{pmatrix} \right\}.$$

This leads to the hybrid projector of Ref. 42. In which case the GLE is of the form

$$\ddot{x}_t = F_e(x_t) - \int_0^t K_H^x(s, x_{t-s}) ds - \int_0^t K_H^v(s) \dot{x}_{t-s} ds + \zeta_t. \quad (36)$$

Next, if we only consider the position observable, namely $\mathcal{O}(\mathbf{X}) = \mathcal{O}_x(\mathbf{X}) = \varphi(\mathbf{q})$, and the function set

$$\mathcal{E}_{ov} = \{h_k : o_1 \mapsto h_k(o_1)\},$$

this leads to an overdamped form of the GLE

$$\dot{x}_t = F_{ov}(x_t) - \int_0^t K_{ov}(s, x_{t-s}) ds + \zeta_t. \quad (37)$$

Similarly, one can consider only the velocity.

After that, let us emphasize that, although kinetic or overdamped equations may, again, be appealing from a physical point of view since many physical laws are associated to these familiar structures, there is a priori no reason to be restricted to equations of order 1 or 2. For instance, considering an observable $\mathcal{O} = (\mathcal{O}_x, \mathcal{O}_v, \mathcal{O}_a)$ with $\mathcal{O}_x(\mathbf{X}) = \varphi(\mathbf{q})$, $\mathcal{O}_v = \mathcal{L}\mathcal{O}_x$ and $\mathcal{O}_a = \mathcal{L}\mathcal{O}_v$ naturally yields a third order effective equation for \mathcal{O}_x , with a structure depending again on the choice of the basis functions.

Finally, in this work we considered basis functions which depend a priori of the whole observable, since we are looking for closed effective dynamics. However, when working on a pair of observables, nothing forbids the choice of a projection on a set of functions which depend only of one of the observables, or even which depend on a different observable than the one we are interested to.

III. VOLTERRA BASED ESTIMATION OF THE MEMORY KERNEL

A. Numerical inversion of the Volterra Integral Equation

Given an all-atom simulation $(\mathbf{X}(t))_{t \in [0, T]}$ (or more generally a set of independent simulations), a practical issue is to parametrize the GLE (2), namely to estimate the coefficients f_k and $g_k(s)$ in (8). This can be done using the Volterra equations (11), when the set \mathcal{E} is finite. Indeed, the averages involved in (11) can be estimated along the trajectory, and then the equations can be numerically inverted using a discretization of the time integral⁴³. In the following, we use the trapezoidal method of Ref. 43, we also point out that this trapezoidal rule suffers from oscillatory error⁴⁴ and requires a smoothing that we implement. The numerical implementation of the inversion of the Volterra equation used in the following is available at <https://github.com/HadrienNU/VolterraBasis> and its documentation is available at <https://volterrabasis.readthedocs.io>.

The inversion of Volterra equation of the first kind is an ill-conditioned problem^{22,45}. Regularization methods can be used to improve the quality of the algorithm but this falls out of scope of this work and we refer to the rich literature on the subject^{45–48}.

Notice that, apart from Volterra-based algorithms, other methods exist to estimate friction kernels^{38,49–51} and could be modified to tackle the position-dependent case. However, here we only consider the former for simplicity and since they are naturally related to the theoretical derivation of the previous section.

B. Lennard-Jones Dimer

In this section, we present some numerical experiments to illustrate this approach (a thorough numerical analysis, including a discussion on the statistical properties of the noise and how it can be generated, on the ability of the effective dynamics to reproduce the dynamics of the initial system, or on other inference methods as in Ref. 52, is out of the scope of this work). The present analysis is based on the molecular dynamics simulation of a Lennard-Jones dimer similar to the one used in our earlier work⁵². Position dependence for the memory kernel in this system has been emphasized in Ref. 53.

This is a 3D system composed of 2 Lennard-Jones (LJ) particles forming a dimer and 510 LJ particles constituting the solvent at reduced temperature $\hat{T} = k_B T / \epsilon = 1$ and reduced density $\hat{\rho} = \rho \sigma^3 = 1$. The two dimer particles interact via a LJ potential with parameters $\epsilon_d = 2$ and $\sigma_d = 1$ (in LJ units) and their distance r is constrained for distance above $r = 3.0$ using a parabolic wall of strength $\kappa = 200$. LJ parameters for dimer-solvent and solvent-solvent interaction are taken as $\epsilon = 1$ and $\sigma = 1$ (in LJ units). The size of the cubic simulation box is 8σ , with periodic boundary conditions in all directions. The dynamics is integrated with a time step of $\Delta t_{MD} = 10^{-3}$ (in LJ units) in the NVE ensemble using the LAMMPS simulation package⁵⁴. We run 500 trajectories with length of $2 \cdot 10^5$ timesteps and CV values are extracted every steps. The observable $\mathcal{O}_x(\mathbf{q})$ is the distance r between the two particles forming the dimer.

We use as set of functions $\{h_k, k \in \mathcal{J}\}$ a set of 14 spline functions of degree 3 with knots non uniformly distributed between $[0.889, 3.297]$ as represented in Fig. 1, the two bounds corresponding to the minimum and maximum value of the distance in the data. More splines function are put on the part of the potential of mean force profile with the most important variation. We refer in the following to this choice of function set as the *splines* GLE.




FIG. 1. (a) Potential of mean force computed from an histogram compared to the integral of the mean force term. We also represent all the function in the splines basis (thin colored line) with the position of the knots marked with black dots. (b) Memory kernel as a function of the distance and time. Horizontal and vertical lines are the position of the cross-section with the same color represented in Fig. 2.

We also study the *linear* GLE of (23) using as function set a linear function and the *minimal* GLE where we use as function set the the PMF as in Eq. (25).

We represent on Fig. 1(a) the potential of mean force obtained via an histogram of the distance or via the integration of the mean force computed via the projection of the acceleration on the spline basis, showing a good agreement between both. The position dependent memory kernel is represented as a 2D plot in Fig. 1(b) and some cross-sections are represented in Fig. 2 along time (a) and space (b).

Comparing cross-sections of the *splines* GLE with cross-sections of other GLEs emphasizes the importance of the basis choice. For the *minimal* GLE most of the memory is concentrated at the boundary (where the potential of mean force is singular) and even if the mean force term is the same than the *splines* GLE, the memory kernel is strongly different from the previous case. Interestingly, using the simple *linear* GLE reconstruct a memory kernel very similar to the one of the *splines* GLE.

Finally, we also study the noise. Indeed, once the mean force and memory kernel have been estimated, the noise can be computed from the molecular dynamics trajectory data by inversion of Eq. (22). Estimation of correlation function between the noise and any function can also be carried out as detailed in the appendix E. In the objective of generating new trajectories using the GLE coarse-grained dynamics, the noise should be modeled, usually as a stationary Gaussian correlated noise^{55–57}. Validity of such models could be then interpreted in the light of noise computed from the trajectory data.




FIG. 2. Cross section of the 2D plot of 1(b) along (a) time and (b) position. Solid line are the cross-section of the *splines* GLE and dashed lines the *minimal* memory kernel computed along the same cross-section. Color matches the lines on Fig. 1. The dot-dashed violet curve on panel(a) is the memory kernel for the *linear* GLE.

Histogram of the value of the noise are plotted on Fig. 3(a) showing similar behavior for all GLE but with non-gaussian tails, a feature that has been noticed on several systems^{37,38}. Notice that with the *splines* GLE, the noise is more Gaussian than with the *linear* GLE or the *minimal* GLE, which illustrates in these two cases the fact that more non-linear features of the dynamics have to be included in the noise.

On Fig. 3(b) we investigate the validity of the FDT by comparing the noise auto-correlation to the constant part of the kernel. The *splines* and *linear* GLE show a good agreement between noise autocorrelation and corresponding coefficient of the memory kernel. The memory kernel of the *minimal* GLE does not have a coefficient corresponding to the noise auto-correlation and is therefore not represented.

Following the discussion of sec. II E 3, the position dependence of the conditional expectation of the variance of the noise as a function of the position at $t = 0$ is plotted on Fig. 4(a). For each trajectories, we bin the value of the noise depending of the position using 50 bins. As expected, the variance of the noise shows a strong dependency in position.

Finally, we study the validity of the position-dependent 2-FDT for the *splines* GLE. We represent on Fig. 4(b) the comparison the memory at three different cross-section with the corresponding 2-FDT curve K_σ of Eq. (31). The disagreement between the curves is explained by the non-zero value of K_ξ in Eq. (30) as illustrated in Fig. 4(c). Indeed, in this case the splines does not form a complete basis of $L^2(\rho, \mathbb{R}^d)$.




FIG. 3. (a) Log of the noise histogram for the *splines* GLE (blue line), the *minimal* GLE (magenta dashed line) and the *linear* GLE (cyan dot dashed line). For comparison, a quadratic curve is fitted to the top of the histogram (dot-dashed yellow thin line). (b) Auto correlation of the noise for the *splines* GLE (blue line), the *minimal* GLE (magenta dashed line) and the linear GLE (cyan dot dashed line) compared to the coefficient $\langle C_0, \mathcal{L}\xi_s \rangle$ of the memory kernel for the spline GLE (blue squares) and the *linear* GLE (cyan circles).

CONCLUSION

We have shown how GLE with a non-linear mean force, a friction kernel linear in the velocity and a Fluctuation-Dissipation Theorem can be rigorously derived within the Mori-Zwanzig formalism, in the Hamiltonian equilibrium settings. However, in contrast to the standard assumption, this requires in general the friction kernel to depend on the position (as confirmed by the numerical experiments). Moreover, we have also emphasized that, even with the position-dependency, this particular form remains arbitrary to some extent, since various other forms may be similarly derived with a suitable choice of observable and of function basis for the projection. The complexity of the model can then be monitored by these choices, a better approximation requiring a larger set of functions, hence of parameters to estimate, leading to a higher sensitivity to the sampling variations. In any cases, the question of obtaining a good approximation cannot be reduced to the choice of the form of the GLE and the estimation of the effective force and the memory kernel: a correct representation of the noise is also necessary, which may require to go beyond space-homogeneous Gaussian processes. In this case, then, this representation cannot be extracted only from the Volterra equations and the 2-FDT, but should rely on a suitable statistical analysis.




FIG. 4. (a) Variance of the noise as a function of the position for the *splines* GLE (blue line), the *minimal* GLE (magenta dashed line) and the *linear* GLE (cyan dot dashed line). (b) Cross section of the memory kernel and $K_\sigma(t, r)$ along time. Solid line are the cross-section of the *splines* memory kernel and cross are the values of $K_\sigma(t, r)$ computed along the same cross-section. (c) Cross section of $K_\xi(t, r)$ along time for the *splines* GLE. For (b) and (c) color matches the lines on Fig. 1.

ACKNOWLEDGMENTS

We thank Tanja Schilling and Fabian Glatzel for enriching discussions on the subject as well as their useful comments on the manuscript. We thank Sara Bonella, Michele Casula, Arthur France-Lanord, Ludovic Goudenège, Fabio Pietrucci, Benjamin Rotenberg, Marco Saitta, Mathieu Salanne, and Rodolphe Vuilleumier for fruitful discussions within the MAE-STRO collaboration.

REFERENCES

- ¹C. Peter and K. Kremer, “Multiscale simulation of soft matter systems – from the atomistic to the coarse-grained level and back,” *Soft Matter* **5**, 4357–4366 (2009).
- ²H. Lei, B. Caswell, and G. E. Karniadakis, “Direct construction of mesoscopic models from microscopic simulations,” *Physical Review E* **81**, 026704 (2010).
- ³F. Legoll and T. Lelièvre, “Effective dynamics using conditional expectations,” *Nonlinearity* **23**, 2131–2163 (2010).
- ⁴W. Zhang, C. Hartmann, and C. Schütte, “Effective dynamics along given reaction coordinates, and reaction rate theory,” *Faraday Discussions* **195**, 365–394 (2017).
- ⁵T. Schilling, “Coarse-Grained Modelling Out of Equilibrium,” arXiv:2107.09972 [cond-mat] (2021), arXiv:2107.09972 [cond-mat].
- ⁶R. Zwanzig, *Nonequilibrium Statistical Mechanics* (Oxford University Press, 2001).
- ⁷A. J. Chorin, O. H. Hald, and R. Kupferman, “Optimal prediction and the Mori–Zwanzig representation of irreversible processes,” *Proceedings of the National Academy of Sciences* **97**, 2968–2973 (2000).
- ⁸A. J. Chorin, O. H. Hald, and R. Kupferman, “Optimal prediction with memory,” *Physica D: Nonlinear Phenomena* **166**, 239–257 (2002).
- ⁹L. Ma, X. Li, and C. Liu, “The derivation and approximation of coarse-grained dynamics from Langevin dynamics,” *The Journal of Chemical Physics* **145**, 204117 (2016).

- ¹⁰S.-H. Chung and M. Roper, “Generalized Langevin Equation: An Introductory Review for Biophysicists,” *Biophysical Reviews and Letters* **14**, 171–196 (2019).
- ¹¹E. Darve, J. Solomon, and A. Kia, “Computing generalized Langevin equations and generalized Fokker–Planck equations,” *Proceedings of the National Academy of Sciences* **106**, 10884–10889 (2009).
- ¹²S. Izvekov, “Microscopic derivation of particle-based coarse-grained dynamics,” *The Journal of Chemical Physics* **138**, 134106 (2013).
- ¹³C. Hijón, P. Español, E. Vanden-Eijnden, and R. Delgado-Buscalioni, “Mori–Zwanzig formalism as a practical computational tool,” *Faraday Discussions* **144**, 301–322 (2009).
- ¹⁴Z. Li, H. S. Lee, E. Darve, and G. E. Karniadakis, “Computing the non-Markovian coarse-grained interactions derived from the Mori–Zwanzig formalism in molecular systems: Application to polymer melts,” *The Journal of Chemical Physics* **146**, 014104 (2017).
- ¹⁵H. S. Lee, S.-H. Ahn, and E. F. Darve, “The multi-dimensional generalized Langevin equation for conformational motion of proteins,” *The Journal of Chemical Physics* **150**, 174113 (2019).
- ¹⁶R. Zwanzig, “Nonlinear generalized Langevin equations,” *Journal of Statistical Physics* **9**, 215–220 (1973).
- ¹⁷H. Mori, “A Continued-Fraction Representation of the Time-Correlation Functions,” *Progress of Theoretical Physics* **34**, 399–416 (1965).
- ¹⁸H. Mori, “Transport, Collective Motion, and Brownian Motion*,” *Progress of Theoretical Physics* **33**, 423–455 (1965).
- ¹⁹G. Ciccotti and J. P. Ryckaert, “On the derivation of the generalized Langevin equation for interacting Brownian particles,” *Journal of Statistical Physics* **26**, 73–82 (1981).
- ²⁰H. Meyer, P. Pelagejcev, and T. Schilling, “Non-Markovian out-of-equilibrium dynamics: A general numerical procedure to construct time-dependent memory kernels for coarse-grained observables,” *EPL (Europhysics Letters)* **128**, 40001 (2020).
- ²¹T. J. Doerries, S. A. M. Loos, and S. H. L. Klapp, “Correlation functions of non-Markovian systems out of equilibrium: Analytical expressions beyond single-exponential memory,” *Journal of Statistical Mechanics: Theory and Experiment* **2021**, 033202 (2021).
- ²²O. F. Lange and H. Grubmüller, “Collective Langevin dynamics of conformational motions in proteins,” *The Journal of Chemical Physics* **124**, 214903 (2006).
- ²³T. Kinjo and S.-a. Hyodo, “Equation of motion for coarse-grained simulation based on microscopic description,” *Phys. Rev. E* **75**, 051109 (2007).
- ²⁴F. Glatzel and T. Schilling, “The interplay between memory and potentials of mean force: A discussion on the structure of equations of motion for coarse grained observables,” *Europhysics Letters* (2021).
- ²⁵B. J. Berne and G. D. Harp, “On the Calculation of Time Correlation Functions,” in *Advances in Chemical Physics* (John Wiley & Sons, Ltd, 1970) pp. 63–227.
- ²⁶M. Berkowitz, J. D. Morgan, D. J. Kouri, and J. A. McCammon, “Memory kernels from molecular dynamics,” *The Journal of Chemical Physics* **75**, 2462–2463 (1981).
- ²⁷J. O. Daldrop, J. Kappler, F. N. Brüning, and R. R. Netz, “Butane dihedral angle dynamics in water is dominated by internal friction,” *Proceedings of the National Academy of Sciences* **115**, 5169–5174 (2018).
- ²⁸V. Klippenstein, M. Tripathy, G. Jung, F. Schmid, and N. F. A. van der Vegt, “Introducing Memory in Coarse-Grained Molecular Simulations,” *The Journal of Physical Chemistry B* (2021), 10.1021/acs.jpcc.1c01120.

- ²⁹C. Ayaz, L. Tepper, F. N. Brünig, J. Kappler, J. O. Daldrop, and R. R. Netz, “Non-markovian modeling of protein folding,” *Proceedings of the National Academy of Sciences* **118** (2021), 10.1073/pnas.2023856118.
- ³⁰Y. Zhu, H. Lei, and C. Kim, “Generalized second fluctuation-dissipation theorem in the nonequilibrium steady state: Theory and applications,” arXiv:2104.05222 [cond-mat, physics:physics] (2021), arXiv:2104.05222 [cond-mat, physics:physics].
- ³¹D. Givon, R. Kupferman, and O. H. Hald, “Existence proof for orthogonal dynamics and the Mori-Zwanzig formalism,” *Israel Journal of Mathematics* **145**, 221–241 (2005).
- ³²H. Meyer, T. Voigtmann, and T. Schilling, “On the non-stationary generalized Langevin equation,” *The Journal of Chemical Physics* **147**, 214110 (2017).
- ³³H. Grabert, *Projection Operator Techniques in Nonequilibrium Statistical Mechanics* (Springer, 2006).
- ³⁴R. Kupferman, “Fractional Kinetics in Kac–Zwanzig Heat Bath Models,” *Journal of Statistical Physics* **114**, 291–326 (2004).
- ³⁵M. Tuckerman, *Statistical mechanics: theory and molecular simulation* (Oxford University Press, 2010).
- ³⁶B. Leimkuhler and M. Sachs, “Ergodic Properties of Quasi-Markovian Generalized Langevin Equations with Configuration Dependent Noise and Non-conservative Force,” in *Stochastic Dynamics Out of Equilibrium*, Springer Proceedings in Mathematics & Statistics, edited by G. Giacomin, S. Olla, E. Saada, H. Spohn, and G. Stoltz (Springer International Publishing, Cham, 2019) pp. 282–330.
- ³⁷H. K. Shin, C. Kim, P. Talkner, and E. K. Lee, “Brownian motion from molecular dynamics,” *Chemical Physics Stochastic Processes in Physics and Chemistry* (in Honor of Peter Hänggi), **375**, 316–326 (2010).
- ³⁸A. Carof, R. Vuilleumier, and B. Rotenberg, “Two algorithms to compute projected correlation functions in molecular dynamics simulations,” *The Journal of Chemical Physics* **140**, 124103 (2014).
- ³⁹G. Jung and F. Schmid, “Fluctuation–dissipation relations far from equilibrium: a case study,” *Soft Matter* **17**, 6413–6425 (2021).
- ⁴⁰H. S. Lee, S.-H. Ahn, and E. F. Darve, “Building a Coarse-Grained Model Based on the Mori-Zwanzig Formalism,” *MRS Online Proceedings Library Archive* **1753** (2015), 10.1557/opl.2015.185.
- ⁴¹W. Zhang, C. Hartmann, and C. Schütte, “Effective dynamics along given reaction coordinates, and reaction rate theory,” *Faraday Discuss.* **195**, 365–394 (2016).
- ⁴²C. Ayaz, B. A. Dalton, and R. R. Netz, “Generalized Langevin Equation with a Non-Linear Potential of Mean Force and Non-Linear Memory Friction From a Hybrid Projection Scheme,” arXiv:2202.01922 [physics] (2022), arXiv:2202.01922 [physics].
- ⁴³P. Linz, “Numerical methods for Volterra integral equations of the first kind,” *The Computer Journal* **12**, 393–397 (1969).
- ⁴⁴J. G. Jones, “On the Numerical Solution of Convolution Integral Equations and Systems of Such Equations,” *Mathematics of Computation* **15**, 131–142 (1961).
- ⁴⁵P. K. Lamm, “A Survey of Regularization Methods for First-Kind Volterra Equations,” in *Surveys on Solution Methods for Inverse Problems*, edited by D. Colton, H. W. Engl, A. K. Louis, J. R. McLaughlin, and W. Rundell (Springer, Vienna, 2000) pp. 53–82.
- ⁴⁶C. T. H. Baker, “A perspective on the numerical treatment of Volterra equations,” *Journal of Computational and Applied Mathematics Numerical Analysis 2000. Vol. VI: Ordinary Differential Equations and Integral Equations*, **125**, 217–249 (2000).

- ⁴⁷J. Biazar, E. Babolian, and R. Islam, “Solution of a system of Volterra integral equations of the first kind by Adomian method,” *Applied Mathematics and Computation* **139**, 249–258 (2003).
- ⁴⁸A. Armand and Z. Gouyandeh, “Numerical solution of the system of Volterra integral equations of the first kind,” *International Journal of Industrial Mathematics* **6**, 27–35 (2014).
- ⁴⁹D. Lesnicki, R. Vuilleumier, A. Carof, and B. Rotenberg, “Molecular Hydrodynamics from Memory Kernels,” *Physical Review Letters* **116**, 147804 (2016).
- ⁵⁰G. Jung, M. Hanke, and F. Schmid, “Iterative Reconstruction of Memory Kernels,” *Journal of Chemical Theory and Computation* **13**, 2481–2488 (2017).
- ⁵¹M. Hanke, “Mathematical analysis of some iterative methods for the reconstruction of memory kernels,” *ETNA - Electronic Transactions on Numerical Analysis* **54**, 483–498 (2021).
- ⁵²H. Vroylandt, L. Goudenège, P. Monmarché, F. Pietrucci, and B. Rotenberg, “Likelihood-based non-Markovian models from molecular dynamics,” *Proceedings of the National Academy of Sciences* **119**, e2117586119 (2022).
- ⁵³J. E. Straub, B. J. Berne, and B. Roux, “Spatial dependence of time-dependent friction for pair diffusion in a simple fluid,” *The Journal of Chemical Physics* **93**, 6804–6812 (1990).
- ⁵⁴S. Plimpton, “Fast parallel algorithms for short-range molecular dynamics,” *Journal of Computational Physics* **117**, 1–19 (1995).
- ⁵⁵K. Lü and J.-D. Bao, “Numerical simulation of generalized Langevin equation with arbitrary correlated noise,” *Physical Review E* **72**, 067701 (2005).
- ⁵⁶C. Maes, S. Safaverdi, P. Visco, and F. van Wijland, “Fluctuation-response relations for nonequilibrium diffusions with memory,” *Physical Review E* **87**, 022125 (2013).
- ⁵⁷J. Schmidt, A. Meistrenko, H. van Hees, Z. Xu, and C. Greiner, “Simulation of stationary Gaussian noise with regard to the Langevin equation with memory effect,” *Physical Review E* **91**, 032125 (2015).

Appendix A: Projection operators

By definition, the orthogonal projection operator on the closure of the space spanned by the set $\mathcal{E} = \{E_k, k \in \mathcal{I}\}$ is the linear operator characterized by the two following properties: first, for all $k \in \mathcal{I}$, $\mathcal{P}_{\mathcal{E}}E_k = E_k$. Second, $\mathcal{P}_{\mathcal{E}}\phi = 0$ for all functions ϕ which are orthogonal to E_k for all $k \in \mathcal{I}$. The idempotency of the projector, namely $\mathcal{P}_{\mathcal{E}}\mathcal{P}_{\mathcal{E}}\phi = \mathcal{P}_{\mathcal{E}}\phi$, follows from those properties.

Let us check that these properties are indeed satisfied by the operator given by Eq.(5). The second one is clear, and for all $k, j \in \mathcal{I}$,

$$\sum_{k' \in \mathcal{I}} G_{k,k'}^{-1} \langle E_{k'}, E_j \rangle = (G^{-1}G)_{k,j} = 1_{k=j},$$

from which $\mathcal{P}_{\mathcal{E}}E_j = E_j$ for all $j \in \mathcal{I}$.

As emphasized before, the projection on various spaces are usually considered. Recall that we consider $E_k = e_k \circ \mathcal{O}$, i.e. in any cases we only consider functions of the observable \mathcal{O} . In that case we can write $\mathcal{P}_{\mathcal{E}}\phi = f_{\phi} \circ \mathcal{O}$ where f_{ϕ} is the function of the observable given by

$$f_{\phi} = \sum_{k \in \mathcal{I}} \sum_{k' \in \mathcal{I}} G_{k,k'}^{-1} \langle E_{k'}, E_j \rangle e_k.$$

- In the case of the projection on a singleton set $\mathcal{E} = \{E_0\}$, the projection becomes

$$f_\phi(z) = \frac{\langle E_0, \phi \rangle}{\langle E_0, E_0 \rangle} e_0(z)$$

Moreover, when the function $e_0(z) = z$ is simply the identity, this is the usual linear Mori projection.

- In the multidimensional case, we can project onto the set of linear functions $e_j : z \mapsto z_j$ returning the j coordinate of z , the projection then reads

$$f_\phi(z) = \sum_{j,j'=1}^d G_{j,j'}^{-1} \langle E_{j'}, \phi \rangle z_j,$$

in particular f_ϕ is a linear function.

- We can also consider the projection upon all functions of $z \in \mathbb{R}^d$. The orthogonal projection is then given by the conditional expectation

$$f_\phi(z) = \mathbb{E}(\phi(\mathbf{X}) \mid \mathcal{O}(\mathbf{X}) = z).$$

Informally, this projector can be obtained from Eq. (5) using for \mathcal{E} the set of Dirac function $\{\delta(\mathcal{O}(\mathbf{X}) - z), z \in \mathbb{R}^d\}$, but choosing $\{e_k, k \in \mathcal{I}\}$ as any Hilbert basis of $L^2(\mathbb{R}^d)$ will give the same result. This is the Zwanzig projection, or "non-linear" projection. Note that "non-linear" refers here to f_ϕ being a non-linear function of z and not to the linearity of the projection operator.

- Finally when we consider an observable that could be separated into a position observable and a velocity observable, as in the main text sec. IID or sec. IIF, we could consider a projection on functions of only the position observable $\varphi(\mathbf{q})$. For instance, the projection upon all function of the position would become for $x \in \mathbb{R}^n$

$$f_\phi(x) = \mathbb{E}(\phi(\mathbf{q}, \mathbf{p}) \mid \varphi(\mathbf{q}) = x).$$

Appendix B: Coarea formula

The projection can be expressed using the coarea formula: for an observable $\phi: \Omega \rightarrow \mathbb{R}^d$, we have

$$\int_{\Omega} \psi(\mathbf{X}) d\mathbf{X} = \int_{\mathbb{R}^d} \int_{\Sigma_z} \psi(\mathbf{X}) \omega_\phi(\mathbf{X}) \sigma_z(d\mathbf{X}) dz$$

for any function ψ on Ω , where $\Sigma_z = \{\mathbf{X} \in \Omega, \phi(\mathbf{X}) = z\}$, σ_z is the Lebesgue measure on Σ_z and $\omega_\phi(\mathbf{X}) = 1/\sqrt{\det((\nabla\phi(\mathbf{X}))^T \nabla\phi(\mathbf{X}))}$ is the inverse of the d -dimensional Jacobian of \mathcal{O} . The measure $\omega_\phi \sigma_z(d\mathbf{X})$ is also denoted by $\delta(\phi(\mathbf{X}) - z) d\mathbf{X}$. Using the notations of sec. IID, we see that the marginal density ρ_φ is given by

$$\rho_\varphi(z) = \int_{\mathbb{R}^D} \int_{\Sigma_z} \omega_\varphi(\mathbf{q}) \rho_{eq}(\mathbf{q}, \mathbf{p}) d\mathbf{p} \sigma_z(d\mathbf{q})$$

where Σ_z corresponds to $\varphi(\mathbf{q}) = z$.

This mean that scalar product between a function g of the observable and a function F can be written as integral over a d -dimensional space

$$\begin{aligned} \int_{\mathbb{R}^{2D}} g(\mathcal{O}(\mathbf{X}))^T \cdot F(\mathbf{X}) \rho_{eq}(\mathbf{X}) d\mathbf{X} &= \int_{\mathbb{R}^d} g(z)^T \cdot \int_{\Sigma_z} F(\mathbf{X}) \omega_{\mathcal{O}}(\mathbf{X}) \rho_{eq}(\mathbf{X}) \sigma_z(d\mathbf{X}) dz \\ &= \int_{\mathbb{R}^d} g(z)^T \cdot F_{\varphi}(z) \rho_{\varphi}(z) dz \\ &= \langle g, F_{\varphi} \rangle_{\varphi} \end{aligned}$$

having introduced the conditional expectation

$$F_{\varphi}(z) = \mathbb{E}(F(\mathbf{X}) \mid \varphi(\mathbf{q}) = z) = \frac{1}{\rho_{\varphi}(z)} \int_{\Sigma_z} F(\mathbf{X}) \omega_{\mathcal{O}}(\mathbf{X}) \rho_{eq}(\mathbf{X}) \sigma_z(d\mathbf{X}).$$

Appendix C: Derivation for the mean force term

Using the skew-symmetry (12) of \mathcal{L} , we have for the mean force term

$$\begin{aligned} \langle B_k, \mathcal{L}\mathcal{O}_v \rangle &= -\langle C_k, \mathcal{O}_v \rangle = -\langle \nabla h_k(\mathcal{O}_x(t)) \mathcal{O}_v, \mathcal{O}_v \rangle \\ &= -\int_{\mathbb{R}^D} \int_{\mathbb{R}^D} \mathcal{O}_v(\mathbf{q}, \mathbf{p})^T \cdot \nabla h_k(\mathcal{O}_x(\mathbf{q}))^T \cdot \mathcal{O}_v(\mathbf{q}, \mathbf{p}) \rho_{eq}(\mathbf{q}, \mathbf{p}) d\mathbf{q} d\mathbf{p} \\ &= -\sum_{j=1}^n \sum_{i=1}^n \int_{\mathbb{R}^D} \int_{\mathbb{R}^D} \partial_{z_j} h_k(\mathcal{O}_x(\mathbf{q}))_i \mathcal{O}_{v,j}(\mathbf{q}, \mathbf{p}) \mathcal{O}_{v,i}(\mathbf{q}, \mathbf{p}) \rho_{eq}(\mathbf{q}, \mathbf{p}) d\mathbf{q} d\mathbf{p}. \end{aligned}$$

From the coarea formula and the definition (17) of the effective mass matrix, we obtain

$$\begin{aligned} \langle B_k, \mathcal{L}\mathcal{O}_v \rangle &= -\int_{\mathbb{R}^n} \sum_{j=1}^n \sum_{i=1}^n \partial_{z_j} h_k(z)_i \left(\int_{\Sigma_z} \int_{\mathbb{R}^D} \mathcal{O}_{v,j}(\mathbf{q}, \mathbf{p}) \mathcal{O}_{v,i}(\mathbf{q}, \mathbf{p}) \omega_{\varphi}(\mathbf{q}) \rho_{eq}(\mathbf{q}, \mathbf{p}) \sigma_z(d\mathbf{q}) \right) d\mathbf{p} dz \\ &= -\frac{1}{\beta} \int_{\mathbb{R}^n} \sum_{j=1}^n \sum_{i=1}^n \partial_{z_j} h_k(z)_i (\rho_{\varphi}(z) M_{\varphi}(z)^{-1})_{ij} dz \\ &= \frac{1}{\beta} \int_{\mathbb{R}^n} \sum_{j=1}^n \sum_{i=1}^n h_k(z)_i \partial_{z_j} (\rho_{\varphi}(z) M_{\varphi}(z)^{-1})_{ij} dz \\ &= \frac{1}{\beta} \int_{\mathbb{R}^n} \sum_{i=1}^n h_k(z)_i \sum_{j=1}^n \left(\partial_{z_j} (M_{\varphi}(z)^{-1})_{ji} + (M_{\varphi}(z)^{-1})_{ji} \partial_{z_j} \ln \rho_{\varphi}(z) \right) \rho_{\varphi}(z) dz \\ &= \frac{1}{\beta} \int_{\mathbb{R}^n} h_k(z)^T \cdot (\nabla \cdot M_{\varphi}(z)^{-1} + M_{\varphi}(z)^{-1} \cdot \nabla \ln \rho_{\varphi}(z)) \rho_{\varphi}(z) dz \end{aligned}$$

which concludes the proof of (19).

Appendix D: Details of the position-dependent 2-FDT

The memory kernel is obtained through the Gram matrix G_c of the function set and the scalar product $\langle C_k, \mathcal{L}\xi_t \rangle$. The Gram matrix of the set $\{c_k, k \in \mathcal{J}\}$ is related to the effective

mass matrix,

$$\begin{aligned}
 G_{C_k, C_{k'}} &= \langle C_k, C_{k'} \rangle = \int_{\mathbb{R}^D} \int_{\mathbb{R}^D} \mathcal{O}_v(\mathbf{q}, \mathbf{p})^T \nabla h_k(\mathcal{O}_x(\mathbf{q}))^T \nabla h_{k'}(\mathcal{O}_x(\mathbf{q})) \mathcal{O}_v(\mathbf{q}, \mathbf{p}) \rho_{eq}(\mathbf{q}, \mathbf{p}) d\mathbf{q} d\mathbf{p} \\
 &= \frac{1}{\beta} \sum_{j=1}^n \sum_{i=1}^n \sum_{l=1}^n \int_{\mathbb{R}^n} \partial_{z_j} h_k(z)_i \partial_{z_i} h_{k'}(z)_l (M_\varphi(z)^{-1})_{ij} \rho_\varphi(z) dz \\
 &= \frac{1}{\beta} \langle \nabla h_k, M_\varphi^{-1} \cdot \nabla h_{k'} \rangle_\varphi.
 \end{aligned}$$

Using the anti-hermitianity of \mathcal{L} , we have for $k \in \mathcal{J}$

$$\begin{aligned}
 \langle C_k, \mathcal{L}\xi_t \rangle &= -\langle \mathcal{L}C_k, \xi_t^v \rangle \\
 &= -\sum_{i=1}^n \sum_{j=1}^n \int_{\mathbb{R}^D} \int_{\mathbb{R}^D} \partial_{z_j} h_k(\mathcal{O}_x(\mathbf{q}))_i \mathcal{L}\mathcal{O}_{v,j} \xi_{t,i}^v(\mathbf{q}, \mathbf{p}) \rho_{eq}(\mathbf{q}, \mathbf{p}) d\mathbf{q} d\mathbf{p} \\
 &\quad - \sum_{i=1}^n \sum_{j=1}^n \sum_{l=1}^n \int_{\mathbb{R}^D} \int_{\mathbb{R}^D} \partial_{z_j} \partial_{z_l} h_k(\mathcal{O}_x(\mathbf{q}))_i \mathcal{O}_{v,l}(\mathbf{q}, \mathbf{p}) \mathcal{O}_{v,j}(\mathbf{q}, \mathbf{p}) \xi_{t,i}^v(\mathbf{q}, \mathbf{p}) \rho_{eq}(\mathbf{q}, \mathbf{p}) d\mathbf{q} d\mathbf{p}.
 \end{aligned}$$

We can then introduce the identity in the form of $1 - \mathcal{P}_\mathcal{E} + \mathcal{P}_\mathcal{E}$

$$\begin{aligned}
 \langle C_k, \mathcal{L}\xi_t \rangle &= -\langle \mathcal{L}C_k, \xi_t^v \rangle \\
 &= -\sum_{i=1}^n \sum_{j=1}^n \int_{\mathbb{R}^D} \int_{\mathbb{R}^D} \partial_{z_j} h_k(\mathcal{O}_x(\mathbf{q}))_i [(1 - \mathcal{P}_\mathcal{E} + \mathcal{P}_\mathcal{E}) \mathcal{L}\mathcal{O}_{v,j}] \xi_{t,i}^v(\mathbf{q}, \mathbf{p}) \rho_{eq}(\mathbf{q}, \mathbf{p}) d\mathbf{q} d\mathbf{p} \\
 &\quad - \sum_{i=1}^n \sum_{j=1}^n \sum_{l=1}^n \int_{\mathbb{R}^D} \int_{\mathbb{R}^D} \partial_{z_j} \partial_{z_l} h_k(\mathcal{O}_x(\mathbf{q}))_i \mathcal{O}_{v,l}(\mathbf{q}, \mathbf{p}) \mathcal{O}_{v,j}(\mathbf{q}, \mathbf{p}) \xi_{t,i}^v(\mathbf{q}, \mathbf{p}) \rho_{eq}(\mathbf{q}, \mathbf{p}) d\mathbf{q} d\mathbf{p} \\
 &= -\sum_{i=1}^n \sum_{j=1}^n \int_{\mathbb{R}^D} \int_{\mathbb{R}^D} \partial_{z_j} h_k(\mathcal{O}_x(\mathbf{q}))_i \xi_{0,j}^v \xi_{t,i}^v(\mathbf{q}, \mathbf{p}) \rho_{eq}(\mathbf{q}, \mathbf{p}) d\mathbf{q} d\mathbf{p} \\
 &\quad - \sum_{i=1}^n \sum_{j=1}^n \int_{\mathbb{R}^D} \int_{\mathbb{R}^D} \partial_{z_j} h_k(\mathcal{O}_x(\mathbf{q}))_i f_b(\mathcal{O}_x(\mathbf{q}))_j \xi_{t,i}^v(\mathbf{q}, \mathbf{p}) \rho_{eq}(\mathbf{q}, \mathbf{p}) d\mathbf{q} d\mathbf{p} \\
 &\quad - \sum_{i=1}^n \sum_{j=1}^n \sum_{l=1}^n \int_{\mathbb{R}^D} \int_{\mathbb{R}^D} \partial_{z_j} \partial_{z_l} h_k(\mathcal{O}_x(\mathbf{q}))_i \mathcal{O}_{v,l}(\mathbf{q}, \mathbf{p}) \mathcal{O}_{v,j}(\mathbf{q}, \mathbf{p}) \xi_{t,i}^v(\mathbf{q}, \mathbf{p}) \rho_{eq}(\mathbf{q}, \mathbf{p}) d\mathbf{q} d\mathbf{p},
 \end{aligned}$$

where we have used that

$$(1 - \mathcal{P}_\mathcal{E}) \mathcal{L}\mathcal{O}_v = \xi_0^v \quad \text{and} \quad \mathcal{P}_\mathcal{E} \mathcal{L}\mathcal{O}_v = f_b \circ \mathcal{O}_x.$$

By definition of the conditional variance of the noise (29) and of the conditional expectation of the noise (26) and (27), we get

$$\begin{aligned}
 \langle C_k, \mathcal{L}\xi_t \rangle &= -\int_{\mathbb{R}^n} \sigma_\varphi(t, z) \cdot \nabla h_k(z) \rho_\varphi(z) dz \\
 &\quad - \int_{\mathbb{R}^n} f_b(z) \cdot \nabla h_k(z) \cdot \bar{\xi}_t(z) \rho_\varphi(z) dz \\
 &\quad - \int_{\mathbb{R}^n} \nabla^2 h_k(z) \bar{\xi}_t^z(z) \rho_\varphi(z) dz.
 \end{aligned}$$

Then a partial integration on the last term of the r.h.s. conclude the derivation of (30).

Appendix E: Projected correlation functions

Correlation function of the noise with an observable $F(\mathbf{X})$ differs from the usual time correlation in that the noise is evolved with the orthogonal dynamics (9). Such projected correlations functions³⁸ are defined as

$$\langle F, \xi_t \rangle = \langle F, e^{t(1-\mathcal{P}\mathcal{E})\mathcal{L}}\xi_0 \rangle.$$

The noise follows from Eq. (8)

$$\xi_t(\mathbf{X}) = \frac{\partial \mathcal{O}(t)}{\partial t} - f(\mathcal{O}(t)) - \int_0^t \sum_k g_k(s) E_k(t-s) ds$$

and the the projected correlation functions can be obtained from

$$\langle F, \xi_t \rangle = \langle F, \frac{\partial \mathcal{O}(t)}{\partial t} - f(\mathcal{O}(t)) \rangle - \int_0^t \sum_k g_k(s) \langle F, E_k(t-s) \rangle ds.$$

Notice that in particular a set of Volterra integral equations of the second kind for the memory kernel can be obtained. Indeed the coefficient of the memory kernel are projected correlation function, since for $\ell \in \mathcal{I}$

$$\sum_k G_{\ell,k} g_k(t) = \langle E_\ell, \mathcal{L}\xi_t \rangle = -\langle \mathcal{L}E_\ell, \xi_t \rangle$$

using the skew-symmetry of \mathcal{L} . This lead to³¹

$$\sum_k G_{\ell,k} g_k(t) = -\left\langle \mathcal{L}E_\ell(0), \frac{\partial \mathcal{O}(t)}{\partial t} \right\rangle + \sum_k f_k \langle \mathcal{L}E_\ell(0), E_k(t) \rangle + \int_0^t \sum_k g_k(s) \langle \mathcal{L}E_\ell(0), E_k(t-s) \rangle ds.$$

From a numerical point of view, we usually compute correlation function using a sum along trajectories sampled every Δt , *i.e.*

$$\langle F, \xi_t \rangle = \sum_m F(\mathbf{X}(m\Delta t))^T \xi_t(\mathbf{X}(m\Delta t)).$$

However, as the noise is evolved with the orthogonal dynamics, we need a relation between $\xi_t(\mathbf{X}(m\Delta t))$ and $\xi_{t+m\Delta t}(\mathbf{X})$. Writing $t = m\Delta t$, we write the noise at time $t + t_0$

$$\xi_{t+t_0}(\mathbf{X}) = e^{t\mathcal{L}} \frac{\partial \mathcal{O}(t_0)}{\partial t} - e^{t\mathcal{L}} f(\mathcal{O}(t_0)) - \int_0^{t+t_0} g(s, \mathcal{O}(t+t_0-s)) ds.$$

Splitting the integral at t and changing the variable s of integration to $u = s - t$ leads to the required relation

$$\xi_{t+t_0}(\mathbf{X}) = \xi_t(\mathbf{X}(t_0)) - \int_0^{t_0} g(u+t, \mathcal{O}(\mathbf{X}(t_0-u))) du.$$

We emphasize that discretizing the integral gives a generalization of the algorithm for the computation of projected correlation function that was discussed in Refs. 38,42,49.