



HAL
open science

The flash-small-pool PCR: how to transform blotting and numerous hybridization steps into a simple denatured PCR

Elodie Dandelot, Geneviève Gourdon

► **To cite this version:**

Elodie Dandelot, Geneviève Gourdon. The flash-small-pool PCR: how to transform blotting and numerous hybridization steps into a simple denatured PCR. *Biotechniques*, 2018, 64 (6), pp.262 - 265. 10.2144/btn-2018-0035 . hal-03753523

HAL Id: hal-03753523

<https://hal.sorbonne-universite.fr/hal-03753523>

Submitted on 18 Aug 2022

HAL is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers.

L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.

The flash-small-pool PCR: how to transform blotting and numerous hybridization steps into a simple denatured PCR

Elodie Dandelot^{1,2} & Geneviève Gourdon^{1,2}

¹Laboratory CTGDM, Inserm UMR1163, 75015 Paris, France; ²Institut Imagine, Université Paris Descartes-Sorbonne Paris Cité, 75015 Paris, France

BioTechniques 64:262-265 (June 2018) 10.2144/btn-2018-0035

Keywords: DM1 • IRDye • microsatellites instability • PCR • trinucleotide repeats

Numerous human diseases are associated with abnormal expansion of unstable trinucleotide repeats (TNRs). TNR instability mechanisms are complex, and remain only partially understood. Small-pool-PCR (SP-PCR) is the reference method to assess TNR instability. SP-PCR amplifies a low number of DNA molecules and is followed by Southern blot. However, SP-PCR remains expensive and time consuming. Here, we describe an optimized SP-PCR that can be done in a day, which reduces cost and experimental biases: the flash-small-pool PCR (FSP-PCR). This method consists of a fluorescent PCR on a few DNA molecules, followed by an alkaline gel electrophoresis revealed with a near infra-red detector system. With reduced experimental steps, cost, and time consumption, microsatellite analysis will become more accessible due to FSP-PCR.

Introduction

An increasing number of human neuromuscular and neurodegenerative diseases are due to unstable trinucleotide repeats (TNRs), including Huntington's disease (with CAG repeats) [1], myotonic dystrophy type 1 (with CTG repeats) [2–4], Fragile X Syndrome (FXS, with CGG repeats) [5] or Friedreich's ataxia (with GAA repeats) [6]. In the general population, repeats remain stable in tissues and through parental transmissions. In patients, the abnormally expanded repeats are unstable and generally biased towards expansions in both germline and somatic cells. In some TNR-associated diseases, such as myotonic dystrophy, larger repeats are associated with more severe symptoms and with an earlier age of onset (anticipation phenomenon) [7]. In somatic cells, TNR instability is age-dependent, tissue-specific and may be associated with symptom progression [8,9]. Therefore, evaluation of TNR instability is critical for prognosis and to better understand these diseases.

Triplets instability appears as a complex result of different mechanisms [7,10]. Development of different models such as mouse, bacterial, yeast, or cell models has given insight about the processes underlying TNR instability but, to date, the dynamic of repeats changes remains only partially understood [10–13]. Further analysis on TNR instability will provide clues for new therapeutic approaches to avoid repeat expansion or to induce TNR contraction down to a non-pathogenic length. Furthermore, accurate TNR instability analysis is necessary to characterize new models and new experimental tools [14–16].

SP-PCR is a powerful method to accurately analyze TNR instability [17–20]. The serial dilutions involved in this method resolve the heterogeneous smear detected by standard PCR experiments. However, current protocols are time consuming, with many steps that can affect the experiment quality such as blotting quality, and hybridization. Moreover, current protocols require many reagents and remain expensive. To eliminate this disadvantage, and to facil-

itate the spread of this high-sensitive method, we have developed an easier and faster method: the flash-small-pool PCR (FSP-PCR) to detect microsatellite repeat-containing PCR products. This protocol comes without biohazard, and does not require a blotting step or hybridization. Moreover, this protocol is faster, feasible in 24 h only, while current procedures take at least 3 days.

In short, this protocol consists of a classical small-pool-PCR performed with one of the primers labeled with near infrared dye (IRD), loaded on a denaturing agarose gel, and analyzed with fluorescent analyzer, here the Odyssey[®] CLx Infrared Imaging System (LI-COR Biosciences – GmbH, NE, USA). In this study, we focused on CTG repeat expansion in DM1 patients.

Material & methods

This protocol has been adapted from previously described procedures [16–18]. The primers used are listed in Table 1. See

METHOD SUMMARY

To analyze more quickly and easily triplet nucleotides repeats, we optimized the commonly used small-pool PCR protocol using a near infrared-labeled primer and alkaline gel electrophoresis. Our protocol gives robust results, and avoids the many experimental reagents and biases of the classic small-pool PCR.

Table 1. List of primers used in flash-small-pool PCR.

Primer name	Primer sequence
Forward primer DMC	5'-AACGGGGCTCGAAGGGTCT-3'
Reverse labeled primer DMBR	5'-IRD700-CGTGGAGGATGGAACACGGGAC-3'

supplementary material for a more detailed protocol.

Restriction endonuclease digestion

All DNA quantifications were done with Qubit® Fluorometer (ThermoFisher Scientific, MA, USA) following manufacturer recommendations. To facilitate DNA accessibility, genomic DNA was digested with 50 U of Hind III endonuclease (Fermentas, MA, USA).

FSP-PCR

Genomic digested DNA was serially diluted ($1.5 \text{ ng.}\mu\text{l}^{-1}$, $300 \text{ pg.}\mu\text{l}^{-1}$, $60 \text{ pg.}\mu\text{l}^{-1}$, $12 \text{ pg.}\mu\text{l}^{-1}$) in deionized milli Rho water with 0.1 M of carrier oligonucleotide (Table 1). PCR mix was prepared as follows (final volume = $7 \mu\text{l}$ per well): $0.2 \mu\text{M}$ of unlabeled-forward primer, $0.2 \mu\text{M}$ of IRD-Reverse primer (Metabion, Planegg, Bavaria, Germany), 1X PCR buffer (ThermoFisher Scientific, ref. SM-0005; MA, USA), $5 \text{ U.}\mu\text{l}^{-1}$ of Thermo Perfect Taq (Integro, Zaandam, The

Netherlands), Deionized milli Rho water up to $6.5 \mu\text{l}$. Each PCR reaction was done with $0.5 \mu\text{l}$ of each serial DNA dilution. PCR program was launched in a Thermal Cycler (Applied Biosystems, CA, USA): one 5-min hold at 96°C , followed by 28 cycles: 96°C (45 s), 68°C (45 s), 70°C (3 min), and a chase at 68°C (1 min) and 70°C (10 min). No PCR optimization was needed because of infra-red labeling (IRD).

Denaturing agarose gel electrophoresis of PCR products

A $20 \times 40 \text{ cm}$ 2.5% agarose gel [19] (50 mM of NaCl and 4 mM of EDTA) was pre-soaked in a cold room for at least 1 h in running buffer (NaOH 30 mM , EDTA 2 mM , volume adjusted with milli Rho water). $2 \mu\text{l}$ of the PCR reactions were used in each well. The ladder used was the 50–1500 bp DNA sizing standard (LI-COR Biosciences – GmbH, USA). The gel electrophoresis was set up as follows 30 min at 140 V, then 2 V/cm (80 V for a $20 \times 40 \text{ cm}$ gel) overnight. Gels were directly observed with the Odyssey® CLx Infrared Imaging System and analyzed with Image Studio™ software (LI-COR Biosciences – GmbH).

Human DNA samples

DNA from DM1 patients were obtained from the Genethon DNA and Cell Bank in collaboration with the DM scope registry. Consent was obtained for the use of DNA samples in this study.

Results & discussion

We provide here an optimized and robust protocol to assess microsatellite instability. While the current protocols with radioactivity or digoxigenin labeling require expensive reagents and days of experimental procedures, this new protocol provides results in a much simpler way (the experimental steps are resumed to one PCR and one agarose gel migration only), and within a decreased procedure time. This protocol has been tested using serial dilution of blood DNA from a DM1




Figure 1. FSP-PCR using blood DNA from patients carrying 150 CTG and non-DM1 controls. (A) The FSP-PCR was performed with successive dilutions of a DM1 patient blood DNA ($1.5 \text{ ng.}\mu\text{l}^{-1}$, $300 \text{ pg.}\mu\text{l}^{-1}$, $60 \text{ pg.}\mu\text{l}^{-1}$ and $12 \text{ pg.}\mu\text{l}^{-1}$). For each dilution, the seven lanes represent seven PCR replicates. PCR products corresponding to normal and mutant alleles are indicated in brackets. Gel agarose percentage: 1.5%. **(B)** FSP-PCR using blood DNA from three DM1 patients. For each patient, seven PCR replicates were performed using the $300 \text{ pg.}\mu\text{l}^{-1}$ dilution. Gel agarose percentage: 1.5%. **(C)** FSP-PCR using blood DNA from three non-DM1 control individuals. For each patient, seven PCR replicates were performed using the $300 \text{ pg.}\mu\text{l}^{-1}$ dilution. Gel agarose percentage: 3%. **(D)** Left panel: SP-PCR (seven replicates using the DNA dilution $300 \text{ pg.}\mu\text{l}^{-1}$) revealed with a (CAG)_n probe labeled with digoxigenin after Southern blot. Right panel: FSP-PCR (seven replicates using the DNA dilution $300 \text{ pg.}\mu\text{l}^{-1}$) revealed directly with Odyssey® CLx Infrared Imaging. L1: 50–1500 bp DNA sizing standard (LI-COR Biosciences – GmbH). L2: Thermo Scientific GeneRuler 100 bp DNA Ladder (ThermoFisher Scientific). L3: DNA Molecular Weight Marker III, DIG-labeled (Roche). DM-1: Myotonic dystrophy type 1; FSP-PCR: Flash-small-pool PCR.




Figure 2. CTG repeat length determination using IRD-labeled leading primer and alkaline gel electrophoresis. PCR with DMC/DMBR primers on 2 ng of DNA extracted from human cerebellum samples with (A) 500 CTG repeats (1645 bp expected) or (B) 900 CTG (2845 bp expected), and (C) mice tail with 1200 CTG (3745 bp expected). L3: Thermo Scientific GeneRuler 1 kb DNA Ladder (ThermoFisher Scientific). The ladder lane was revealed with ethidium bromide while the rest of the gel with the PCR products was revealed with Odyssey® CLx Infrared Imaging.

patient carrying 150 CTG repeats. Figure 1A shows clearly the CTG repeat somatic mosaicism in the different dilutions as observed in classical small-pool PCR [16]. FSP-PCR was repeated with blood DNA from three different DM1 patients carrying between 150 and 200 CTG repeats. PCR amplification using 0.5 μl of the 300 μg . μl^{-1} dilution shows again clearly the CTG repeat mosaicism resulting from somatic instability (Figure 1B and Supplementary Figure S1). As expected, no somatic mosaicism was detected in non-DM1 individuals (Figure 1C and Supplementary Figure S1). We can note different repeat sizes for normal alleles between individuals, reflecting *DMPK* normal allele polymorphisms [21]. In order to verify the efficiency of FSP-PCR, we compared the CTG repeat expansion mosaicism using FSP-PCR and classical SP-PCR (Figure 1D) [18]. The magnitude of the CTG repeat instability was comparable

using both methods showing normal sensitivity with FSP-PCR. We also used various DNA samples carrying different CTG repeat expansions including DNA extracted from DM1 cerebellum samples (with 500 and 900 CTG repeats) and tail DNA from the DMSXL mouse model carrying 1200 CTG repeats. Using 2 ng of DNA, we were able to detect up to 1200 CTG repeats (Figure 2). The results obtained demonstrate that the protocol is useful for detecting a wide range of CTG repeat lengths both in human (blood and tissues) and in mice. Since no optimization was necessary in the original PCR program used with classical unlabeled primers, we assume that this protocol can be easily adapted to any usual PCR protocol [17–20].

Recently, it has been reported that ethidium bromide modifies the mobility of CAG•CTG alternative DNA structures generated by PCR during electrophoresis [22]. Using a denaturing gel and fluorescent primers, this protocol overcomes the possible migration bias resulting from secondary DNA structures formed during PCR. Moreover, we used here IRD700 labeling, as it is empirically known to provide sharper detection than the IRD800 labeling [23]. To our knowledge, there are no IRD-labeled ladders with bands higher than 1500 bp. This problem can be easily overcome by using a usual standard and performing a TBE 1X/ethidium bromide bath after analysis with the Odyssey CLx Infrared Imaging System: fluorescent and UV pictures can be overlaid in computational post-treatment (Figure 2). Another solution can be to create your own labeled ladder as already described elsewhere [24].

In conclusion, the FSP-PCR protocol using fluorescent PCR, an alkaline gel electrophoresis, and a near Infra-red detector system is a highly sensitive, easy, reliable, and faster method for assessing microsatellite instability.

Acknowledgements

The authors thank S.Tome and the DM-Scope registry (especially Dr. G. Bassez and C. Dogan) for their help in obtaining DNA samples.

Author contributions

E.D. designed and performed the experiments and prepared the manuscript. G.G. supervised the study and corrected the manuscript.

Financial & competing interests disclosure

This study was supported by grants from AFM-Téléthon (France, AFM-Telethon grant n° 19757 to G. G.), INSERM (France), Université Paris Descartes (France) as well as a PhD fellowship from Ministère Français de la Recherche et Technologie (to E.D.). The Imagine Institute received a state subsidy managed by the National Research Agency under the “Investments for the Future” program (ANR-10-IAHU-01). The authors have no other relevant affiliations or financial involvement with any organization or entity with a financial interest in or financial conflict with the subject matter or materials discussed in the manuscript apart from those disclosed.

No writing assistance was utilized in the production of this manuscript.

Ethical conduct of research

The authors state that they have obtained appropriate institutional review board approval or have followed the principles outlined in the Declaration of Helsinki for all human or animal experimental investigations. In addition, for investigations involving human subjects, informed consent has been obtained from the participants involved.

Open access

This work is licensed under the Creative Commons Attribution 4.0 License. To view a copy of this license, visit <http://creativecommons.org/licenses/by/4.0/>

Supplementary data

To view the supplementary data that accompany this paper please visit the journal website at: www.future-science.com/doi/full/10.4155/btn-2018-0035

References

Papers of special note have been highlighted as:
• of interest; •• of considerable interest


1. The Huntington's Disease Collaborative Research Group. A novel gene containing a trinucleotide repeat that is expanded and unstable on Huntington's disease chromosomes. *Cell* 72, 971–983 (1993).
2. Aslanidis C, Jansen G, Amemiya C *et al.* Cloning of the essential myotonic dystrophy region and mapping of the putative defect. *Nature* 355, 548–551 (1992).
3. Brook JD, McCurrach ME, Harley HG *et al.* Molecular basis of myotonic dystrophy: expansion of a trinucleotide (CTG) repeat at the 3' end of a transcript encoding a protein kinase family member. *Cell* 68, 799–808 (1992).
4. Fu YH, Pizzuti A, Fenwick RG Jr, King J, Rajnarayan S. An unstable triplet repeat in a gene related to myotonic muscular dystrophy. *Science* 255, 1256–1258 (1992).
5. Ashley CT, Sutcliffe JS, Kunst CB *et al.* Human and murine FMR-1: alternate splicing and translational initiation downstream of the CGG-repeat. *Nat. Gene.* 4, 244–251 (1993).
6. Campuzano V, Montermini L, Moltò MD *et al.* Friedreich's ataxia: autosomal recessive disease caused by an intronic GAA triplet repeat expansion. *Science* 271, 1423–1427 (1996).
7. Pearson CE, Nichol Edamura K, Cleary JD. Repeat instability: mechanisms of dynamic mutations. *Nat. Rev. Genet.* 6(10), 729–742 (2005).
8. Morales F, Couto JM, Higham CF *et al.* Somatic instability of the expanded CTG triplet repeat in myotonic dystrophy type 1 is a heritable quantitative trait and modifier of disease severity. *Human Mol. Gen.* 21(16), 3558–3567 (2012).
9. Illarioshkin SN, Igarashi S, Onodera O *et al.* Trinucleotide repeat length and rate of progression of Huntington's disease. *Neurology* 36(4), 630–635 (1994).
10. López castel A, Cleary JD, Pearson CE. Repeat instability as the basis for human diseases and as a potential target for therapy. *Nat. Rev. Mol. Cell Biol.* 11(3), 165–170 (2010).
11. Usdin K, House NCM, Freudenreich CH. Repeat instability during DNA repair: Insights from model systems. *Crit. Rev. Biochem. Mol. Biol.* 50(2), 142–167 (2015).
12. Gomes-Pereira M, Cooper TA, Gourdon G. Myotonic dystrophy mouse models: towards rational therapy development. *Trends Mol. Med.* 17(9), 10.1016 (2011).
13. Brouwer JR, Willemsen R, Oostra BA. Microsatellite repeat instability and neurological disease. *BioEssays* 31(1), 71–83 (2009).
14. Polyzos A, McMurray C. Close encounters: Moving along bumps, breaks, and bubbles on expanded trinucleotide tracts. *DNA Repair* 56, 144–155 (2017).
15. Cinesi C, Aeschbach L, Yang B, Dion V. Contracting CAG/CTG repeats using the CRISPR-Cas9 nickase. *Nat. Commun.* 7, 13272 (2016).
16. Monckton DG, Wong LJ, Ashizawa T, Caskey CT. Somatic mosaicism, germline expansions, germline reversions and intergenerational reductions in myotonic dystrophy males: small pool PCR analyses. *Hum. Mol. Gene.* 4(1), 1–8 (1995).
•• First description of the Small-Pool PCR method.
17. Gomes-Pereira M, Bidichandani SI, Monckton DG. Analysis of unstable triplet repeats using small-pool polymerase chain reaction. In: *Trinucleotide Repeat Protocols. Methods in Molecular Biology*[™]. Kohwi Y (Ed.). Humana Press, New York City, NY, USA (2004).
• Full description of the Small-Pool PCR method.
18. Tomé S, Nicole A, Gomes-Pereira M, Gourdon G. Non-radioactive detection of trinucleotide repeat size variability. *PLoS Currents* 6 March 2014 Edition 1 (2014).
19. Drouin R, Gao S, Holmquist GP. Agarose gel electrophoresis for DNA damage analysis. In: *Technologies for Detection of DNA Damage and Mutations*. Pfeifer GP (Ed.). Springer, Boston, MA, USA (1996).
20. Aeschbach L, Dion V. Minimizing carry-over PCR contamination in expanded CAG/CTG repeat instability applications. *Sci. Rep.* 7(1), 18026 (2017).
21. Zerylnick C, Torroni A, Sherman SL, Warren ST. Normal variation at the myotonic dystrophy locus in global human populations. *Am. J. Hum. Genet.* 56, 123–130 (1995).
22. Gomes-Pereira M, Monckton DG. Ethidium bromide modifies the agarose electrophoretic mobility of CAG•CTG alternative DNA structures generated by PCR. *Front. Cell. Neurosci.* 11, 153 (2017).
23. Ying B-W, Fourmy D, Yoshizawa S. Substitution of the use of radioactivity by fluorescence for biochemical studies of RNA. *RNA* 13(11), 2042–2050 (2007).
24. Gupta S, Charakana C, Sreelakshmi Y, Sharma R. Fluorescent dye labeled DNA size standards for molecular mass detection in visible/infrared range. *BMC Res Notes* 4, 12 (2011).

Received: 9 January 2018; Accepted for publication: 9 May 2018

Address correspondence to Geneviève Gourdon, Laboratory CTGDM, Inserm UMR1163, 75015 Paris, France. E-mail: genevieve.gourdon@inserm.fr

To purchase reprints of this article, contact s.cavana@future-science.com

Capture Full-Length Genes for phased sequencing and structural analysis.



sageHLS[™] HMW Library System

- flank genomic regions with CRISPR/Cas9
- target regions up to 500kb
- collect multiple targets