



**HAL**  
open science

## **Ex Vivo Test for Measuring Complement Attack on Endothelial Cells: From Research to Bedside**

Marie-Sophie Meuleman, Anna Duval, Véronique Fremeaux-Bacchi, Lubka T Roumenina, Sophie Chauvet

► **To cite this version:**

Marie-Sophie Meuleman, Anna Duval, Véronique Fremeaux-Bacchi, Lubka T Roumenina, Sophie Chauvet. Ex Vivo Test for Measuring Complement Attack on Endothelial Cells: From Research to Bedside. *Frontiers in Immunology*, 2022, 13, pp.860689. 10.3389/fimmu.2022.860689 . hal-03775681

**HAL Id: hal-03775681**

**<https://hal.sorbonne-universite.fr/hal-03775681v1>**

Submitted on 12 Sep 2022

**HAL** is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers.

L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.

1 ***Ex vivo* test for measuring complement attack on endothelial cells: from**  
2 **research to bedside**

3

4 **Marie-Sophie Meuleman<sup>1</sup>, Anna Duval<sup>1</sup>, Véronique Fremeaux-Bacchi<sup>1</sup>, Lubka T Roumenina<sup>1</sup>,**  
5 **Sophie Chauvet<sup>1</sup>**

6 <sup>1</sup> Centre de Recherche des Cordeliers, INSERM, Sorbonne Université, Université de Paris, Paris,  
7 France

8 **\* Correspondence:**

9 Marie-Sophie MEULEMAN

10 Cordeliers Research Center, INSERM UMRS 1138;

11 15 rue de l'Ecole de Médecine; 75006 Paris, France

12 e-mail: marie-sophie.meuleman@inserm.fr

13

14 **Keywords: Complement; Endothelial Cells; Diagnostics; Prognostics; Therapeutics; Kidney**  
15 **injury, Nephrology, Explorations**

16 Number of words: 5553

17 Number of figures: 4

18 Number of tables: 2

19

20 **Abstract**

21 As part of the innate immune system, the complement system plays a key role in defense against  
22 pathogens and in host cell homeostasis. This enzymatic cascade is rapidly triggered in the presence of  
23 activating surfaces. Physiologically, it is tightly regulated on host cells to avoid uncontrolled activation  
24 and self-damage. In cases of abnormal complement dysregulation/overactivation, the endothelium is  
25 one of the primary targets.

26 Complement has gained momentum as a research interest in the last decade because its dysregulation  
27 has been implicated in the pathophysiology of many human diseases. Thus, it appears to be a promising  
28 candidate for therapeutic intervention.

29 However, detecting abnormal complement activation is challenging. In many pathological conditions,  
30 complement activation occurs locally in tissues. Standard routine exploration of the plasma  
31 concentration of the complement components shows values in the normal range. The available tests to

32 demonstrate such dysregulation with diagnostic, prognostic, and therapeutic implications are limited.  
33 There is a real need to develop tools to demonstrate the implications of complement in diseases and to  
34 explore the complex interplay between complement activation and regulation on human cells. The  
35 analysis of complement deposits on cultured endothelial cells incubated with pathologic human serum  
36 holds promise as a reference assay. This *ex vivo* assay most closely resembles the physiological context.  
37 It has been used to explore complement activation from sera of patients with atypical hemolytic uremic  
38 syndrome, malignant hypertension, elevated liver enzymes low platelet syndrome, sickle cell disease,  
39 pre-eclampsia, and others. In some cases, it is used to adjust the therapeutic regimen with a  
40 complement-blocking drug. Nevertheless, an international standard is lacking, and the mechanism by  
41 which complement is activated in this assay is not fully understood. Moreover, primary cell culture  
42 remains difficult to perform, which probably explains why no standardized or commercialized assay  
43 has been proposed. Here, we review the diseases for which endothelial assays have been applied. We  
44 also compare this test with others currently available to explore complement overactivation. Finally,  
45 we discuss the unanswered questions and challenges to overcome for validating the assays as a tool in  
46 routine clinical practice.

47

### 48 **Introduction**

49 As part of the complex innate immune surveillance system, the complement system plays a key role in  
50 defense against pathogens and in host homeostasis. This enzymatic cascade is rapidly triggered in the  
51 presence of activating surfaces, such as bacteria or apoptotic necrotic cells. However, the cascade is  
52 highly physiologically regulated on host cells to avoid self-aggression. The endothelium is one of the  
53 primary targets of complement dysregulation. There is increasing evidence of complement  
54 implications in the pathophysiology of many human diseases. Many complement-blocking therapeutics  
55 are under development, and some are already available in clinical practice. Nevertheless, detection of  
56 abnormal functioning complement is challenging, because in many pathological conditions C3 and C4  
57 plasma levels, the two main biomarkers of complement activation, remain within normal ranges. The  
58 available tests to demonstrate such overactivation with diagnostic, prognostic, and therapeutic  
59 implications are limited. Methods are poorly standardized, and only a few have functional value.  
60 Therefore, there is a need to develop a robust and standardized tool for identifying infraclinical  
61 complement activation.

62 The final objective is to allow better pathophysiologically based therapeutic management of patients.  
63 The analysis of complement deposits on cultured endothelial cells (EC) incubated with patient serum  
64 holds promise as a reference assay. This approach has been used to explore complement activation in  
65 the sera of patients with atypical hemolytic uremic syndrome (aHUS), malignant hypertension,  
66 hemolysis, elevated liver enzymes, and low platelet (HELLP) syndrome, sickle cell disease (SCD), and  
67 pre-eclampsia. In some cases, adjusting the complement-blocking drugs has been considered.  
68 Nevertheless, the international standard for this test is lacking, and the mechanism by which  
69 complement is activated in this assay is not fully understood.

70 After a brief summary of the complement cascade, we present the mechanisms of complement  
71 activation and how they contribute to cell damage in several human diseases. We then provide an  
72 overview of the tests currently available to explore complement overactivation in routine practice.  
73 Finally, through a comparative analysis of the available endothelial assays for complement exploration,  
74 we discuss the unanswered questions and challenges to overcome to validate the study of complement  
75 deposition on cultured EC as a tool in routine clinical practice.

76

### 77 **The complement system in health and disease**

78 The complement system plays a key role in cell homeostasis, inflammation, and defense against  
79 pathogens. It is the first line of defense. The system comprises more than 30 soluble and membrane-  
80 bound proteins. Three different pathways lead to complement activation: the classical (CP), lectin (LP),  
81 and alternative (AP) pathways. When activated, these serine protease cascades converge to the  
82 formation of two enzymes, C3 convertase and C5 convertase, allowing the generation of the main  
83 effectors of this system: anaphylatoxins (C3a and C5a), opsonin (C3b/iC3b), and the membrane attack  
84 complex (MAC) (C5b-9). CP and LP are initiated by the recognition of pathogen-associated molecular  
85 patterns or damage-associated molecular patterns by pattern-recognition molecules (C1q and mannose-  
86 binding lectin). Conversely, AP is constantly activated at a low level in the fluid phase, generating a  
87 small quantity of C3b. In the presence of an activating surface (apopto-necrotic or bacterial), C3b  
88 covalently binds to the surface, and thus, initiates cell surface C3 convertase formation (C3bBb) and  
89 the AP amplification loop. To avoid self-aggression, AP is highly regulated in the fluid phase and on  
90 the host cell surface by soluble (factor H (FH), factor I (FI)) and membrane-bound regulators

91 (membrane cofactor protein (MCP) or CD46, complement receptor 1 (CR1) or CD35, decay  
92 accelerating factor or CD55, and CD59). In humans, deficiencies in complement regulatory proteins  
93 are associated with rare diseases, such as aHUS, C3 glomerulopathy (C3G), and paroxysmal nocturnal  
94 hemoglobinuria (PNH). However, complement activation triggered by different pathophysiological  
95 processes that overwhelm the capacity of regulation has been increasingly described in a wide spectrum  
96 of diseases.

97

### 98 **Complement implication in diseases**

99 While AP overactivation is the central mechanism of cell and tissue injury in complementopathies  
100 (aHUS, C3G, and PNH), complement is crucial to tissue injury in a wide variety of diseases. These  
101 include age-related macular degeneration (AMD), antibody-mediated rejection (ABMR),  
102 cryoglobulinemic vasculitis (CV), IgA nephropathy (IgAN), systemic lupus erythematosus (SLE), anti-  
103 phospholipid syndrome (APS), ANCA-associated vasculitis (AAV), rheumatoid arthritis (RA),  
104 HELLP syndrome, pre-eclampsia, myasthenia gravis (MG), neuromyelitis optica spectrum disorder  
105 (NMOSD), SCD, and rhabdomyolysis-induced acute kidney injury (RIAKI). To a lesser extent,  
106 complement seems to be involved in an increasing spectrum of human pathological conditions, such  
107 as inflammatory disorders, ischemia/reperfusion, cancer, degenerative disorders (e.g., Alzheimer's  
108 disease, atherosclerosis), and more recently, viral infections that include COVID-19 (1,2) (**Figure 1**).

109 Complementopathies are characterized by a specific cell target of AP-mediated damage. In aHUS and  
110 PNH, AP dysregulation occurs on the cell membrane, EC surface or platelets (3) and erythrocyte  
111 surface (4). In C3G, overactivation of C3 and C5 convertases may occur in the fluid phase or locally  
112 within the glomeruli, where the targeted surface remains to be determined (suggestions include  
113 glomerular EC and mesangial cells). AP dysregulation is a central pathophysiological mechanism in  
114 these diseases. It can be related to innate or acquired abnormalities in complement components, mainly  
115 regulators (FH, FI, or MCP) or C3 convertase components (C3 or FB) (5–16).

116 In diseases with major complement contributions, complement activation can be triggered by one or  
117 another pathway. In CV (17) and ABMR (18), activation occurs through CP in the presence of immune  
118 complexes (IC). In cryoglobulinemia (type II), IC are composed of IgM with rheumatoid factor activity  
119 associated with polyclonal IgG. In ABMR, IC are composed of IgG and donor HLA molecules.  
120 Conversely, despite the disease being triggered by the presence of IC, AP appears to be essential for  
121 disease development in mouse models of RA (19,20) and SLE (21–23). This activation can be  
122 enhanced by apoptotic and necrotic cells due to prior damage (24) or by proteins of the extracellular  
123 matrix (ECM) from damaged cartilage in RA (25). In IgAN, AP (26), and LP (27) activation is  
124 mediated by polymeric IgA. In vitro, a correlation was found between C3 cleavage products (iC3b,  
125 C3c, C3dg) and IgA-A-IgG IC levels, suggesting that IC-containing IgA may act as a surface for  
126 soluble AP activation (28). In AAV, AP may be activated by neutrophil extracellular traps, thus  
127 amplifying complement activation and damage of EC (29). Finally, a disease-specific soluble factor  
128 has been implicated in complement activation. Free heme renders EC more sensitive to complement  
129 activation in SCD (30), aHUS (31), and RIAKI (32). In vitro, thrombin induces C5 cleavage in C5a in  
130 APS (33).

131 Complement activation does not arise from a unique mechanism but can be triggered in several ways  
132 according to the disease pathophysiology. Identification of the precise mechanisms of complement  
133 activation will help determine different potential therapeutic targets within the cascade.

## Measuring complement attack on endothelial cells

134 Complement activation contributes to cell and tissue injuries in different ways. First, it promotes  
135 inflammatory cell recruitment mainly in CV (34), ABMR (35), AMD (36), SLE (37) and RA. C5a and  
136 its receptor C5aR are involved in neutrophil recruitment (38–40) and endothelial activation (41) in  
137 AAV. Complement activation can promote specific disease processes. Thus, MAC can directly affect  
138 collagenase production by synovial fibroblasts in RA (42). In IgAN, mesangial cells exposed to  
139 complement activation and C3 deposition promote phenotypic conversion to a more synthetic and  
140 proliferative state (43). In AMD, C3a and C5a promote choroidal and C5a induces vascular endothelial  
141 growth factor secretion by retinal pigment epithelium (36). In pre-eclampsia, it has been suggested that  
142 the binding of C5a to C5aR expressed on trophoblasts contributes to the acquisition of their anti-  
143 angiogenic phenotype (44).

144 The complement system can also act as an amplifier for other molecules involved in injury. The  
145 C5a/C5aR axis participates in neutrophil recruitment and activation, which in turn can induce  
146 complement activation in AAV (39). C5a induces tissue factor expression by neutrophils, leading to  
147 factor X activation and thrombin generation, which in turn cleaves C5 into C5a in APS (33).

148 Ultimately, several triggers of complement activation and effectors may contribute to cell and tissue  
149 damage in heterogeneous human diseases. The identification of specific triggers of complement  
150 activation and fine pathophysiological mechanisms resulting in cell and tissue complement-mediated  
151 injury is needed to determine the best therapeutic target within the cascade. Complement inhibitor anti-  
152 C5 monoclonal antibody (eculizumab, and more recently its long-acting form, ravulizumab) is the gold  
153 standard in two complementopathies, aHUS and PNH, and has obtained Food and Drug Administration  
154 (FDA) approval for MG and NMO. Avacopan is a C5aR1 antagonist that has also been approved  
155 by the FDA for patients with AAV, another disease with a major complement contribution.  
156 Understanding the detailed mechanisms of complement activation and complement-mediated damage  
157 is necessary to guide the prescription of new complement inhibitors.

### 158 **Overview of the tests exploring complement activation**

#### 159 Quantification of complement components

160 Currently available tests mainly consist in quantification of individual complement components or  
161 activation products.

- 162 ○ For the quantification of individual complement proteins in plasma, various types of immunoassays  
163 are used to determine the concentration of individual complement components. The most common  
164 is nephelometry. Polyclonal antibodies to component are added in excess of the sample and bind  
165 to their target. Quantification is performed by passing a light beam through the sample, which is  
166 distorted by the IC that have formed (45).
- 167 ○ Quantification of complement activation products corresponding to cleavage fragments or  
168 complement protein complexes (C3a, C3dg, C4a, C4d, Ba, Bb, C5a, C3bBbP, MASP2, and sC5b-  
169 9) is possible. Several assays have been described, mostly based on the recognition of a neoepitope  
170 of the complement component in an enzyme-linked immunosorbent assay (ELISA) format. Thus,  
171 C4a and C4d reflect CP/LP activation, Ba, Bb, and C3bBbP reflect AP activation, MASP2 is a key  
172 enzyme in LP activation (46) and increasing soluble C5b-9 reflects TP activation (47). C3a and  
173 C5a are common to the three activation pathways.
- 174 ○ Detection of auto-Abs (anti-FH, FB, C3b, C3bBb, and C1q) targeting complement proteins can be  
175 performed using ELISA (15)

176 Functional assays

- 177 • Quantification of complement function is used to explore the activity of a pathway or the entire  
178 cascade.
- 179 ○ In hemolytic assays, CP activation can be assessed by incubating patient sera with sheep  
180 erythrocytes coated with rabbit anti-sheep red blood cell antibodies (48). In this assay, termed the  
181 CH50 assay, C1q binds to immunoglobulins, initiates the formation of CP C3 convertase, and leads  
182 to MAC assembly and erythrocyte lysis. Hemoglobin release is determined to calculate the number  
183 of hemolytic sites per cell. Activation through AP can be assessed using rabbit or guinea pig  
184 erythrocytes, which are activators of human AP, incubated with patient serum added to ethylene  
185 glycol-bis( $\beta$ -aminoethyl ether) (EGTA), which chelates Ca<sup>2+</sup> and inhibits activation via CP and  
186 LP (49). This hemolytic assay is termed the AP50 assay.
- 187 ○ Liposomes coated with an activator can be used in a similar manner to CH50 assays (50). The main  
188 difference is the readout, which consists of the unquenching of a fluorescent dye and not the lysis  
189 of erythrocytes.
- 190 ○ Assays based on ELISA method can also be used to explore the function of the three pathways.  
191 Microtiter plate wells are coated with recognition structures specific to each pathway (IgM for CP,  
192 mannan or acetylated bovine serum albumin for LP, and LPS for AP). Patient serum is added and  
193 incubated under conditions in which only one pathway is operative at any given time; the other two  
194 pathways are blocked. Finally, activation capacity is detected through the formation of the C5b-9  
195 complex by monoclonal antibodies targeting a neo-epitope in complex-bound C9 (51).

196

- 197 • Different hemolytic assays have been developed to explore specific steps of the AP.
- 198 ○ Sanchez-Corral et al. (52) developed a hemolytic assay to study FH functional defects in aHUS.  
199 The assay relies on the knowledge that sheep erythrocytes are highly sialylated and favor FH  
200 binding, whereas their membrane complement regulators are incompatible with human  
201 complement proteins. Therefore, they are protected from complement lysis due to the binding of  
202 human FH to their surface. In the assay, sheep erythrocytes are incubated with human plasma in  
203 Mg-EGTA buffer, allowing activation of AP only. Normal plasma does not induce lysis, whereas  
204 aHUS plasma with FH functional defects (mutations or autoantibodies) induces lysis under these  
205 conditions (52).
- 206 ○ Hemolytic assays can also be used to study the stabilization of cell-bound AP convertases (53).  
207 This assay has been used to detect C3Nef in C3G cells. Sheep erythrocytes bearing C3 convertase  
208 C3bBb (generated by exposure of sheep erythrocytes bearing C3b to FB and FD) were incubated  
209 with patient IgG. C3Nef activity correlates with residual C3bBb hemolytic sites, and lysis is  
210 developed by the addition of rat serum.

211

- 212 • Staining of tissue sections for the deposition of complement activation products can provide  
213 information about local complement activation in tissue. This can be performed by  
214 immunohistochemistry or immunofluorescence (54). For example, this technique has been used  
215 to study C5b-9 deposition in the skin of patients with aHUS (55).

216 These tests allow only the characterization of a specific molecule or step of the complement cascade.  
217 To reproduce human pathological conditions and their complexity, several authors have proposed the  
218 use of an *ex vivo* endothelial assay. The assay detects and quantifies complement component deposition

## Measuring complement attack on endothelial cells

219 on the EC surface after incubation with human serum. The EC surface is used as the regulating surface.  
220 The objective is to detect abnormal complement deposition that could result from either complement  
221 overactivation exceeding the capacity of regulation, or from a defect in complement regulation in fluid  
222 or on the EC surface. The *ex vivo* endothelial assay is presented in **Figure 2**.

223 We next discuss the advantages and limits of this functional approach.

### 224 **Study of complement deposition on cultured EC**

#### 225 Heterogeneity of endothelial cells populations and their complement regulation

226 EC line blood vessels and constitute an active regulatory organ that has been implicated in vascular  
227 homeostasis, permeability regulation, vasomotor tone, angiogenesis, and diapedesis of immune cells  
228 (56). As first barrier between the blood and interstitium it is in constant equilibrium with the  
229 environment. Thus, heterogeneity in the structure and function of EC is a core property of the  
230 endothelium, allowing diverse vascular functions and regional specificity (57,58). This diversity can  
231 be partially explained by a distinct transcriptional profile (59) in relation to neighboring cells (60).  
232 Hence, EC from different blood vessels have distinct and dynamic expression profiles of complement  
233 components and regulators, which may explain the different susceptibility and specific organ tropism  
234 observed in some complement-mediated diseases (61).

235 At a steady state, EC can produce most complement components and express high levels of  
236 complement regulators on their membranes (**Table S1**). Under inflammatory conditions, complement  
237 component production and regulatory protein expression are modified (**Table 1**). In addition to the  
238 steady state, the modulation of complement protein expression under inflammatory conditions differs  
239 according to the EC type and probably contributes to a specific damage mediated by AP and the  
240 different organ tropisms observed in complement-mediated diseases. Sartain et al. demonstrated that  
241 resting or tumor necrosis factor (TNF)-stimulated brain microvascular EC expressed higher levels of  
242 regulatory molecules (FH, FI, CD46, CD55, and THBD), generated lower levels of C3a and C4a, and  
243 enhanced lower degree AP activation (measured by lower Ba generation) than human renal glomerular  
244 EC (HRGEC) (62). The authors also demonstrated a slight increase in CD46 expression, decrease in  
245 thrombomodulin (TM), and increase in C3 and FB transcription in HRGEC exposed to TNF (63).  
246 These results agree with the prior demonstrations of an increase in C3 and FB production by human  
247 umbilical vein EC (HUVEC) exposed to TNF (64), increased FH transcription and production by  
248 HUVEC exposed to interferon (INF) gamma (65), increased C2, FH, FB, and C1inh transcription, and  
249 decreased C3 production by HUVEC exposed to INF gamma (66). May et al. compared the properties  
250 of four EC types (HRGEC, glomerular EC (GEnC), human microvascular EC (HMEC), and HUVEC)  
251 in the resting state and after overnight exposure to heme (67). While there was no difference in  
252 expression of regulatory factors (MCP, CD55, TM) at resting state, after overnight heme exposure, C3  
253 deposits on glomerular EC were greater than on other EC. This was associated with, and possibly  
254 explained by, weaker FH binding and TM upregulation and lower upregulation of heme-oxygenase 1  
255 (cytoprotective heme-degrading enzyme) compared to HUVEC. Moreover, HUVEC, but not EC, of  
256 glomerular origin were protected from complement deposition after re-challenge with heme (**Table**  
257 **S2**).

258 EC used for *ex vivo* experiments comprise two types: conditionally immortalized EC (CI-EC) and  
259 primary EC (**Table 2**). Primary EC can be difficult to isolate and maintain in culture, and have  
260 a limited lifespan. Moreover, differences in the genetic background of individual donors can  
261 lead to interexperimental variability. In particular, inter-individual heterogeneity in



262 complement regulator expression at the EC surface cannot be excluded. CI-EC has been  
263 developed to overcome these difficulties. HMEC-1 and CI-GEnC are HMEC and GEnCs,  
264 respectively, that have been transfected with SV40 large T antigen (68,69). EA.hy926 cells  
265 were obtained by fusing HUVEC with A549 cells obtained from human lung carcinoma (70).  
266 The EA.hy926 cells were used to generate glycosylphosphatidylinositol-anchored complement  
267 regulatory protein-deficient cells when treated with phosphatidylinositol-specific  
268 phospholipase C. These cells have been used along with the PIGA-mutant TF-1 to study  
269 complement deposits by confocal microscopy and flow cytometry after incubation with serum  
270 from patients with thrombotic microangiopathy (TMA), this test was called the modified Ham  
271 test (71). After incubation with serum from aHUS patients, cell surface C5b-9 deposits were  
272 reportedly higher than after incubation with thrombotic thrombocytopenic purpura (TTP)  
273 serum. Therefore, this test has been considered a tool to distinguish aHUS from TTP. It is  
274 important to note that sC5b-9, reflecting terminal pathway activation and regulation, is elevated  
275 under both aHUS and TTP plasma conditions (72,73). One possibility is that both conditions  
276 are associated with complement activation. However, in aHUS, complement overactivation  
277 exceeds alternative and terminal pathway regulation, leading to C5b-9 deposits. In contrast, in  
278 TTP, complement activation is counterbalanced by complement regulation, leading to sC5b-9  
279 release, but not C5b-9 deposits in the modified Ham test.

280 Micro- or macrovascular origin of the EC tissue lineages also needs to be considered. Complement-  
281 mediated EC injury demonstrates specific cell tropism according to pathophysiological processes. In  
282 HUS and TTP, microvascular EC of dermal, renal, and cerebral origin are more sensitive to apoptosis,  
283 whereas microvascular EC of pulmonary and hepatic origin and macrovascular EC are resistant (74).  
284 Distinct sensitivity of EC to complement attack has also been explored in aHUS and heme exposure.  
285 The demonstration of a distinct EC response in terms of complement regulator expression after a trigger  
286 (here heme) was proposed to partially explain the kidney tropism in this disease (67).

287 HUVEC are primary macrovascular EC isolated from human umbilical cords. These are the most  
288 frequently used cells for *ex vivo* assays (75). If tissue specificity is required, HRGEC (76) or GEnCs  
289 (77), which are both isolated from human glomeruli, can be used. More recently, the use of blood  
290 outgrowth EC obtained from the differentiation of circulating marrow-derived endothelial progenitor  
291 cells isolated from peripheral blood has been proposed (78).

### 292 Comparative analysis of the available endothelial assays

293 These tests consist of the quantification of complement activation products (C3 activation fragments  
294 and C5b-9) deposits on EC by immunofluorescence (IF) measured by confocal microscopy or flow  
295 cytometry (fluorescence-activated cell sorting, FACS) after incubation with a serum sample of interest.  
296 Different protocols have been proposed to study complement activation on the EC surface in several  
297 pathological conditions, including aHUS (31,79–89), TMA of other etiologies (90,91), HELLP  
298 syndrome and pre-eclampsia (92), C3G (15,93), lupus nephritis (LN) (94,95), APS (96,97), SCD (30),  
299 hemolytic anemia (98) and hyperhemolytic transfusion reaction without hemoglobinopathy (99).

300 The general procedure of the *ex vivo* assay and the different protocols are presented in **Figure 3**.

301 *To pre-activate or not pre-activate EC?*

302 Resting EC or EC pre-activated by cytokines, ADP, or heme can be used (**Table S3**) to provide  
303 additional information.

## Measuring complement attack on endothelial cells

304 When resting HUVEC were incubated with aHUS FB mutants added to FB-depleted normal human  
305 serum (NHS), enhanced C3b/iC3b-fragment deposition as measured by an anti-C3c-reacting antibody  
306 was observed (79). The same result was obtained for some cases when aHUS patient serum was  
307 incubated with resting HUVEC (31,80). Nevertheless, incubation with NHS depleted in FB and  
308 reconstituted with other aHUS FB variants (81) or incubation with aHUS serum from patients carrying  
309 some C3 or FH variants (31,80) may be insufficient to induce C3 or C5b-9 deposits. When quiescent  
310 HMEC-1 were incubated with aHUS serum from patients carrying mutations in FH, FI, C3, or  
311 FH/CFHR1 hybrid, enhanced C3c or C5b-9 deposition was reported only if serum was collected during  
312 the acute phases of the disease and not after reaching remission (83,90). Furthermore, deposits on  
313 quiescent HMEC-1 are better correlated with relapse risk during the tapering or discontinuation of  
314 eculizumab (86).

315 To increase test sensitivity, the authors proposed pre-activating EC. Modifications in surface-bound  
316 protein expression enable complement activation. This was achieved in the case of P-selectin  
317 expression on HMEC-1 pre-activated with ADP, LPS, or thrombin (83) or P-selectin expression on  
318 HUVEC or GEnC pre-activated with heme (31,81,100), which could allow C3b binding and C3  
319 convertase formation. Enhanced formation of C3 fragments by TNF/IFN pre-activation and C5b-9  
320 deposition by ADP pre-activation on HUVEC or HMEC-1 cells was described after incubation of these  
321 cells with serum from asymptomatic carriers of mutations in AP regulatory proteins or C3 (80,83). The  
322 normal range was established when pre-activated EC were incubated with sera from healthy donors.  
323 In addition, serum from healthy family members without the mutation was within the normal range in  
324 this assay (80).

325 *What kind of blood samples might be incubated with EC?*

326 Serum has been used as the source of complement proteins in the vast majority of the tests described  
327 above. One limitation of these tests, particularly when deposits are detected by IF, is the variation in  
328 the results, reportedly from 30% to 52% when activated HMEC-1 were incubated with serum collected  
329 at the acute phase of aHUS (92). To reduce this variation, Palomo et al. proposed the use of activated  
330 plasma, which refers to citrated plasma mixed 1:1 with a control serum pool. Using this approach, the  
331 authors derived a coefficient of variation of 9% to 18% (92). C3 consumption by the patient or loss of  
332 C3 activity during the pre-analytical phase are also potential factors responsible for this variation (101).  
333 Finally, for all complement assays and to avoid in vitro complement activation, proper blood collection  
334 and processing must be achieved (102). Processing of plasma or serum sample must be performed  
335 within a few hours of collection, with storage at  $-80^{\circ}\text{C}$  and defrosting immediately before use to avoid  
336 repeated freezing and thawing.

337 To explore the functional consequences of autoantibodies against C3 and properdin in SLE, Vasilev et  
338 al. and Radanova et al. incubated HUVEC with NHS supplemented with purified IgG from patients  
339 positive for such autoantibodies (94,95). Using this strategy, complement deposition on EC can be  
340 directly ascribed to the addition of autoantibodies to NHS. The same approach was applied for anti-  
341 C3b/FB autoantibodies in patients with C3G (15). To understand the mechanism behind complement  
342 deposits on EC from patients with SCD, microvesicles from normal or patient-derived erythrocytes  
343 were added to normal serum to model the disease condition. Enhanced binding of the C3 activation  
344 products was demonstrated (30,103).

345 *Which controls are relevant?*

346 Most often, NHS is used as a negative control (15,30,31,79–84) (**Table S4**). An important aspect to  
347 consider is the inter-individual variability in deposits induced by normal sera. FACS analysis has  
348 revealed that this variability was relatively low when sera from 50 healthy donors were tested (80).  
349 However, this is a concern, particularly when deposits are detected by IF. This has not been directly  
350 reported, but has been suggested by the use of pooled sera in more recent papers (86,92) and our own  
351 experience. Aiello et al. reported that C3 and C5b-9 deposits obtained after a single healthy subject  
352 serum (N=12) incubation range from 0.5 to 1.5 fold increase of stained surface area compared to pooled  
353 serum (from 10 healthy donors) run in parallel (88).

354 Several authors did not use any positive controls for their experiments (83,84,86,91,92). The  
355 comparison was only made with the deposits obtained with negative controls. It might be interesting  
356 to position the results on a scale. Positive controls with published data are FH or FI depleted NHS  
357 (15,80,82) or normal serum supplemented with blocking anti-FH antibodies targeting the N-terminus  
358 or C-terminus (31,81) or with FH19-20, corresponding to the two last domains of FH, able to compete  
359 with the full FH protein for cell surface binding (98).

360 The main issues with this type of assay are the lack of validated international standards as well as  
361 standardized positive and negative controls. The variability of the results in samples from healthy  
362 donors needs to be studied extensively to determine the appropriate cutoff. In addition, the impact of  
363 C3 or other complement protein consumption in the patient and the influence of the pre-analytical  
364 phase must be determined to avoid false positive and false negative results.

365 *Which deposits should be measured?*

366 The objective of these tests is to demonstrate and explore complement overactivation or dysregulation  
367 on the EC surface after incubation with blood samples of interest. This is enabled by quantification of  
368 the deposition of complement component products resulting from activation or regulation. C3c (a  
369 common epitope to C3, C3(H<sub>2</sub>O), C3b, and iC3b) (which reflects C3 convertase activity and the early  
370 phase of the complement cascade) can be detected by polyclonal anti-C3c antibody. Antibody targeting  
371 C5b-9 reveals the final step of the cascade. When a signal is detected on the cell membrane, it can be  
372 assumed that the detected fragment is C3b or iC3b covalently attached to the surface. Nevertheless,  
373 heme-activated EC and likely ADP-activated EC (104,105) express P-selectin, which recruits C3b,  
374 C3(H<sub>2</sub>O), and a C3(H<sub>2</sub>O)-like form of C3 generated after contact with heme (31,100). Properdin also  
375 binds to heme-exposed or stressed EC, promoting complement activation in a similar manner without  
376 covalent C3b binding (98). This is an additional mechanism for amplification of complement activation  
377 on the EC surface. C5b-9 deposits may be more relevant in identifying dysregulation at any step.  
378 Nevertheless, early dysregulation can induce C3 activation fragment deposits without C5b-9 formation  
379 because of TP regulation. C5b-9 is readily detectable by IF but is much more difficult to detect by  
380 FACS because of the weak shifts of the peaks. To test for CP participation, the presence of C4d-positive  
381 deposits was also investigated (83,90). Staining can also be performed under the same conditions for  
382 von Willebrand Factor, C5aR1, P-selectin, and others (88).

383 *Evaluation of activated pathways*

384 The test can be modified to assess which complement pathway is activated in given pathological  
385 settings. The test can be performed under different conditions to avoid CP and LP contributions, which  
386 include C2 (31) or C1q (106) depleted NHS, addition of SCR1 (88) or Mg-EGTA buffer (31,79,94,95).  
387 EGTA chelates Ca<sup>2+</sup>, which is crucial for CP and LP activation, whereas AP depends on Mg<sup>2+</sup>. If AP  
388 has to be inhibited, FB-depleted NHS can be used. These reagents are applicable for test conditions,

## Measuring complement attack on endothelial cells

389 where the activating factor is added externally to the serum (i.e., IgG, heme, microvesicles, etc.). When  
390 patient samples are used directly, the same effect can be achieved by inhibiting C1q, C4, FB, or  
391 properdin with blocking antibodies, protein constructs, or small molecules, if available (98).  
392 Quantification of complement activation products (split fragments generated by cleavage of  
393 complement components or protein complexes when activated components bind their target (i.e., C3a,  
394 C4a, Ba, Bb, C5a, and sC5b-9) in the supernatant might be an additional element to study complement  
395 cascade activation.

396 *Which techniques are used for detection and quantification?*

397 The two main detection techniques commonly used are FACS and IF. HUVEC pre-activated with heme  
398 and then incubated with NHS or aHUS serum showed results similar by FACS or IF detection(31). IF  
399 directly analyzes deposits on EC grown on slides. FACS requires a cell detachment step before  
400 staining, with the potential risk of losing a part of the deposit signal. In contrast, as mentioned by  
401 Gavriilaki et al., obtaining quantitative data by IF requires confocal microscopy and further analysis  
402 using specialized software (71). When IF is used, the area occupied by fluorescent staining in fields  
403 systematically digitized along the surface is quantified. The quantified results expressed as the mean  
404 of the square number of pixels per field are compared with the negative control (83,84,86,90,92).  
405 Considering the number of EC on which fluorescence has been measured and the staining intensity  
406 might appear relevant. In contrast, FACS allows the rapid and objective quantification of deposits.

407 *What are the functional consequences of such deposits?*

408 If enhancement of complement fragment deposits on EC is interpreted as pathogenic, the functional  
409 consequences of such deposits must be questioned. Lactate dehydrogenase release from EC reflects  
410 cell damage. This release can be measured in the cell culture supernatant (106). Analysis of  
411 complement deposits can also be associated with a cell viability assay, corresponding to a colorimetric  
412 assay based on cleavage of the WST-1 tetrazolium salt by mitochondrial dehydrogenases in viable cells  
413 (71). Cellular integrity can be verified by May-Grunwald Giemsa staining (89). Direct cell death rarely  
414 occurs under these experimental conditions. Experiments testing cell activation status by complement  
415 overactivation have not been reported in the literature and are needed to further understand the impact  
416 of complement on endothelial injury. Analysis of transcriptomic modifications in EC exposed to  
417 complement deposits under several conditions could also be of interest.

418

419 Clinical and therapeutic relevance of the obtained results

420 The *ex vivo* EC assay, consisting in the quantification of complement activation products (C3 activation  
421 fragments or C5b-9) deposits on EC (by IF measured on confocal microscopy or FACS), after  
422 incubation with a serum sample of interest, was first used for specific characterization of complement  
423 component abnormalities (79–82,84) or exploration of mechanisms implicated in EC injury (31) in the  
424 main complementopathy, aHUS. The assay was then used to demonstrate and explore complement  
425 activation and participation in the pathophysiology of several diseases, including C3G (15,93), HELLP  
426 syndrome and pre-eclampsia (92,107), TMA associated with severe hypertension (90,108), drug-  
427 induced TMA (109), SCD (30), hemolytic anemia (98,99), SLE (94,95), and APS (97). Demonstration  
428 of increasing complement deposits on EC incubated with pathological sera is not sufficient to  
429 determine what is responsible for complement activation at the EC surface. Modulation of the test

430 conditions can help in detailing complement activation. This was the case when complement activation  
431 was inhibited by the addition of hemopexin to the sera of patients with SCD (30).

432 Noris et al. and Galbusera et al. also proposed the use of this *ex vivo* EC test to monitor eculizumab  
433 therapy in patients with aHUS (83,86). During eculizumab tapering or discontinuation, disease relapse  
434 preceded or was associated with an increase in C5b-9 deposits on resting HMEC-1 in all patients. In  
435 contrast, only one patient without relapse showed increased deposits (86). In clinical practice, CH50 is  
436 the only routine test used to monitor eculizumab therapy. CH50 is reportedly strongly suppressed in  
437 patients receiving eculizumab according to the standard protocol. However, CH50 does not allow  
438 monitoring of eculizumab dosage tapering or discontinuation, as it is not well correlated with relapse  
439 risk (83,86). Eculizumab therapy monitoring using the Wieslab® complement system screen (110) or  
440 the modified Ham test (111) has also been proposed. Thus, the *ex vivo* EC assay could represent a test  
441 of interest to a better personalized complement-blocking therapy, but first needs to be more  
442 standardized.

443 This test can also be used to better classify and assess the prognosis of specific diseases. This is the  
444 case for hypertensive TMA, as Timmermans et al. demonstrated in a cohort of hypertension  
445 emergencies associated with TMA(108). The authors demonstrated a statistical association between  
446 increased C5b-9 deposition in the EC *ex vivo* test and kidney survival. Moreover, they reported an  
447 improvement in renal function for those with increased deposits treated with eculizumab. The authors  
448 proposed a classification of TMA-hypertensive emergency based on the EC *ex vivo* test (108).

449 Finally, many new anti-complement drugs targeting specific steps of the cascade have been under  
450 development in recent years (112). A standardized and validated assay to study complement activation  
451 could be a useful tool in their development.

452

### 453 **Discussion and Conclusion**

454 The increasing demonstration of complement involvement in the pathophysiology of many human  
455 diseases has mandated the development of tools to finely explore complement activation. Complement  
456 is a complex enzymatic cascade that is highly regulated in constant interplay with its environment. The  
457 current arsenal for complement exploration does not provide functional characterization and does not  
458 report on the complex interplay between complement and its environment, particularly the cell surface.

459 The development of tests with these capabilities could allow for a deeper exploration of the  
460 mechanisms of complement activation in several diseases. This information could inform the  
461 development of a complement blocking therapeutic strategy based on pathophysiological mechanisms.

462 *Ex vivo* complement activation on EC represents a promising tool for demonstrating and exploring  
463 complement activation. It not only recapitulates complex complement cascade regulation *in vivo*, but  
464 also allows modification of several steps of the experimental procedure to characterize complement  
465 activation mechanisms.

466 However, there are still unanswered questions hindering broad used. The first is the variability in the  
467 results and the inter-individual variability in deposits induced by normal sera. Comprehension of the  
468 precise mechanism responsible for complement deposition in this assay would improve its better use.  
469 The second issue is to standardize the main steps of the procedure to improve the interexperimental  
470 comparison.

471 The use of such a test could be multiple, including molecular functional characterization, disease  
472 pathophysiology exploration, prognosis classification, complement targeting drug development, and  
473 complement therapeutic monitoring. The use of standardized conditions will expand the field of this  
474 promising tool.

### 475 **Authors Contributions**

476 MSM, SC, and LR conceptualized and conceived the manuscript. MSM drafted the manuscript,  
477 including the literature search, reading, and writing. SC, VFB, AD, and LR edited and critically  
478 evaluated the manuscript. All authors have contributed to the manuscript and approved the submitted  
479 version.

### 480 **Conflict of Interest**

481 The authors declare that the review was conducted in the absence of any commercial or financial  
482 relationships that could be construed as potential conflicts of interest.

483

### 484 **References**

- 485 1. Magro C, Mulvey JJ, Laurence J, Seshan S, Crowson AN, Dannenberg AJ, Salvatore S, Harp  
486 J, Nuovo GJ. Docked SARS CoV-2 Proteins within the Cutaneous and Subcutaneous  
487 Microvasculature and their Role in the Pathogenesis of Severe COVID-19. *Hum Pathol* (2020) doi:  
488 10.1016/j.humpath.2020.10.002
- 489 2. Magro C, Mulvey JJ, Laurence J, Seshan S, Crowson AN, Dannenberg AJ, Salvatore S, Harp  
490 J, Nuovo GJ. Docked SARS CoV-2 Proteins within the Cutaneous and Subcutaneous  
491 Microvasculature and their Role in the Pathogenesis of Severe COVID-19. *Hum Pathol* (2020) doi:  
492 10.1016/j.humpath.2020.10.002
- 493 3. Fakhouri F, Zuber J, Frémeaux-Bacchi V, Loirat C. Haemolytic uraemic syndrome. *The*  
494 *Lancet* (2017) **390**:681–696. doi: 10.1016/S0140-6736(17)30062-4
- 495 4. Brodsky RA. Paroxysmal nocturnal hemoglobinuria. *Blood* (2014) **124**:2804–2811. doi:  
496 10.1182/blood-2014-02-522128
- 497 5. Roumenina LT, Loirat C, Dragon-Durey M-A, Halbwachs-Mecarelli L, Sautes-Fridman C,  
498 Frémeaux-Bacchi V. Alternative complement pathway assessment in patients with atypical HUS. *J*  
499 *Immunol Methods* (2011) **365**:8–26. doi: 10.1016/j.jim.2010.12.020
- 500 6. Heinen S, Józsi M, Hartmann A, Noris M, Remuzzi G, Skerka C, Zipfel PF. Hemolytic  
501 Uremic Syndrome: A Factor H Mutation (E1172Stop) Causes Defective Complement Control at the  
502 Surface of Endothelial Cells. *J Am Soc Nephrol* (2007) **18**:506–514. doi: 10.1681/ASN.2006091069
- 503 7. Dragon-Durey M-A. Heterozygous and Homozygous Factor H Deficiencies Associated with  
504 Hemolytic Uremic Syndrome or Membranoproliferative Glomerulonephritis: Report and Genetic  
505 Analysis of 16 Cases. *J Am Soc Nephrol* (2004) **15**:787–795. doi:  
506 10.1097/01.ASN.0000115702.28859.A7



- 507 8. Lehtinen MJ, Rops AL, Isenman DE, van der Vlag J, Jokiranta TS. Mutations of Factor H  
508 Impair Regulation of Surface-bound C3b by Three Mechanisms in Atypical Hemolytic Uremic  
509 Syndrome. *J Biol Chem* (2009) **284**:15650–15658. doi: 10.1074/jbc.M900814200
- 510 9. Manuelian T, Hellwage J, Meri S, Caprioli J, Noris M, Heinen S, Jozsi M, Neumann HPH,  
511 Remuzzi G, Zipfel PF. Mutations in factor H reduce binding affinity to C3b and heparin and surface  
512 attachment to endothelial cells in hemolytic uremic syndrome. *J Clin Invest* (2003) **111**:1181–1190.  
513 doi: 10.1172/JCI16651
- 514 10. Kavanagh D, Richards A, Noris M, Hauhart R, Liszewski MK, Karpman D, Goodship JA,  
515 Fremeaux-Bacchi V, Remuzzi G, Goodship THJ, et al. Characterization of mutations in complement  
516 factor I (CFI) associated with hemolytic uremic syndrome. *Mol Immunol* (2008) **45**:95–105. doi:  
517 10.1016/j.molimm.2007.05.004
- 518 11. Bienaime F, Dragon-Durey M-A, Regnier CH, Nilsson SC, Kwan WH, Blouin J, Jablonski  
519 M, Renault N, Rameix-Welti M-A, Loirat C, et al. Mutations in components of complement  
520 influence the outcome of Factor I-associated atypical hemolytic uremic syndrome. *Kidney Int* (2010)  
521 **77**:339–349. doi: 10.1038/ki.2009.472
- 522 12. Fremeaux-Bacchi V, Moulton EA, Kavanagh D, Dragon-Durey M-A, Blouin J, Caudy A,  
523 Arzouk N, Cleper R, Francois M, Guest G, et al. Genetic and Functional Analyses of Membrane  
524 Cofactor Protein (CD46) Mutations in Atypical Hemolytic Uremic Syndrome. *J Am Soc Nephrol*  
525 (2006) **17**:2017–2025. doi: 10.1681/ASN.2005101051
- 526 13. Dragon-Durey M-A, Loirat C, Cloarec S, Macher M-A, Blouin J, Nivet H, Weiss L, Fridman  
527 WH, Frémeaux-Bacchi V. Anti-Factor H Autoantibodies Associated with Atypical Hemolytic  
528 Uremic Syndrome. *J Am Soc Nephrol* (2005) **16**:555–563. doi: 10.1681/ASN.2004050380
- 529 14. Blanc C, Togarsimalemath SK, Chauvet S, Le Quintrec M, Moulin B, Buchler M, Jokiranta  
530 TS, Roumenina LT, Fremeaux-Bacchi V, Dragon-Durey M-A. Anti-Factor H Autoantibodies in C3  
531 Glomerulopathies and in Atypical Hemolytic Uremic Syndrome: One Target, Two Diseases. *J*  
532 *Immunol* (2015) **194**:5129–5138. doi: 10.4049/jimmunol.1402770
- 533 15. Marinozzi MC, Roumenina LT, Chauvet S, Hertig A, Bertrand D, Olagne J, Frimat M,  
534 Ulinski T, Deschênes G, Burtey S, et al. Anti-Factor B and Anti-C3b Autoantibodies in C3  
535 Glomerulopathy and Ig-Associated Membranoproliferative GN. *J Am Soc Nephrol* (2017) **28**:1603–  
536 1613. doi: 10.1681/ASN.2016030343
- 537 16. Iatropoulos P, Noris M, Mele C, Piras R, Valoti E, Bresin E, Curreri M, Mondo E, Zito A,  
538 Gamba S, et al. Complement gene variants determine the risk of immunoglobulin-associated MPGN  
539 and C3 glomerulopathy and predict long-term renal outcome. *Mol Immunol* (2016) **71**:131–142. doi:  
540 10.1016/j.molimm.2016.01.010
- 541 17. Sansonno D, Gesualdo L, Manno C, Schena FP, Dammacco F. Hepatitis C virus-related  
542 proteins in kidney tissue from hepatitis C virus-infected patients with cryoglobulinemic  
543 membranoproliferative glomerulonephritis. *Hepatology* (1997) **25**:1237–1244. doi:  
544 10.1002/hep.510250529
- 545 18. Stegall MD, Chedid MF, Cornell LD. The role of complement in antibody-mediated rejection  
546 in kidney transplantation. *Nat Rev Nephrol* (2012) **8**:670–678. doi: 10.1038/nrneph.2012.212

## Measuring complement attack on endothelial cells

- 547 19. Banda NK, Takahashi K, Wood AK, Holers VM, Arend WP. Pathogenic Complement  
548 Activation in Collagen Antibody- Induced Arthritis in Mice Requires Amplification by the  
549 Alternative Pathway. *J Immunol* (2007) **179**:4101–4109. doi: 10.4049/jimmunol.179.6.4101
- 550 20. Banda NK, Thurman JM, Kraus D, Wood A, Carroll MC, Arend WP, Holers VM. Alternative  
551 Complement Pathway Activation Is Essential for Inflammation and Joint Destruction in the Passive  
552 Transfer Model of Collagen-Induced Arthritis. *J Immunol* (2006) **177**:1904–1912. doi:  
553 10.4049/jimmunol.177.3.1904
- 554 21. Watanabe H, Garnier G, Circolo A, Wetsel RA, Ruiz P, Holers VM, Boackle SA, Colten HR,  
555 Gilkeson GS. Modulation of Renal Disease in MRL/ *lpr* Mice Genetically Deficient in the  
556 Alternative Complement Pathway Factor B. *J Immunol* (2000) **164**:786–794. doi:  
557 10.4049/jimmunol.164.2.786
- 558 22. Elliott MK, Jarmi T, Ruiz P, Xu Y, Holers VM, Gilkeson GS. Effects of complement factor D  
559 deficiency on the renal disease of MRL/*lpr* mice. *Kidney Int* (2004) **65**:129–138. doi: 10.1111/j.1523-  
560 1755.2004.00371.x
- 561 23. Sekine H, Kinser TTH, Qiao F, Martinez E, Paulling E, Ruiz P, Gilkeson GS, Tomlinson S.  
562 The benefit of targeted and selective inhibition of the alternative complement pathway for  
563 modulating autoimmunity and renal disease in MRL/*lpr* mice. *Arthritis Rheum* (2011) **63**:1076–1085.  
564 doi: 10.1002/art.30222
- 565 24. Nauta AJ, Trouw LA, Daha MR, Tijssma O, Nieuwland R, Schwaeble WJ, Gingras AR,  
566 Mantovani A, Hack EC, Roos A. Direct binding of C1q to apoptotic cells and cell blebs induces  
567 complement activation. *Eur J Immunol* (2002) **32**:1726. doi: 10.1002/1521-  
568 4141(200206)32:6<1726::AID-IMMU1726>3.0.CO;2-R
- 569 25. Sjöberg AP, Manderson GA, Mörgelin M, Day AJ, Heinegård D, Blom AM. Short leucine-  
570 rich glycoproteins of the extracellular matrix display diverse patterns of complement interaction and  
571 activation. *Mol Immunol* (2009) **46**:830–839. doi: 10.1016/j.molimm.2008.09.018
- 572 26. Russell MW, Mansa B. Complement-Fixing Properties of Human IgA Antibodies Alternative  
573 Pathway Complement Activation by Plastic-Bound, But Not Specific Antigen-Bound, IgA. *Scand J*  
574 *Immunol* (1989) **30**:175–183. doi: 10.1111/j.1365-3083.1989.tb01199.x
- 575 27. Roos A, Bouwman LH, van Gijlswijk-Janssen DJ, Faber-Krol MC, Stahl GL, Daha MR.  
576 Human IgA Activates the Complement System Via the Mannan-Binding Lectin Pathway. *J Immunol*  
577 (2001) **167**:2861–2868. doi: 10.4049/jimmunol.167.5.2861
- 578 28. Maillard N, Boerma LJ, Hall S, Huang ZQ, Mrug M, Moldoveanu Z, Julian B, Renfrow MB,  
579 Novak J. L'analyse protéomique de complexes immuns artificiels IgA1-IgG révèle une association  
580 avec des formes activées de C3. *Néphrologie Thérapeutique* (2014) **10**:279. doi:  
581 10.1016/j.nephro.2014.07.347
- 582 29. Schreiber A, Rousselle A, Becker JU, von Mässenhausen A, Linkermann A, Kettritz R.  
583 Necroptosis controls NET generation and mediates complement activation, endothelial damage, and  
584 autoimmune vasculitis. *Proc Natl Acad Sci* (2017) **114**:E9618–E9625. doi:  
585 10.1073/pnas.1708247114



- 586 30. Roumenina LT, Chadebech P, Bodivit G, Vieira-Martins P, Grunenwald A, Boudhabhay I,  
587 Poillerat V, Pakdaman S, Kiger L, Jouard A, et al. Complement activation in sickle cell disease:  
588 Dependence on cell density, hemolysis and modulation by hydroxyurea therapy. *Am J Hematol*  
589 (2020) **95**:456–464. doi: 10.1002/ajh.25742
- 590 31. Frimat M, Tabarin F, Dimitrov JD, Poitou C, Halbwachs-Mecarelli L, Fremeaux-Bacchi V,  
591 Roumenina LT. Complement activation by heme as a secondary hit for atypical hemolytic uremic  
592 syndrome. *Blood* (2013) **122**:282–292. doi: 10.1182/blood-2013-03-489245
- 593 32. Boudhabhay I, Poillerat V, Grunenwald A, Torset C, Leon J, Daugan MV, Lucibello F, El  
594 Karoui K, Ydee A, Chauvet S, et al. Complement activation is a crucial driver of acute kidney injury  
595 in rhabdomyolysis. *Kidney Int* (2020)S0085253820312448. doi: 10.1016/j.kint.2020.09.033
- 596 33. Ritis K, Doumas M, Mastellos D, Micheli A, Giaglis S, Magotti P, Rafail S, Kartalis G,  
597 Sideras P, Lambris JD. A Novel C5a Receptor-Tissue Factor Cross-Talk in Neutrophils Links Innate  
598 Immunity to Coagulation Pathways. *J Immunol* (2006) **177**:4794–4802. doi:  
599 10.4049/jimmunol.177.7.4794
- 600 34. Sansonno D, Dammacco F. Hepatitis C virus, cryoglobulinaemia, and vasculitis: immune  
601 complex relations. *Lancet Infect Dis* (2005) **5**:227–236. doi: 10.1016/S1473-3099(05)70053-0
- 602 35. Loupy A, Lefaucheur C. Antibody-Mediated Rejection of Solid-Organ Allografts. *N Engl J*  
603 *Med* (2018) **379**:1150–1160. doi: 10.1056/NEJMra1802677
- 604 36. Nozaki M, Raisler BJ, Sakurai E, Sarma JV, Barnum SR, Lambris JD, Chen Y, Zhang K,  
605 Ambati BK, Baffi JZ, et al. Drusen complement components C3a and C5a promote choroidal  
606 neovascularization. *Proc Natl Acad Sci* (2006) **103**:2328–2333. doi: 10.1073/pnas.0408835103
- 607 37. Botto M, Dell’Agnola C, Bygrave AE, Thompson EM, Cook HT, Petry F, Loos M, Pandolfi  
608 PP, Walport MJ. Homozygous C1q deficiency causes glomerulonephritis associated with multiple  
609 apoptotic bodies. *Nat Genet* (1998) **19**:56–59. doi: 10.1038/ng0598-56
- 610 38. Dick J, Gan P-Y, Ford SL, Odobasic D, Alikhan MA, Loosen SH, Hall P, Westhorpe CL, Li  
611 A, Ooi JD, et al. C5a receptor 1 promotes autoimmunity, neutrophil dysfunction and injury in  
612 experimental anti-myeloperoxidase glomerulonephritis. *Kidney Int* (2018) **93**:615–625. doi:  
613 10.1016/j.kint.2017.09.018
- 614 39. Schreiber A, Xiao H, Jennette JC, Schneider W, Luft FC, Kettritz R. C5a Receptor Mediates  
615 Neutrophil Activation and ANCA-Induced Glomerulonephritis. *J Am Soc Nephrol* (2009) **20**:289–  
616 298. doi: 10.1681/ASN.2008050497
- 617 40. Camous L, Roumenina L, Bigot S, Brachemi S, Frémeaux-Bacchi V, Lesavre P, Halbwachs-  
618 Mecarelli L. Complement alternative pathway acts as a positive feedback amplification of neutrophil  
619 activation. *Blood* (2011) **117**:1340–1349. doi: 10.1182/blood-2010-05-283564
- 620 41. Foreman KE, Vaporciyan AA, Bonish BK, Jones ML, Johnson KJ, Glovsky MM, Eddy SM,  
621 Ward PA. C5a-induced expression of P-selectin in endothelial cells. *J Clin Invest* (1994) **94**:1147–  
622 1155. doi: 10.1172/JCI117430
- 623 42. Jahn B, Von Kempis J, Krämer KL, Filsinger S, Hänsch GM. Interaction of the terminal

## Measuring complement attack on endothelial cells

- 624 complement components C5b-9 with synovial fibroblasts: binding to the membrane surface leads to  
625 increased levels in collagenase-specific mRNA. *Immunology* (1993) **78**:329–334.
- 626 43. Wan J-X, Fukuda N, Endo M, Tahira Y, Yao E-H, Matsuda H, Ueno T, Matsumoto K.  
627 Complement 3 is involved in changing the phenotype of human glomerular mesangial cells. *J Cell*  
628 *Physiol* (2007) **213**:495–501. doi: 10.1002/jcp.21129
- 629 44. Ma Y, Kong L-R, Ge Q, Lu Y-Y, Hong M-N, Zhang Y, Ruan C-C, Gao P-J. Complement 5a-  
630 mediated trophoblasts dysfunction is involved in the development of pre-eclampsia. *J Cell Mol Med*  
631 (2017) doi: 10.1111/jcmm.13466
- 632 45. Ekdahl KN, Persson B, Mohlin C, Sandholm K, Skattum L, Nilsson B. Interpretation of  
633 Serological Complement Biomarkers in Disease. *Front Immunol* (2018) **9**: doi:  
634 10.3389/fimmu.2018.02237
- 635 46. Elhadad S, Chapin J, Copertino D, Van Besien K, Ahamed J, Laurence J. MASP2 levels are  
636 elevated in thrombotic microangiopathies: association with microvascular endothelial cell injury and  
637 suppression by anti-MASP2 antibody narsoplimab. *Clin Exp Immunol* (2021) **203**:96–104. doi:  
638 10.1111/cei.13497
- 639 47. Mollnes TE, Lea T, Froland SS, Harboe M. Quantification of the Terminal Complement  
640 Complex in Human Plasma by an Enzyme-Linked Immunosorbent Assay Based on Monoclonal  
641 Antibodies against a Neoantigen of the Complex. *Scand J Immunol* (1985) **22**:197–202. doi:  
642 10.1111/j.1365-3083.1985.tb01871.x
- 643 48. Mayer MM. On the destruction of erythrocytes and other cells by antibody and complement.  
644 *Cancer Res* (1961) **21**:1262–1269.
- 645 49. Platts-Mills TA, Ishizaka K. Activation of the alternate pathway of human complements by  
646 rabbit cells. *J Immunol Baltim Md 1950* (1974) **113**:348–358.
- 647 50. Yamamoto S, Kubotsu K, Kida M, Kondo K, Matsuura S, Uchiyama S, Yonekawa O, Kanno  
648 T. Automated homogeneous liposome-based assay system for total complement activity. *Clin Chem*  
649 (1995) **41**:586–590.
- 650 51. Seelen MA, Roos A, Wieslander J, Mollnes TE, Sjöholm AG, Wurzner R, Loos M, Tedesco  
651 F, Sim RB, Garred P, et al. Functional analysis of the classical, alternative, and MBL pathways of the  
652 complement system: standardization and validation of a simple ELISA. *J Immunol Methods* (2005)  
653 **296**:187–198. doi: 10.1016/j.jim.2004.11.016
- 654 52. Sanchezcorral P. Functional analysis in serum from atypical Hemolytic Uremic Syndrome  
655 patients reveals impaired protection of host cells associated with mutations in factor H. *Mol Immunol*  
656 (2004) **41**:81–84. doi: 10.1016/j.molimm.2004.01.003
- 657 53. Daha MR, Fearon DT, Austen KF. C3 nephritic factor (C3NeF): stabilization of fluid phase  
658 and cell-bound alternative pathway convertase. *J Immunol Baltim Md 1950* (1976) **116**:1–7.
- 659 54. Mori H, Cardiff RD. “Methods of Immunohistochemistry and Immunofluorescence:  
660 Converting Invisible to Visible.” In: Ursini-Siegel J, Beauchemin N, editors. *The Tumor*  
661 *Microenvironment*. New York, NY: Springer New York (2016). p. 1–12 doi: 10.1007/978-1-4939-

662 3801-8\_1

- 663 55. Magro CM, Momtahan S, Mulvey JJ, Yassin AH, Kaplan RB, Laurence JC. Role of the skin  
664 biopsy in the diagnosis of atypical hemolytic uremic syndrome. *Am J Dermatopathol* (2015) **37**:349–  
665 356; quiz 357–359. doi: 10.1097/DAD.0000000000000234
- 666 56. Sturtzel C. “Endothelial Cells,.” In: Sattler S, Kennedy-Lydon T, editors. *The Immunology of*  
667 *Cardiovascular Homeostasis and Pathology*. Cham: Springer International Publishing (2017). p. 71–  
668 91 doi: 10.1007/978-3-319-57613-8\_4
- 669 57. Aird WC. Endothelial Cell Heterogeneity. *Cold Spring Harb Perspect Med* (2012)  
670 **2**:a006429–a006429. doi: 10.1101/cshperspect.a006429
- 671 58. Dumas SJ, Meta E, Borri M, Luo Y, Li X, Rabelink TJ, Carmeliet P. Phenotypic diversity and  
672 metabolic specialization of renal endothelial cells. *Nat Rev Nephrol* (2021) **17**:441–464. doi:  
673 10.1038/s41581-021-00411-9
- 674 59. Chi J-T, Chang HY, Haraldsen G, Jahnsen FL, Troyanskaya OG, Chang DS, Wang Z,  
675 Rockson SG, van de Rijn M, Botstein D, et al. Endothelial cell diversity revealed by global  
676 expression profiling. *Proc Natl Acad Sci* (2003) **100**:10623–10628. doi: 10.1073/pnas.1434429100
- 677 60. Jambusaria A, Hong Z, Zhang L, Srivastava S, Jana A, Toth PT, Dai Y, Malik AB, Rehman J.  
678 Endothelial heterogeneity across distinct vascular beds during homeostasis and inflammation. *eLife*  
679 (2020) **9**: doi: 10.7554/eLife.51413
- 680 61. Roumenina LT, Rayes J, Frimat M, Fremeaux-Bacchi V. Endothelial cells: source, barrier,  
681 and target of defensive mediators. *Immunol Rev* (2016) **274**:307–329. doi: 10.1111/imr.12479
- 682 62. Sartain SE, Turner NA, Moake JL. Brain microvascular endothelial cells exhibit lower  
683 activation of the alternative complement pathway than glomerular microvascular endothelial cells. *J*  
684 *Biol Chem* (2018) **293**:7195–7208. doi: 10.1074/jbc.RA118.002639
- 685 63. Sartain SE, Turner NA, Moake JL. TNF Regulates Essential Alternative Complement  
686 Pathway Components and Impairs Activation of Protein C in Human Glomerular Endothelial Cells. *J*  
687 *Immunol* (2016) **196**:832–845. doi: 10.4049/jimmunol.1500960
- 688 64. Kawakami Y, Watanabe Y, Yamaguchi M, Haruko Sakaguchi, Kono I, Ueki A. TNF- $\alpha$   
689 stimulates the biosynthesis of complement C3 and factor B by human umbilical cord vein endothelial  
690 cells. *Cancer Lett* (1997) **116**:21–26. doi: 10.1016/S0304-3835(97)04737-X
- 691 65. Brooimans RA, van der Ark AA, Buurman WA, van Es LA, Daha MR. Differential  
692 regulation of complement factor H and C3 production in human umbilical vein endothelial cells by  
693 IFN- $\gamma$  and IL-1. *J Immunol Baltim Md 1950* (1990) **144**:3835–3840.
- 694 66. Lappin DF, Guc D, Hill A, McShane T, Whaley K. Effect of interferon- $\gamma$  on complement  
695 gene expression in different cell types. *Biochem J* (1992) **281**:437–442. doi: 10.1042/bj2810437
- 696 67. May O, Merle NS, Grunenwald A, Gnemmi V, Leon J, Payet C, Robe-Rybkin T, Paule R,  
697 Delguste F, Satchell SC, et al. Heme Drives Susceptibility of Glomerular Endothelium to  
698 Complement Overactivation Due to Inefficient Upregulation of Heme Oxygenase-1. *Front Immunol*

699 (2018) **9**: doi: 10.3389/fimmu.2018.03008

700 68. Ades EW, Candal FJ, Swerlick RA, George VG, Summers Susan, Bosse DC, Lawley TJ.  
701 HMEC-1: Establishment of an Immortalized Human Microvascular Endothelial Cell Line. *J Invest*  
702 *Dermatol* (1992) **99**:683–690. doi: 10.1111/1523-1747.ep12613748

703 69. Satchell SC, Tasman CH, Singh A, Ni L, Geelen J, von Ruhland CJ, O’Hare MJ, Saleem MA,  
704 van den Heuvel LP, Mathieson PW. Conditionally immortalized human glomerular endothelial cells  
705 expressing fenestrations in response to VEGF. *Kidney Int* (2006) **69**:1633–1640. doi:  
706 10.1038/sj.ki.5000277

707 70. Edgell CJ, McDonald CC, Graham JB. Permanent cell line expressing human factor VIII-  
708 related antigen established by hybridization. *Proc Natl Acad Sci* (1983) **80**:3734–3737. doi:  
709 10.1073/pnas.80.12.3734

710 71. Gavriilaki E, Yuan X, Ye Z, Ambinder AJ, Shanbhag SP, Streiff MB, Kickler TS, Moliterno  
711 AR, Sperati CJ, Brodsky RA. Modified Ham test for atypical hemolytic uremic syndrome. *Blood*  
712 (2015) **125**:3637–3646. doi: 10.1182/blood-2015-02-629683

713 72. Réti M, Farkas P, Csuka D, Rázsó K, Schlammadinger Á, Udvardy ML, Madách K, Domján  
714 G, Bereczki C, Reusz GS, et al. Complement activation in thrombotic thrombocytopenic purpura. *J*  
715 *Thromb Haemost JTH* (2012) **10**:791–798. doi: 10.1111/j.1538-7836.2012.04674.x

716 73. Bettoni S, Galbusera M, Gastoldi S, Donadelli R, Tentori C, Spartà G, Bresin E, Mele C,  
717 Alberti M, Tortajada A, et al. Interaction between Multimeric von Willebrand Factor and  
718 Complement: A Fresh Look to the Pathophysiology of Microvascular Thrombosis. *J Immunol* (2017)  
719 **199**:1021–1040. doi: 10.4049/jimmunol.1601121

720 74. Mitra D, Jaffe EA, Weksler B, Hajjar KA, Soderland C, Laurence J. Thrombotic  
721 thrombocytopenic purpura and sporadic hemolytic-uremic syndrome plasmas induce apoptosis in  
722 restricted lineages of human microvascular endothelial cells. *Blood* (1997) **89**:1224–1234.

723 75. Jaffe EA, Nachman RL, Becker CG, Minick CR. Culture of Human Endothelial Cells  
724 Derived from Umbilical Veins. IDENTIFICATION BY MORPHOLOGIC AND IMMUNOLOGIC  
725 CRITERIA. *J Clin Invest* (1973) **52**:2745–2756. doi: 10.1172/JCI107470

726 76. McGinn S, Poronnik P, Gallery ED, Pollock CA. A method for the isolation of glomerular and  
727 tubulointerstitial endothelial cells and a comparison of characteristics with the human umbilical vein  
728 endothelial cell model. *Nephrology* (2004) **9**:229–237. doi: 10.1111/j.1440-1797.2004.00254.x

729 77. van Setten PA, van Hinsbergh VWM, van der Velden TJAN, van de Kar NCAJ, Vermeer M,  
730 Mahan JD, Assmann KJM, van den Heuvel LPWJ, Monnens LAH. Effects of TNF $\alpha$  on  
731 verocytotoxin cytotoxicity in purified human glomerular microvascular endothelial cells. *Kidney Int*  
732 (1997) **51**:1245–1256. doi: 10.1038/ki.1997.170

733 78. Martin-Ramirez J, Hofman M, van den Biggelaar M, Hebbel RP, Voorberg J. Establishment  
734 of outgrowth endothelial cells from peripheral blood. *Nat Protoc* (2012) **7**:1709–1715. doi:  
735 10.1038/nprot.2012.093

736 79. Roumenina LT, Jablonski M, Hue C, Blouin J, Dimitrov JD, Dragon-Durey M-A, Cayla M,

- 737 Fridman WH, Macher M-A, Ribes D, et al. Hyperfunctional C3 convertase leads to complement  
738 deposition on endothelial cells and contributes to atypical hemolytic uremic syndrome. *Blood* (2009)  
739 **114**:2837–2845. doi: 10.1182/blood-2009-01-197640
- 740 80. Roumenina LT, Frimat M, Miller EC, Provot F, Dragon-Durey M-A, Bordereau P, Bigot S,  
741 Hue C, Satchell SC, Mathieson PW, et al. A prevalent C3 mutation in aHUS patients causes a direct  
742 C3 convertase gain of function. *Blood* (2012) **119**:4182–4191. doi: 10.1182/blood-2011-10-383281
- 743 81. Marinozzi MC, Vergoz L, Rybkine T, Ngo S, Bettoni S, Pashov A, Cayla M, Tabarin F,  
744 Jablonski M, Hue C, et al. Complement Factor B Mutations in Atypical Hemolytic Uremic  
745 Syndrome—Disease-Relevant or Benign? *J Am Soc Nephrol* (2014) **25**:2053–2065. doi:  
746 10.1681/ASN.2013070796
- 747 82. Schramm EC, Roumenina LT, Rybkine T, Chauvet S, Vieira-Martins P, Hue C, Maga T,  
748 Valoti E, Wilson V, Jokiranta S, et al. Mapping interactions between complement C3 and regulators  
749 using mutations in atypical hemolytic uremic syndrome. *Blood* (2015) **125**:2359–2369. doi:  
750 10.1182/blood-2014-10-609073
- 751 83. Noris M, Galbusera M, Gastoldi S, Macor P, Banterla F, Bresin E, Tripodo C, Bettoni S,  
752 Donadelli R, Valoti E, et al. Dynamics of complement activation in aHUS and how to monitor  
753 eculizumab therapy. *Blood* (2014) **124**:1715–1726. doi: 10.1182/blood-2014-02-558296
- 754 84. Valoti E, Alberti M, Tortajada A, Garcia-Fernandez J, Gastoldi S, Besso L, Bresin E,  
755 Remuzzi G, Rodriguez de Cordoba S, Noris M. A Novel Atypical Hemolytic Uremic Syndrome–  
756 Associated Hybrid *CFHR1/CFH* Gene Encoding a Fusion Protein That Antagonizes Factor H–  
757 Dependent Complement Regulation. *J Am Soc Nephrol* (2015) **26**:209–219. doi:  
758 10.1681/ASN.2013121339
- 759 85. Rigothier C, Delmas Y, Roumenina LT, Contin-Bordes C, Lepreux S, Bridoux F, Goujon JM,  
760 Bachelet T, Touchard G, Frémeaux-Bacchi V, et al. Distal Angiopathy and Atypical Hemolytic  
761 Uremic Syndrome: Clinical and Functional Properties of an Anti-Factor H IgA $\lambda$  Antibody. *Am J*  
762 *Kidney Dis* (2015) **66**:331–336. doi: 10.1053/j.ajkd.2015.03.039
- 763 86. Galbusera M, Noris M, Gastoldi S, Bresin E, Mele C, Breno M, Cuccarolo P, Alberti M,  
764 Valoti E, Piras R, et al. An Ex Vivo Test of Complement Activation on Endothelium for  
765 Individualized Eculizumab Therapy in Hemolytic Uremic Syndrome. *Am J Kidney Dis* (2019) **74**:56–  
766 72. doi: 10.1053/j.ajkd.2018.11.012
- 767 87. Madden I, Roumenina LT, Langlois-Meurinne H, Guichoux J, Llanas B, Frémeaux-Bacchi V,  
768 Harambat J, Godron-Dubrasquet A. Hemolytic uremic syndrome associated with *Bordetella pertussis*  
769 infection in a 2-month-old infant carrying a pathogenic variant in complement factor H. *Pediatr*  
770 *Nephrol* (2019) **34**:533–537. doi: 10.1007/s00467-018-4174-1
- 771 88. Aiello S, Gastoldi S, Galbusera M, Ruggenti PL, Portalupi V, Rota S, Rubis N, Liguori L,  
772 Conti S, Tironi M, et al. C5a and C5aR1 are key drivers of microvascular platelet aggregation in  
773 clinical entities spanning from aHUS to COVID-19. *Blood Adv* (2021) **bloodadvances.2021005246**.  
774 doi: 10.1182/bloodadvances.2021005246
- 775 89. Piras R, Iatropoulos P, Bresin E, Todeschini M, Gastoldi S, Valoti E, Alberti M, Mele C,  
776 Galbusera M, Cuccarolo P, et al. Molecular Studies and an ex vivo Complement Assay on

## Measuring complement attack on endothelial cells

- 777 Endothelium Highlight the Genetic Complexity of Atypical Hemolytic Uremic Syndrome: The Case  
778 of a Pedigree With a Null CD46 Variant. *Front Med* (2020) **7**:579418. doi:  
779 10.3389/fmed.2020.579418
- 780 90. Timmermans SAMEG, Abdul-Hamid MA, Potjewijd J, Theunissen ROMFIH, Damoiseaux  
781 JGMC, Reutelingsperger CP, van Paassen P, on behalf of the Limburg Renal Registry. C5b9  
782 Formation on Endothelial Cells Reflects Complement Defects among Patients with Renal  
783 Thrombotic Microangiopathy and Severe Hypertension. *J Am Soc Nephrol* (2018) **29**:2234–2243.  
784 doi: 10.1681/ASN.2018020184
- 785 91. Blasco M, Martínez-Roca A, Rodríguez-Lobato LG, Garcia-Herrera A, Rosiñol L, Castro P,  
786 Fernández S, Quintana LF, Cibeira MT, Bladé J, et al. Complement as the enabler of carfilzomib-  
787 induced thrombotic microangiopathy. *Br J Haematol* (2020) doi: 10.1111/bjh.16796
- 788 92. Palomo M, Blasco M, Molina P, Lozano M, Praga M, Torramade-Moix S, Martinez-Sanchez  
789 J, Cid J, Escolar G, Carreras E, et al. Complement Activation and Thrombotic Microangiopathies.  
790 *Clin J Am Soc Nephrol* (2019) **14**:1719–1732. doi: 10.2215/CJN.05830519
- 791 93. Chauvet S, Roumenina LT, Bruneau S, Marinozzi MC, Rybkine T, Schramm EC, Java A,  
792 Atkinson JP, Aldigier JC, Bridoux F, et al. A Familial C3GN Secondary to Defective C3 Regulation  
793 by Complement Receptor 1 and Complement Factor H. *J Am Soc Nephrol* (2016) **27**:1665–1677. doi:  
794 10.1681/ASN.2015040348
- 795 94. Vasilev VV, Noe R, Dragon-Durey M-A, Chauvet S, Lazarov VJ, Deliyska BP, Fremeaux-  
796 Bacchi V, Dimitrov JD, Roumenina LT. Functional Characterization of Autoantibodies against  
797 Complement Component C3 in Patients with Lupus Nephritis. *J Biol Chem* (2015) **290**:25343–  
798 25355. doi: 10.1074/jbc.M115.647008
- 799 95. Radanova M, Mihaylova G, Ivanova D, Daugan M, Lazarov V, Roumenina L, Vasilev V.  
800 Clinical and functional consequences of anti-properdin autoantibodies in patients with lupus  
801 nephritis. *Clin Exp Immunol* (2020) doi: 10.1111/cei.13443
- 802 96. Chaturvedi S, Braunstein EM, Yuan X, Yu J, Alexander A, Chen H, Gavriilaki E, Alluri R,  
803 Streiff MB, Petri M, et al. Complement activity and complement regulatory gene mutations are  
804 associated with thrombosis in APS and CAPS. *Blood* (2020) **135**:239–251. doi:  
805 10.1182/blood.2019003863
- 806 97. Timmermans S, Damoiseaux J, Reutelingsperger C, van Paassen P. More About Complement  
807 in the Antiphospholipid Syndrome. *Blood* (2020) doi: 10.1182/blood.2020005171
- 808 98. Chen JY, Galwankar NS, Emch HN, Menon SS, Cortes C, Thurman JM, Merrill SA, Brodsky  
809 RA, Ferreira VP. Properdin Is a Key Player in Lysis of Red Blood Cells and Complement Activation  
810 on Endothelial Cells in Hemolytic Anemias Caused by Complement Dysregulation. *Front Immunol*  
811 (2020) **11**: doi: 10.3389/fimmu.2020.01460
- 812 99. Cid J, Fernández J, Palomo M, Blasco M, Bailó N, Diaz-Ricart M, Lozano M.  
813 Hyperhemolytic Transfusion Reaction in Non-Hemoglobinopathy Patients and Terminal  
814 Complement Pathway Activation: Case Series and Review of the Literature. *Transfus Med Rev*  
815 (2020) doi: 10.1016/j.tmr.2020.06.002



- 816 100. Merle NS, Paule R, Leon J, Daugan M, Robe-Rybkin T, Poillerat V, Torset C, Frémeaux-  
817 Bacchi V, Dimitrov JD, Roumenina LT. P-selectin drives complement attack on endothelium during  
818 intravascular hemolysis in TLR-4/heme-dependent manner. *Proc Natl Acad Sci* (2019) **116**:6280–  
819 6285. doi: 10.1073/pnas.1814797116
- 820 101. Roumenina LT, Loirat C, Dragon-Durey M-A, Halbwachs-Mecarelli L, Sautes-Fridman C,  
821 Frémeaux-Bacchi V. Alternative complement pathway assessment in patients with atypical HUS. *J*  
822 *Immunol Methods* (2011) **365**:8–26. doi: 10.1016/j.jim.2010.12.020
- 823 102. Mollnes TE, Garred P, Bergseth G. Effect of time, temperature and anticoagulants on in vitro  
824 complement activation: consequences for collection and preservation of samples to be examined for  
825 complement activation. *Clin Exp Immunol* (1988) **73**:484–488.
- 826 103. Merle NS, Grunenwald A, Rajaratnam H, Gnemmi V, Frimat M, Figueres M-L, Knockaert S,  
827 Bouzekri S, Charue D, Noe R, et al. Intravascular hemolysis activates complement via cell-free heme  
828 and heme-loaded microvesicles. *JCI Insight* (2018) **3**: doi: 10.1172/jci.insight.96910
- 829 104. del Conde I, Cruz MA, Zhang H, López JA, Afshar-Kharghan V. Platelet activation leads to  
830 activation and propagation of the complement system. *J Exp Med* (2005) **201**:871–879. doi:  
831 10.1084/jem.20041497
- 832 105. Morigi M, Galbusera M, Gastoldi S, Locatelli M, Buelli S, Pezzotta A, Pagani C, Noris M,  
833 Gobbi M, Stravalaci M, et al. Alternative Pathway Activation of Complement by Shiga Toxin  
834 Promotes Exuberant C3a Formation That Triggers Microvascular Thrombosis. *J Immunol* (2011)  
835 **187**:172–180. doi: 10.4049/jimmunol.1100491
- 836 106. Noone DG, Riedl M, Pluthero FG, Bowman ML, Liszewski MK, Lu L, Quan Y, Balgobin S,  
837 Schneppenheim R, Schneppenheim S, et al. Von Willebrand factor regulates complement on  
838 endothelial cells. *Kidney Int* (2016) **90**:123–134. doi: 10.1016/j.kint.2016.03.023
- 839 107. Youssef L, Miranda J, Blasco M, Paules C, Crovetto F, Palomo M, Torramade-Moix S,  
840 García-Calderó H, Tura-Ceide O, Dantas AP, et al. Complement and coagulation cascades activation  
841 is the main pathophysiological pathway in early-onset severe preeclampsia revealed by maternal  
842 proteomics. *Sci Rep* (2021) **11**:3048. doi: 10.1038/s41598-021-82733-z
- 843 108. Timmermans SAMEG, Wérion A, Damoiseaux JGMC, Morelle J, Reutelingsperger CP, van  
844 Paassen P. Diagnostic and Risk Factors for Complement Defects in Hypertensive Emergency and  
845 Thrombotic Microangiopathy. *Hypertension* (2020) **75**:422–430. doi:  
846 10.1161/HYPERTENSIONAHA.119.13714
- 847 109. Blasco M, Martínez-Roca A, Rodríguez-Lobato LG, Garcia-Herrera A, Rosiñol L, Castro P,  
848 Fernández S, Quintana LF, Cibeira MT, Bladé J, et al. Complement as the enabler of carfilzomib-  
849 induced thrombotic microangiopathy. *Br J Haematol* (2020) doi: 10.1111/bjh.16796
- 850 110. Volokhina EB, van de Kar NCAJ, Bergseth G, van der Velden TJAM, Westra D, Wetzels  
851 JFM, van den Heuvel LP, Mollnes TE. Sensitive, reliable and easy-performed laboratory monitoring  
852 of eculizumab therapy in atypical hemolytic uremic syndrome. *Clin Immunol Orlando Fla* (2015)  
853 **160**:237–243. doi: 10.1016/j.clim.2015.05.018
- 854 111. Merrill SA, Brittingham ZD, Yuan X, Moliterno AR, Sperati CJ, Brodsky RA. Eculizumab

## Measuring complement attack on endothelial cells

855 cessation in atypical hemolytic uremic syndrome. *Blood* (2017) **130**:368–372. doi: 10.1182/blood-  
856 2017-02-770214

857 112. Mastellos DC, Ricklin D, Lambris JD. Clinical promise of next-generation complement  
858 therapeutics. *Nat Rev Drug Discov* (2019) **18**:707–729. doi: 10.1038/s41573-019-0031-6

859



861 **Table 1: Production of distinct complement components and expression of regulators according**  
 862 **to endothelial cell type after stimulation**

		TNF	INF gamma	IL1 beta	Heme
<b>HRGEC</b>	C3	↑	↑	→*	
	C4	→*	↑	→*	
	C5	→*		→*	
	FB	↑		→*	
	FD	→*		→*	
	Properdin	↓*		→*	
	FH	→		→*	
	FI			→*	
	TM	↓		↑	
	CD46	↑		→	↓
	CD55	→		→	↓
	CD59	→		→	
	E-selectine	↑	→		
	C3aR	↑			
	C5aR	0			
<b>BMVEC</b>	C3	↑			
	C4	→*			
	C5	→*			
	FB	↑			
	FD	→*			
	Properdin	↓*			
	FH	→			
	CD46	↑			
	CD55	→			
	C3aR	↑			
	C5aR	0			
<b>HMEC</b>	E-selectine	↑	→		
	C3		→		
	C4		↑		
	CD46				↓
	CD55				→
<b>HUVEC</b>	C2		↑		
	C3	↑	→/↓	↑	
	C4			→*	
	C5			→*	
	FB	↑	↑	↑*	
	FD			→*	
	Properdin			→*	
	FH		↑	↓	
	FI			→*	
	TM	↓		→	
	CD46	↑		→	↓
	CD55	↑		↑	↓
	CD59	→		→	→
	E-selectin	↑	→		
	P-selectin				↑
	C1-inh		↑		

## Measuring complement attack on endothelial cells

863 The data presented here are mainly concerned with protein expression. \* denotes transcriptomic data. For details, please  
864 refer to **Table S2**. Abbreviations: BMVEC, brain microvascular endothelial cells; HMEC, human microvascular endothelial  
865 cells; HRGEC, human renal glomerular endothelial cells; HUVEC, human umbilical vein endothelial cells

866

867

868 **Table 2: Endothelial cells used for *ex vivo* experiments**

	Conditionally immortalized	Primary
Macrovascular		<b>HUVEC</b>
Microvascular	<b>CI-GEnC</b> <b>HMEC-1</b>	<b>BMVEC</b> <b>HRGEC</b> <b>BOEC</b>

869 Abbreviations: BMVEC, brain microvascular endothelial cells; BOEC, blood outgrowth endothelial cells; CI-GEnC,  
870 conditionally immortalized human glomerular endothelial cells; HMEC, human microvascular endothelial cells; HRGEC,  
871 human renal glomerular endothelial cells; HUVEC, human umbilical vein endothelial cells

872

## 873 **Figures Legends**

### 874 **Figure 1: Complement implication in human diseases**

875 Complement dysregulation has been implicated in the pathophysiology of several human diseases.  
876 Complementopathies in which alternative pathway dysregulation is the central mechanism of cell and  
877 tissue injury are represented in red. Conditions in which the complement system has been demonstrated  
878 to contribute significantly to tissue injury are represented in pink. Other diseases in which complement  
879 plays an accessory role are represented in gray.

### 880 **Figure 2: Concept of *ex vivo* complement deposition on endothelial cells**

881 An *ex vivo* endothelial assay was developed to reproduce human pathological conditions and their  
882 complexity. The assay consists of the detection and quantification of complement component  
883 deposition on the cultured endothelial cells (EC) surface after incubation with human serum. The EC  
884 surface was used as the regulatory surface. (A): In serum from healthy individuals, the alternative  
885 pathway is active at low levels but tightly regulated in the fluid phase by regulators, resulting in a very  
886 low level of complement activation product deposition on the EC surface. The detection of an increased  
887 complement deposition when incubation is performed with pathological serum (B) could result in  
888 either i) complement overactivation that overwhelms EC capacity of regulation (orange) or ii) defect  
889 in complement regulation in fluid or solid phase. Both are induced by tested human serum incubated  
890 with EC. Orange arrows represent some mechanisms involved in complement overactivation in serum:  
891 (1) the participation of a coactivation of classical/alternative pathway due to pathological  
892 immunoglobulins, immune complexes, or lectin pathway activation by polymeric IgA in IgA  
893 nephropathy, (2) an increase in the formation of fluid phase C3 convertases in the presence of heme or  
894 fluid phase activating surface, and the stabilization of C3 (3) or C5 (4) convertases by pathological  
895 immunoglobulins, such as C3 and C5 nephritic factors. Red crosses represent potential defects in  
896 alternative complement pathway regulation in the fluid phase (1) and on the cell surface (2, 3). These  
897 defects in complement regulation could be the consequence of inhibition of the main alternative  
898 pathway regulator FH due to anti-factor H antibodies (such as in aHUS), a lack of function, or a  
899 quantitative deficiency of FH and FI due to pathological genetic variants. \*CR1: weak expression of  
900 CR1 on endothelial cells Abbreviations: CR1: complement receptor 1 (CD35), FB: factor B, FD: factor  
901 D, FH: factor H, FI: factor I, FP: properdin, MCP: membrane cofactor protein.

### 902 **Figure 3: Comparative analysis of different protocols used for the *ex vivo* complement** 903 **activation test with endothelial cells**

904 1: The *ex vivo* test for measuring complement attack on endothelial cells can be performed on different  
905 endothelial cells, including human dermal microvascular endothelial cells (HMEC-1), human  
906 umbilical vein endothelial cells (HUVEC), blood outgrowth endothelial cells (BOEC), and glomerular  
907 endothelial cells (GEnC). 2: Cultured EC are then used at their resting state or after an activation by  
908 either ADP, heme, LPS, TNF/INF gamma, or apoptonecrotic cells. 3: EC are incubated with sample of  
909 interest. Either serum or activated plasma (consisting of patient citrated plasma mixed 1:1 with control  
910 serum pool) or normal human serum with addition of the protein of interest (e.g., IgG). Complement  
911 activation can be modulated in by addition of sCR1, anti-C5 antibody, anti-FH antibody, anti-properdin  
912 antibody, or EGTA-Mg buffer. 4: Complement activation products are then revealed by fluorescent  
913 tagged antibody. Antibody directed again C3c or C5b9 can be used. According to the context, staining  
914 for other molecules have been proposed and include IgG, P-selectin, vWF, and CD31. 5: Quantification

## Measuring complement attack on endothelial cells

915 is then performed using immunofluorescence scanning, flow cytometry, or ELISA. 6: Controls are  
916 required and vary according to the protocols.

### 917 **Figure 4: Current and future application fields of the *ex vivo* complement activation test on** 918 **endothelial cells**

919 There is a wide range of potential applications of *ex vivo* complement activation tests in endothelial  
920 cells. Currently used to decipher *in vitro* complement pathophysiology in research, a standardized test  
921 would represent a promising tool in clinical and therapeutic fields, paving the way for tailored medicine  
922 in complementopathies.

923