



HAL
open science

Validation of the Performance of A1HPV6, a Triage Blood Test for the Early Diagnosis and Prognosis of SARS-CoV-2 Infection

Pauline Maisonnasse, Thierry Poynard, Mehdi Sakka, Sepideh Akhavan, Romain Marlin, Valentina Peta, Olivier Deckmyn, Nesrine Braham Ghedira, Yen Ngo, Marika Rudler, et al.

► To cite this version:

Pauline Maisonnasse, Thierry Poynard, Mehdi Sakka, Sepideh Akhavan, Romain Marlin, et al.. Validation of the Performance of A1HPV6, a Triage Blood Test for the Early Diagnosis and Prognosis of SARS-CoV-2 Infection. *Gastro Hep Advances*, 2022, 1 (3), pp.393–402. 10.1016/j.gastha.2021.12.009 . hal-03793486

HAL Id: hal-03793486

<https://hal.sorbonne-universite.fr/hal-03793486>

Submitted on 19 Dec 2022

HAL is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers.

L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.



Distributed under a Creative Commons Attribution - NonCommercial - NoDerivatives 4.0 International License

ORIGINAL RESEARCH—CLINICAL

Validation of the Performance of A1HPV6, a Triage Blood Test for the Early Diagnosis and Prognosis of SARS-CoV-2 Infection



Pauline Maisonnasse,^{1,*} Thierry Poynard,^{2,*} Mehdi Sakka,³ Sepideh Akhavan,⁴ Romain Marlin,¹ Valentina Peta,^{2,5} Olivier Deckmyn,⁵ Nesrine Braham Ghedira,³ Yen Ngo,⁵ Marika Rudler,⁶ Sylvie van der Werf,^{7,8} Stephane Marot,⁴ Dominique Thabut,⁶ Harry Sokol,⁹ Chantal Housset,² Alain Combes,¹⁰ Roger Le Grand,^{1,*} and Patrice Cacoub,^{11,*} for the ProCop Study Group

¹Center for Immunology of Viral, Auto-immune, Hematological and Bacterial diseases (IMVA-HB/IDMIT), Université Paris-Saclay, Inserm, CEA, Fontenay-aux-Roses, France; ²Assistance Publique-Hôpitaux de Paris (AP-HP), INSERM, Centre de Recherche Saint-Antoine (CRSA), Institute of Cardiometabolism and Nutrition (ICAN), Sorbonne Université, Paris, France; ³Department of Metabolic Biochemistry, AP-HP Pitié-Salpêtrière, Sorbonne Université, Paris, France; ⁴Department of Virology, AP-HP Pitié-Salpêtrière, Sorbonne Université, Paris, France; ⁵BioPredictive, Research, Paris, France; ⁶Department of Hepatology, AP-HP Pitié-Salpêtrière, Sorbonne Université, Paris, France; ⁷Department of Virology, CNRS UMR 3569, Institut Pasteur, Paris, France; ⁸National Reference Center for Respiratory Viruses, Institut Pasteur, Paris, France; ⁹Gastroenterology Department, INSERM, Centre de Recherche Saint Antoine, Sorbonne Université, Paris, France; ¹⁰Department of Intensive Care Unit, AP-HP Pitié-Salpêtrière, Sorbonne Université, Paris, France; and ¹¹Department of Internal Medicine and Clinical Immunology, AP-HP Pitié-Salpêtrière, Sorbonne Université, Paris, France

BACKGROUND AND AIMS: Apolipoprotein A1 (A1) and haptoglobin (HP) serum levels are associated with the spread and severity of severe acute respiratory syndrome coronavirus 2 (SARS-CoV-2) infection. We have constructed and validated a multivariable risk calculator (A1HPV6) integrating A1, HP, alpha2-macroglobulin, and gamma glutamyl transferase to improve the performances of virological biomarkers. **METHODS:** In a prospective observational study of hospitalized patients with nonsevere SARS-CoV-2 infection, A1HPV6 was constructed in 127 patients and validated in 116. The specificity was assessed in 7482 controls representing the general population. The primary diagnostic endpoint was the area under the receiver operating characteristic curve in patients with positive SARS-CoV-2 PCR. The primary prognostic endpoint was the age-and-sex adjusted risk of A1HPV6 to predict patients with WHO-stage > 4 ($W > 4$) severity. We assessed the kinetics of the A1HPV6 components in a nonhuman primate model (NHP), from baseline to 7 days (D7) after SARS-CoV-2 infection. **RESULTS:** The area under the receiver operating characteristic curve for A1HPV6 was 0.99 (95% CI 0.97–0.99) in the validation subset, which was not significantly different from that in the construction subset, 0.99 (0.99–0.99; $P = .80$), like for sensitivity 92% (85–96) vs 94% (88–97; $P = .29$). A1HPV6 was associated with $W > 4$, with a significant odds ratio of 1.3 (1.1–1.5; 0.002). In NHP, A1 levels decreased ($P <$

.01) at D2 and normalized at D4; HP levels increased at D2 and peaked at D4. In patients, A1 concentration was very low at D2 vs controls ($P < .01$) and increased at D14 ($P < .01$) but was still lower than controls; HP increased at D2 and remained elevated at D14. **CONCLUSION:** These results validate the diagnostic and prognostic performances of A1HPV6. Similar kinetics of apolipoprotein A1, HP, and alpha-2-macroglobulin were observed in the NHP model. ClinicalTrials.gov number, NCT01927133.

Keywords: COVID-19; Nonhuman Primate Model; Apolipoprotein A1; Haptoglobin

Introduction

The pandemic of the respiratory disease (COVID-19) associated with the novel coronavirus (severe acute respiratory syndrome coronavirus 2 [SARS-CoV-2]) has highlighted the need for biomarkers that detect different risks, that is, the risk of infection before the exposure, named “predisposing biomarkers”, the prognostic factors during the exposure, named “acute phase biomarker”, and the risk of sequelae after exposure, named “sequelae bio-

*These authors contributed equally to this work.

Abbreviations used in this paper: A1, Apolipoprotein A1; A2M, alpha2-macroglobulin; ALT, alanine aminotransferase; Dpi, days after infection; GGT, gamma glutamyl transpeptidase; HDL-C, high-density lipoprotein cholesterol; HP, haptoglobin; ICU, intensive care unit; NHP, non-human primate model; SAA, serum amyloid A.

Most current article

Copyright © 2022 The Authors. Published by Elsevier Inc. on behalf of the AGA Institute. This is an open access article under the CC BY-NC-ND license (<http://creativecommons.org/licenses/by-nc-nd/4.0/>).

2772-5723

<https://doi.org/10.1016/j.gastha.2021.12.009>

marker".¹⁻³ Among simple available blood biomarkers, 2 proteins associated with cell repair, apolipoprotein-A1 (A1) and haptoglobin (HP), could be accurate components of such multianalyte risk markers.

Several prospective population-based studies have shown that a low level of A1 as well as an associated low level of high-density lipoprotein cholesterol were associated with a significant risk of hospitalization 10 years later, confirming their importance as a predisposing biomarker.⁴⁻⁷ Several prospective studies in COVID-19 hospitalized patients have shown that A1 alone or associated with HP also had significant diagnostic and short-term prognostic value for the acute phase of infection.⁸⁻¹² Biologically, A1 interacts with lipid rafts on cellular membranes that are enriched in immune cell receptors such as toll-like receptors on macrophages, T-cell receptors, and B-cell receptors, which may all modulate immune responses.⁹⁻¹³ HDL also has immunomodulatory, antithrombotic, and antioxidant effects that could provide important clarity why genetically determined levels of high-density lipoprotein cholesterol, but not low-density lipoprotein cholesterol, provide a protective effect against infectious diseases.^{4,12}

The increase in serum HP levels associated with SARS-CoV-2 infection was expected based on its well-known acute-phase protein profile.¹⁴⁻¹⁶ HP is the most abundant protein among those modulated in SARS-CoV-2 infection.¹⁷ Unlike A1, there is no association between serum HP levels before SARS-CoV-2 exposure or genetic polymorphisms.^{3,14} However, A1 and HP interact during the acute phase response to infection.¹⁴⁻¹⁶ A1 collaborates with HP to downregulate hemoglobin-redox activity.^{15,16} It also facilitates the uptake of hemoglobin by interacting with the HDL-scavenger receptor B type 1 (SR-B1) displayed on macrophages and hepatocytes.¹⁶

The interaction of the SARS-CoV-2 S1 spike protein with HDL cholesterol could also participate in the transient decrease of A1 by its main transporter during infection. The SARS-CoV-2 S protein binds to cholesterol, followed by an enhanced attachment via SR-B1, which facilitates the SARS-CoV-2 entry. SR-B1 and host cell entry are initiated through interactions with the angiotensin-converting enzyme 2.¹⁸

In a previous study, we described the temporal association between A1 and HP in patients with nonalcoholic fatty liver disease with the incidence of SARS-CoV-2-infected cases in the United States. The significant association between serum A1 levels and SARS-CoV-2 infection persisted after stratification of the main confounding factors.³ The value of combining these 2 proteins was based on the high sensitivity and specificity of the combination in the general population. The kinetics of a transient low A1 and high HP were not observed in 4 million FibroTest (FibroSure in USA) results since 2001, which combine these 2 proteins to assess the stage of liver fibrosis.³

The primary endpoint of this study was to assess the diagnostic value of A1HPV6, a multianalyte SARS-CoV-2 risk marker integrating A1 and HP; alpha-2-macroglobulin

(A2M), a marker of liver fibrosis; and the prognostic value of gamma glutamyl transpeptidase (GGT), a sensitive marker of liver injury.³ We report the prospective construction of A1HPV6 during the first wave of COVID-19, the "construction subset", and the internal validation in the "validation-subset" during the following 2 waves.

The second aim was to assess the prognostic value of A1HPV6 to predict WHO stage > 4 ($W > 4$) severity COVID-19 disease in patients hospitalized in an internal medicine department (Table 1), who did not require direct admission to the intensive care unit (ICU). To confirm the value of A1HPV6 components, we compared the kinetics observed in the patients during hospitalization to those observed in a nonhuman primate (NHP) mild-disease model.^{19,20} This is a unique model to assess the impact of SARS-CoV-2 on biomarkers in the first 14 days postinfection (Dpi), which is not feasible in humans.

Finally, to identify the risk of false positive or false negative A1HPV6 values and its components,^{21,22} we assessed the A1HPV6 components in 3 extremely severe ICU patients with persistence of SARS-CoV-2 infection and multiple repeated samples.

Patients and Methods

Patients

Five subsets were analyzed, a "construction-subset," a "validation-subset," a "prognostic-subset," a "controls-subset," and a "kinetic-subset" combining all the positive SARS-CoV-2 PCR cases admitted in the internal medicine department, plus 3 severe cases with persistent plasma infection followed in the ICU, and controls from the general population (Figure 1).

Ethics

The prospective observational study in COVID-19 patients was approved by CER-Sorbonne University IRB, CER-2020-14, with signed informed consent. All the previously published patient analyses from retrospective databases were non-interventional studies, without supplementary blood samples, and were exempt from IRB review (NCT01927133). This study was performed according to the principles of the Declaration of Helsinki. All authors had access to the study data and reviewed and approved the final manuscript.

Construction, Validation, and Prognostic Subsets

The construction-subset was performed between January and June 2020 and the validation-subset between October 2020 and May 2021. The methods and characteristics of patients included in the construction-subset have already been described elsewhere, as well as the univariate performance of A1 and HP.³ The inclusion criteria were the same in both subsets. Patients aged 18 years or older and hospitalized in the internal medicine department were eligible for the study if they had confirmed SARS-CoV-2 infection (positive on RT-PCR or typical chest CT scan). Very severe patients directly admitted in ICU were not eligible, as well as patients with serum assessments not contemporaneous, defined as more than 14 days before or after admission.

Table 1. Characteristics of COVID-19 Patients of the Construction and Validation Subsets to Assess Sensitivity of A1HPV6

Characteristics	Construction subset	Validation subset	P value	All
Number n (%)	127	116		243
Male sex	83 (65.4)	74 (63.4)	.90	157 (65.0)
Median age (IQR) year	71 (57–81)	67 (53–77)	.03	69 (56–78)
Age category			.22	
<50 y	16 (12.6)	21 (18.1)		37 (15.2)
50 to <70 y	45 (35.4)	47 (40.5)		92 (37.9)
≥70 y	66 (52.0)	48 (41.4)		114 (46.9)
Geographic origin			.03	
Caucasian	75 (59.1)	61 (52.6)		136 (56.0)
North African, Middle East	36 (28.4)	25 (21.6)		61 (25.1)
Other (Subsaharan, Asian)	16 (12.6)	30 (25.9)		46 (18.9)
Severity WHO stages				
<5	40 (31.5)	62 (53.5)	<.001 ^a	102 (42.0)
0–2 not hospitalized	2 (1.8)	0 (0)		2 (0.8)
3 hospitalized without oxygen	3 (2.4)	25 (21.6)		28 (11.5)
4 oxygen support mask	35 (27.6)	37 (31.9)		72 (29.6)
5–8	87 (68.5)	54 (46.6)		141 (58.0)
5 high flow or ventilation	64 (50.4)	29 (25.0)		93 (38.3)
6–7 Invasive oxygen support	8 (6.3)	3 (2.3)		11 (4.5)
8 death	15 (11.8)	22 (19.0)		37 (15.2)
Coexisting conditions				
Obesity (BMI ≥ 30)	27 (21.3)	35 (30.2)	.14	62 (25.5)
Hypertension	71 (55.9)	59 (50.9)	.44	130 (53.5)
Diabetes type 2	33 (26.0)	29 (25.0)	.88	62 (25.5)
Dyslipidaemia	44 (34.7)	37 (32.0)	.68	81 (33.3)
Stage liver fibrosis (FibroTest)			.15	
F0F1F2	117 (92.1)	100 (89.3)		217 (89.3)
F3F4 (cirrhosis)	10 (7.9)	16 (13.8)		26 (10.7)
Initial presentation				
Anosmia	15 (11.8)	13 (11.2)	1.00	28 (11.5)
Diarrhea	28 (22.1)	18 (15.5)	.25	46 (18.9)
Apolipoprotein ≤ 1.25g/L	115 (90.6)	105 (90.5)	1.00	220 (90.5)
Median laboratory (IQR)				
Apolipoprotein-A1 g/L	0.86 (0.71–1.03)	0.88 (0.67–1.07)	.96	0.87 (0.69–1.05)
Haptoglobin g/L	3.16 (2.22–4.08)	3.17 (1.91–4.11)	.78	3.17 (2.15–4.09)
Alpha-2 macroglobulin g/L	1.41 (1.18–1.97)	1.56 (1.20–2.00)	.46	1.50 (1.19–1.98)
GGT IU per liter	50 (30–117)	79 (40–134)	.02	63 (33–125)
ALT IU per liter	32 (23–63)	43.5 (24–83.3)	.02	36 (24–67)
Total bilirubin micromol/L	7 (5–11)	7 (5–10)	.96	7 (5–10)
Time clinic-PCR (d)	6 (2–11)	6 (0–11)	.82	6 (1–11)
Time clinic-serum sample (d)	9 (5–14)	8 (3–14)	.06	9 (4–14)
Time clinic-last news (d)	12 (8–18)	13 (7–17)	.90	12 (8–17)

ALT, alanine aminotransferase; BMI, body mass index; IQR, interquartile range.

^aP value severity WHO, stages <5 vs from 5 to 8.

The prognostic analysis was performed in all patients by combining the construction and validation subsets and called the “prognostic-subset” (Figure 1).

Kinetic-Subsets

The kinetics were analyzed in the “kinetics-subset,” which included the prospective integrated prognostic-subset and retrospectively severe cases admitted directly to the ICU (ICU patients).

In non-ICU patients, all cases with at least 3 repeated samples were analyzed. The aim was to describe the kinetics of A1HPV6 and its components, 8 and 14 days after hospital

admission, that is around 16 and 24 days postinfection (Dpi). Only kinetics in NHP were able to assess the kinetics during the first 14 Dpi.

The severe cases were identified as ICU patients with persistent SARS-CoV-2 viremia and at least 9 samples repeated frozen plasma samples. The severe cases allowed us to identify the risks of A1HPV6 false negatives and its components, such as organ failure and aggressive treatment.²²

Controls Subset to Assess the Specificity of A1HPV6

We previously collected 5 cohorts, called “specificity-cohorts,” which were used to retrospectively validate the




Figure 1. Flow chart of patients and controls.

specificity of A1 and HP in a large group of subjects without COVID-19 or pneumonia (Table A1). Specificity was assessed in the group of healthy volunteers that was representative of the French population before 2019.²³

Biochemical and Virological Methods

Measurements were all performed on fresh prospectively collected serum or plasma in the biochemistry unit of the AHPV-PSL hospital. A1, HP, A2M, GGT, alanine aminotransferase and bilirubin were assessed according to BioPredictive (Paris, France) analytical recommendations.^{3,24} Serum amyloid A (SAA) was performed by the ELISA method (Life Diagnostic, SAA-3). The virological methods used to diagnose SARS-CoV-2 in respiratory samples are described in File A2.

Kinetic Study in NHPs

Briefly, 4 female and 3 male cynomolgus macaques and 2 female rhesus macaques, aged 3–6 years, were originating from Mauritian and Chinese Accreditation of Laboratory Animal Care-certified breeding centers,^{9,25} respectively, and with the related NHP references described in File A2.

Statistical Methods

The criterion for main prognostic endpoint was the accurate prediction of severe stage WHO 5–8 (Table 1) by A1HPV6 assessed by the odds ratio using regression analysis adjusted for the subset group. A1HPV6 values were also compared between WHO severities using median tests values and the percentage of high-risk A1HPV6 by Fisher Exact test. The variability of repeated samples of A1HPV6 and its components was assessed in the kinetic subsets by repeated analysis of variance using the Time F-Ratio. We assumed that in humans, the median presymptomatic infectious period across studies varied from <1 to 4 days.²⁶ The different repeated means were compared to the first assessment at baseline as control by Dunnett's 2-sided multiple-comparison test, and the mean differences significance was compared by Tukey-Kramer's all pairs simultaneous confidence intervals.

Results

A1HPV6 Diagnostic Performances

Patients Included. A total of 292 patients with suspected of COVID-19 infection were preincluded, 136 in the construction subset between January and June 2020, and 156 in the validation subset between October 2020 and May 2021. After noninclusion of 49 patients with no positive PCR or for noncontemporaneous assessment of A1HPV6 components, a total of 243 patients were included, 127 in the construction subset and 116 in the validation subset (Figure 1 and Table 1). Forty out of 156 patients (25.6%) were not eligible, due to the absence of contemporaneous serum samples, vs 9 out of 136 patients (6.6%) preincluded in the construction subset ($P < .001$) (Table A2).

Several characteristics were significantly different in the validation subset compared with the construction subset, including lower median age (interquartile range; P value) 67 (53–77) years vs 71 (57–81; $P = .03$) years, a lower percentage of patients with European geographic origin (53% vs 59%; $P = .03$), and less need for oxygen support (78% vs 96%; $P < .001$).

Construction and Validation of A1HPV6 (Table 2). A1HPV6 was built including the 127 patients in the construction subset and the 7482 control patients in the general population, using logistic regression with and HP, GGT, and A2M, all adjusted for age and gender (patent pending).

The area under the receiver operating characteristic curve in the construction subset ($n = 127$) was 0.994 (0.982–0.998) with a sensitivity of 0.976 (0.933–0.995) and a specificity of 0.959 (0.955–0.964) using the cutoff of A1HPV6 ≥ 0.01 to define a high risk of infection.

The area under the receiver operating characteristic curve for the diagnosis of COVID-19 was 0.989 (0.971–0.991) in the validation subset ($n = 116$), with a sensitivity of 0.957 (0.902–0.986), which was nonsignificantly lower than that in the construction subset, with the

same specificity of 0.959 (0.955–0.964) because the same control subset was used.

Prognostic Performances

In the pooled population of patients with COVID-19, the prognostic value of A1HPV6 was significant on univariate analysis, with an odds ratio = 2.10 (1.27–3.50; $P = .004$) and a similar odds ratio of 2.07 (1.2–3.5; $P = .006$) in the construction subset after adjustment by regression analysis (Table 2). The prevalence of high-risk A1HPV6 was 99.3% in the 141 patients with severe WHO stages 5–8, which was greater than the 93.1% in the 102 patients with nonsevere WHO stages. A post-hoc analysis including geographical origins as the independent variable did not change the prognostic significance of A1HPV6, odds ratio = 2.14 (1.28–3.57; $P = .004$) (Table 2).

Kinetic Studies

Non-ICU Patients. A total of 222 sera were assessed in 59 patients (Figure 2). No significant difference was between the mean A1HPV6 values on D7 and D14 (mean difference -0.04 ; -0.13 to 0.06 ; $P = .64$) compared with D2 values. The risk was only low in one case on D2 and in 4 cases thereafter (panel A). The A1 mean decreased from 1.24 on D2 to 0.98 g/L on D14 for a difference of 0.26 g/L (0.19–0.33; $P < .001$) (panel B). The HP means decreased from 4.15 on D2 to 2.99 g/L on D14, for a difference of 1.16 g/L (0.85–1.47; $P < .001$) (panel C). The kinetics of bilirubin and GGT were described in panels D and E, respectively. The A2M decreased during hospitalization from 2.04 on D2 to 1.85 and 1.77 g/L on D7 and D14, respectively. On D14, the mean difference was 0.27 g/L (0.20–0.34; $P < .001$) (panel F).

ICU Patients. Three ICU patients with persistent SARS-CoV-2 plasma viremia for more than 2 months who later died had 9 aliquots available each (Table A2). The

repeated mean A1HPV6 values were not significantly different during follow-up compared with values at admission (D0) (Figure 3; panel A). The mean A1 on D0 and D80 were below 1 g/L D2, which were extremely low compared with normal values (>1.15 g/L), with a significant transient increase around D15 (panel B). HP values were severely decreased, with several values below detectable levels (0.08 g/L) during follow-up (panel C) and with a temporal association with cardiac arrests and extracorporeal membrane oxygenation (Table A2). Total bilirubin increased on D5 (panel D; $P = .004$). The mean GGT means was elevated from D5 to D20 (panel E, $P = .03$). A2M values were not significantly different (panel F). As expected in the presence of cardiac arrest, the 3 proteins were at the limit of detection on D2 in patient #2 (blue line).

Kinetic Study in NHPs (Figure A2)

There was no significant change in A1 in NHP from 7 to 26 Dpi (panel A), unlike in COVID-19 patients (black box) whose values were lower ($P < .001$) than those in control (blue box) COVID-19 patients before, during, and after recovery in all repeated samples (panel B). HP peaked at 7 Dpi in NHP (panel C), close to the peak of nasopharyngeal SARS-CoV-2 viral loads and was still elevated at 20 Dpi.¹⁹ Changes in HP were similar in patients, with a peak at D14 which then returned to normal values at D60 (panel D).

Kinetic Study in NHPs, During the First Week After Infection (Figure 4)

A1 mean values fluctuated significantly (repeated time F-ratio = 5.1; $P = .008$) with lowest values on 2 and 3 Dpi and returned to baseline at 4 and 6–26 Dpi (panel A). It is interesting that we were not able to see this early decrease in the initial study because previous assessments were performed at 7 Dpi. The mean HP increased from 1.00 g/L at

Table 2. A1HPV6 Performances

A1HPV6 outcome	Construction	Validation	Significant difference
Prevalence (%) of high risk	127/7609 (1.7)	116/7598 (1.5)	Not applicable
Diagnostic performance			
Area under ROC curve	0.994 (0.982–0.998)	0.989 (0.971–0.996)	<0.05
Sensitivity % (95% CI)	97.6 (93.3–95.9)	95.7 (90.2–98.6)	<0.05
Specificity	95.9 (95.5–96.4)	95.9 (95.5–96.4)	Not applicable same controls
Negative predictive value	99.9 (99.8–100)	99.9 (99.8–100)	<0.05
Positive predictive value	29.0 (24.7–33.5)	26.8 (22.6–31.3)	<0.05
Prognostic performance	Construction and validation pooled		Significance
WHO severe vs nonsevere	Stages 1–4 vs 5–8, n = 243		
Log A1HPV6 n/median/SD	87/–0.01/0.28 vs 40/–0.22/0.85		<0.001
Odds ratio univariate (95% CI)	2.10 (1.27–3.50)		0.004
Odds ratio subset (validation vs construction) adjusted	2.07 (1.24–3.48)		0.006
Odds ratio geographical origin adjusted	2.14 (1.28–3.57)		0.004

CI, confidence interval; ROC, receiver operating characteristics; SD, standard deviation; WHO, World Health Organization.




Figure 2. Kinetic in non-ICU patients. (A) A1HPV6. The dotted black line is the cutoff ($\text{Log} = -2.1$) defining the high risk of SARS-CoV-2 infection. (B) ApoA1. The dotted black line is the lower normal value (1.15 g/L). (C) Haptoglobin. The dotted black line is the upper normal value (3.60 g/L). (D) Total bilirubin. The dotted black line is the upper normal value (11.1 $\mu\text{mol/L}$). (E) GGT. The dotted black line is the upper normal value (62 IU/L). (F) A2M. The dotted black line is the upper normal value (2.76 g/L).

baseline to 0.60 g/L at 2 Dpi2 to, for a 0.40 g/L difference (0.14–0.66; $P = .002$) (panel B). The mean GGT decreased later ($P = .04$) (panel C). The mean A2M decreased regularly from 1.29 g/L at baseline to 1.04 g/L at 4 Dpi, with a difference of 0.25 g/L (0.16–0.33; $P < .001$) (panel D).

Mean CRP values did not fluctuate significantly during the week (panel E). SARS-CoV-2 viral loads in the trachea peaked at 2–3 Dpi 3, 8×10^5 copies/mL (1.9–5.4; $P < .001$) (panel F). Mean SAA values increased significantly very early from 195 to 1704 mg/L ($P = .04$) (Figure A3).

Discussion

We constructed and prospectively validated A1HPV6, a multivariate risk calculator for the diagnosis of SARS-CoV-2 infection in hospitalized patients who did not require ICU at admission. We confirmed the prognostic value of A1HPV6 for the severity of SARS-CoV-2 infection, defined as COVID-19 WHO stages >4 . To further our understanding of the 3 liver proteins included in A1HPV6, that is, A1, HP, and A2M, we evaluated their kinetics during hospitalization in non-ICU patients and in some ICU patients with persistent infection. Finally, we analyzed the kinetics of these components in a NHP model, which revealed a temporal

association with viral load during infection, including viral recovery.

Diagnostic Performances

Based on the results of our previous analyses of A1 and HP demonstrating their independent diagnostic value in patients with SARS-CoV-2 infection,³ we constructed A1HPV6 by adding A2M and GGT, 2 other independent risk factors, to these proteins. A2M is a well-validated liver fibrosis biomarker in patients with chronic liver disease.^{27,28} Recently a large cohort study showed that chronic liver disease plays a significant role in the burden of mechanical ventilation in severe COVID-19 patients.²⁹ GGT is a sensitive marker of liver injury³⁰ and an independent prognostic risk factor in patients hospitalized for COVID-19.^{3,30} The main limitations were the relatively small size of the 2 subsets, with a non-contemporaneous control population, and the absence of independent external validation. Another limitation was the higher percentage of noneligible patients in the validation subset due to the too late serum assessment of tests' components. However, post-hoc comparisons found only a significant increase in patients with type-2 diabetes among the noneligible patients (18 out of 40, 45%, vs 29 out of 116,




Figure 3. Kinetic in ICU patients with persistent plasmatic viremia. (A) A1HPV6. The dotted black line is the cutoff (Log = -2.1) defining the high risk of SARS-CoV-2 infection. (B) ApoA1. The dotted black line is the lower normal value (1.15 g/L). (C) Haptoglobin. The dotted black line is the upper normal value (3.60 g/L). (D) Total bilirubin. The dotted black line is the upper normal value (17.1 $\mu\text{mol/L}$). (E) GGT. The dotted black line is the upper normal value (62 IU/L). (F) A2M. The dotted black lines are the upper (2.76 g/L) and the lower (1.10 g/L) normal values.

25%; $P = .02$) compared with eligible patients. Most of the other characteristics were nonsignificant between the groups. The other significant differences were all expected in patients with late assessment as related to the recovery profile of the kinetics analyses: higher levels of A1, lower levels of HP, and a lower level of A1HPV6 score (Table A2).

However, the strengths include an assessment of the risk of false positive and negative in the A1HPV6 components since 2001 in a large number of patients including multi-racial ethnic populations.^{23,24,28} The kinetics of A1HPV6 and its components in ICU patients with persistent viremia (Figure 3) strongly suggests that the diagnostic value of HP, A1, and A2M should not be assessed in association with cardiac arrest or extracorporeal membrane oxygenation to avoid false positive or negative conclusions.

Prognostic Performance

We used the COVID-19 WHO severity staging system to compare the performance of A1HPV6 with that of other blood tests. The main limitations were the small sample size, the limited number of events, and the short follow-up

during hospitalization, which prevents a powerful multivariate analysis. There were 3 significant differences, in the validation subset vs the construction subset: median age 3 years younger, twice more patients from Asian and Subsaharan origin, and twice less high flow or ventilation. However, the A1HPV6 score already included age as a covariable, and at the inclusion, there were 3 significant differences, in the validation vs construction subset: median age 3 years younger, twice more patients from Asian and Subsaharan origin, and twice less high flow or ventilation. One regression analysis included the subset as an independent variable (validation vs construction 2.07 (1.24–3.48); $P = .006$), without a significant change in the prognostic performance of A1HPV6. A second regression analysis included geographical origin as an independent variable, which also did not change the prognostic significance of A1HPV6, odds ratio = 2.14 (1.28–3.57; $P = .004$) (Table 2).

Protein Kinetics

This is the first time that similar results have been identified for liver protein kinetics in patients and in an NHP




Figure 4. Early kinetics in Cynomolgus model. (A) ApoA1. (B) Haptoglobin. (C) GGT. (D) A2M. (E) CRP. (F) Tracheal SARS-CoV-2 viral loads.

model. Despite the limited number of cases, the numerous repeated sera analyses support the early kinetics of these liver proteins. In this study, early measurement of A1 during the first week of infection highlights the sensitivity of this protein to SARS-CoV-2 infection. Of note, in our initial NHP study, there was no significant variation detectable between 7 Dpi and 26 Dpi (Figure A2 panel A). However, an analysis of earlier samples shows a significant decrease in A1 on 2 Dpi associated with a peak in the virus load in the tracheal samples (Figure 4; panels A and F). On 6 Dpi, A1 returned to baseline while only traces of the virus were found in tracheal samples.

The second original result was the early sensitivity of serum HP (Figure 4, panel B), which was found to be more sensitive than CRP (Figure 4, panel E), the standard acute phase protein in human, or SAA (Figure A3). HP could also be better than CRP to evaluate inflammation recovery because of its sustained duration in SARS-CoV-2 infection. HP curves were similar in humans and NHP with abnormal values between 2 and 25 Dpi (Figure 2, panel C; Figure 3, panel C; Figure 4, panel B).

The third original and unexpected result was the highly significant decrease in A2M in both patients (Figure 2, panel F) and NHP (Figure 4, panel D). One

hypothesis in line with these results is that A2M protects the endothelium from a variety of potentially harmful intravascular proteases during SARS-CoV-2 infection.³¹ Our findings are also in line with the hypothesis that children are to some extent protected by higher A2M levels from severe COVID-19.³² These results and hypotheses encourage new studies of A2M in COVID-19 patients. The interpretation of A2M variability must take into account the age of patients.

Finally, A1HPV6 score added diagnostic and prognostic information before, during, and after SARS-CoV-2 infection. These data could improve the appropriateness of vaccines, antivirals, and anti-inflammatory prescriptions.

In conclusion, these results validate the diagnostic and prognostic performances of A1HPV6 in patients infected by SARS-CoV-2, not hospitalized in ICU. These results are reinforced by similar kinetics of A1, HP, and A2M observed in the macaque model.

Supplementary Materials

Material associated with this article can be found in the online version at <https://doi.org/10.1016/j.gastha.2021.12.009>.

References

- Whetton AD, Preston GW, Abubeker S, et al. Proteomics and informatics for understanding phases and identifying biomarkers in COVID-19 disease. *J Proteome Res* 2020;19:4219–4232.
- Demichev V, Tober-Lau P, Lemke O, et al. A time-resolved proteomic and prognostic map of COVID-19. *Cell Syst* 2021;12:780–794.e7.
- Poynard T, Deckmyn O, Rudler M, et al. Performance of serum apolipoprotein-A1 as a sentinel of Covid-19. *PLoS One* 2020;15:e0242306.
- Trinder M, Walley KR, Boyd JH, et al. Causal inference for genetically determined levels of high-density lipoprotein cholesterol and risk of infectious disease. *Arterioscler Thromb Vasc Biol* 2020;40:267–278.
- Madsen CM, Varbo A, Tybjærg-Hansen A, et al. U-shaped relationship of HDL and risk of infectious disease: two prospective population-based cohort studies. *Eur Heart J* 2018;39:1181–1190.
- Scalsky RJ, Chen YJ, Desai K, et al. Baseline cardiometabolic profiles and SARS-CoV-2 infection in the UK Biobank. *PLoS One* 2021;16:e0248602.
- Lassale C, Hamer M, Hernáez Á, et al. Association of pre-pandemic high-density lipoprotein cholesterol with risk of COVID-19 hospitalisation and death: the UK Biobank cohort study. *Prev Med Rep* 2021;23:101461.
- Sun JT, Chen Z, Nie P, et al. Lipid profile features and their associations with disease severity and mortality in patients with COVID-19. *Front Cardiovasc Med* 2020;7:584987.
- Begue F, Tanaka S, Mouktadi Z, et al. Altered high-density lipoprotein composition and functions during severe COVID-19. *Sci Rep* 2021;11:2291.
- Geyer PE, Arend FM, Doll S, et al. High-resolution serum proteome trajectories in COVID-19 reveal patient-specific seroconversion. *EMBO Mol Med* 2021;13:e14167.
- Zhu Z, Yang Y, Fan L, et al. Low serum level of apolipoprotein A1 may predict the severity of COVID-19: a retrospective study. *J Clin Lab Anal* 2021;35:e23911.
- Sorokin AV, Karathanasis SK, Yang ZH, et al. COVID-19-associated dyslipidemia: implications for mechanism of impaired resolution and novel therapeutic approaches. *FASEB J* 2020;34:9843–9853.
- Cochran BJ, Ong KL, Manandhar B, et al. Apolipoprotein-A1: a protein with multiple therapeutic functions. *Curr Atheroscler Rep* 2021;23:11.
- Delanghe JR, De Buyzere ML, Speeckaert MM. Genetic polymorphisms in the host and COVID-19 infection. *Adv Exp Med Biol* 2021;1318:109–118.
- Spagnuolo MS, Cigliano L, Abrescia P. The binding of haptoglobin to apolipoprotein A1: influence of hemoglobin and concanavalin A. *Biol Chem* 2003;384:1593–1596.
- Du R, Winarsih I, Ho B, et al. Lipid-free apolipoprotein A-I exerts an antioxidative role against cell-free hemoglobin. *Am J Clin Exp Immunol* 2012;1:33–48.
- Barberis E, Vanella VV, Falasca M, et al. Circulating exosomes are strongly involved in SARS-CoV-2 infection. *Front Mol Biosci* 2021;8:632290.
- Wei C, Wan L, Yan Q, et al. HDL-scavenger receptor B type 1 facilitates SARS-CoV-2 entry. *Nat Metab* 2020;2:1391–1400.
- Sokol H, Contreras V, Maisonnasse P, et al. SARS-CoV-2 infection in nonhuman primates alters the composition and functional activity of the gut microbiota. *Gut Microbes* 2021;13:1–19.
- Gonçalves A, Maisonnasse P, Donati F, et al. SARS-CoV-2 viral dynamics in non-human primates. *PLoS Comput Biol* 2021;17:e1008785.
- Tanaka S, De Tymowski C, Assadi M, et al. Lipoprotein concentrations over time in the intensive care unit COVID-19 patients: results from the ApoCOVID study. *PLoS One* 2020;15:e0239573.
- Tanaka S, De Tymowski C, Zappella N, et al. Lipoprotein concentration in patients requiring extracorporeal membrane oxygenation. *Sci Rep* 2021;11:17225.
- Poynard T, Lebray P, Ingiliz P, et al. Prevalence of liver fibrosis and risk factors in a general population using non-invasive biomarkers (FibroTest). *BMC Gastroenterol* 2010;10:40.
- Poynard T, Munteanu M, Deckmyn O, et al. Applicability and precautions of use of liver injury biomarker FibroTest. A reappraisal at 7 years of age. *BMC Gastroenterol* 2011;11:39.
- Maisonnasse P, Aldon Y, Marc A, et al. COVA1-18 neutralizing antibody protects against SARS-CoV-2 in three preclinical models. *Nat Commun* 2021;12:6097.
- Byrne AW, McEvoy D, Collins AB, et al. Inferred duration of infectious period of SARS-CoV-2: rapid scoping review and analysis of available evidence for asymptomatic and symptomatic COVID-19 cases. *BMJ Open* 2020;10:e039856.
- Naveau S, Poynard T, Benattar C, et al. Alpha-2-macroglobulin and hepatic fibrosis. Diagnostic interest. *Dig Dis Sci* 1994;39:2426–2432.
- Poynard T, Munteanu M, Deckmyn O, et al. Validation of liver fibrosis biomarker (FibroTest) for assessing liver fibrosis progression: proof of concept and first application in a large population. *J Hepatol* 2012;57:541–548.
- Mallet V, Beeker N, Bouam S, et al. Prognosis of French COVID-19 patients with chronic liver disease: a national retrospective cohort study for 2020. *J Hepatol* 2021;75:848–855.
- Adamidi ES, Mitsis K, Nikita KS. Artificial intelligence in clinical care amidst COVID-19 pandemic: a systematic review. *Comput Struct Biotechnol J* 2021;19:2833–2850.
- Becker CG, Harpel PC. alpha-2-Macroglobulin on human vascular endothelium. *J Exp Med* 1976;144:1–9.
- Seitz R, Gürtler L, Schramm W. Thromboinflammation in COVID-19: can α_2 -macroglobulin help to control the fire? *J Thromb Haemost* 2021;19:351–354.

Received November 15, 2021. Accepted December 29, 2021.

Correspondence:

Address correspondence to: Thierry Poynard, MD, PhD, Hepatology Groupe Hospitalier Pitié-Salpêtrière, 57 Bd Hôpital, Paris 75013, France. e-mail: thierry@poynard.com.

Authors' Contributions:

Pauline Maisonnasse: Data curation, equal; investigation, equal; writing – original draft, equal. Thierry Poynard: Conceptualization, lead; data curation, equal; formal analysis, equal; funding acquisition, equal; investigation, equal;

methodology, lead; project administration, lead; resources, equal; supervision, lead; validation, lead; visualization, lead; writing – original draft, lead; writing – review & editing, lead. Mehdi Sakka: Investigation, equal; resources: equal; validation: equal. Sepideh Akhavan: Data curation, equal; investigation, equal; resources, equal; validation, equal. Romain Marlin: Data curation, equal; investigation, equal. Valentina Peta: Data curation, equal; resources, equal; references review & editing, equal. Olivier Deckmyn: Conceptualization, equal; data curation, equal; methodology, equal; software, lead; validation, equal; visualization, equal; writing – original draft, equal; writing – review & editing, equal. Nesrine Braham Ghedira: Investigation, equal; resources, equal; validation, equal. Yen Ngo: Data curation, equal; investigation, equal; validation, equal. Marika Rudler: Data curation, equal; investigation, equal; resources, equal. Sylvie van der Werf: Data curation, equal; investigation, equal; validation, equal. Stephane Marot: Data curation, equal; investigation, equal; resources, equal; validation, equal. Dominique Thabut: Project administration, equal; resources, equal; supervision, equal; validation, equal. Harry Sokol: Formal analysis, equal; supervision, equal; validation, equal. Chantal Housset: Formal analysis, equal; supervision, equal; validation, equal; writing – original draft, equal. Alain Combes: Data curation, equal; investigation, equal; resources, equal. Roger Le Grand: Conceptualization, equal; formal analysis, equal; investigation, equal; supervision, equal; validation, lead; writing – original draft, equal. Patrice Cacoub: Conceptualization, lead; formal analysis, lead; funding acquisition, equal; investigation, equal; project administration, lead; resources,

equal; supervision, lead; validation, lead; visualization, lead; writing – original draft, equal; writing review & editing, equal.

Conflicts of Interest:

These authors disclose the following: Thierry Poynard is the inventor of FibroTest and founder of BioPredictive, the patents belong to the public organization Assistance Publique-Hôpitaux deParis. Olivier Deckmyn, Valentina Peta, and Yen Ngo are full-time employees of BioPredictive. The remaining authors disclose no conflicts.

Funding:

This study was supported by Grant EIT health ProCoP 20879 to Pr Patrice Cacoub APHP France.

Ethical Statement:

The corresponding author, on behalf of all authors, jointly and severally, certifies that their institution has approved the protocol for any investigation involving humans or animals and that all experimentation was conducted in conformity with ethical and humane principles of research.

Data Transparency Statement:

Data, analytic methods, and study materials of this nonintervantional study will be made available to other researchers, at thierry@poynard.com.