



HAL
open science

Effect of 10 UV filters on the brine shrimp *Artemia salina* and the marine microalgae *Tetraselmis* sp

Evane Thorel, Fanny Clergeaud, Lucie Jaugeon, Alice Rodrigues, Julie Lucas,
Didier Stien, Philippe Lebaron

► **To cite this version:**

Evane Thorel, Fanny Clergeaud, Lucie Jaugeon, Alice Rodrigues, Julie Lucas, et al.. Effect of 10 UV filters on the brine shrimp *Artemia salina* and the marine microalgae *Tetraselmis* sp. 2022. hal-03796512

HAL Id: hal-03796512

<https://hal.sorbonne-universite.fr/hal-03796512v1>

Preprint submitted on 4 Oct 2022

HAL is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers.

L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.

Effect of 10 UV filters on the brine shrimp *Artemia salina* and the marine microalgae *Tetraselmis* sp.

Evane Thorel, Fanny Clergeaud, Lucie Jaugeon, Alice M. S. Rodrigues, Julie Lucas, Didier Stien, Philippe Lebaron*

Sorbonne Université, CNRS USR3579, Laboratoire de Biodiversité et Biotechnologie Microbienne, LBBM, Observatoire Océanologique, 66650, Banyuls-sur-mer, France.

*corresponding author : E-mail address : lebaron@obs-banyuls.fr

Abstract

The presence of pharmaceutical and personal care products' (PPCPs) residues in the aquatic environment is an emerging issue due to their uncontrolled release, through grey water, and accumulation in the environment that may affect living organisms, ecosystems and public health. The aim of this study is to assess the toxicity of benzophenone-3 (BP-3), bis-ethylhexyloxyphenol methoxyphenyl triazine (BEMT), butyl methoxydibenzoylmethane (BM), methylene bis-benzotriazolyl tetramethylbutylphenol (MBBT), 2-Ethylhexyl salicylate (ES), diethylaminohydroxybenzoyl hexyl benzoate (DHHB), diethylhexyl butamido triazone (DBT), ethylhexyl triazone (ET), homosalate (HS), and octocrylene (OC) to marine organisms from two major trophic levels including autotrophs (*Tetraselmis sp.*) and heterotrophs (*Artemia salina*). In general, EC₅₀ results show that both HS and OC are the most toxic for our tested species, followed by a significant effect of BM on *Artemia salina* but only at high concentrations (1 mg/L) and then an effect of ES, BP3 and DHHB on the metabolic activity of the microalgae at 100 µg/L. BEMT, DBT, ET, MBBT had no effect on the tested organisms, even at high concentrations (2mg/L). OC toxicity represent a risk for those species since it is observed at concentrations only 15 to 90 times

higher than the highest concentrations reported in the natural environment and HS toxicity is for the first time reported on microalgae and was very important on *Tetraselmis sp.* at concentrations close to the natural environment concentrations.

Keywords

UV-filters

Toxicity tests

Marine microalgae

Artemia salina

Marine environment

1 **Introduction**

2

3 In recent decades, the sunscreen production and skin application have continuously
4 increased to protect against sunlight exposure damages and to prevent skin cancer risk (Azoury and
5 Lange, 2014; Waldman and Grant-Kels, 2019). Sunscreen products contain many chemical
6 ingredients including UltraViolet (UV) filters, the aim of which is to absorb or reflect UVA and/or
7 UVB radiations ranging from 280 to 400 nm (Sánchez-Quiles and Tovar-Sánchez, 2015).

8 More than 50 different UV filters are currently on the market (Shaath, 2010). Despite the
9 fact that the use of these compounds is subject to different regulations around the world, UV filters
10 are regularly detected in various aquatic environmental compartments including lakes, rivers,
11 surface marine waters and sediments. These chemicals can enter in the marine environment in two
12 ways, either indirectly from wastewater treatment plants effluent or directly from swimming or
13 recreational activities (Giokas et al., 2007). Furthermore, their lipophilic nature results in low
14 solubility in water, high stability and tendency to bioaccumulate (Gago-Ferrero et al., 2015; Vidal-
15 Liñán et al., 2018).

16 In recent years, many ecotoxicological studies have focused on the impact of organic UV
17 filters on various trophic levels, from microalgae to fish, passing by corals. Several studies have
18 demonstrated that some of these compounds can disrupt survival (Chen et al., 2018; He et al., 2019;
19 Paredes et al., 2014), behavior (Araújo et al., 2018; Barone et al., 2019), growth (Mao et al., 2017;
20 Paredes et al., 2014; Sieratowicz et al., 2011), development (Giraldo et al., 2017; Torres et al.,
21 2016), metabolism (Esperanza et al., 2019; Seoane et al., 2017; Stien et al., 2019), gene expression
22 (Gao et al., 2013; Zucchi et al., 2011) and reproduction (Araújo et al., 2018; Coronado et al., 2008;
23 Kaiser et al., 2012) in various species. It should be noted that the majority of toxicological studies
24 on organic UV filters were conducted on BP3, EMC and 4-MBC.

25 The adoption and implementation of the European legislation on the registration, evaluation,
26 authorization and restriction of chemicals (REACH) required several additional ecotoxicity data

27 promoting the use of invertebrates as models for toxicity assays (European Commission, 2007).
28 Brine shrimps *Artemia* spp. (here *A. salina*) are readily available worldwide and easy to breed. They
29 have therefore been frequently used as test organism in ecotoxicity assays, and *A. salina* was chosen
30 to investigate the toxicity of UV filters in this study (Caldwell et al., 2003; Libralato, 2014; Nunes
31 et al., 2006).

32 The green algae *Dunaliella tertiolecta* is commonly used for chronic algal toxicity testing.
33 The Haptophyta *Isochrysis galbana* is interesting too as it is widely cultured as food for the bivalve
34 industry. Green algae such as *Chlorella* and *Tetraselmis* sp., belonging to the phylum Chlorophyta,
35 have also been frequently exploited in toxicity assays. *Tetraselmis* have been used before to study
36 the toxic effect of several antibiotics and PCPs, but also the UV filter BP-3 (Seoane et al., 2017,
37 2014). This is why this algae was used in this study.

38 Previous studies about the toxic effects of different pollutants on microalgae physiology
39 demonstrate that flow cytometry (FCM) can be an alternative to standard algal population-based
40 endpoints, since it allows a rapid, quantitative and simultaneous measurement of multiple responses
41 to stress in individual cells (Esperanza et al., 2019; Hadjoudja et al., 2009; Prado et al., 2015;
42 Seoane et al., 2017). Seoane et al., (2017) showed that the most sensitive parameters are the
43 metabolic activity and cytoplasmic membrane potential. Therefore, we decided to use FCM to
44 analyze the toxicity of UV filters on *Tetraselmis* cells.

45

46 **2. Materials and methods**

47

48 *2.1 Test substances and experimental solutions*

49 The UV filters benzophenone-3 (BP-3), bis-ethylhexyloxyphenol methoxyphenyl triazine
50 (BEMT), butyl methoxydibenzoylmethane (BM), and methylene bis-benzotriazolyl
51 tetramethylbutylphenol (MBBT) were purchased from Sigma-Aldrich (Saint-Quentin Fallavier,
52 France). 2-Ethylhexyl salicylate (ES), diethylaminohydroxybenzoyl hexyl benzoate (DHBB),

53 diethylhexyl butamido triazone (DBT), ethylhexyl triazone (ET), homosalate (HS), and octocrylene
54 (OC) were provided by Pierre Fabre Laboratories.

55 Before each toxicity test, and due to the low water solubility of the compounds, stock
56 solutions at 1 mg/ml were prepared by dissolving each UV filters in dimethyl sulfoxide (DMSO,
57 Sigma-Aldrich, purity >99%). These solutions were diluted in order to add the same amount of
58 DMSO to all samples and to obtain exposure concentrations ranging from 20 ng/L to 2 mg/L for *A.*
59 *salina*, and 10 µg/L to 1 mg/L for *Tetraselmis* sp.. The lower concentrations tested were roughly
60 those reported in natural ecosystems. DMSO concentration in the experiments was always 2.5 %
61 (v/v). A DMSO control (2.5 % v/v) and a blank control were also included. The blank control was
62 artificial seawater for *A. salina* and growth medium for *Tetraselmis* sp..

63

64 2.2 *Artemia salina* mortality test

65

66 *A. salina* cysts were purchased from AquarHéak Aquaculture (Ars-en-Ré, France) and
67 stored at 4 °C. Dried cysts were hatched in a constantly aerated transparent «V» hatching incubator
68 filled with 500 mL of artificial seawater (ASW) at a salinity of 37 g/L, prepared with Instant Ocean
69 salt (Aquarium Systems, Sarrebourg, France). Incubation was carried out for 48 h, at 25 °C under
70 continuous light the first 24 hours. A 12:12 h light regime was then applied until the nauplii reached
71 the instar II-III stage. Ten nauplii were transferred into 5 ml glass tubes filled with ASW (2 mL)
72 contingently supplemented with DMSO and UV filters. The tubes were incubated at 25 °C under a
73 12:12 h light regime. The experiments were performed in sextuplicate. During the exposure period,
74 there was no aeration and the nauplii were not fed. The mortality rate was estimated after 48 h by
75 counting the dead nauplii under binocular. Organisms with no swimming activity or movement of
76 appendices for 10 s even after mechanical stimulation with a Pasteur pipette were counted as dead.
77 The tests were considered valid if the control's average mortality rate was < 20%.

78

79 2.3 *Tetraselmis* sp. toxicity test

80

81 2.3.1 *Experimental procedure*

82 *Tetraselmis* sp. (RCC500) was purchased from the Roscoff culture collection and was
83 grown in filtered (pore size: 0.22 μm) and autoclaved seawater enriched with a 50-fold diluted f/2
84 medium (Sigma–Aldrich). The culture was maintained under controlled conditions at 18 °C (\pm 1
85 °C) with a photon flux of 70 $\mu\text{mol photons.m}^{-2}.\text{s}^{-1}$ under a dark:light cycle of 12:12 h. Toxicity tests
86 were conducted in 150 mL Erlenmeyer flasks containing 50 mL of culture. Algae cells in
87 exponential growth phase were used as inoculum and the initial cell density was 5.10^4 cells/mL.
88 Three replicates per UV filter concentration were performed. After 7 days of exposure, different
89 morphological and physiological cells properties were monitored *via* flow cytometry (FCM).
90 Analyzed parameters were granularity, relative cell volume, chlorophyll a fluorescence, esterase
91 activity, and growth. The control experiment was a *Tetraselmis* sp. culture supplemented with
92 DMSO (2.5%).

93

94 2.3.2 *Flow cytometry (FCM) analyses*

95 Aliquots were collected after 7 days of exposition to be analyzed in a FACSCanto II flow
96 cytometer (Becton Dickinson, Franklin Lakes, New Jersey, USA) equipped with an air-cooled
97 argon laser (488 nm, 15 mW). To characterize the microalgae population, and to exclude non-algal
98 particles, the forward scatter (FSC, an estimation of cell size) and side scatter (SSC, an estimation
99 of granularity) dot-plots were used before each measurement. The flow rate of the cytometer was
100 set to low (acquisition time: 1 min).

101 The data recorded by FCM were measured either directly (autofluorescence, granularity,
102 size) or indirectly by the use of fluorochromes (esterase activity). Cellular density was determined
103 using Becton Dickinson Trucount™ 10 μm beads for calibration, as described by Pecqueur et al.
104 (2011). Growth rate (μ), expressed as day^{-1} were calculated using the following equation: $\mu =$

105 $(\ln(N_t) - \ln(N_0)) / (t - t_0)$, where N_t is the cell density at time t and N_0 is initial cell density. Chlorophyll
106 a natural autofluorescence was measured and detected in the FL3 channel (nm). Relative cell
107 volume (or size) and granularity were directly estimated with the forward light scatter (FSC
108 channel) and with the side scatter channel (SSC), respectively. To determine the metabolic activity
109 based on the esterase activity study, cells were stained with the fluorochrome Chemchrom V6 (10-
110 fold diluted in ChemSol B26 buffer– Biomérieux, France) at 1 % final concentration, and
111 incubated for 15 min at room temperature in the dark before analysis. Reading was performed with
112 the FL1 channel (nm). All cytometry data were analyzed using BD FACSDiva (Becton Dickinson).
113 Results were expressed as percentage of variation relative to control (100 %).

114

115 *2.4. Statistical analysis*

116

117 Results are reported as mean and standard deviation (SD), calculated from the 3 or 6
118 replicates. For both tests and all the parameters measured, differences between controls and nominal
119 concentrations of UV filter were analyzed using R software, by one-way analysis of variance
120 (ANOVA) followed by post-hoc Tukey HSD tests for pairwise comparisons. In all cases,
121 significance was accepted when $p < 0.05$. Dose–response curves, LD/LC₅₀-values were estimated
122 by a log(agonist) vs. response - Variable slope (four parameters) regression model in GraphPad
123 Prism 5.

124

125 **3. Results and discussion**


126

127 *3.1 Effects on Artemia salina mortality*

128

129 The toxicity of several organic UV filters on the marine crustacean *Artemia salina* (Nauplii
130 Instar II III) was determined after 48 h of exposure by counting dead larvae (Fig. 1). At the highest

131 concentration tested (2 mg/L), HS, BM, and OC demonstrated a significant effect on Nauplii
132 survival ($p < 0.05$) with mortality values reaching $54 \pm 16\%$, $64 \pm 19\%$ and $88 \pm 16\%$, respectively.
133 At lower concentrations of these filters no significant effect was detected. For BP3, BEMT, MBBT,
134 ES, DHHB, DBT and ET no toxicity was observed, even at the highest concentration.
135



136

137

138 **Fig 1** Mortality rate of *A. salina* exposed to the 10 UV filters at 6 concentrations. Boxes delineate
139 the minimal and maximal values, and the vertical line is the median of six replicates. Significance
140 levels relative to control determined by ANOVA followed by the Tukey's multiple comparison test:
141 *** $p < 0.001$. Results were not significant unless otherwise stated. For BM, HS and OC, the LD₅₀
142 is reported on the figure.

143


144 Our results indicate that among the different UV filters tested in this study, OC was the most toxic
145 molecule showing the lowest LD₅₀ value (0.6 mg/L), followed by BM and HS (1.8 mg/L and 2.4
146 mg/L, respectively). Environmental HS and BM concentrations reported so far are at least 500
147 times lower than LD₅₀, with values always lower than 3 µg/L (Fagervold et al., 2019; Sánchez-
148 Quiles and Tovar-Sánchez, 2015; Tsui et al., 2014; Ramos et al., 2015). OC concentrations in
149 coastal waters are higher and can reach 9 µg/L (Langford and Thomas, 2008; Tsui et al., 2014;
150 Ramos et al, 2015). This is the first report showing OC toxicity on *Artemia salina*. We also
151 observed a concentration-dependent increase in mortality of *Artemia* with respect to the control.
152 This is congruent with the toxicity observed, at lower concentration, on coral (50 µg/L, (Stien et al.,
153 2019), urchin, mussel and algae (40-80 µg/L, (Giraldo et al., 2017). OC also affects the
154 developmental process in zebrafish (Blüthgen et al., 2014). Here the LD₅₀ on *A. salina* is
155 approximately 90 times higher than the highest OC concentrations in marine waters reported so far.
156 It should be mentioned as well that OC concentrations in the 50-100 µg/kg range have been
157 frequently reported in sediments (Gago-Ferrero et al., 2011; Kameda et al., 2011), in which case
158 OC may affect benthic crustaceans.

159

160

161 *3.2 Effects on Tetraselmis sp.*

162



163

164

165

166

167

168

169

170

171

172

173

174

175

176

177

178

179

Fig 2 Relative A) growth rate, B) granularity, C) cell volume, D) fluorescence and E) metabolic activity of exposed *Tetraselmis* compared to control, set to 100 %. The boxes delineate the minimal and maximal values. The vertical line in the boxes is at mean. Significance levels relative to negative control determined by ANOVA followed by the Tukey's multiple comparison test: *** $p < 0.001$, ** $p < 0.01$, * $p < 0.05$. Results were not significant unless otherwise stated. N/A: not applicable, the data could not be obtained due to extensive cell death.

3.2.1 Growth rate and EC_{50} values

After 7 days of exposure, HS, BP3, and ES induced a significant decrease of algae growth (Fig. 2A). The growth rate of algae exposed to ES at 1 mg/L decreased by 24 % compared to control ($p < 0.05$). For BP-3 we observed a concentration dependent decrease in growth, which was statistically significant at 100 µg/L ($p < 0.01$) and 1 mg/L ($p < 0.001$). At 1 mg/L, the growth rate was negative, which translates a decreased cell concentration compared to t_0 . The 7-days LC_{50} value for BP3 was 143 µg/L. BP3 LC_{50} values on several microalgae species have been reported previously to be

180 roughly in the 100 µg/L to 20 mg/L range (Esperanza et al., 2019; Mao et al., 2017; Pablos et al.,
181 2015; Zhong et al., 2019). Seawater BP3 concentrations in the µg/L range have been frequently
182 reported in the literature (Gago-Ferrero et al., 2015; Tsui et al., 2014). Extremely high values of 1.4
183 and 0.6 mg/L have been recorded in the U.S. Virgin Islands (Downs et al., 2016). Finally, the most
184 important decline was with UV filter HS. No algal cells were detected in the presence of HS at 100
185 µg/L and 1 mg/L. LC₅₀ with HS was estimated at 74 µg/L, while it has been shown that HS
186 concentration in aquatic environments can reach ~3 µg/L (Tsui et al. 2014; Ramos et al., 2015). OC
187 induced a slight but significant increase of the growth rate at 1mg/L with a decrease in the
188 metabolic activity as determined by esterase activity. This increased metabolic activity was not
189 explained. Similar differences in the response of different physiological parameters were already
190 reported by Esperanza et al (2019) for the toxicity of BP3 in the microalgae *Chlamydomonas*
191 *reinhardtii*. The growth rate of *Tetraselmis* was not affected with BEMT, MBBT, DHHB, DBT, ET
192 and BM, even at 1 mg/L.

193

194 3.2.2 Impact on cell morphology

195

196 Three of the UV filters induced cell morphological alterations (Fig. 2B/C). Cells cultured in
197 the presence of BM have experienced a significant increase in cell volume and granularity at 1
198 mg/L ($p < 0.05$). This concentration is 1,000 to 10,000 times higher than the few concentrations
199 reported in the field (Fagervold et al., 2019; Sánchez-Quiles and Tovar-Sánchez, 2015; Tsui et al.,
200 2014). According to environmental concentrations of other UV filters, one can assume that the
201 effective concentration of 1 mg/L is probably higher than any BM environmental concentration, but
202 this remains to be confirmed. BP3 caused a dose dependent increase of relative cell volume at 100
203 µg/L and above, reaching up to 129 % of control cell volume at 1 mg/L ($p < 0.001$). Meanwhile,
204 this UV filter induced a significant 38 % granularity decrease at 1 mg/L ($p < 0.001$). With reference
205 to the environmental concentrations of BP3 (see above), this filter should exert a significant impact

206 on phytoplankton communities. These results are congruent with what was recently reported by
207 Zhong et al. (2019) on *Arthrospira* sp. With HS, cell volume and granularity could not be measured
208 at 100 and 1000 $\mu\text{g/L}$. However, a significant cell volume decrease was observed at 10 $\mu\text{g/L}$ of HS
209 (-11% , $p < 0.001$). In our experiment, HS No Observed Effect Concentration (NOEC) was
210 therefore lower than 10 $\mu\text{g/L}$, *i.e.*, within the same order of magnitude than the highest water
211 column concentration reported so far by Tsui et al. (2014). Again, it is expected that HS should
212 affect microalgae communities in bathing areas. No significant effect was observed for BEMT,
213 MBBT, ES, DHHB, DBT, ET, HS and OC.

214

215 3.2.3 Impact on autofluorescence

216

217 The results of FCM analysis revealed that several UV filters significantly reduced
218 chlorophyll a (Chl a) cell fluorescence (Fig 4D). The decrease was significant with BM (-13% , $p <$
219 0.01) and ES (-15% , $p < 0.05$) at 1 mg/L. A strong dose-dependent autofluorescence inhibition was
220 observed upon exposure to BP3 at concentrations of 100 $\mu\text{g/L}$ ($p < 0.001$) and above. Inhibition
221 reached 78 % at the highest dose. Again, autofluorescence could not be measured in cells treated
222 with HS at 100 and 1000 $\mu\text{g/L}$ due to the important cell degradation at these concentrations. No
223 significant effect was observed for BEMT, MBBT, DHHB, DBT, ET and OC.

224

225 3.2.4 Impact on cell metabolic activity

226

227 Metabolic activity was determined by estimating the relative esterase activity in exposed
228 cells compared to control ones. It was measured by CV6 staining and highlighted significant
229 decreased metabolic activities with half of the tested UV filters (Fig 2E). Algae exposed to ES and
230 HS experienced a decreased esterase activity at 10 $\mu\text{g/L}$ UV filter. The effect of BP3, DHHB and
231 OC was significant at 100 $\mu\text{g/L}$ and above. A similar decreased esterase activity was reported for *C.*

232 *reinhardtii* exposed to BP3, although at concentrations in the mg/L range (Esperanza et al., 2019;
233 Seoane et al., 2017). For DHHB, the effect was only observed for the microalgae and for the
234 esterase activity but not for other parameters. Therefore, the environmental risk cannot be estimated
235 since natural concentrations have never been reported for this filter. No significant effect was
236 observed for BEMT, MBBT, DBT, ET and BM.

237

238 **4. Conclusion**

239

240 The present work demonstrates that several filters exert toxicity on *A. salina* and *Tetraselmis*
241 sp. HS was the most toxic UV filter for the microalgae. LC₅₀ was 74 µg/L and significant adverse
242 effects were recorded at the lowest concentration tested (10 µg/L). HS was also toxic for *A. salina*,
243 although at much higher concentrations (LD₅₀ 2.4 mg/L). Overall, HS NOEC was lower than 10
244 µg/L. Its Lowest Observed Effect Concentration (LOEC) was 10 µg/L or below. HS concentrations
245 up to 3 µg/L have been reported in the natural environment in which case HS may represent a
246 potential risk for marine phytoplankton communities. Further research is needed to investigate on
247 HS toxicity with a larger diversity of phytoplankton species.

248 OC was toxic on both models with a dose-dependent effect on the microalgae. OC
249 significantly altered *Tetraselmis* sp. metabolic activity at 100 µg/L. On *A. salina*, LD₅₀ was 610
250 µg/L. Overall, OC LOEC was 100 µg/L with these models. The toxicity of OC occurred at
251 concentrations only 90 (*A. salina*) and 15 (*Tetraselmis* sp.) times higher than the highest
252 environmental concentrations reported so far. These results confirm the toxicity of OC on marine
253 organisms. BM was toxic towards the brine shrimp at high concentrations with a LD₅₀ of 1.84 mg/L
254 and had little effect on the microalgae. BM LOEC was 1 mg/L. The toxic concentrations reported
255 here may be high enough. BM might not have any effect on marine ecosystems, although the
256 occurrence of this filter should be monitored in a large range of ecosystems to better estimate its
257 natural concentrations. ES (LOEC 10 µg/L), BP3 and DHHB (LOEC 100 µg/L) had a significant

258 impact on the microalgae metabolic activity and had little effect on *A. salina*.

259 Overall, this research supports the need of establishing environmental quality standards for
260 UV-filters based on toxicity testing with key marine organisms, as well as identifying and reducing
261 environment input sources for the toxic ones. There are still many filters for which environmental
262 concentrations are missing to better estimate the potential environmental risk of their occurrence in
263 coastal ecosystems. It is probably important also to design new user's and environment friendly UV
264 filters.

265

266 **Acknowledgements**

267 We thank the BIO2MAR and the BIOPIC platforms from the Observatoire Océanologique
268 de Banyuls for providing technical support and access to instrumentation.

269

270 **Funding sources**

271 This work was carried out with the financial support of the Pierre Fabre company [grant
272 number 2018-02108]

273

274 **References**

275

276 Araújo, M.J., Rocha, R.J.M., Soares, A.M.V.M., Benedé, J.L., Chisvert, A., Monteiro, M.S., 2018.

277 Effects of UV filter 4-methylbenzylidene camphor during early development of *Solea*

278 *senegalensis* Kaup, 1858. Sci. Total Environ. 628–629, 1395–1404.

279 <https://doi.org/10.1016/j.scitotenv.2018.02.112>

280 Azoury, S.C., Lange, J.R., 2014. Epidemiology, risk factors, prevention, and early detection of

281 melanoma. Surg. Clin. North Am., Melanoma 94, 945–962.

282 <https://doi.org/10.1016/j.suc.2014.07.013>

- 283 Barone, A.N., Hayes, C.E., Kerr, J.J., Lee, R.C., Flaherty, D.B., 2019. Acute toxicity testing of
284 TiO₂-based vs. oxybenzone-based sunscreens on clownfish (*Amphiprion ocellaris*). Environ.
285 Sci. Pollut. Res. 26, 14513–14520. <https://doi.org/10.1007/s11356-019-04769-z>
- 286 Blüthgen, N., Meili, N., Chew, G., Odermatt, A., Fent, K., 2014. Accumulation and effects of the
287 UV-filter octocrylene in adult and embryonic zebrafish (*Danio rerio*). Sci. Total Environ.
288 476–477, 207–217. <https://doi.org/10.1016/j.scitotenv.2014.01.015>
- 289 Caldwell, G.S., Bentley, M.G., Olive, P.J.W., 2003. The use of a brine shrimp (*Artemia salina*)
290 bioassay to assess the toxicity of diatom extracts and short chain aldehydes. Toxicon 42,
291 301–306. [https://doi.org/10.1016/S0041-0101\(03\)00147-8](https://doi.org/10.1016/S0041-0101(03)00147-8)
- 292 Chen, L., Li, X., Hong, H., Shi, D., 2018. Multigenerational effects of 4-methylbenzylidene
293 camphor (4-MBC) on the survival, development and reproduction of the marine copepod
294 *Tigriopus japonicus*. Aquat. Toxicol. 194, 94–102.
295 <https://doi.org/10.1016/j.aquatox.2017.11.008>
- 296 Coronado, M., De Haro, H., Deng, X., Rempel, M.A., Lavado, R., Schlenk, D., 2008. Estrogenic
297 activity and reproductive effects of the UV-filter oxybenzone (2-hydroxy-4-methoxyphenyl-
298 methanone) in fish. Aquat. Toxicol. 90, 182–187.
299 <https://doi.org/10.1016/j.aquatox.2008.08.018>
- 300 Downs, C.A., Kramarsky-Winter, E., Segal, R., Fauth, J., Knutson, S., Bronstein, O., Ciner, F.R.,
301 Jeger, R., Lichtenfeld, Y., Woodley, C.M., Pennington, P., Cadenas, K., Kushmaro, A.,
302 Loya, Y., 2016. Toxicopathological effects of the sunscreen UV filter, oxybenzone
303 (benzophenone-3), on coral planulae and cultured primary cells and its environmental
304 contamination in Hawaii and the U.S. Virgin Islands. Arch. Environ. Contam. Toxicol. 70,
305 265–288. <https://doi.org/10.1007/s00244-015-0227-7>
- 306 Esperanza, M., Seoane, M., Rioboo, C., Herrero, C., Cid, Á., 2019. Differential toxicity of the UV-
307 filters BP-3 and BP-4 in *Chlamydomonas reinhardtii*: A flow cytometric approach. Sci.
308 Total Environ. 669, 412–420. <https://doi.org/10.1016/j.scitotenv.2019.03.116>

- 309 European Commission, 2007. Chemicals Legislation | Internal Market, Industry, Entrepreneurship
310 and SMEs [WWW Document]. URL
311 https://ec.europa.eu/growth/sectors/chemicals/legislation_en (accessed 12.3.19).
- 312 Fagervold, S.K., Rodrigues, A.M.S., Rohée, C., Roe, R., Bourrain, M., Stien, D., Lebaron, P., 2019.
313 Occurrence and environmental distribution of 5 UV filters during the summer season in
314 different water bodies. *Water. Air. Soil Pollut.* 230, 172–172.
- 315 Gago-Ferrero, P., Díaz-Cruz, M.S., Barceló, D., 2015. UV filters bioaccumulation in fish from
316 Iberian river basins. *Sci. Total Environ.* 518–519, 518–525.
317 <https://doi.org/10.1016/j.scitotenv.2015.03.026>
- 318 Gago-Ferrero, P., Díaz-Cruz, M.S., Barceló, D., 2011. Fast pressurized liquid extraction with in-cell
319 purification and analysis by liquid chromatography tandem mass spectrometry for the
320 determination of UV filters and their degradation products in sediments. *Anal. Bioanal.*
321 *Chem.* 400, 2195–2204. <https://doi.org/10.1007/s00216-011-4951-1>
- 322 Gao, L., Yuan, T., Zhou, C., Cheng, P., Bai, Q., Ao, J., Wang, W., Zhang, H., 2013. Effects of four
323 commonly used UV filters on the growth, cell viability and oxidative stress responses of the
324 *Tetrahymena thermophila*. *Chemosphere* 93, 2507–2513.
325 <https://doi.org/10.1016/j.chemosphere.2013.09.041>
- 326 Giraldo, A., Montes, R., Rodil, R., Quintana, J.B., Vidal-Liñán, L., Beiras, R., 2017.
327 Ecotoxicological evaluation of the UV filters ethylhexyl dimethyl *p*-aminobenzoic acid and
328 octocrylene using marine organisms *Isochrysis galbana*, *Mytilus galloprovincialis* and
329 *Paracentrotus lividus*. *Arch. Environ. Contam. Toxicol.* 72, 606–611.
330 <https://doi.org/10.1007/s00244-017-0399-4>
- 331 Hadjoudja, S., Vignoles, C., Deluchat, V., Lenain, J.-F., Le Jeune, A.-H., Baudu, M., 2009. Short
332 term copper toxicity on *Microcystis aeruginosa* and *Chlorella vulgaris* using flow
333 cytometry. *Aquat. Toxicol.* 94, 255–264. <https://doi.org/10.1016/j.aquatox.2009.07.007>

- 334 He, T., Tsui, M.M.P., Tan, C.J., Ma, C.Y., Yiu, S.K.F., Wang, L.H., Chen, T.H., Fan, T.Y., Lam,
335 P.K.S., Murphy, M.B., 2019. Toxicological effects of two organic ultraviolet filters and a
336 related commercial sunscreen product in adult corals. *Environ. Pollut.* 245, 462–471.
337 <https://doi.org/10.1016/j.envpol.2018.11.029>
- 338 Kaiser, D., Sieratowicz, A., Zielke, H., Oetken, M., Hollert, H., Oehlmann, J., 2012.
339 Ecotoxicological effect characterisation of widely used organic UV filters. *Environ. Pollut.*
340 163, 84–90. <https://doi.org/10.1016/j.envpol.2011.12.014>
- 341 Kameda, Y., Kimura, K., Miyazaki, M., 2011. Occurrence and profiles of organic sun-blocking
342 agents in surface waters and sediments in Japanese rivers and lakes. *Environ. Pollut.* 159,
343 1570–1576. <https://doi.org/10.1016/j.envpol.2011.02.055>
- 344 Langford, K.H., Thomas, K.V., 2008. Inputs of chemicals from recreational activities into the
345 Norwegian coastal zone. *J. Environ. Monit.* 10, 894–898. <https://doi.org/10.1039/B806198J>
- 346 Libralato, G., 2014. The case of *Artemia* spp. in nanoecotoxicology. *Mar. Environ. Res.* 101, 38–43.
347 <https://doi.org/10.1016/j.marenvres.2014.08.002>
- 348 Mao, F., He, Y., Kushmaro, A., Gin, K.Y.-H., 2017. Effects of benzophenone-3 on the green alga
349 *Chlamydomonas reinhardtii* and the cyanobacterium *Microcystis aeruginosa*. *Aquat.*
350 *Toxicol.* 193, 1–8. <https://doi.org/10.1016/j.aquatox.2017.09.029>
- 351 Nunes, B.S., Carvalho, F.D., Guilhermino, L.M., Van Stappen, G., 2006. Use of the genus *Artemia*
352 in ecotoxicity testing. *Environ. Pollut.* 144, 453–462.
353 <https://doi.org/10.1016/j.envpol.2005.12.037>
- 354 Pablos, M.V., García-Hortigüela, P., Fernández, C., 2015. Acute and chronic toxicity of emerging
355 contaminants, alone or in combination, in *Chlorella vulgaris* and *Daphnia magna*. *Environ.*
356 *Sci. Pollut. Res.* 22, 5417–5424. <https://doi.org/10.1007/s11356-015-4119-1>
- 357 Paredes, E., Perez, S., Rodil, R., Quintana, J.B., Beiras, R., 2014. Ecotoxicological evaluation of
358 four UV filters using marine organisms from different trophic levels *Isochrysis galbana*,

- 359 *Mytilus galloprovincialis*, *Paracentrotus lividus*, and *Siriella armata*. *Chemosphere* 104,
360 44–50. <https://doi.org/10.1016/j.chemosphere.2013.10.053>
- 361 Pecqueur, D., Vidussi, F., Fouilland, E., Le Floc'h, E., Mas, S., Roques, C., Salles, C., Tournoud,
362 M.-G., Mostajir, B., 2011. Dynamics of microbial planktonic food web components during a
363 river flash flood in a Mediterranean coastal lagoon. *Hydrobiologia* 673, 13–27.
364 <https://doi.org/10.1007/s10750-011-0745-x>
- 365 Prado, R., García, R., Rioboo, C., Herrero, C., Cid, Á., 2015. Suitability of cytotoxicity endpoints
366 and test microalgal species to disclose the toxic effect of common aquatic pollutants.
367 *Ecotoxicol. Environ. Saf.* 114, 117–125. <https://doi.org/10.1016/j.ecoenv.2015.01.021>
- 368 Sánchez-Quiles, D., Tovar-Sánchez, A., 2015. Are sunscreens a new environmental risk associated
369 with coastal tourism? *Environ. Int.* 83, 158–170.
370 <https://doi.org/10.1016/j.envint.2015.06.007>
- 371 Seoane, M., Esperanza, M., Rioboo, C., Herrero, C., Cid, Á., 2017. Flow cytometric assay to assess
372 short-term effects of personal care products on the marine microalga *Tetraselmis suecica*.
373 *Chemosphere* 171, 339–347. <https://doi.org/10.1016/j.chemosphere.2016.12.097>
- 374 Seoane, M., Rioboo, C., Herrero, C., Cid, Á., 2014. Toxicity induced by three antibiotics commonly
375 used in aquaculture on the marine microalga *Tetraselmis suecica* (Kylin) Butch. *Mar.*
376 *Environ. Res.* 101, 1–7. <https://doi.org/10.1016/j.marenvres.2014.07.011>
- 377 Shaath, N.A., 2010. Ultraviolet filters. *Photochem. Photobiol. Sci.* 9, 464–469.
378 <https://doi.org/10.1039/B9PP00174C>
- 379 Sieratowicz, A., Kaiser, D., Behr, M., Oetken, M., Oehlmann, J., 2011. Acute and chronic toxicity
380 of four frequently used UV filter substances for *Desmodesmus subspicatus* and *Daphnia*
381 *magna*. *J. Environ. Sci. Health Part A* 46, 1311–1319.
382 <https://doi.org/10.1080/10934529.2011.602936>
- 383 Stien, D., Clergeaud, F., Rodrigues, A.M.S., Lebaron, K., Pillot, R., Romans, P., Fagervold, S.,
384 Lebaron, P., 2019. Metabolomics reveal that octocrylene accumulates in *Pocillopora*

- 385 *damicornis* tissues as fatty acid conjugates and triggers coral cell mitochondrial dysfunction.
386 Anal. Chem. 91, 990–995. <https://doi.org/10.1021/acs.analchem.8b04187>
- 387 Torres, T., Cunha, I., Martins, R., Santos, M.M., 2016. Screening the toxicity of selected personal
388 care products using embryo bioassays: 4-MBC, propylparaben and triclocarban. Int. J. Mol.
389 Sci. 17, 1762. <https://doi.org/10.3390/ijms17101762>
- 390 Tsui, M.M.P., Leung, H.W., Wai, T.-C., Yamashita, N., Taniyasu, S., Liu, W., Lam, P.K.S.,
391 Murphy, M.B., 2014. Occurrence, distribution and ecological risk assessment of multiple
392 classes of UV filters in surface waters from different countries. Water Res. 67, 55–65.
393 <https://doi.org/10.1016/j.watres.2014.09.013>
- 394 Vidal-Liñán, L., Villaverde-de-Sáa, E., Rodil, R., Quintana, J.B., Beiras, R., 2018. Bioaccumulation
395 of UV filters in *Mytilus galloprovincialis* mussel. Chemosphere 190, 267–271.
396 <https://doi.org/10.1016/j.chemosphere.2017.09.144>
- 397 Waldman, R.A., Grant-Kels, J.M., 2019. The role of sunscreen in the prevention of cutaneous
398 melanoma and nonmelanoma skin cancer. J. Am. Acad. Dermatol. 80, 574-576.e1.
399 <https://doi.org/10.1016/j.jaad.2018.06.069>
- 400 Zhong, X., Downs, C.A., Che, X., Zhang, Z., Li, Yiman, Liu, B., Li, Q., Li, Yuting, Gao, H., 2019.
401 The toxicological effects of oxybenzone, an active ingredient in sunscreen personal care
402 products, on prokaryotic alga *Arthrospira* sp. and eukaryotic alga *Chlorella* sp. Aquat.
403 Toxicol. 216, 105295. <https://doi.org/10.1016/j.aquatox.2019.105295>
- 404 Zucchi, S., Oggier, D.M., Fent, K., 2011. Global gene expression profile induced by the UV-filter
405 2-ethyl-hexyl-4-trimethoxycinnamate (EHMC) in zebrafish (*Danio rerio*). Environ. Pollut.
406 159, 3086–3096. <https://doi.org/10.1016/j.envpol.2011.04.013>
- 407