



**HAL**  
open science

# Enantiopure Cyclometalated Rh(III) and Ir(III) Complexes Displaying Rigid Configuration at Metal Center: Design, Structures, Chiroptical Properties and Role of the Iodide Ligand

Antoine Groué, Jean-Philippe Tranchier, Geoffrey Gontard, Marion Jean, Nicolas Vanthuyne, Hani Amouri

► **To cite this version:**

Antoine Groué, Jean-Philippe Tranchier, Geoffrey Gontard, Marion Jean, Nicolas Vanthuyne, et al.. Enantiopure Cyclometalated Rh(III) and Ir(III) Complexes Displaying Rigid Configuration at Metal Center: Design, Structures, Chiroptical Properties and Role of the Iodide Ligand. *Chemistry*, 2022, 4 (1), pp.156-167. 10.3390/chemistry4010014 . hal-03811858

**HAL Id: hal-03811858**

<https://hal.sorbonne-universite.fr/hal-03811858v1>

Submitted on 12 Oct 2022

**HAL** is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers.

L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.

1 Article

 2 **Enantiopure cyclometalated Rh(III) and Ir(III) complexes displaying rigid configuration at metal center: Design, structures,**  
 3 **chiroptical properties and role of the iodide ligand**

 5 Antoine Groue,<sup>1</sup> Jean Philippe Tranchier,<sup>1</sup> Geoffrey Gontard,<sup>1</sup> Marion Jean,<sup>2</sup> Nicolas Vanthuyne,<sup>2</sup> and Hani Amouri  
 6 <sup>1,\*</sup>

 7 <sup>1</sup> Sorbonne Université- Campus Pierre et Marie Curie, Institut Parisien de Chimie Moléculaire (IPCM) UMR  
 8 CNRS 8232, 4 place Jussieu, 75252 Paris cedex 05, France; antoinegroue@gmail.com ;  
 9 jean-philippe.tranchier@sorbonne-universite.fr ;geoffrey.gontard@sorbonne-universite.fr; ha-  
 10 ni.amouri@sorbonne-universite.fr .

 11 <sup>2</sup> Aix Marseille Univ, CNRS, Centrale Marseille, iSm2, Marseille, France.; marion.jean@univ-amu.fr; nico-  
 12 las.vanthuyne@univ-amu.fr.

13 \* Correspondence: HA : hani.amouri@sorbonne-universite.fr

 14 *Dedicated in memoriam to Professor Alexander von Zelewsky*

 16 **Abstract:** Enantiopure *N*-heterocyclic carbene half-sandwich metal complexes of the general formula [Cp\*M(C<sup>∧</sup>C:)] (M = Rh, Ir; C<sup>∧</sup>C: = NI-NHC; NI-H = Naphthalimide; NHC = *N*-heterocyclic carbene) are reported. The rhodium compound was obtained as a single isomer displaying six membered metallacycle and was resolved on chiral column chromatography to the corresponding enantiomers (*S*)-[Cp\*Rh(C<sup>∧</sup>C:)] (*S*)-**2** and (*R*)-[Cp\*Rh(C<sup>∧</sup>C:)] (*R*)-**2**. The iridium congener however, furnishes a pair of regioisomers, which were resolved into (*S*)-[Cp\*Ir(C<sup>∧</sup>C:)] (*S*)-**3** and (*R*)-[Cp\*Ir(C<sup>∧</sup>C:)] (*R*)-**3** and (*S*)-[Cp\*Ir(C<sup>∧</sup>C:)] (*S*)-**4** and (*R*)-[Cp\*Ir(C<sup>∧</sup>C:)] (*R*)-**4**. These regioisomers differ from each other, only by the size of the metallacycle; Five-membered for **3** and six-membered for **4**. The molecular structures of (*S*)-**2** and (*S*)-**4** are reported. Moreover, the chiroptical properties of these compounds are presented and discussed. These compounds display exceptional stable configurations at the metal center in solution with enantiomerization barrier Δ*G*<sup>‡</sup> up to 124 kJ/mol. This is because the nature of the naphthalimide-NHC clamp ligand and the iodide ligand contribute to their configuration's robustness. In contrast to related complexes reported in the literature, which are often labile in solution.

 26 **Citation:** Lastname, F.; Lastname, F.  
 27 Lastname, F. Title. *Chemistry* **2022**, *4*, 28  
 28 Firstpage–Lastpage.  
 29 <https://doi.org/10.3390/xxxxx>

 30 Academic Editor: Firstname  
 31 Lastname

Received: date

Accepted: date

32 Published: date

 33 **Publisher's Note:** MDPI stays  
 34 neutral with regard to jurisdictional  
 35 claims in published maps and  
 36 institutional affiliations.

 37 **Copyright:** © 2022 by the authors.  
 38 Submitted for possible open access  
 39 publication under the terms and  
 40 conditions of the Creative Commons  
 41 Attribution (CC BY) license  
 42 (<https://creativecommons.org/licenses/by/4.0/>).

 30 **Keywords:** Chiral resolution; Enantiopure; Configurational stability; Circular dichroism.

 32 **1. Introduction**

 33 Chirality is an ever-fascinating topic and occurs in many fields of science.[1-4] In the  
 34 area of transition metal complexes Brunner [5,6], Gladysz [7,8], von Zelewsky[9],  
 35 Meggers[10], Constable [11] and others[12,13] have made great contributions to the ad-  
 36 vance and comprehension of the elements that control the chirality at metal centers at the  
 37 molecular and supramolecular levels.[11,14] For instance octahedral iridium complexes  
 38 displaying helical chirality (*Δ*, *Λ*) show a stable configuration at the metal center.[15-19]  
 39 Coordination and organometallic complexes with planar chirality show also stable con-  
 40 figuration.[20-25] In contrast, half-sandwich rhodium and iridium complexes with piano  
 41 stool geometry, displaying central chirality are labile in solution, as demonstrated by  
 42 Brunner and co-workers and others.[26-29] More recently efforts were devoted to using  
 43 strongly coordinated *N*-heterocyclic carbene (NHC) ligands to increase the stability at the  
 44 metal center, however, only a few examples were reported.[30-32]


We recently described the synthesis of some half-sandwich iridium complexes displaying stable configuration at metal center.[32] Such compounds contain Cp\*Ir complexes in which the NHC unit is directly attached to a naphthalimide (NI) molecule. The latter chelates strongly the metal center; moreover an iodide ligand completes the coordination sphere around the metal center. Iodide is known to act as a strong  $\sigma$ -donor and a weak-leaving group[33], thus we anticipated such a combination of ligands should bring rigidity to the configuration of the metal center. In this work we extended our synthetic procedures to the related rhodium congener. Interestingly only one regioisomer complex is obtained, displaying a six-membered metallacycle. The iridium complexes are obtained as two regioisomers in roughly 1: 1 ratio as a mixture of five-membered and six membered iridacycles (Scheme 1).

The racemic complexes were all resolved using chiral column chromatography. The chiroptical optical properties were studied and the configurational stability were investigated by measuring the enantiomerization barrier at T = 60°C for these complexes and were found to be up to  $\Delta G^\ddagger_{\text{enantiomerization}} = 124$  kJ/mole, suggesting that these compounds display high configurational stability at room temperature.

## 2. Results and Discussions

### 2.1 Synthesis and characterization

The target compounds **2-4** were prepared by treatment of two equivalents of the imidazolium salt (**1**) with the chloro-bridged dimeric precursor  $[\text{Cp}^*\text{M}(\mu\text{-Cl})\text{Cl}]_2$  in the presence of  $\text{Cs}_2\text{CO}_3$  and  $\text{NBu}_4\text{I}$ , in  $\text{CH}_3\text{CN}$  under reflux overnight. When  $[\text{Cp}^*\text{Rh}(\mu\text{-Cl})\text{Cl}]_2$  was used as starting material only one single rhodium carbene compound identified as  $[\text{Cp}^*\text{Rh}(\text{NI-NHC})\text{I}]$  (**2**) (NI-H = Naphthalimide) was obtained in good yield as red microcrystalline solid after eluting on a column chromatography. The  $^1\text{H-NMR}$  of **2** recorded in  $\text{CDCl}_3$  showed the presence of a singlet assigned to  $\text{Cp}^*\text{Rh}$  protons at  $\delta 1.51$  ppm and the aromatic protons of the naphthalimide and NHC moiety appeared in the range of  $\delta 7.1$  to  $\delta 8.61$  ppm.



Scheme 1. Preparation of the cyclometalated half-sandwich complexes **2-4**.

Starting with iridium precursor  $[\text{Cp}^*\text{Ir}(\mu\text{-Cl})\text{Cl}]_2$ , under similar experimental conditions, two regioisomeric complexes **3-4** were obtained. These compounds display the same chemical composition but differ only with the size of the iridacycle. The formation of the two isomers results from the different modes of metallacyclization; (a) whether the cyclization process takes place at the arene attached directly to the NHC moiety to give the five membered metallacycle. (b) Metallacyclization occurs at the other arene ring generating the six-membered metallacycle. Both isomers **3** and **4** were obtained in 1: 1 ratio. The  $^1\text{H-NMR}$  of **4** recorded in  $\text{CDCl}_3$  displayed a singlet at  $\delta 1.54$  ppm, assigned to the  $\text{Cp}^*\text{Ir}$  protons, this upfield shift is a diagnosis to the formation of a six-membered metallacycle.

The aromatic protons of the naphthalimide and NHC moiety appeared in the range of  $\delta$  7.1 to  $\delta$  8.57 ppm. These data are similar to those obtained for complex **2**. The  $^1\text{H-NMR}$  of compound **3**, on the other hand, displayed a singlet at  $\delta$  1.96 ppm to the  $\text{Cp}^*\text{Ir}$  protons, which is downfield with respect to the six-membered metallacycle complexes **2** and **4**. Moreover the aromatic protons of the naphthalimide and NHC moiety appeared downfield in the range 7.1 to 9.2 ppm and displayed a different pattern. Furthermore, 2D COSY, ROESY, HSQC, and HMBC experiments carried out on complexes **3** and **4** allowed us to fully characterize both isomers (Figures S1-S2, 2D ROESY Spectra of **3** and **4**). In addition, the identity of these molecules was also confirmed by determining the molecular structures of the enantiopure complexes *S*-**2** and *S*-**4** (vide infra).




Figure 1. Optically active molecules (*S*)-**2**, (*S*)-**3** and (*S*)-**4**/ (*R*)-**2**, (*R*)-**3** and (*R*)-**4** described in this work and priority rule used ( $\text{I} > \text{Cp} > \text{C-Carbene} > \text{C-arene}$ ) to assign the absolute configuration.[34,35]

## 2.2 Chiral resolution of complexes $[\text{Cp}^*\text{M}(\text{C}^{\wedge}\text{C})\text{I}]$ ( $M = \text{Rh}$ , (**2**); $M = \text{Ir}$ (**3-4**) and chiroptical properties

Half-sandwich rhodium and iridium *N*-heterocyclic carbene complexes (**2-4**) were resolved on the chiral stationary phase (Experimental details are given in the SI). For the rhodium complex, Chiralpak IF column was chosen using heptane/ethanol/ dichloromethane (50/30/20) mixture as eluent. The related iridium complexes **3-4** were resolved using (*S,S*)-Whelk-O1 chiral column and using the same solvent mixture as eluent. Remarkably all complexes were stable during resolution process highlighting their robustness and were obtained with  $> 98\%$  ee. The CD spectra of the enantiomers of **2**, **3**, and **4** are given in Figures 2, 3, and 4. The optical rotations of both enantiomers were monitored as well (experimental details of their resolution are given in the SI).




Figure 2. CD spectra of (*S*)-2 (green) and (*R*)-2 (red) recorded in CH<sub>3</sub>CN at 0.251mM.

The CD curves of both enantiomers (Figure 2) displayed an excellent mirror image relationship. For instance, the CD trace of *S*-2 displayed a positive band at 200 nm ( $\Delta\epsilon = +42$ ) and a negative band at 265 nm ( $\Delta\epsilon = -42$ ), followed by a broad positive band at 390 nm ( $\Delta\epsilon = +36$ ) and a negative band at 445 nm ( $\Delta\epsilon = -16$ ). The first eluted complex (CD green trace) was crystallized and provided suitable crystals for x-ray diffraction study. The x-ray molecular structure confirmed the identity of the complex (vide infra). Grati-fyingly all enantiopure compounds displayed stable configuration in strongly coordinated CH<sub>3</sub>CN solution and did not epimerize upon standing for several days in solution as demonstrated by their CD curves (vide infra).

On the other hand, the CD curves of both enantiomers of 3 displayed again an excellent mirror image patterns confirming the enantiomeric relationship between the two complexes (*S*-3 and *R*-3). It is noteworthy that the positive and negative Cotton bands above 350 nm appeared weaker than those observed for enantiomers of 2 and 4 suggesting perhaps the influence of the size of the five- or six-membered metallacycles (vide infra)




Figure 3. CD spectra of *S*-3 (green) and *R*-3 (red) recorded in CH<sub>3</sub>CN at 0.24mM.

The CD spectra of both enantiomers of 4 are shown in Figure 4. The traces showed opposite image absorption bands, confirming the enantiomeric relationship. Moreover, the absorption profile appears to be comparable to those of complex 2 displaying a six membered metallacycle. For instance, the CD trace of (*S*)-4 showed a negative cotton band at 267 nm ( $\Delta\epsilon = -26$ ), and a broad positive band at 371 nm ( $\Delta\epsilon = +27$ ) comparable to those observed for (*S*)-2. Crystals of the first eluted sample (*S*)-4 were obtained for x-ray diffraction. The structure is presented and discussed in the next section.




Figure 4. CD spectra of (*S*)-4 (green) and (*R*)-4 (red) recorded in CH<sub>3</sub>CN at 0.24mM.

2.3 Molecular structures of *S*-2 and *S*-4

Single crystals of (*S*)-2 were obtained by slow diffusion of diethyl ether into a CH<sub>2</sub>Cl<sub>2</sub> solution of the complex. X-ray crystallography confirmed the proposed structure of complex (*S*)-2 (Figure 5).




Figure 5. Molecular structure of (*S*)-2 with thermal ellipsoids drawn at the 30% probability level, hydrogen atoms were omitted for clarity. The asymmetric unit comprises two molecules with the same configuration and almost identical conformations, differing only in their nBu-groups. One molecule was arbitrarily chosen for representation. Selected average bond distances (Å) and angles (deg): Rh1-I1 2.699(2), Rh1-C1 1.983(5), C1-N1 1.371(7), N1-C5 1.414(6), C5-C10 1.426(7), C10-C11 1.427(7), C11-Rh1 2.026(8), Rh1-C21 2.154(5), Rh1-C22 2.248(8), Rh1-C23 2.289(5), Rh1-C24 2.276(5), Rh1-C25 2.248(5), C1-Rh1-C11 86.5(3), Rh1-C1-N1 126.0(6), Rh1-C11-C10 122.2(12), C1-N1-C5 125.8(4), C11-C10-C5 123.9(5), N1-C5-C10 120.0(4).

The structure shows the formation of the six-membered rhodacycle, which adopts a boat conformation. The absolute configuration around the metal center was determined to be (*S*) by the refinement of the Flack *x* parameter (Table 1). The metal center is chelated by the (C<sup>^</sup>C) NHC-naphthalimide ligand, is symmetrically bound to η-Cp\*. Finally, the iodide ligand completes the coordination sphere around the rhodium and confers a distorted tetrahedral geometry. The iridium complex (*S*)-4 was found to be isostructural to that of the rhodium complex (*S*)-2 (Figure 6). We note however the nBu-group of the naphthalimide points downward opposite to that observed for (*S*)-2. The absolute configuration of the complex was assigned by refining the Flack *x* parameter (Table 1).




Figure 6. Molecular structure of (*S*)-4 with thermal ellipsoids drawn at the 30% probability level, hydrogen atoms were omitted for clarity. Selected bond distances (Å) and angles (deg): Ir1-I1 2.702(1), Ir1-C1 1.982(5), C1-N1 1.367(6), N1-C5 1.430(6), C5-C10 1.422(7), C10-C11 1.422(7), C11-Ir1

2.053(4), Ir1-C21 2.168(5), Ir1-C22 2.230(5), Ir1-C23 2.271(4), Ir1-C24 2.279(4), Ir1-C25 2.246(4), C1-Ir1-C11 86.6(2), Ir1-C1-N1 125.1(3), Ir1-C11-C10 121.0(4), C1-N1-C5 125.8(4), C11-C10-C5 123.8(4), N1-C5-C10 119.7(4).

#### 2.4 Determination of the enantiomerization barriers and configurational stability

To evaluate the stability of the enantiopure complex, (*R*)-**2** was dissolved in acetonitrile, heated at 60°C and then analyzed by chiral HPLC. Complex **2** was chemically stable but underwent racemization. Monitoring of the enantiomeric excess along time revealed a clean first-order reaction and allowed to determine an enantiomerization barrier of 115.9 kJ/mol. Half-life time was extrapolated at 25°C by considering zero enantiomerization entropy: 130 days for **2** at 25°C in acetonitrile. So, enantiopure **2** can be handled and evaporated without any risk of enantiopurity loss. Similar racemization kinetics studies performed on (*S*)-**3** and (*R*)-**4**, provided the enantiomerization barriers in acetonitrile, to be 124.4 kJ/mol and 122.3 kJ/mol, respectively. These values confirm the chemical robustness of these complexes and allowed the estimation of their half-life times at 25°C, to be 11 and 5 years for **3** and **4**, respectively. Highlighting the role of the metallacycle size 5-membered versus 6-membered. The configurational stability is weaker in ethanol, with lower enantiomerization barriers of 4 kJ/mol (see Supporting Information for details). The racemization is faster with rhodium than with iridium, and in protic solvents, this facilitate the iodide decoordination as one might expect. These results underline the importance of iodine on the stabilization of the chiral configuration at the metal center. For comparison purpose, we note that *Brunner and co-workers* have studied the half sandwich complexes [Cp\*M(N<sup>+</sup>N<sup>-</sup>)Cl] (M = Rh, Ir) where N<sup>+</sup>N<sup>-</sup> is the anion of (+)-2-N-[(*S*)-1-phenylethylpyrrolcarbaldimine] and showed that such compounds are configurationally unstable at metal centers. Time dependent integration of <sup>1</sup>H NMR signals of the complexes revealed that the epimerization is a first-order reaction. The half-lives in CD<sub>2</sub>Cl<sub>2</sub> solution at -50°C were 19.4 min for the rhodium complex and 30.6 min for iridium compound.[36]

#### 2.5 Discussion

As mentioned in the introduction of this manuscript, half-sandwich metal complexes continue to attract attention due to their importance in medicinal chemistry[37,38], materials science[39,40] and catalysis.[41-43] Of prime importance is the control of the configurational stability at the metal center. For instance Brunner[26], White[27] and others [29] demonstrated that half sandwich metal complexes of rhodium and iridium racemize in solution, because they display labile configuration at metal center.

More recently efforts were made to use strongly donor NHC ligands[44,45] associated with helicene[31] or other elements of chirality to increase the configurational rigidity at the metal center with some success. Thus, our complexes presented in this work display only one stereogenic element of chirality. This is a rare example of chiral half-sandwich metal complexes displaying exceptional rigidity at the metal center. The data obtained from the determination of the free energy of enantiomerization confirm this behavior.


We feel this unique class of complexes owe their rigid configuration to the nature of the NI-NHC (C<sup>+</sup>C<sup>-</sup>) ligand that acts as a clamp and strongly chelates the metal center. Moreover, iodide ligand is known to act as a strong donor ligand but a weak leaving group[46], which permits to lock of the configuration at the metal center in these complexes. Therefore, our judicious choice to combine strongly NHC-chelating (C<sup>+</sup>C<sup>-</sup>) ligands and iodide generate novel class of chiral cyclometalated complexes with exceptional configuration at the metal center which might show interesting applications in variety of fields, spanning from medical chemistry, organometallic catalysis to materials science.

### 3. Materials and Methods

#### 2.1 General synthetic procedure

All synthetic manipulations were carried out under argon using Schlenk tube techniques. The  $^1\text{H}$  and  $^{13}\text{C}$ , and spectra were recorded in  $\text{CD}_2\text{Cl}_2$  and  $\text{CD}_3\text{CN}$  using a Bruker Avance 400 and 300 NMR spectrometer at 400.13 and 100.61 and 76.31 MHz respectively. The azolium salt [IMZ-NI-H][I] (**1**) was prepared according to our procedure published in the literature.[47-49]


**Synthesis of 2.** In a dry Schlenk tube, imidazolium salt (**1**) (150 mg, 0.33 mmol),  $[\text{Cp}^*\text{Rh}(\mu\text{Cl})\text{Cl}]_2$  (60 mg, 0.11 mmol),  $\text{Cs}_2\text{CO}_3$  (82 mg, 0.253 mmol) and  $\text{NBu}_4\text{I}$  (243 mg, 0.66 mmol) were mixed in 15 mL of distilled and degassed acetonitrile, the solution was refluxed overnight. The resulting solution was cooled at r.t., filtered to remove a precipitate and washed with distilled acetonitrile. After evaporation of the solvents, the crude solid was purified by chromatography column ( $\text{SiO}_2$ ) with a mixture of  $\text{CH}_2\text{Cl}_2$  / Acetone (99/1). The desired fractions were then concentrated to give product **2** as orange microcrystalline solid in 65% yield.



$^1\text{H}$  NMR (300 MHz,  $\text{CDCl}_3$ )  $\delta$  8.61 (dd,  $J = 8.0, 2.1$  Hz, 2H,  $\text{H}_{9,11\text{napht}}$ ), 8.21 (d,  $J = 7.8$  Hz, 1H,  $\text{H}_{12\text{napht}}$ ), 7.78 (d,  $J = 2.4$  Hz, 1H,  $\text{H}_{3\text{imid}}$ ), 7.65 (d,  $J = 8.0$  Hz, 1H,  $\text{H}_{4\text{napht}}$ ), 7.28 (q,  $J = 2.3$  Hz, 1H,  $\text{H}_{2\text{imid}}$ ), 4.20 (t,  $J = 7.7$  Hz, 2H,  $\text{H}_{17}$ ), 4.06 (d,  $J = 2.2$  Hz, 3H,  $\text{H}_{16}$ ), 1.76 (t,  $J = 8.2$  Hz, 2H,  $\text{H}_{18}$ ), 1.51 (s, 15H,  $\text{CH}_3$   $\text{Cp}^*$ ), 1.28 (m, 2H,  $\text{H}_{19}$ ), 1.01 (m,  $J = 6.4, 5.7, 3.8$  Hz, 3H,  $\text{H}_{20}$ ).

$^{13}\text{C}$  NMR (101 MHz,  $\text{CD}_2\text{Cl}_2$ )  $\delta$  178.02 ( $\text{C}_{14}$ ), 177.58 ( $\text{C}_{15}$ ), 165.12 ( $\text{C}_1$ ), 163.22 ( $\text{C}_{10}$ ), 144.34 ( $\text{C}_{11}$ ), 141.09 ( $\text{C}_5$ ), 129.84 ( $\text{C}_9$ ), 128.66 ( $\text{C}_6$ ), 128.28 ( $\text{C}_7$ ), 125.07 ( $\text{C}_2$ ), 122.44 ( $\text{C}_8$ ), 119.86 ( $\text{C}_3$ ), 116.92 ( $\text{C}_{13}$ ), 112.96 ( $\text{C}_4$ ), 99.55 ( $\text{C}_q$   $\text{Cp}^*$ ), 41.29 ( $\text{C}_{16}$ ), 39.85 ( $\text{C}_{17}$ ), 30.27 ( $\text{C}_{18}$ ), 20.43 ( $\text{C}_{19}$ ), 13.65 ( $\text{C}_{20}$ ), 9.37 ( $\text{CH}_3$   $\text{Cp}^*$ ). Anal. calcd. for **2** C 51.67, H 4.77, N 6.03; Found: C 51.99, H 4.77, N 5.72.


**Synthesis of 3 and 4.** In a dry Schlenk tube, imidazolium salt (**1**) (88 mg, 0.185 mmol),  $[\text{Cp}^*\text{Ir}(\mu\text{Cl})\text{Cl}]_2$  (50 mg, 0.061 mmol),  $\text{Cs}_2\text{CO}_3$  (46 mg, 0.14 mmol) and  $\text{NBu}_4\text{I}$  (137 mg, 0.37 mmol) were mixed in 15 mL of distilled and degassed acetonitrile, the solution was refluxed overnight. The resulting solution was cooled at r.t., filtered to remove a precipitate and washed with distilled acetonitrile. After evaporation of the solvents, the crude solid was purified by chromatography column ( $\text{SiO}_2$ ) with a mixture of  $\text{CH}_2\text{Cl}_2$  / Acetone (99/1). The desired fractions were then concentrated to give products **3** and **4** in almost 1/1 ratio.



**3** : A crystalline red powder. Yield 32 mg (33 %).  $^1\text{H}$  NMR (500 MHz,  $\text{CDCl}_3$ )  $\delta$  9.09 (s, 1H,  $\text{H}_9\text{napht}$ ), 8.48 (dd,  $J = 8.7; 0.9$  Hz, 1H,  $\text{H}_{10\text{napht}}$ ), 8.45 (dd,  $J = 7.3; 0.9$  Hz, 1H,  $\text{H}_{12\text{napht}}$ ), 8.07 (d,  $J = 2.3$  Hz, 1H,  $\text{H}_{3\text{imid}}$ ), 7.71 (dd,  $J = 8.7, 7.3$  Hz, 1H,  $\text{H}_{11\text{napht}}$ ), 7.19 (d,  $J = 2.3$  Hz, 1H,  $\text{H}_{2\text{imid}}$ ), 4.23 - 4.15 (m, 2H,  $\text{H}_{17}$ ), 3.98 (s, 3H,  $\text{H}_{16}$ ), 1.96 (s, 15H,  $\text{CH}_3$   $\text{Cp}^*$ ), 1.76-1.70 (m, 2H,  $\text{H}_{18}$ ), 1.50 - 1.42 (m, 2H,  $\text{H}_{19}$ ), 0.98 (t,  $J = 7.4$  Hz, 3H,  $\text{H}_{20}$ )  $^{13}\text{C}$  NMR (125 MHz, Chloroform- $d$ )  $\delta$  169.4 ( $\text{C}_1$ ), 165.0 ( $\text{C}_{14}$ ), 164.3 ( $\text{C}_{15}$ ), 145.9 ( $\text{C}_5$ ), 143.4 ( $\text{C}_9$ ), 142.3 ( $\text{C}_4$ ), 128.0 ( $\text{C}_{12}$ ), 127.0 ( $\text{C}_7$ ),



265 125.8 (C<sub>11</sub>), 125.3 (C<sub>10</sub>), 123.5 (C<sub>13</sub>), 122.2 (C<sub>2</sub>), 120.0 (C<sub>6</sub>), 118.8 (C<sub>3</sub>), 117.9 (C<sub>8</sub>), 93.0 (C<sub>q</sub>.Cp\*),  
 266 40.3 (C<sub>17</sub>), 38.4 (C<sub>16</sub>), 30.4 (C<sub>18</sub>), 20.6 (C<sub>19</sub>), 14.0 (C<sub>20</sub>), 10.5 (CH<sub>3</sub>.Cp\*). Anal. calcd. for 3 C  
 267 45.74, H 4.35, N 5.21; Found: C 45.71, H 4.39, N 5.08.  
 268



269  
 270  
 271 **4**: A crystalline red powder. Yield 38 mg (39%). <sup>1</sup>H NMR (500 MHz, CDCl<sub>3</sub>) δ 8.57 (d,  
 272 *J* = 8.1 Hz, 1H, H<sub>9</sub><sub>naph</sub>), 8.38 (d, *J* = 7.8 Hz, 1H, H<sub>11</sub><sub>naph</sub>), 8.16 (d, *J* = 7.8 Hz, 1H, H<sub>12</sub><sub>naph</sub>), 7.69  
 273 (d, *J* = 2.3 Hz, 1H, H<sub>3</sub><sub>imid</sub>), 7.54 (d, *J* = 8.1 Hz, 1H, H<sub>4</sub><sub>naph</sub>), 7.22 (d, *J* = 2.3 Hz, 1H, H<sub>2</sub><sub>imid</sub>), 4.18  
 274 – 4.14 (m, 2H, H<sub>17</sub>), 3.98 (s, 3H, H<sub>16</sub>), 1.76–1.70 (m, 2H, H<sub>18</sub>), 1.54 (s, 15H, CH<sub>3</sub> Cp\*), 1.50 –  
 275 1.42 (m, 2H, H<sub>19</sub>), 0.98 (t, *J* = 7.4 Hz, 3H, H<sub>20</sub>). <sup>13</sup>C NMR (125 MHz, CDCl<sub>3</sub>) δ 165.4 (C<sub>14</sub>),  
 276 163.8 (C<sub>15</sub>), 158.5 (C<sub>1</sub>), 158.4 (C<sub>10</sub>), 145.5 (C<sub>11</sub>), 141.6 (C<sub>5</sub>), 130.2 (C<sub>12</sub>), 129.8 (C<sub>9</sub>), 128.5 (C<sub>6</sub>),  
 277 128.3 (C<sub>7</sub>), 124.0 (C<sub>2</sub>), 122.2 (C<sub>8</sub>), 118.8 (C<sub>3</sub>), 116.5 (C<sub>13</sub>), 112.8 (C<sub>4</sub>), 93.7 (C<sub>q</sub>.Cp\*), 40.9 (C<sub>16</sub>),  
 278 40.1 (C<sub>17</sub>), 30.4 (C<sub>18</sub>), 20.6 (C<sub>19</sub>), 14.0 (C<sub>20</sub>), 9.4 (CH<sub>3</sub>.Cp\*). Anal. calcd. for 4 C 45.74, H 4.35,  
 279 N 5.21; Found: C 45.65, H 4.30, N 5.09.  
 280  
 281

282 **X-Ray crystal structure determination.** A single crystal was selected, mounted and  
 283 transferred into a cold nitrogen gas stream. Intensity data was collected with a Bruker  
 284 Kappa-APEX2 system using micro-source Cu-Kα radiation. Unit-cell parameters deter-  
 285 mination, data collection strategy, integration and absorption correction were carried out  
 286 with the Bruker APEX2 suite of programs. The structure was solved with SHELXT and  
 287 refined anisotropically by full-matrix least-squares methods with SHELXL using WinGX.  
 288 Absolute structure was determined by anomalous scattering effects analysis and chemi-  
 289 cal absolute configuration was then deduced. The structures were deposited at the  
 290 Cambridge Crystallographic Data Centre with numbers CCDC 2145150 and 2145151  
 291 which can be obtained free of charge via [www.ccdc.cam.ac.uk](http://www.ccdc.cam.ac.uk).

292 Table 1. Summary of crystallographic, data collection and refinement parameters for  
 293 (S)-[Cp\*<sup>+</sup>Rh(C<sup>−</sup>C:)] (S)-2 and (S)-[Cp\*<sup>+</sup>Ir(C<sup>−</sup>C:)] (S)-4.

Compound	(S)-[Cp* <sup>+</sup> Rh(C <sup>−</sup> C:)] (S)-2	(S)-[Cp* <sup>+</sup> Ir(C <sup>−</sup> C:)] (S)-4
Empirical formula	C <sub>30</sub> H <sub>33</sub> I RhN <sub>3</sub> O <sub>2</sub>	C <sub>30</sub> H <sub>33</sub> I Ir N <sub>3</sub> O <sub>2</sub>
Formula weight	697.40	786.69
Crystal system	Monoclinic	Orthorhombic
Space group	P 2 <sub>1</sub>	P 2 <sub>1</sub> 2 <sub>1</sub> 2 <sub>1</sub>
Unit cell dimensions	a = 17.2315(4) Å	a = 8.0943(2) Å
	b = 10.6098(2) Å	b = 16.2456(4) Å
	c = 17.2604(4) Å	c = 21.3531(5) Å
	α = 90°	α = 90°
	β = 119.616(1)°	β = 90°
	γ = 90°	γ = 90°
Volume	2743.34(11) Å <sup>3</sup>	2807.86(12) Å <sup>3</sup>
Z	4	4
Temperature	200(1) K	200(1) K
Wavelength	1.54178 Å	1.54178 Å

$\theta$ range for data collection	5.10° to 66.62°	4.96° to 66.64°
Reflections (all / independent)	34037 / 9689	15376 / 4959
R(int)	2.54%	2.28%
Completeness	99.7%	99.8%
Data / parameters / restraints	4959 / 340 / 0	9689 / 679 / 1
R1 [ $I > 2\sigma(I)$ ]	2.08%	1.79%
wR2 (all data)	5.31%	4.55%
Flack parameter	-0.015(2)	0.001(3)

294  
295  
296  
297  
298  
299  
300  
301  
302  
303  
304  
305  
306  
307  
  
308  
  
309  
  
310  
  
311  
312  
313  
314  
315  
316  
  
317  
318  
319  
320  
321  
  
322  
323  
  
324  
  
325

#### 4. Conclusions

In this work we reported a novel procedure to prepare half sandwich piano stool complexes of rhodium and iridium. These complexes were resolved using chiral column chromatography into the corresponding enantiomers. The enantiomerization barrier was measured in acetonitrile and ethanol and was up to 124 kJ/mol at 60°C. This data suggests that our compounds display rigid configuration at the metal center in solution. Our choice to combine a strong chelating (C<sup>^</sup>C:) ligand with a strong donor/weak leaving iodide ligand[33,46] generates complexes with strong configuration at the metal center. Due to the presence of an organic chromophore, these complexes might be used to prepare novel luminescent materials or as photocatalysts; the results will be disclosed in due course.

#### Supporting Materials.

**Author Contributions:** The synthesis and spectroscopic characterization of the new complexes were performed by Antoine Groué and Jean-philippe Tranchier. The X-ray structural determination was carried out by Geoffrey Gontard. Dr. M. Jean and Dr. N. Vanthuyne performed the chiral separation of the enantiomers and determined the optical rotation and CD spectra. All authors participated in the discussion. Preparation and writing of the manuscript were made by Hani Amouri who also directed the project

**Data Availability Statement:** In this section, please provide details regarding where data supporting reported results can be found, including links to publicly archived datasets analyzed or generated during the study. Please refer to suggested Data Availability Statements in section "MDPI Research Data Policies" at <https://www.mdpi.com/ethics>. If the study did not report any data, you might add "Not applicable" here.

**Acknowledgments:** This work was supported by the CNRS, by the Sorbonne Université, campus Pierre et Marie Curie which we gratefully acknowledged.

**Conflicts of Interest:** The authors declare no conflict of interest.

## References

- 326  
327  
328
- 329 1. von Zelewsky, A. *Stereochemistry of Coordination Compounds*. Wiley: Chichester, U.K., 1996; p 254 pp.
  - 330 2. Constable, E.C. Through a Glass Darkly-Some Thoughts on Symmetry and Chemistry. *Symmetry-Basel* **2021**, *13*.
  - 331 3. Collet, A.; Crassous, J.; Dutasta, J.P.; Guy, L. *Molécules Chirales: Stéréochimie et Propriétés*. EDP Sciences: 2006.
  - 332 4. Amouri, H.; Gruselle, M. *Chirality in Transition Metal Chemistry: Molecules, Supramolecular Assemblies and Materials*. Wiley:  
333 Chichester, U.K., 2008.
  - 334 5. Brunner, H. Optical Activity at an Asymmetrical Manganese Atom. *Angew. Chem. Int. Ed.* **1969**, *8*, 382-383.
  - 335 6. Brunner, H.; Tsuno, T. Ligand Dissociation: Planar or Pyramidal Intermediates? *Acc. Chem. Res.* **2009**, *42*, 1501-1510.
  - 336 7. Merrifield, J.H.; Strouse, C.E.; Gladysz, J.A. Synthesis, Optical Resolution, and Absolute-Configuration of  
337 Pseudotetrahedral Organorhenium Complexes (Eta-C5H5)Re(NO)(PPh3)(X). *Organometallics* **1982**, *1*, 1204-1211.
  - 338 8. Mukherjee, T.; Ghosh, S.K.; Wititsuwannakul, T.; Bhuvanesh, N.; Gladysz, J.A. Chiral-at-Metal Ruthenium Complexes  
339 with Guanidinobenzimidazole and Pentaphenylcyclopentadienyl Ligands: Synthesis, Resolution, and Preliminary  
340 Screening as Enantioselective Second Coordination Sphere Hydrogen Bond Donor Catalysts. *Organometallics* **2020**, *39*,  
341 1163-1175.
  - 342 9. Mamula, O.; von Zelewsky, A. Supramolecular coordination compounds with chiral pyridine and polypyridine ligands  
343 derived from terpenes. *Coord. Chem. Rev.* **2003**, *242*, 87-95.
  - 344 10. Huang, X.; Meggers, E. Asymmetric photocatalysis with bis-cyclometalated rhodium complexes. *Acc. Chem. Res.* **2019**, *52*,  
345 833-847.
  - 346 11. Constable, E.C. Stereogenic metal centres - from Werner to supramolecular chemistry. *Chem. Soc. Rev.* **2013**, *42*, 1637-1651.
  - 347 12. Li, L.P.; Yao, S.Y.; Ou, Y.L.; Wei, L.Q.; Ye, B.H. Diastereoselective Synthesis and Photophysical Properties of  
348 Bis-Cyclometalated Ir(III) Stereoisomers with Dual Stereocenters. *Organometallics* **2017**, *36*, 3257-3265.
  - 349 13. Li, L.P.; Peng, H.L.; Ye, B.H. Thermodynamic Resolution and Enantioselective Synthesis of C-2-Symmetric Bis-sulfoxides  
350 Based on Chiral Iridium(III) Complexes. *Inorg. Chem.* **2019**, *58*, 12245-12253.
  - 351 14. Ehnbon, A.; Ghosh, S.K.; Lewis, K.G.; Gladysz, J.A. Octahedral Werner complexes with substituted ethylenediamine  
352 ligands: a stereochemical primer for a historic series of compounds now emerging as a modern family of catalysts. *Chem.*  
353 *Soc. Rev.* **2016**, *45*, 6799-6811.
  - 354 15. Schaffner-Hamann, C.; von Zelewsky, A.; Barbieri, A.; Barigelletti, F.; Muller, G.; Riehl, J.P.; Neels, A. Diastereoselective  
355 formation of chiral tris-cyclometalated iridium (III) complexes: Characterization and photophysical properties. *J. Am.*  
356 *Chem. Soc.* **2004**, *126*, 9339-9348.
  - 357 16. Coughlin, F.J.; Westrol, M.S.; Oyler, K.D.; Byrne, N.; Kraml, C.; Zysman-Colman, E.; Lowry, M.S.; Bernhard, S. Synthesis,  
358 separation, and circularly polarized luminescence studies of enantiomers of iridium(III) luminophores. *Inorg. Chem.* **2008**,  
359 *47*, 2039-2048.
  - 360 17. Chepelin, O.; Ujma, J.; Wu, X.; Slawin, A.M.Z.; Pitak, M.B.; Coles, S.J.; Michel, J.; Jones, A.C.; Barran, P.E.; Lusby, P.J.  
361 Luminescent, Enantiopure, Phenylatopyridine Iridium-Based Coordination Capsules. *J. Am. Chem. Soc.* **2012**, *134*,  
362 19334-19337.
  - 363 18. Damas, A.; Moussa, J.; Rager, M.N.; Amouri, H. Chiral octahedral bimetallic assemblies with  $\Delta$ -TRISPHAT as counter  
364 anion: design, anion metathesis, and Cp\*Ir as a probe for chiral recognition. *Chirality* **2010**, *22*, 889-895.
  - 365 19. Groue, A.; Montier-Sorkine, E.; Cheng, Y.P.; Rager, M.N.; Jean, M.; Vanthuyne, N.; Crassous, J.; Lopez, A.C.; Moncada,  
366 A.S.; Barbieri, A., *et al.* Enantiopure, luminescent, cyclometalated Ir(III) complexes with N-heterocyclic

- 367 carbene-naphthalimide chromophore: design, vibrational circular dichroism and TD-DFT calculations. *Dalton Trans.* **2022**,  
368 *51*, 2750-2759.
- 369 20. Djukic, J.-P.; Hijazi, A.; Flack, H.D.; Bernardinelli, G. Non-racemic (scalemic) planar-chiral five-membered metallacycles:  
370 routes, means, and pitfalls in their synthesis and characterization. *Chem. Soc. Rev.* **2008**, *37*, 406-425.
- 371 21. Amouri, H.; Thouvenot, R.; Gruselle, M. Differentiation of planar chiral enantiomers of [Cp\*M(2-alkyl-phenoxo)][BF<sub>4</sub>] {M  
372 = Rh, Ir} by the trisphat anion. *C. R. Chim.* **2002**, *5*, 257-262.
- 373 22. Amouri, H.; Caspar, R.; Gruselle, M.; Guyard-Duhayon, C.; Boubekeur, K.; Lev, D.A.; Collins, L.S.B.; Grotjahn, D.B. Chiral  
374 Recognition and Resolution Mediated by p-p Interactions: Synthesis and X-ray Structure of  
375 trans-[(Sp,Sp)-bis(CpRu)-carbazolyl][D-Trisphat]. *Organometallics* **2004**, *23*, 4338-4341.
- 376 23. Dubarle-Offner, J.; Axet, M.R.; Chamoreau, L.M.; Amouri, H.; Cooksy, A.L. Enantiomerically Pure, Planar Chiral Cp\*Ru  
377 Complexes: Synthesis, Molecular Structures, DFT and Coordination Properties. *Organometallics* **2012**, *31*, 4429-4434.
- 378 24. Moussa, J.; Chamoreau, L.M.; Amouri, H. Planar Chiral Iridium Complexes with the -TRISPHAT Anion: Toward the First  
379 Enantiopure o-Quinone Methide pi-Complex. *Chirality* **2013**, *25*, 449-454.
- 380 25. Puig, E.; Gontard, G.; Rager, M.N.; Amouri, H. Optically active Pt-terpyridyl coordination assemblies derived from planar  
381 chiral metathioligands. *Inorg. Chimica. Acta.* **2021**, *517*.
- 382 26. Brunner, H. Stability of the metal configuration in chiral-at-metal half-sandwich compounds. *Eur. J. Inorg. Chem.* **2001**,  
383 *905-912*.
- 384 27. Pettinari, C.; Pettinari, R.; Fianchini, M.; Marchetti, F.; Skelton, B.W.; White, A.H. Syntheses, structures, and reactivity of  
385 new pentamethylcyclopentadienyl-rhodium(III) and -iridium(III) 4-acyl-5-pyrazolonate complexes. *Inorg. Chem.* **2005**, *44*,  
386 *7933-7942*.
- 387 28. Mimassi, L.; Guyard-Duhayon, C.; Rager, M.N.; Amouri, H. Chiral recognition and resolution of the enantiomers of  
388 supramolecular triangular hosts: Synthesis, circular dichroism, NMR, and X-ray molecular structure of [Li subset  
389 of(R,R,R)-{Cp\*Rh(5-chloro-2,3-dioxypyridine)}(3)][Delta-TRISPHAT]. *Inorg. Chem.* **2004**, *43*, 6644-6649.
- 390 29. Feghali, E.; Barloy, L.; Issenhuth, J.T.; Karmazin-Brelot, L.; Bailly, C.; Pfeffer, M. Cyclometalation of  
391 (2R,5R)-2,5-Diphenylpyrrolidine and 2-Phenyl-2imidazoline Ligands with Half-Sandwich Iridium(III) and Rhodium(III)  
392 Complexes. *Organometallics* **2013**, *32*, 6186-6194.
- 393 30. Corberan, R.; Sanau, M.; Peris, E. Highly stable Cp\*-Ir(III) complexes with N-heterocyclic carbene ligands as C-H  
394 activation catalysts for the deuteration of organic molecules. *J. Am. Chem. Soc.* **2006**, *128*, 3974-3979.
- 395 31. Hellou, N.; Jahier-Diallo, C.; Basle, O.; Srebro-Hooper, M.; Toupet, L.; Roisnel, T.; Caytan, E.; Roussel, C.; Vanthuyne, N.;  
396 Autschbach, J., *et al.* Electronic and chiroptical properties of chiral cycloiridiated complexes bearing helicenic NHC ligands.  
397 *Chem. Commun.* **2016**, *52*, 9243-9246.
- 398 32. Groue, A.; Tranchier, J.-P.; Rager, M.-N.; Gontard, G.; Jean, M.; Vanthuyne, N.; Pearce, H.R.; Cooksy, A.L.; Amouri, H.  
399 Unique Class of Enantiopure N-Heterocyclic Carbene Half-Sandwich Iridium(III) Complexes with Stable Configurations:  
400 Probing Five-Membered versus Six-Membered Iridacycles. *Inorg. Chem.* **2019**, *58*, 2930-2933.
- 401 33. Fagnou, K.; Lautens, M. Halide effects in transition metal catalysis. *Angew. Chem. Int. Ed.* **2002**, *41*, 26-47.
- 402 34. Stanley, K.; Baird, M.C. *J. Am. Chem. Soc.* **1975**, *97*, 6598-6599.
- 403 35. Merrifield, J.H.; Fernandez, J.M.; Buhro, W.E.; Gladysz, J.A. Cleavage of the Rhenium Methyl Bond of  
404 (Eta-C<sub>5</sub>H<sub>5</sub>)Re(NO)(PPh<sub>3</sub>)(CH<sub>3</sub>) by Protic and Halogen Electrophiles - Stereochemistry at Rhenium. *Inorg. Chem.* **1984**, *23*,  
405 *4022-4029*.
- 406 36. Brunner, H.; Kollnberger, A.; Burgemeister, T.; Zabel, M. Optically active transition metal complexes - Part 125.  
407 Preparation and epimerization of chiral-at-metal pentamethylcyclopentadienyl-rhodium(III) and iridium(III)  
408 half-sandwich complexes. *Polyhedron* **2000**, *19*, 1519-1526.

- 409 37. Liu, Z.; Habtemariam, A.; Pizarro, A.M.; Fletcher, S.A.; Kisova, A.; Vrana, O.; Salassa, L.; Bruijninx, P.C.A.; Clarkson, G.J.;  
410 Brabec, V., *et al.* Organometallic half-sandwich iridium anticancer complexes. *J. Med. Chem.* **2011**, *54*, 3011-3026.
- 411 38. Liu, Z.; Sadler, P.J. Organoiridium Complexes: Anticancer Agents and Catalysts. *Acc. Chem. Res.* **2014**, *47*, 1174-1185.
- 412 39. Grotjahn, D.B.; Brown, D.B.; Martin, J.K.; Marelius, D.C.; Abadjian, M.-C.; Tran, H.N.; Kalyuzhny, G.; Vecchio, K.S.; Specht,  
413 Z.G.; Cortes-Llamas, S.A., *et al.* Evolution of Iridium-Based Molecular Catalysts during Water Oxidation with Ceric  
414 Ammonium Nitrate. *J. Am. Chem. Soc.* **2011**, *133*, 19024-19027.
- 415 40. Schulze, M.; Steffen, A.; Wurthner, F. Near-IR Phosphorescent Ruthenium(II) and Iridium(III) Perylene Bisimide Metal  
416 Complexes. *Angew. Chem. Int. Ed.* **2015**, *54*, 1570-1573.
- 417 41. Specht, Z.G.; Cortes-Llamas, S.A.; Tran, H.N.; van Niekerk, C.J.; Rancudo, K.T.; Golen, J.A.; Moore, C.E.; Rheingold, A.L.;  
418 Dwyer, T.J.; Grotjahn, D.B. Enabling Bifunctionality and Hemilability of N-Heteroaryl NHC Complexes. *Chem. Eur. J.* **2011**,  
419 *17*, 6606-6609.
- 420 42. Gomez-Lopez, J.L.; Chavez, D.; Parra-Hake, M.; Royappa, A.T.; Rheingold, A.L.; Grotjahn, D.B.; Miranda-Soto, V.  
421 Synthesis and reactivity of bis(protic N-heterocyclic carbene)iridium(III) complexes. *Organometallics* **2016**, *35*, 3148-3153.
- 422 43. Peris, E. Smart N-Heterocyclic Carbene Ligands in Catalysis. *Chem. Rev.* **2018**, *118*, 9988-10031.
- 423 44. Thongpaen, J.; Schmid, T.E.; Toupet, L.; Dorcet, V.; Mauduit, M.; Basle, O. Directed ortho C-H borylation catalyzed using  
424 Cp\*Rh(iii)-NHC complexes. *Chem. Commun.* **2018**, *54*, 8202-8205.
- 425 45. Shibata, T.; Hashimoto, H.; Kinoshita, I.; Yano, S.; Nishioka, T. Unprecedented diastereoselective generation of  
426 chiral-at-metal, half sandwich Ir(III) and Rh(III) complexes via anomeric isomerism on "sugar-coated" N-heterocyclic  
427 carbene ligands. *Dalton Trans.* **2011**, *40*, 4826-4829.
- 428 46. Zhang, W.Y.; Banerjee, S.; Hughes, G.M.; Bridgewater, H.E.; Song, J.I.; Breeze, B.; Clarkson, G.J.; Coverdale, J.P.C.;  
429 Sanchez-Cano, C.; Ponte, F., *et al.* Ligand-centred redox activation of inert organoiridium anticancer catalysts. *Chem. Sci.*  
430 **2020**, *11*, 5466-5480.
- 431 47. Lanoe, P.H.; Chan, J.; Gontard, G.; Monti, F.; Armaroli, N.; Barbieri, A.; Amouri, H. Deep-Red Phosphorescent Iridium(III)  
432 Complexes with Chromophoric N-Heterocyclic Carbene Ligands: Design, Photophysical Properties, and DFT Calculations.  
433 *Eur. J. Inorg. Chem.* **2016**, 1631-1634.
- 434 48. Lanoe, P.H.; Najjari, B.; Hallez, F.; Gontard, G.; Amouri, H. N-Heterocyclic Carbene Coinage Metal Complexes Containing  
435 Naphthalimide Chromophore: Design, Structure, and Photophysical Properties. *Inorganics* **2017**, *5*.
- 436 49. Lanoe, P.H.; Chan, J.; Groue, A.; Gontard, G.; Jutand, A.; Rager, M.N.; Armaroli, N.; Monti, F.; Barbieri, A.; Amouri, H.  
437 Cyclometalated N-heterocyclic carbene iridium(III) complexes with naphthalimide chromophores: a novel class of  
438 phosphorescent heteroleptic compounds. *Dalton Trans.* **2018**, *47*, 3440-3451.
- 439