



HAL
open science

Each Pregnancy Linearly Changes Immune Gene Expression in the Blood of Healthy Women Compared with Breast Cancer Patients

Eiliv Lund, Aurelie Nakamura, Igor Snapkov, Jean-Christophe Thalabard, Karina Standahl Olsen, Lars Holden, Marit Holden

► **To cite this version:**

Eiliv Lund, Aurelie Nakamura, Igor Snapkov, Jean-Christophe Thalabard, Karina Standahl Olsen, et al. Each Pregnancy Linearly Changes Immune Gene Expression in the Blood of Healthy Women Compared with Breast Cancer Patients. *Clinical Epidemiology*, 2018, 10, pp.931–940. 10.2147/CLEP.S163208 . hal-03833313

HAL Id: hal-03833313

<https://hal.sorbonne-universite.fr/hal-03833313v1>

Submitted on 2 Nov 2022

HAL is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers.

L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.

Each pregnancy linearly changes immune gene expression in the blood of healthy women compared with breast cancer patients

Eiliv Lund^{1,2}
 Aurelie Nakamura^{3,4}
 Igor Snapkov¹
 Jean-Christophe Thalabard⁵
 Karina Standahl Olsen¹
 Lars Holden⁶
 Marit Holden⁶

¹Department of Community Medicine, UiT The Arctic University of Norway, Tromsø, Norway; ²The Cancer Registry of Norway, Oslo, Norway; ³Department of Social Epidemiology, Pierre Louis Institute of Epidemiology and Public Health, Sorbonne University, INSERM, Paris, France; ⁴French School of Public Health (EHESP), Doctoral Network, Rennes, France; ⁵MAP5, UMR CNRS 8145, Université Paris Descartes, Sorbonne Paris Cite, Paris, France; ⁶Norwegian Computing Center, Oslo, Norway

Background: There is a large body of evidence demonstrating long-lasting protective effect of each full-term pregnancy (FTP) on the development of breast cancer (BC) later in life, a phenomenon that could be related to both hormonal and immunological changes during pregnancies. In this work, we studied the pregnancy-associated differences in peripheral blood gene expression profiles between healthy women and women diagnosed with BC in a prospective design.

Methods: Using an integrated system epidemiology approach, we modeled BC incidence as a function of parity in the Norwegian Women and Cancer (NOWAC) cohort (165,000 women) and then tested the resulting mathematical model using gene expression profiles in blood in a nested case-control study (460 invasive case-control pairs) of women from the NOWAC postgenome cohort. Lastly, we undertook a gene set enrichment analysis for immunological gene sets.

Results: A linear trend fitted the dataset precisely showing an 8% decrease in risk of BC for each FTP, independent of stratification on other risk factors and lasting for decades after a woman's last FTP. Women with six children demonstrated 48% reduction in the incidence of BC compared to nulliparous. When we looked at gene expression, we found that 756 genes showed linear trends in cancer-free controls (false discovery rate [FDR] 5%), but this was not the case for any of the genes in BC cases. Gene set enrichment analysis of immunologic gene sets (C7 collection in Molecular Signatures Database) revealed 215 significantly enriched human gene sets (FDR 5%).

Conclusion: We found marked differences in gene expression and enrichment profiles of immunologic gene sets between BC cases and healthy controls, suggesting an important protective effect of the immune system on BC risk.

Keywords: breast cancer, Norwegian Women and Cancer Study, gene expression, parity, semi-allograft, hormones, pregnancy

Introduction

Each pregnancy represents a unique and remarkable challenge to the mother's immune system, as her body must accept and carry a fetus to term without immune rejection. Immunological tolerance during pregnancy is necessary for women's bodies to accept the fetus.^{1,2} This obligate tolerance has been observed for many years, and because of it, each fetus can be seen as a different semi-allograft. Interestingly, "immune evasion", which consists of some similar immunological changes, was included as another hallmark of cancer,³ and researchers have recognized some common characteristics between these two highly different biological processes.⁴⁻⁷ Looking at this, it seems paradoxical that increasing number of full-term pregnancies reduces the risk of breast cancer (BC) in women.^{8,9} Some studies have even shown a protective effect up to eight

Correspondence: Eiliv Lund
 Department of Community Medicine,
 UiT The Arctic University of Norway,
 9037 Tromsø, Norway
 Tel +47 77 64 4816
 Email elu000@post.uit.no

or nine children.⁸ However, neither spontaneous nor induced abortions confer the same risk reduction.^{10,11}

The role of hormones is well established in both pregnancy and tumorigenesis; however, the influence of hormonal fluctuations associated with pregnancy on the cancer development in a long-term perspective remains unclear.^{12,13} High, changing levels of hormones such as estrogens, gestagens, and human chorionic hormone initiate and support immunological tolerance during pregnancy,¹⁴ whereas the high levels of estrogens and progestins found in oral contraceptives and hormone replacement therapy are classified as carcinogens.¹⁵

We have previously shown that blood gene expression profiles differ between women with BC and BC-free controls, both at time of diagnosis¹⁶ and during the years prior to diagnosis.¹⁷ Additionally, in a recent study, we compared gene expression in blood and breast tumor tissue.¹⁸ We revealed some similarities in expression profiles mainly for immunogenic tumors; however, we demonstrated that there is no obligate common gene expression pattern.

Here, we aim to explore changes in blood gene expression in women with BC and cancer-free controls according to number of full-term pregnancies (FTPs).

Methods

Norwegian Women and Cancer (NOWAC) study

The NOWAC study¹⁹ is a national population-based study in which a random sample of women living in Norway was recruited starting in 1991. Among the 172,748 women invited to participate, 165,227 women replied to at least one questionnaire and make up the study population. Death and emigration status of these women were updated through linkage to the Cause of Death Registry at Statistics Norway and Central Population Registry, respectively. Information on parity was taken from the NOWAC questionnaires. The date of entry into the present study was set as the date of the first questionnaire answered. The exit date from NOWAC was determined either by the date of cancer diagnosis or end of follow-up (2013/12/31). The NOWAC postgenome cohort consisted of 48,692 women randomly sampled from the whole NOWAC study, born in 1943–1957, with a blood sample buffered to preserve the gene expression profile (PAXgene Blood RNA system; Preanalytix/Qiagen, Hilden, Germany). The NOWAC postgenome biobank contained blood samples from women taken by local physicians and mailed overnight to the study center in Tromsø. The blood was collected in 2003–2006.²⁰ BC cases that were diagnosed in 2003–2009 were identified through linkage to the National Cancer Registry of Norway.

Altogether 546 cases were identified prospectively, ie, developed BC after blood sample donation.

For each BC case, a control from the NOWAC postgenome cohort was assigned, matched by time of blood sampling and year of birth, to be analyzed together with the case. The controls were used to establish the average (mean) gene expression in individuals without cancer and to serve as the basis for exposure-adjusted analyses. The expression level of a gene not involved in the carcinogenic process varied depending on day-to-day changes in exposures such as environment and nutrition, resulting in random fluctuations of the difference in gene expression between cases and matched controls around a population-average constant over time. On the other hand, the difference in the expression of genes related to different stages of the carcinogenic process varied over time in a nonrandom way, thus exhibiting a nonrandom trend. Moreover, the changes in the expression of genes related to the carcinogenesis could be complicated by other effects caused by exposure to carcinogens.

Laboratory procedures

All extraction and microarray services were provided by the Genomics Core Facility, Norwegian University of Science and Technology, Trondheim, Norway. To control for technical variability such as different batches of reagents and kits, day-to-day variations, microarray production batches, and effects related to different laboratory operators, each case–control pair was kept together throughout all extraction, amplification, and hybridization procedures. RNA extraction was performed using the PAXgene Blood mRNA Isolation kit (Preanalytix/Qiagen) according to the manufacturer's instructions. RNA quality and purity were assessed using the NanoDrop ND 8000 Spectrophotometer (Thermo Fisher Scientific, Waltham, MA, USA) and Agilent Bioanalyzer (Agilent Technologies, Santa Clara, CA, USA), respectively. RNA amplification was performed on 96-well plates using 300 ng of total RNA and the Illumina TotalPrep-96 RNA Amplification Kit (Ambio, Inc., Austin, TX, USA). The amplification procedure consisted of reverse transcription with a T7 promoter and ArrayScript, followed by a second-strand synthesis. In vitro transcription with T7 RNA polymerase using a biotin–nucleoside triphosphates mix produced biotinylated cRNA. All case–control pairs were run on either the IlluminaHumanAWG-6 version 3 expression bead chips or the HumanHT-12 version 4 expression bead chips. Outliers were excluded after visual examination of dendrograms, principal component analysis (PCA) plots and density plots. Individuals who were considered borderline outliers were

excluded if their laboratory quality measures were below given thresholds (RNA integrity number value <7 , 260/280 ratio <2 , 260/230 ratio <1.7 , and $50 < \text{RNA} < 500$).

Statistical methods

The first stage of the analysis was to explore the relationship between full-term pregnancies and BC in the whole NOWAC cohort. Based on person-years, the percentage change in incidence rates of BC for each additional child was calculated by using a Poisson linear regression. Each covariate was analyzed separately in a stratified analysis with test for interactions using the Wald test. A Cox proportional semiparametric hazard analysis with attained age as follow-up was used for estimating the overall effect of potential confounders.

Then, gene expression profiles were analyzed for BC cases and controls using microarray technology. The dataset with 30,046 probes for each individual was preprocessed as previously described.²¹ The dataset was background corrected using negative control probes, \log_2 transformed using a variance stabilizing technique,²² and quantile normalized. Data from the two Illumina chips (HumanWG-6 v3 and HumanHT-12 v4) were combined on identical nucleotide universal identifiers.²³ We retained probes present in at least 70% of the individuals. If a gene was represented with more than one probe, the average expression level of the probes was used as the expression for the gene, resulting in a dataset with 8155 genes. The probes were translated to genes using the lumiHumanIDMapping database.²⁴ Finally, the differences in the \log_2 gene expression for each case–control pair were computed and used in the statistical analyses. We then excluded 79 case–control pairs in which the case was diagnosed with in situ BC and seven pairs in which the controls were diagnosed with BC before the end of follow-up in 2014. The final preprocessed dataset included 460 case–control pairs.

A nested case–control design was chosen in order to reduce batch effects in the laboratory. In PCA plots, we observed no batch effects in the data that were obtained after we computed differences in \log_2 gene expression between cases and controls. We used the Bioconductor R-package Limma (linear models for microarrays) to identify the genes that were influenced by parity.²⁵ In the linear model, the responses were the differences in the \log_2 gene expression for each case–control pair, while we included the parity of the control and the parity of the case as covariates. In the analyses, we merged parities 1–3 and 4–6 so that the parity data consisted of three different values: 0, 1–3, and 4–6. The merging was made in order to reduce the effect of the highest parities. The distribution of parities for cases and controls

included in this study is shown in Table S1. We identified gene sets that were influenced by parity using Limma in the same way as we did for individual genes by using enrichment scores for gene sets instead of differences in the \log_2 gene expressions as responses in the linear model. The enrichment scores for gene sets were obtained from the differences in \log_2 gene expressions using the Bioconductor R-package gene set variation analysis (GSVA).²⁶

Ethics approval and consent to participate

The NOWAC study was approved by the Norwegian Data Inspectorate and the Regional Ethical Committee of North Norway (REK). The study was conducted in compliance with the Declaration of Helsinki, and all participants gave written informed consent. The linkages of the NOWAC database to national registries such as the Cancer Registry of Norway and registries on death and emigration were approved by the Directorate of Health. The women were informed about these linkages. Furthermore, the collection and storing of human biological material was approved by the REK in accordance with the Norwegian Biobank Act. Women were informed in the letter of introduction that the blood samples would be used for gene expression analyses.

Results

Reduction of BC risk after full-term pregnancies

Our first step was to estimate the mathematical relationship between parity and BC incidence in the complete NOWAC cohort, which consisted of 165,227 participants and 6536 registered incident BC cases identified through linkage to the Cancer Registry of Norway. Increasing parity was related to less BC in mothers, less use of hormone replacement therapy, larger body mass index (BMI), earlier first birth, more breast feeding, and less use of oral contraceptives (Table 1). Among both premenopausal and postmenopausal women, a regular decrease in incidence rates was found (Figure 1 and Table S2), and in linear regression analyses of BC incidence in the complete NOWAC cohort, each new pregnancy conferred a strong linear reduction in the absolute risk (8.0%; Figure 1). This led to an observed reduction of 48% for a woman with six children.

Differences in BC incidence rates were of the same order between nulliparous and uniparous women and for women with five or six children, illustrating an additive effect. The linearity of BC incidence was equally clear regardless of menopausal status, showing the persistent effect of age,

but postmenopausal women had higher BC incidence rates (Figure 1). Stratification on all risk factors (current use of hormone replacement therapy, BMI, age at first birth, breastfeeding, ever use of oral contraceptives, and maternal history of BC) revealed the same linear patterns (Table S3). There were no significant interactions between different strata of the risk factors (none in the Wald test). Using the Cox's multivariate hazard model adjusted for the same risk factors (taken from Table S3), the crude estimated relative risk (RR) in women with six children versus nulliparous women was 0.50 (95% CI 0.37–0.66) and adjusted RR was 0.49 (95% CI 0.36–0.68), similar to the ones derived from Poisson regression analysis (Table S4).

Table 1 Description of the entire study population according to parity

Characteristics	Parity (number of children)			
	0	1–2	3–4	5–6
Maternal history of BC	5.9	5.2	5.2	4.5
Current use of hormone replacement therapy	17.4	17.1	14.7	12.7
BMI <25 kg/m ²	58.7	58.5	53.7	43.8
Age at first birth <25 years	NA	53.2	70.8	82.2
Years of lactation				
Never	100.0	7.8	4.1	3.1
<2 years	NA	88.3	69.5	47.8
2+	NA	3.8	26.4	49.1
Ever use of oral contraceptives	47.6	59.6	52.7	37.5

Note: Numbers represent a percentage of women within the corresponding parity category.

Abbreviations: BC, breast cancer; BMI, body mass index; NA, data not available.

Differential pregnancy-associated blood gene expression between cases and controls

This linear relationship independent of other covariates was then used to test the hypothesis that blood gene expression of some genes increased or decreased linearly according to parity.

An overall analysis without parity status showed no significant differences in expressed genes between cases and controls. When we added one parity variable for the cases and one for the controls in the model, we found a significant linear relationship between log gene expression and parity for 756 genes in BC-free controls, but we found no such relationship in BC cases (false discovery rate [FDR] 5%). Among the controls, 96 of 756 significant genes had at least 5% change per child (91 were upregulated and five were downregulated) and four genes had at least 10% change per child (all were upregulated). A description of potential carcinogenic effects in BC of the ten top significant genes showed (Table 2) that four of them are tumor promoter genes and one is a tumor suppressor gene; however, the function of five significant genes was unknown.

Gene set enrichment analysis

Table 3 demonstrates the results of gene set enrichment analysis. We identified 588 gene sets from the C7 collection in Molecular Signatures Database (MSigDB; <http://software.broadinstitute.org/gsea/index.jsp>)²⁷ that were significantly enriched when the parity of the controls varied (FDR 5%).




Figure 1 BC incidence rates in the NOWAC study, 1991–2013.

Note: 95% CIs and linear trends according to parity (A) and parity and menopausal status (B; n = 165,227, BC cases = 6,536, PY = 2,390,385).

Abbreviations: BC, breast cancer; CI, confidence interval; NOWAC, Norwegian Women and Cancer; PY, person-years.

Table 2 Top 10 single genes ranked according to *q*-value with logFC and their role in BC

Gene ID	logFC ^a	<i>q</i> -value ^b	Role in BC	Function
<i>CNOT8</i>	0.064	0.013	TP	Involved in translation and transcription regulation
<i>ZRANB2</i>	0.140	0.013	NA	Protein regulating alternative splicing; exact function is unknown
<i>TFRC</i>	0.080	0.013	TP	Required for transfer of iron into cells from transferrin
<i>CD47</i>	0.105	0.013	TP	Involved in many cellular processes including proliferation, migration, and apoptosis
<i>PSORS1C1</i>	-0.039	0.013	NA	Function is not known. Mutations are associated with psoriasis, systemic sclerosis, and rheumatoid arthritis
<i>EML4</i>	0.073	0.013	NA	Protein involved in microtubule formation
<i>FKBP1</i>	-0.027	0.013	TS	A wide range of functions including cell cycle regulation, trafficking, and angiogenesis inhibition
<i>HACL1</i>	0.044	0.013	NA	Exact role is unclear. Catalyzes a carbon-carbon cleavage reaction. Involved in lipid metabolism
<i>PPP1R2</i>	0.093	0.017	NA	Serine/threonine phosphatase. Function is not known
<i>LPINI</i>	0.075	0.017	TP	Protein controlling the metabolism of fatty acids. Transcriptional coactivator

Notes: ^alogFC is the estimated log-fold change in gene expression when the parity increases with one. ^b*q*-value is an FDR-adjusted *p*-value.

Abbreviations: BC, breast cancer; FDR, false discovery rate; NA, data not available; TP, tumor promoter; TS, tumor suppressor.

Table 3 Gene set enrichment analyses for all gene sets included in MSigDB

Collection name	All gene sets			Significant gene sets (FDR 5%)		
	Total	Human	Other	Total	Human	Other
C1: positional gene sets	326	326	0	75	75	0
C2: curated gene sets	4729	3750	979	280	229	51
C3: motif gene sets	836	835	1	22	0	22
C4: computational gene sets	858	857	1	217	217	0
C5: gene ontology gene sets	6166	6162	4	0	0	0
C6: oncogenic signatures	189	144	45	42	35	7
C7: immunologic signatures	4872	1844	3028	588	215	373

Note: In non-human gene sets, mice are the dominant species (<http://software.broadinstitute.org/gsea/msigdb/index.jsp> MSigDB v5.2). Copyright ©2018. Dove Medical Press. Reproduced from Subramanian A, Tamayo P, Mootha VK, et al. Gene set enrichment analysis: a knowledge-based approach for interpreting genome-wide expression profiles. *Proceedings of the National Academy of Sciences*. 2005;102(43):15545–50.27.

Abbreviations: FDR, false discovery rate; MSigDB, Molecular Signatures Database.

Experimentally produced gene sets were submitted to MSigDB from researchers using both in vivo and in vitro materials, as well as both human cells or tissues and animal models. Of our 588 enriched gene sets, 215 were derived from human data and 373 were derived mostly from mouse data.

Detailed characterization of the top 10 gene sets from each species showed a discrepancy in the tissues analyzed: blood and cord blood in humans versus bone marrow and spleen in mice (Tables 4 and S5). We had information on gender for only one of the top 10 human gene sets and four of the mouse gene sets (after contacting main authors of the associated publications). None of the MSigDB gene sets from either humans or mice included information on the number of full-term pregnancies (Table 4).

Discussion

We suggest that our unexpected finding of no changes in gene expression levels from circulating immune-competent cells in BC cases as opposed to parity-dependent changes in healthy women might be linked to the long-term immune-related

protection against BC conferred by semi-allograft experience of each pregnancy. Our findings demonstrating the 8% decrease in BC risk for each birth correspond well to a number of previously published studies.^{8,9}

The systems epidemiology approach we used is based on an integrated analysis in the same cohort; first, hypotheses describing the mathematical relationship between parity and BC incidence in the whole NOWAC study are explored. The findings were that the linear relationship between parity and BC could be described as independent of other risk factors. Then, this hypothesis was tested using gene expression profiles obtained from the postgenome biobank of NOWAC. Consequently, we reduced the number of multiple testing in the gene expression analysis and the potential for false-positive findings.

In a recent analysis of interactions between gene expression in BC tissues and blood samples nested in the NOWAC postgenome study,¹⁸ we found that genes expressed in tumors and used as markers are not necessarily expressed in a similar manner in blood. Not surprisingly, tumors were enriched

Table 4 Top 10 gene sets for humans and mice with information on experimental design

geneSetID	scoreDiff ^a	FDR (q-value)	Source	Immune system	Parity	Cells	Gender
Human							
GSE3982	0.023	0.027	Cord blood	I	NA	Macrophages	NA
GSE2770	-0.026	0.027	Cord blood	A	NA	CD4+ T cells	NA
GSE16385	0.047	0.027	Blood	I	NA	Monocytes	NA
GSE1460	0.034	0.027	Cord blood	A	NA	CD4+ T cells	NA
			Blood				
GSE13411	-0.035	0.027	Spleen	A	NA	B cells	M, F
GSE2770	0.027	0.027	Cord blood	A	NA	CD4+ T cells	NA
GSE29618	-0.029	0.027	Blood	I	NA	DC	NA
GSE17974	0.028	0.027	Cord blood	A	NA	CD4+ T cells	NA
GSE2770	-0.028	0.027	Cord blood	A	NA	CD4+ T cells	NA
GSE29615	0.038	0.027	Blood	I, A	NA	PBMCs	NA
Mouse							
GSE17721	0.018	0.027	Bone marrow	I	NA	DC	F
GSE14769	0.032	0.027	Bone marrow	I	NA	Macrophages	NA
GSE3691	0.037	0.027	Various tissues	I	NA	DC	F
GSE37301	0.035	0.027	Bone marrow	I, A	NA	HSCs, CLPCs	NA
GSE32034	0.028	0.027	Various tissues	I	NA	Monocytes	M
GSE17721	0.026	0.027	Bone marrow	I	NA	DC	F
GSE21063	0.036	0.027	Spleen	A	NA	B cells	NA
GSE11924	-0.029	0.027	Spleen	A	NA	CD4+ T cells	NA
GSE28237	0.035	0.027	Spleen	A	NA	B cells	NA
GSE13547	0.034	0.027	Spleen	A	NA	B cells	NA

Note: ^ascoreDiff is the estimated change in enrichment score when the parity increases with one.

Abbreviations: A, adaptive immune system; CLPCs, common lymphoid progenitor cells; DC, dendritic cells; F, female; FDR, false discovery rate; HSCs, hematopoietic stem cells; I, innate immune system; M, male; NA, data not available; PBMCs, peripheral blood mononuclear cells.

for genes related to cancer hallmarks, while in blood, the genes represented either general cellular or immune-related processes.

Currently, there is no unifying concept explaining mechanism of pregnancy-related BC protection. Yet, four main theories may be underlined: decreased number of mammary stem cells, changes in circulating hormone levels, differentiation of breast epithelial cells, and altered estrogen responsiveness.²⁸ Notably, most of these theories are based on studies focusing on the local changes in the mammary gland, while we used blood samples as the source of data, thus looking at systemic processes.

The single-gene analysis revealed a set of genes linearly changing expression after pregnancies in controls. Almost all of these genes were upregulated. This parity-dependent effect was nonsignificant when we compared all cases and controls without taking parity status into account. A possible explanation could be the low parity in the study population with two children as the median value. Of the top ten significant genes, four upregulated genes were known to be related to breast carcinogenesis.^{29–32} Similar to our results, one study found lower levels of TFRC in plasma of BC cases compared to

controls.³³ However, the overall meaning of these expression changes in healthy women is uncertain.

Previously, Rotunno et al³⁴ demonstrated the significant enrichment of inflammatory and immune response genes in parous women compared to nulliparous. In our gene set enrichment analysis, we observed significant enrichment of numerous gene sets related to the immune system (C7 collection in MSigDB). It is nearly impossible to make an unambiguous conclusion from these results due to the complexity of gene sets' data and lack of essential information in gene sets' description (eg, we were unable to ascertain the gender of blood donors for the majority of human experiments even after contacting the authors of publications). Nevertheless, we observed a clear interspecific difference between components of the immune system: the gene sets were related to (in mice, most of the gene sets were of innate immunity origin, while in humans, they were of adaptive origin) and between the sources of the cells for the experiments. While the latter can be explained by technical and ethical considerations, the former raises yet another concern on the validity of results obtained in animal models.^{35–39}

Pregnancy and cancer share multiple characteristics including proliferation, invasion, neoangiogenesis, and

modulation of local and systemic immune response.⁴ Indeed, in our study, we observed that women with a high parity have a higher expression of tumor promoter genes and a lower expression of tumor suppressor genes; therefore, we propose that the body might experience each pregnancy as a “pseudo cancer”. It is possible to suggest that some pregnancy-related processes may have common genetic signatures and protein expression profile with initial stages of breast tumor formation. Being securely controlled by a variety of local and systemic mechanisms during pregnancy, the immune system does not respond to the changes in breast tissue, while storing memory and effector cells. Later, these cells can initiate rapid and extensive response against cancer cells during the early stages of carcinogenesis. This phenomenon of long-term storage of various subsets of antigen-experienced immune cells after pregnancies has previously been described.^{40,41} Slight expressional differences in each pregnancy expand the pool of antigen-specific immune cells, thus increasing protective effect with the number of pregnancies. Recently, Krause et al⁴² published an experimental study confirming our hypothesis.

Our findings of an impaired immune surveillance in BC patients opposed to healthy women may generate novel approaches to immunotherapy based on the natural mechanisms involved in amplified protection against BC with repeated pregnancies. Deciphering of these mechanisms could significantly improve response rate of BC patients to immunotherapy, which at present remains poor.^{43–45}

The major weakness of the study is the lack of statistical power for the gene expression analyses. However, this was compensated by the strict hypothesis testing approach. The strength of the design is the complete follow-up on cancer, emigration, and mortality based on a unique national birth number. The external validity of the parity data in NOWAC was found to be excellent in a linkage study based on information from the fertility register in Statistics Norway.⁴⁶

Conclusion

To our knowledge, this is the first study revealing the involvement of the immune system in parity-associated BC protection with the use of systems epidemiology approach. Owing to descriptive study design, it is not possible to hypothesize the exact mechanisms responsible for the pregnancy-associated protection from BC using data obtained in this work. Our focus on systemic rather than on local changes may provide the scientific community with a fresh view on the biology of BC and may become a foundation for future research.

Data sharing statement

The datasets are stored at the European Genome-phenome Archive with accession number EGAD00010001062.

Acknowledgments

We are thankful to and impressed by the women who donated blood to the NOWAC postgenome cohort. Bente Augdal, Merete Albertsen, and Knut Hansen were responsible for all infrastructure and administrative issues. The funders had no role in the design of the study; collection, analyses, and interpretation of the data; writing of the manuscript; or decision to submit for publication. This study was supported by a grant from the European Research Council (ERC-AdG 232997 TICE). Some of the data in this article are from the Cancer Registry of Norway. The Cancer Registry of Norway is not responsible for the analysis or interpretation of the data presented. Microarray service was provided by the Genomics Core Facility, Norwegian University of Science and Technology, and NMC – a national technology platform supported by the functional genomics program (FUGE) of the Research Council of Norway.

Author contributions

EL, JT, and MH carried out conception and design of the study. EL, LH, and MH were involved in the development of methodology. EL, LH, and MH contributed toward acquisition of data. EL, AN, IS, JT, KSO, LH, and MH performed analysis and interpretation of data (eg, statistical analysis, biostatistics, and computational analysis). EL, IS, JT, KSO, LH, and MH carried out writing, review, and/or revision of the manuscript. EL, IS, KSO, LH, and MH provided administrative, technical, or material support. EL conducted study supervision. All authors contributed toward data analysis, drafting and critically revising the paper, gave final approval of the version to be published, and agree to be accountable for all aspects of the work.

Disclosure

The authors report no conflicts of interest in this work.

References

1. Erlebacher A. Mechanisms of T cell tolerance towards the allogeneic fetus. *Nat Rev Immunol.* 2012;13(1):23–33.
2. Williams Z. Inducing tolerance to pregnancy. *N Engl J Med.* 2012;367(12):1159–1161.
3. Hanahan D, Weinberg RA. Hallmarks of cancer: the next generation. *Cell.* 2011;144(5):646–674.
4. Holtan SG, Creedon DJ, Haluska P, Markovic SN. Cancer and pregnancy: parallels in growth, invasion, and immune modulation and implications for cancer therapeutic agents. *Mayo Clin Proc.* 2009;84(11):985–1000.

5. Curigliano G. Immunity and autoimmunity: revising the concepts of response to breast cancer. *Breast*. 2011;20:S71–S74.
6. Elinav E, Nowarski R, Thaiss CA, Hu B, Jin C, Flavell RA. Inflammation-induced cancer: crosstalk between tumours, immune cells and microorganisms. *Nat Rev Cancer*. 2013;13(11):759–771.
7. Zhao H, Ozen M, Wong RJ, Stevenson DK. Heme oxygenase-1 in pregnancy and cancer: similarities in cellular invasion, cytoprotection, angiogenesis, and immunomodulation. *Front Pharmacol*. 2014;5:295.
8. Lund E. Childbearing in marriage and mortality from breast cancer in Norway. *Int J Epidemiol*. 1990;19(3):527–531.
9. Collaborative Group on Hormonal Factors in Breast Cancer. Breast cancer and breastfeeding: collaborative reanalysis of individual data from 47 epidemiological studies in 30 countries, including 50302 women with breast cancer and 96973 women without the disease. *Lancet*. 2002;360(9328):187–195.
10. Reeves GK, Kan SW, Key T, et al. Breast cancer risk in relation to abortion: results from the EPIC study. *Int J Cancer*. 2006;119(7):1741–1745.
11. Guo J, Huang Y, Yang L, et al. Association between abortion and breast cancer: an updated systematic review and meta-analysis based on prospective studies. *Cancer Causes Control*. 2015;26(6):811–819.
12. Tal R, Taylor HS, Burney RO, Mooney SB, Giudice LC. *Endocrinology of Pregnancy*. South Dartmouth, MA: MDText.com; 2000.
13. Henderson BE, Ponder BAJ, Ross RK. *Hormones, Genes, and Cancer*. Oxford; New York: Oxford University Press; 2003.
14. Robinson DP, Klein SL. Pregnancy and pregnancy-associated hormones alter immune responses and disease pathogenesis. *Horm Behav*. 2012;62(3):263–271.
15. IARC. Hormonal Contraception and Post-menopausal Hormonal Therapy. IARC Monographs on the Evaluation of Carcinogenic Risks to Humans Volume 72. Lyon: IARC; 1999.
16. Lund E, Holden L, Bøvelstad H, et al. A new statistical method for curve group analysis of longitudinal gene expression data illustrated for breast cancer in the NOWAC postgenome cohort as a proof of principle. *BMC Med Res Methodol*. 2016;16(1):28.
17. Dumeaux V, Ursini-Siegel J, Flatberg A, et al. Peripheral blood cells inform on the presence of breast cancer: a population-based case-control study. *Int J Cancer*. 2015;136(3):656–667.
18. Dumeaux V, Fjukstad B, Fjosne HE, et al. Interactions between the tumor and the blood systemic response of breast cancer patients. *PLoS Comput Biol*. 2017;13(9):e1005680.
19. Lund E, Dumeaux V, Braaten T, et al. Cohort profile: the Norwegian women and cancer study – NOWAC – Kvinner og kreft. *Int J Epidemiol*. 2007;37(1):36–41.
20. Dumeaux V, Børresen-Dale A-L, Frantzen J-O, Kumle M, Kristensen VN, Lund E. Gene expression analyses in breast cancer epidemiology: the Norwegian women and cancer postgenome cohort study. *Breast Cancer Res*. 2008;10(1):R13.
21. Holden M, Holden L. *Statistical Analysis of Gene Expression in Blood Before Diagnosis of Breast Cancer*. NR note, SAMBA/07/16; 2016. Available from: <https://www.nr.no/files/samba/smbi/note2016SAMBA0716BreastCancer.pdf>. Accessed July 5, 2018.
22. Lin SM, Du P, Huber W, Kibbe WA. Model-based variance-stabilizing transformation for Illumina microarray data. *Nucleic Acids Res*. 2008;36(2):e11.
23. Du P, Kibbe WA, Lin SM. nuID: a universal naming schema of oligonucleotides for Illumina, Affymetrix, and other microarrays. *Biol Direct*. 2007;2(1):16.
24. Du P, Feng G, Kibbe W, Lin S. *lumiHumanIDMapping: Illumina Identifier mapping for Human*. 2016. R package version 1.10.1. Available from: <https://bioconductor.org/packages/release/data/annotation/html/lumiHumanIDMapping.html>. Accessed July 5, 2018.
25. Smyth GK. Linear models and empirical Bayes methods for assessing differential expression in microarray experiments. *Stat Appl Genet Mol Biol*. 2004;3(1):1–25.
26. Hänzelmann S, Castelo R, Guinney J. GSEA: gene set variation analysis for microarray and RNA-Seq data. *BMC Bioinformatics*. 2013;14(1):7.
27. Subramanian A, Tamayo P, Mootha VK, et al. Gene set enrichment analysis: a knowledge-based approach for interpreting genome-wide expression profiles. *Proc Natl Acad Sci USA*. 2005;102(43):15545–15550.
28. Britt K, Ashworth A, Smalley M. Pregnancy and the risk of breast cancer. *Endocr Relat Cancer*. 2007;14(4):907–933.
29. Aslam A, Mittal S, Koch F, Andrau JC, Winkler GS. The Ccr4-NOT deadenylase subunits CNOT7 and CNOT8 have overlapping roles and modulate cell proliferation. *Mol Biol Cell*. 2009;20(17):3840–3850.
30. Yang YH, Markus MA, Mangs AH, Raitskin O, Sperling R, Morris BJ. ZRANB2 localizes to supraspliceosomes and influences the alternative splicing of multiple genes in the transcriptome. *Mol Biol Rep*. 2013;40(9):5381–5395.
31. Daniels TR, Delgado T, Rodriguez JA, Helguera G, Penichet ML. The transferrin receptor part I: biology and targeting with cytotoxic antibodies for the treatment of cancer. *Clin Immunol*. 2006;121(2):144–158.
32. Zhang H, Lu H, Xiang L, et al. HIF-1 regulates CD47 expression in breast cancer cells to promote evasion of phagocytosis and maintenance of cancer stem cells. *Proc Natl Acad Sci U S A*. 2015;112(45):E6215–E6223.
33. Buas MF, Rho JH, Chai X, Zhang Y, Lampe PD, Li CI. Candidate early detection protein biomarkers for ER+/PR+ invasive ductal breast carcinoma identified using pre-clinical plasma from the WHI observational study. *Breast Cancer Res Treat*. 2015;153(2):445–454.
34. Rotunno M, Sun X, Figueroa J, et al. Parity-related molecular signatures and breast cancer subtypes by estrogen receptor status. *Breast Cancer Res*. 2014;16(4):R74.
35. Beura LK, Hamilton SE, Bi K, et al. Normalizing the environment recapitulates adult human immune traits in laboratory mice. *Nature*. 2016;532(7600):512–516.
36. Anisimov VN, Ukraintseva SV, Yashin AI. Cancer in rodents: does it tell us about cancer in humans? *Nat Rev Cancer*. 2005;5(10):807–819.
37. Beckers J, Wurst W, de Angelis MH. Towards better mouse models: enhanced genotypes, systemic phenotyping and envirotype modelling. *Nat Rev Genet*. 2009;10(6):371–380.
38. [No authors listed]. Of men, not mice. *Nat Med*. 2013;19(4):379.
39. Mak IW, Evaniew N, Ghert M. Lost in translation: animal models and clinical trials in cancer treatment. *Am J Transl Res*. 2014;6(2):114–118.
40. Rowe JH, Ertelt JM, Xin L, Way SS. Pregnancy imprints regulatory memory that sustains anergy to fetal antigen. *Nature*. 2012;490(7418):102–106.
41. Mor G, Aldo P, Alvero AB. The unique immunological and microbial aspects of pregnancy. *Nat Rev Immunol*. 2017;17(8):469–482.
42. Krause AL, Schuetz F, Boudewijns M, et al. Parity improves anti-tumor immunity in breast cancer patients. *Oncotarget*. 2017;8(62):104981–104991.
43. Vonderheide RH, Domchek SM, Clark AS. Immunotherapy for breast cancer: what are we missing? *Clin Cancer Res*. 2017;23(11):2640–2646.
44. Salatino M, Girotti MR, Rabinovich GA. Glycans pave the way for immunotherapy in triple-negative breast cancer. *Cancer Cell*. 2018;33(2):155–157.
45. Tolba MF, Omar HA. Immunotherapy, an evolving approach for the management of triple negative breast cancer: converting non-responders to responders. *Crit Rev Oncol Hematol*. 2018;122:202–207.
46. Lund E, Kumle M, Braaten T, et al. External validity in a population-based national prospective study – the Norwegian Women and Cancer Study (NOWAC). *Cancer Causes Control*. 2003;14(10):1001–1008.

Supplementary materials

Table S1 Distribution of parities for the cases and controls

	Parities							Sum
	0	1	2	3	4	5	6	
Cases	69	67	208	83	29	4	0	460
Controls	74	62	174	116	26	5	3	460

Table S2 BC cases, PY, and incidence rates per 100,000 PY, according to menopausal status and parity

Status	Parity						
	0	1	2	3	4	5	6
Premenopause							
Cases	168	219	688	358	86	13	2
PY	7,6189	101,736	356,276	212,648	55,512	12,065	2744
Rates	220.50	215.26	193.11	168.35	154.92	107.75	72.88
Postmenopause							
Cases	535	618	2082	1254	372	95	27
PY	132,262	172,437	636,545	420,991	144,901	43,189	13,662
Rates	404.50	358.39	327.08	297.87	256.73	219.97	197.62
Total							
Cases	703	837	2770	1612	458	108	29
PY	20,8451	274,173	992,821	633,639	200,413	55,253	16,407
Rates	337.25	305.28	279.00	254.40	228.53	195.46	176.76

Note: Data from the entire NOWAC cohort.

Abbreviations: BC, breast cancer; NOWAC, Norwegian Women and Cancer; PY, person-years.

Table S3 Tests for linear trend between strata of risk factors

Stratification	% decrease per child	Intercept	β	Pr > t β
Percentage change per child				
Overall	8	334.34	-26.84	<0.001
Stratified according to:				
Premenopause	11	236.4	-24.86	<0.001
Postmenopause	9	398.29	-34.57	<0.001
With mother history of BC	10	631.76	-61.68	0.01
Without mother history of BC	8	317.59	-24.56	<0.001
Age at first birth <25 years	7	305.73	-22.4	0.002
Age at first birth \geq 25 years	6	341.39	-20.93	0.06
BMI <25 kg/m ²	9	337.08	-28.81	<0.001
BMI \geq 25 kg/m ²	8	333.51	-25.09	<0.001
Use of OC	6	323.64	-20.76	<0.001
No use of OC	9	340.42	-30.21	<0.001
No lactation	11	340.34	-35.78	0.022
Years of lactation: 1-2	9	340.61	-30.33	0.002
Years of lactation: 3+	14	488.44	-68.55	0.017
HRT (only postmenopause)	9	530.49	-49.27	0.009
No HRT (only postmenopause)	9	377.31	-32.77	<0.001

Note: Data from the entire NOWAC cohort.

Abbreviations: BC, breast cancer; BMI, body mass index; HRT, hormone replacement therapy; NOWAC, Norwegian Women and Cancer; OC, oral contraceptives; Pr, probability.

Table S4 Age-adjusted and multivariate-adjusted Cox proportional hazard ratios and 95% CIs for BC according to parity in the entire NOWAC cohort

Parity	BC	
	Crude	Adjusted ^a
1	0.91 (0.82–1.00)	0.86 (0.77–0.96)
2	0.81 (0.75–0.88)	0.75 (0.68–0.83)
3	0.73 (0.68–0.80)	0.68 (0.61–0.76)
4	0.63 (0.56–0.71)	0.59 (0.51–0.67)
5	0.52 (0.43–0.64)	0.49 (0.39–0.62)
6+	0.50 (0.37–0.66)	0.49 (0.36–0.68)

Note: ^aAdjustment for maternal history of BC (yes, no), BMI (<25, 25+ kg/m²), current use of hormone replacement therapy (yes, no), years of lactation (0, 1–2, 2+), and use of oral contraceptives (ever used oral contraceptives, never used contraceptives).

Abbreviations: BC, breast cancer; BMI, body mass index; CI, confidence interval; NOWAC, Norwegian Women and Cancer.

Table S5 Full names of gene sets described in Table 4

Human

GSE3982_CTRL_VS_LPS_4H_MAC_UP
 GSE2770_UNTREATED_VS_ACT_CD4_TCELL_48H_DN
 GSE16385_MONOCYTE_VS_12H_ROSIGLITAZONE_TREATED_MACROPHAGE_UP
 GSE1460_CORD_VS_ADULT_BLOOD_NAIVE_CD4_TCELL_DN
 GSE13411_IGM_VS_SWITCHED_MEMORY_BCELL_UP
 GSE2770_UNTREATED_VS_TGFB_AND_IL12_TREATED_ACT_CD4_TCELL_6H_UP
 GSE29618_LAIV_VS_TIV_FLU_VACCINE_DAY7_PDC_DN
 GSE17974_2.5H_VS_72H_IL4_AND_ANTI_IL12_ACT_CD4_TCELL_UP
 GSE2770_TGFB_AND_IL4_VS_TGFB_AND_IL12_TREATED_ACT_CD4_TCELL_2H_UP
 GSE29615_CTRL_VS_DAY3_LAIV_IFLU_VACCINE_PBMC_DN

Mouse

GSE17721_LPS_VS_PAM3CSK4_4H_BMDM_DN
 GSE14769_UNSTIM_VS_240MIN_LPS_BMDM_UP
 GSE3691_IFN_PRODUCING_KILLER_DC_VS_PLASMACYTOID_DC_SPLEEN_DN
 GSE37301_HEMATOPOIETIC_STEM_CELL_VS_COMMON_LYMPHOID_PROGENITOR_DN
 GSE32034_UNTREATED_VS_ROSIGLITAZONE_TREATED_LY6C_HIGH_MONOCYTE_DN
 GSE17721_LPS_VS_PAM3CSK4_1H_BMDM_UP
 GSE21063_WT_VS_NFATC1_KO_BCELL_DN
 GSE11924_TFH_VS_TH17_CD4_TCELL_UP
 GSE28237_EARLY_VS_LATE_GC_BCELL_UP
 GSE13547_CTRL_VS_ANTI_IGM_STIM_ZFX_KO_BCELL_2H_DN

Clinical Epidemiology

Publish your work in this journal

Clinical Epidemiology is an international, peer-reviewed, open access, online journal focusing on disease and drug epidemiology, identification of risk factors and screening procedures to develop optimal preventative initiatives and programs. Specific topics include: diagnosis, prognosis, treatment, screening, prevention, risk factor modification,

Submit your manuscript here: <https://www.dovepress.com/clinical-epidemiology-journal>

systematic reviews, risk and safety of medical interventions, epidemiology and biostatistical methods, and evaluation of guidelines, translational medicine, health policies and economic evaluations. The manuscript management system is completely online and includes a very quick and fair peer-review system, which is all easy to use.

Dovepress