



HAL
open science

A Hamilton-Jacobi Approach to Evolution of Dispersal

King-Yeung Lam, Yuan Lou, Benoît Perthame

► **To cite this version:**

King-Yeung Lam, Yuan Lou, Benoît Perthame. A Hamilton-Jacobi Approach to Evolution of Dispersal. Communications in Partial Differential Equations, In press. hal-03835868

HAL Id: hal-03835868

<https://hal.sorbonne-universite.fr/hal-03835868>

Submitted on 1 Nov 2022

HAL is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers.

L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.

A Hamilton-Jacobi Approach to Evolution of Dispersal

King-Yeung Lam¹, Yuan Lou², and Benoît Perthame³

¹Department of Mathematics, Ohio State University, Columbus, OH, USA,
lam.184@math.ohio-state.edu

²School of Mathematical Sciences, CMA-Shanghai, Shanghai Jiao Tong University,
Shanghai 200240, China, yuanlou@sjtu.edu.cn

³Sorbonne Université, CNRS, Université de Paris, Inria, Laboratoire Jacques-Louis
Lions, F-75005 Paris, France, benoit.perthame@sorbonne-universite.fr

Abstract

The evolution of dispersal is a classical question in evolutionary biology, and it has been studied in a wide range of mathematical models. A selection-mutation model, in which the population is structured by space and a phenotypic trait, with the trait acting directly on the dispersal (diffusion) rate, was formulated by Perthame and Souganidis [Math. Model. Nat. Phenom. 11 (2016), 154–166] to study the evolution of random dispersal towards the evolutionarily stable strategy. For the rare mutation limit, it was shown that the equilibrium population concentrates on a single trait associated to the smallest dispersal rate. In this paper, we consider the corresponding evolution equation and characterize the asymptotic behaviors of the time-dependent solutions in the rare mutation limit, under mild convexity assumptions on the underlying Hamiltonian function.

AMS Subject Class. [2010]: 35B25, 35F21, 35K57, 92D15

Keywords: evolution of dispersal, nonlocal PDE, constrained Hamilton-Jacobi equation, effective fitness, principal Floquet bundle

1 Introduction

Recently, various mathematical models for evolutionary biology have been developed with the theories of adaptive dynamics, competition/selection equations, deterministic or stochastic. Here, we are interested in the specific case of the evolution of dispersal in a bounded domain and explain, in a continuous setting, the selection of the slowest. We introduce the framework in a general setting without space before describing the full model.

1.1 Continuous modeling without space

A convenient modeling background, based on nonlocal Lotka-Volterra parabolic equations, is considered in [2, 27, 33]. It is expressed under the form

$$\begin{cases} \epsilon \partial_t \tilde{n}_\epsilon = \epsilon^2 \Delta_z \tilde{n}_\epsilon + \tilde{n}_\epsilon R(z, \tilde{\rho}_\epsilon(t)) & \text{for } z \in \mathbb{R}^N, t > 0, \\ \tilde{\rho}_\epsilon(t) = \int_{\mathbb{R}^N} \tilde{n}_\epsilon(z, t) dz & \text{for } t > 0, \\ \tilde{n}_\epsilon(z, 0) = \tilde{n}_{\epsilon, in}(z) & \text{for } z \in \mathbb{R}^N. \end{cases} \quad (1.1)$$

The model (1.1) describes the dynamics of a population with density $\tilde{n}_\epsilon(z, t)$, which is structured by a physiological trait $z \in \mathbb{R}^N$. The population dynamics of (1.1) is driven by both mutation and selection. The mutation process is modeled by the Laplacian Δ_z , and the selection is

expressed by the dependence of the (sign-changing) growth rate of an individual, given by $R(z, \tilde{\rho}_\epsilon(t))$, on the trait z and the limiting total population $\tilde{\rho}_\epsilon(t)$. We refer to [9] for a derivation of this type of equations from individual based stochastic models. A striking property of (1.1) is that, as $\epsilon \rightarrow 0$, the generic solution of this equation concentrates as a moving Dirac mass, i.e.

$$\tilde{n}_\epsilon(t) \approx \tilde{\rho}(t)\delta(z - \bar{z}(t)) \quad \text{in distribution,} \quad (1.2)$$

where $\tilde{\rho}(t) = \lim_{\epsilon \rightarrow 0} \tilde{\rho}_\epsilon(t)$ and the trajectory $\bar{z}(t)$ can be interpreted biologically as the dominant trait at the time t . Furthermore, the trajectory $\bar{z}(t)$ can be inferred from the total population $\tilde{\rho}(t)$ via the relation

$$R(\bar{z}(t), \tilde{\rho}(t)) = 0 \quad \text{a.e.} \quad (1.3)$$

and the rare mutation limit of the WKB-ansatz $\tilde{u}_\epsilon(z, t) := -\epsilon \log \tilde{n}_\epsilon(z, t)$ and the total population $\tilde{\rho}(t)$ satisfy, in the viscosity sense, the following Hamilton-Jacobi equation with a nonnegativity constraint.

$$\begin{cases} \partial_t \tilde{u} + |\partial_z \tilde{u}|^2 + R(z, \tilde{\rho}(t)) = 0 & \text{for } z \in \mathbb{R}^N, t > 0, \\ \inf_z \tilde{u}(\cdot, t) = u(\bar{z}(t), t) = 0 & \text{for } t > 0, \\ \tilde{u}(z, 0) = \tilde{u}_{in}(z) & \text{for } z \in \mathbb{R}^N. \end{cases} \quad (1.4)$$

The key to understanding the evolutionary dynamics $\bar{z}(t)$ is thus contained in the question of uniqueness of solution $(\tilde{u}, \tilde{\rho})$ of (1.4), which was investigated in [7, 30, 33].

The model (1.1) enables a rigorous derivation of the so-called *canonical equation*, which has been formally proposed in the framework of adaptive dynamics to describe the trait evolution. See also [11, 22, 27, 33]. In [7], it is further proved that $\tilde{\rho}(t)$ is a nondecreasing function. Since the dominant trait satisfies (1.3), this leads to the conclusion that evolution favors the traits that can retain the proliferative advantage when the level of the total population is high.

1.2 Discrete modeling with spatial structure

In many biological situations, however, the proliferative advantage is not so directly linked to the physiological trait in question. This is the case, in particular, in the study of evolution of dispersal where individuals adopt different dispersal strategies without an apparent proliferative advantage. An interesting question in this direction is the selection of random dispersal rate [19]. For K interacting populations with densities $n_i(x, t)$, $1 \leq i \leq K$, competing for a common resource $m(x)$ and differing only in their dispersal rates α_i , their population dynamics can be described by the following competition system introduced in [15]: for $1 \leq i \leq K$,

$$\partial_t n_i = \alpha_i \Delta_x n_i + n_i \left(m(x) - \sum_{j=1}^K n_j \right) + \sum_{j=1}^K M_{ij} n_j \quad \text{for } x \in D, t > 0. \quad (1.5)$$

In this context, these K species are (i) subject to a mutation with switching rate M_{ij} from phenotype j to i ; (ii) moving randomly with diffusion rates α_i and (iii) competing for a common resource. The proliferative advantage of an individual is influenced by the way the individual moves and utilizes resources. For the two-species case without mutation ($M_{ij} \equiv 0$), a well-known result due to Dockery et al. [15] says that if $m(x)$ is non-constant, and the populations are subject to the no-flux boundary conditions, then the species with the smaller dispersal rate always exclude the faster competitor. The corresponding question for system (1.5) consisting of three or more species remains open; see [8] for recent progress.

The system (1.5) describes the competition among finitely many phenotypes with different dispersal rates, denoted by α_i . In the next section, we will introduce a mathematical model with infinitely many phenotypes, parameterized by a one-dimensional trait variable z . Each z corresponds to a phenotype with dispersal rate $\alpha(z)$, which is a continuous function of z .

1.3 The continuous model for evolution of dispersal

In this paper, we consider a nonlocal reaction-diffusion equation, introduced in [34], which can be viewed as the extension of the discrete trait model (1.5) to the continuous trait setting. The main modeling assumptions are as follows.

- (i) The dispersal rate of an individual is a positive function of its phenotypic trait z , that we denote by $\alpha(z)$;
- (ii) A Fisher-type Lotka-Volterra growth rate with spatially heterogeneous carrying capacity $m(x)$ and limitation by the total population at the same spatial location; and
- (iii) Rare mutation acting on the phenotypic trait variable and modeled by a diffusion with covariance $\sqrt{2}\epsilon$.

More precisely, we study the asymptotic behavior, as $\epsilon \rightarrow 0$, of the density function $n_\epsilon = n_\epsilon(x, z, t) : D \times I \times [0, T] \rightarrow [0, \infty)$, of the nonlocal, nonlinear problem

$$\begin{cases} \epsilon \partial_t n_\epsilon = \alpha(z) \Delta_x n_\epsilon + n_\epsilon (m(x) - \rho_\epsilon(x, t)) + \epsilon^2 \partial_z^2 n_\epsilon & \text{for } x \in D, z \in I, t > 0, \\ \partial_\nu n_\epsilon = 0 & \text{for } x \in \partial D, z \in I, t > 0, \\ \partial_z n_\epsilon = 0 & \text{for } x \in D, z \in \partial I, t > 0, \\ n_\epsilon(x, z, 0) = n_{\epsilon, 0}(x, z) & \text{for } x \in D, z \in I, \end{cases} \quad (1.6)$$

where

$$\rho_\epsilon(x, t) := \int_I n_\epsilon(x, z, t) dz, \quad x \in D, t > 0. \quad (1.7)$$

Here D represents a bounded spatial domain in \mathbb{R}^N with smooth boundary ∂D ; ν denotes the outward unit normal vector on ∂D ; $\Delta_x = \sum_{i=1}^N \partial_{x_i}^2$ is the Laplace operator in x variables; $I = (a, b) \subset \mathbb{R}$ is a bounded open interval with $a < b$; $\rho_\epsilon(x, t)$ is the population density of the entire species at location x and time t . Without loss of generality we assume $|D| = 1$ and $|I| = 1$, i.e., D and I are both of unit Lebesgue measure. The interest of this problem is that the fitness is controlled by a combination between the dispersal $\alpha(z)$ and the competition for resources expressed by $m(x) - \rho_\epsilon(x, t)$. It is important that the resource $m(x)$ satisfies

(M) $m \in C^\beta(\bar{D})$ for some $\beta \in (0, 1)$, and it is a nonconstant, positive function.

On the one hand, if **(M)** does not hold, i.e. $m(x)$ is a constant and $\alpha(z) = z$, then it can be shown that the constant steady state solution is globally asymptotically stable among all nonnegative nontrivial solutions [24]. On the other hand, under the assumption **(M)** and some mild assumptions, it can be shown that, for all ϵ sufficiently small, (1.6) has a positive steady state solution $\tilde{n}_\epsilon(x, z)$ concentrated at the trait where α attains the minimum value [25, 34]. Under suitable assumptions, the positive steady state $\tilde{n}_\epsilon(x, z)$ is shown to be unique and locally asymptotically stable [24]. We will enforce hypothesis **(M)** throughout the rest of the paper.

Our goal is to show that, for the evolution of dispersal, the same concentration effect holds around the fittest trait $\bar{z}(t)$ as expressed in (1.2).

1.4 Assumptions and Main Results

Definition 1.1 (Invasion exponent).

For each $z_1, z_2 \in I$, let $\lambda(z_1, z_2)$ be the smallest eigenvalue of the linear equation

$$\begin{cases} \alpha(z_1) \Delta_x \phi + (m - \theta_{z_2}) \phi + \lambda \phi = 0 & \text{for } x \in D, \\ \partial_\nu \phi = 0 & \text{for } x \in \partial D, \end{cases} \quad (1.8)$$

where, for each $z \in I$, $\theta_z(x)$ is the unique positive solution of

$$\begin{cases} \alpha(z) \Delta_x \theta_z + (m - \theta_z) \theta_z = 0 & \text{for } x \in D, \\ \partial_\nu \theta_z = 0 & \text{for } x \in \partial D. \end{cases} \quad (1.9)$$

Notice that for $z_1 = z_2 =: z$, we have $\lambda(z, z) = 0$, which corresponds to $\phi = \theta_z$. Throughout the paper we impose the following assumptions:

(H1) $\alpha : I \rightarrow \mathbb{R}_+$ is smooth and is chosen such that $0 < \inf_I \alpha \leq \sup_I \alpha < +\infty$ and

$$(\exists K_\lambda > 0) \quad 2K_\lambda \leq \partial_{z_1}^2 \lambda(z_1, z_2) \leq \frac{2}{K_\lambda}, \quad \text{for all } (z_1, z_2) \in I \times I,$$

and

$$\partial_{z_1} \lambda(a, a) < 0 \quad \text{and} \quad \partial_{z_1} \lambda(b, b) > 0,$$

(H2) Let $u_\epsilon(x, z, t) = -\epsilon \log n_\epsilon(x, z, t)$, then the initial condition satisfies

$$\left\| u_\epsilon(x, z, 0) - V_0(z) - \frac{1}{2} \epsilon \log \epsilon \right\|_{C^2(\bar{D} \times \bar{I})} = O(\epsilon),$$

where $V_0(z) \in C^\infty(\bar{I})$ is a non-negative function such that for some $K_0 > 0$,

$$\partial_z^2 V_0(z) > 2K_0 \quad \text{for } z \in I \quad \text{and} \quad \inf_I V_0(z) = V_0(\bar{z}_0) = 0 \quad \exists \bar{z}_0 \in \text{Int } I.$$

Remark 1.2. **(H1)** ensures that the trajectory of the dominant trait remains continuous, and bounded away from ∂I , for $t \geq 0$. See Section 6 for an explicit example of α so that **(H1)** can be explicitly verified¹. We leave the general case for future work.

The following uniqueness result, which seems to be of independent interest, plays a critical role in characterizing the solution trajectories of (1.6). Its proof, which uses the convexity of $\lambda(\cdot, z_2)$, is presented in Appendix C.

Proposition 1.3 (Constrained Hamilton-Jacobi eq.). *Suppose that **(H1)** - **(H2)** hold.*

- (i) *There exists a unique viscosity solution $(V(z, t), \bar{z}(t)) \in W^{1,\infty}(I \times [0, \infty)) \times W^{1,\infty}([0, \infty))$ to the following constrained Hamilton-Jacobi equation:*

$$\begin{cases} \partial_t V + |\partial_z V|^2 - \lambda(z, \bar{z}(t)) = 0 & \text{for } z \in I, t > 0, \\ \partial_z V(z, t) = 0 & \text{for } z \in \partial I, t > 0, \\ V(z, 0) = V_0(z) & \text{for } z \in I, \\ \inf_{z \in I} V(z, t) = 0 & \text{for } t > 0, \end{cases} \quad (1.10)$$

where $V_0(z)$ is given in **(H2)**.

- (ii) $\bar{z} \in C^1([0, \infty))$ and for each $t \geq 0$, $V(z, t) = 0$ if and only if $z = \bar{z}(t)$. Furthermore,

$$V(z, t) = \frac{1}{2} \partial_z^2 V(\bar{z}(t), t) |z - \bar{z}(t)|^2 + o(|z - \bar{z}(t)|^2)$$

and for each $T > 0$ there exists $K_3 > 0$ such that $K_3^{-1} \leq \partial_z^2 V(\bar{z}(t), t) \leq K_3$ for $t \in [0, T]$.

- (iii) $\bar{z}(t)$ satisfies the ODE with coefficient $\sigma(t) := \partial_z^2 V(\bar{z}(t), t)$

$$\frac{d}{dt} \bar{z}(t) = -\frac{\partial_{z_1} \lambda(\bar{z}(t), \bar{z}(t))}{\sigma(t)}, \quad \text{for } t > 0, \quad (1.11)$$

with the initial data $\bar{z}(0) = \bar{z}_0$, where $\bar{z}_0 = \text{argmin } V_0$.

Next, we state our main theorem.

¹When convexity assumption fails, then in general the dominant trait has jump discontinuities and belongs to the class of BV functions for which the uniqueness of the Hamilton-Jacobi equation (1.10) is more subtle; See [7].

Theorem 1.4 (Dynamics of the fittest trait). *Assume (H1) - (H2). For each $T > 0$, as $\epsilon \rightarrow 0$,*

$$n_\epsilon(x, z, t) \rightarrow \delta_0(z - \bar{z}(t))\theta_{\bar{z}(t)}(x) \quad \text{in distribution sense in } \bar{D} \times \bar{I} \times (0, T], \quad (1.12)$$

where $\bar{z}(t) \rightarrow \operatorname{argmin} \alpha(z)$ as $t \rightarrow \infty$. In fact, we have

$$-\epsilon \log n_\epsilon(x, z, t) \rightarrow V(z, t) \quad \text{in } C(\bar{D} \times \bar{I} \times [0, T]), \quad (1.13)$$

and

$$\rho_\epsilon(x, t) \rightarrow \theta_{\bar{z}(t)}(x) \quad \text{in } C_{loc}(\bar{D} \times (0, T]), \quad (1.14)$$

where $V(z, t)$ and $\bar{z}(t)$ are given by Proposition 1.3; and, for $z \in I$, $\theta_z(x)$ is the unique positive solution of (1.9).

Remark 1.5. One can also replace the hypothesis (H1) - (H2) by

(H1') Suppose $V_0(z)$ and $\alpha(z)$ are chosen such that (1.10) has a unique viscosity solution $(V(z, t), \bar{z}(t)) \in W^{1,\infty}(D \times (0, \infty)) \times W^{1,\infty}(0, \infty)$ such that the conclusions (ii) and (iii) of Proposition 1.3 hold.

For instance, our conclusion holds for the case $\alpha(z) > 0$ being periodic in z , and the Neumann condition being replaced by a periodic condition in z , provided that α has a non-degenerate minimum attained at \hat{z} , and such that $V_0(z) = K(z - \bar{z}_0)^2$, provided $|\hat{z} - \bar{z}_0| \ll 1$ and $K \gg 1$.

1.5 Biological interpretation

The Darwinian evolution of a quantitative trait is the combined effects of two biological processes: (i) mutations generating variations in the trait value; and (ii) selection via relative reproductive fitness, resulting from ecological interactions between individuals and their environment. The framework of adaptive dynamics [13, 17] is based on the assumption of separation of timescales between the mutation/evolutionary and selection/ecological processes. One important advance of this theory, due to Dieckmann and Law [12], is the formal derivation of the so-called canonical equation of adaptive dynamics: An ordinary differential equation that gives the rate of change over time of the expected trait value in a monomorphic population. In previous works [14, 27], the canonical equation was rigorously derived in case the ecological interaction can be described by ODEs, i.e., the fitness function is explicitly given in terms of the trait z . This is not the case, however, for evolution of dispersal, for which the incorporation of spatial structure in the model is essential and causes considerable mathematical difficulties. See [6, 23, 29]. Also, using a continuous trait explains the ‘‘accelerating waves’’ which have been actively studied recently, [3, 4, 5, 20].

In this paper, we define an effective Hamiltonian $H_\epsilon(z, t)$ by improving the existing theory of the principal Floquet bundle for parabolic problems. This Hamiltonian can be viewed as a fitness function of the trait z interacting with the environment at time t (as described by $m(x) - \rho_\epsilon(x, t)$). One achievement of this paper is to show rigorously that, in a suitable timescale, the fitness function $H_\epsilon(z, t)$ converges to $\lambda(z, \bar{z}(t))$ in the rare mutation limit, where $\bar{z}(t)$ is the fittest trait at time t and, in this environment set by the fittest trait, the function $\lambda(z, \bar{z}(t))$ is the relative fitness of other trait z and is defined implicitly by an elliptic linear eigenvalue problem arising from the pairwise ecological interaction between two traits. The canonical equation (1.11) can then be rigorously derived. It is interesting to note that the mean trait $\bar{z}(t)$ and the variance $\partial_{zz}V(z, t)|_{z=\bar{z}(t)}$ does not satisfy a closed system of ODEs.

While our previous works [24, 25, 34] characterized the unique evolutionarily stable strategy (ESS) by solving the steady state problem, our present work on the time-dependent problem describes the approach of the dominant trait to the ESS in the evolutionary timescale.

1.6 A heuristic presentation of the analytical approach

The WKB-ansatz defines u_ϵ by

$$n_\epsilon(x, z, t) = e^{-\frac{u_\epsilon(x, z, t)}{\epsilon}},$$

where the rate function $u_\epsilon(x, z, t)$ satisfies the equation

$$\partial_t u_\epsilon - \frac{\alpha(z)}{\epsilon} \Delta_x u_\epsilon + \frac{\alpha(z)}{\epsilon^2} |\nabla_x u_\epsilon|^2 - \epsilon \partial_z^2 u_\epsilon + |\partial_z u_\epsilon|^2 + m(x) - \rho_\epsilon(x, t) = 0. \quad (1.15)$$

Whereas the large coefficients of the spatial derivatives suggests that $u(z, t) = \lim_{\epsilon \rightarrow 0} u_\epsilon(x, z, t)$ is constant in x in the limit, the equation (1.15) itself depends non-trivially on the spatial variables, through the terms $m(x) - \rho_\epsilon(x, t)$.

To obtain the limiting Hamilton-Jacobi equation (which is supposed to be free of the x -dependence), the perturbed test function method has been invented in [16] by considering

$$n_\epsilon(x, z, t) = \Phi_\epsilon(x, z, t) e^{-\frac{v_\epsilon(x, z, t)}{\epsilon}},$$

i.e. $v_\epsilon(x, z, t) = u_\epsilon + \epsilon \log \Phi_\epsilon(x, z, t)$. To eliminate the x -dependence in (1.15), we construct the corrector² $\Phi_\epsilon(x, z, t) > 0$ as the *normalized principal Floquet bundle* that satisfies, for each z ,

$$\epsilon \partial_t \Phi_\epsilon - \alpha(z) \Delta_x \Phi_\epsilon = (m(x) - \rho_\epsilon(x, t) + H_\epsilon(z, t)) \Phi_\epsilon, \quad \int_D \Phi_\epsilon(x, z, t) dx = 1. \quad (1.16)$$

We can build (see Appendix B) the normalizing factor $t \mapsto H_\epsilon(z, t)$ so that $\Phi_\epsilon(x, z, t)$ satisfies the mass 1 constraint (and is bounded in $C(\bar{D} \times [0, \infty))$ thanks to the Harnack inequality). Denoting $\varphi_\epsilon := -\log \Phi_\epsilon$, the equation of v_ϵ can be written as

$$\begin{aligned} \partial_t v_\epsilon - \frac{\alpha(z)}{\epsilon} \Delta_x v_\epsilon + \frac{\alpha(z)}{\epsilon^2} |\nabla_x v_\epsilon|^2 + 2 \frac{\alpha(z)}{\epsilon} \nabla_x v_\epsilon \nabla_x \varphi_\epsilon - \epsilon \partial_z^2 v_\epsilon + |\partial_z v_\epsilon|^2 + 2\epsilon \partial_z v_\epsilon \partial_z \varphi_\epsilon \\ = H_\epsilon(z, t) - \epsilon^2 (\partial_z^2 \varphi_\epsilon + |\partial_z \varphi_\epsilon|^2) \quad \text{for } x \in D, z \in I, t > 0. \end{aligned} \quad (1.17)$$

Provided that $\varphi_\epsilon \in L^\infty$ and it has bounded derivatives in the trait variable³, we can show by comparison that $v_\epsilon(x, z, t) \approx \tilde{V}_\epsilon(z, t)$, where the approximate solution $\tilde{V}_\epsilon(z, t)$ can be obtained by solving the following Hamilton-Jacobi equation which is free of x -dependence:

$$\partial_t \tilde{V}_\epsilon + |\partial_z \tilde{V}_\epsilon|^2 = H_\epsilon(z, t). \quad (1.18)$$

Therefore, we can approximate for a given time t the dominant trait, which maximizes $z \mapsto n_\epsilon(x, z, t)$, by the value $\bar{z}_\epsilon(t)$ where \tilde{V}_ϵ attains its minimum. In fact, we can determine this approximate trajectory $\bar{z}_\epsilon(t)$ from $H_\epsilon(z, t)$ as well. To this end, use the uniqueness of convex solutions $(V_\epsilon(z, t), \bar{z}_\epsilon(t))$ in the class $W^{1, \infty}(I \times (0, T)) \times W^{1, \infty}(0, T)$ of the following Hamilton-Jacobi equation with a constraint, due to Mirrahimi and Roquejoffre [30] (for the uniqueness in $W^{1, \infty}(I \times (0, T)) \times BV(0, T)$, see also [7] and Appendix C):

$$\begin{cases} \partial_t V_\epsilon + |\partial_z V_\epsilon|^2 = H_\epsilon(z, t) - H_\epsilon(\bar{z}_\epsilon(t), t) & \text{for } z \in I, t \in [0, T], \\ \inf_z V_\epsilon(\cdot, t) = 0 & \text{for } t \in [0, T], \\ V_\epsilon(z, 0) = V_0(z) & \text{for } z \in I. \end{cases} \quad (1.19)$$

²The elliptic eigenfunction and eigenvalues (i.e. by omitting the term $\epsilon \partial_t \Phi_\epsilon$ in (1.16)) are not adopted here. The reason is that the lack of regularity in time for ρ_ϵ will then render it quite difficult to estimate $\epsilon \partial_t \Phi_\epsilon$, which only goes to zero in some weak, average sense.

³We need to show the terms $-\epsilon^2 (\partial_z^2 \varphi_\epsilon + |\partial_z \varphi_\epsilon|^2) \rightarrow 0$ uniformly. In general, the Cauchy problem (1.16) does not guarantee smoothness with respect to z , due to the dependence on initial data and the timescale ϵt . We will therefore make a canonical choice of an appropriate eternal solution of (1.16). This requires *a priori* Hölder estimates of $\rho_\epsilon(x, t)$, as well as an improvement upon existing theory of principal Floquet bundle (Appendix B).

Hence, we have defined the approximate trajectory $\bar{z}_\epsilon(t)$ and effective Hamiltonian $H_\epsilon(z, t)$ in terms of the quantity $\rho_\epsilon(x, t)$.⁴ Now, having proved that the population has dominant trait at $\bar{z}_\epsilon(t)$, we can further determine that as $\epsilon \rightarrow 0$,

$$|\rho_\epsilon(x, t) - \theta_{\bar{z}_\epsilon(t)}(x)| \rightarrow 0. \quad (1.20)$$

Since $H_\epsilon(z, t)$ depends smoothly in terms of $\rho_\epsilon(x, t)$, (1.20) and the separation of timescales imply that the effective Hamiltonian $H_\epsilon(z, t)$ converges to the invasion exponent in (1.8), i.e.,

$$|H_\epsilon(z, t) - \lambda(z, \bar{z}_\epsilon(t))| \rightarrow 0.$$

Since $\lambda(z_1, z_2) = 0$ when $z_1 = z_2$, we have in fact

$$H_\epsilon(z, t) - H_\epsilon(\bar{z}(t), t) \approx \lambda(z, \bar{z}(t)) - \lambda(\bar{z}(t), \bar{z}(t)) = \lambda(z, \bar{z}(t)). \quad (1.21)$$

This, in turn, allows us to pass to the limit in the constrained Hamilton-Jacobi equation (1.19), resulting in (1.10) in Proposition 1.3.

However, for the uniqueness argument to work on (1.19), which is crucial in the definition of approximate trajectory $\bar{z}_\epsilon(t)$, we need

$$\partial_z^2 H_\epsilon(\bar{z}(t), t) > 0.$$

This is indeed the case in a small time interval $[0, \delta_1]$, as one can show, by the separation of time-scale between the evolutionary dynamics in z and ecological dynamics of reaching the spatial equilibrium $\theta_{\bar{z}_0}$, that $\rho_\epsilon(x, t)$ does not deviate too much from $\theta_{\bar{z}_0}(x)$ in the time interval $[0, \delta_1]$. This implies the effective Hamiltonian $H_\epsilon(x, t)$ is in the proximity of $\lambda(z, \bar{z}_0)$, which has the relevant convexity in z . This is done in Corollary 4.6. In this way, the argument above is valid and delivers that

$$H_\epsilon(z, t) \rightarrow \lambda(z, \bar{z}(t)) \quad \text{and} \quad \rho_\epsilon(x, t) \rightarrow \theta_{\bar{z}(t)}(x) \quad \text{for } 0 \leq t \leq \delta_1.$$

By carefully examining the smooth dependence of the effective Hamiltonian H_ϵ and corrector Φ_ϵ on z and ρ_ϵ , we establish a uniform lower bound for the time-step δ_1 for which the above argument can be applied. Iterating step by step in time, we can prove the convergence over the time interval $[0, T]$ for all $T > 0$.

A different mutation-selection model involving a spatial variable is also studied in [23]. In the setting of that work, the rate function u_ϵ can be shown to be uniformly convex in z *a priori*. This is not the case in our setting. Also, the quasi-steady state approximation is used in that paper, i.e., the elliptic eigenvalue problem instead of the parabolic principal Floquet bundle problem (1.16) in choosing the corrector Φ_ϵ . In our setting ρ_ϵ does not have enough *a priori* regularity in time, so our definition of corrector via the parabolic problem affords the needed additional time regularity.

A similar kind of result has also been obtained for the corresponding model with age structure [32], but with a different strategy. Therein the corrector Φ_ϵ was defined by a couple of nonlinear mappings and the main analysis was devoted to showing the uniform boundedness of the corrector Φ_ϵ . In contrast, here we adopt a relatively direct approach by defining the corrector Φ_ϵ directly to be the (bounded) normalized principal Floquet bundle for parabolic problem (1.16). This is made possible thanks to our new *a priori* estimates on the quantity $\rho_\epsilon(x, t)$.

⁴At this point $u_\epsilon(x, z, t) \approx v_\epsilon(x, z, t) \approx V_\epsilon(z, t) + \int_0^t H_\epsilon(\bar{z}_\epsilon(s), s) ds$. However, by the uniform positive upper and lower bounds of $\|\rho_\epsilon(\cdot, t)\|_{L^1(D)}$, we deduce that $\int_0^t H_\epsilon(\bar{z}_\epsilon(s), s) ds \rightarrow 0$ uniformly. Hence, we have $v_\epsilon(x, z, t) \approx V_\epsilon(z, t)$.

1.7 Organization of the Paper

In Section 2, we establish some *a priori* estimates of the solution $n_\epsilon(x, z, t)$ and its integrated version $\rho_\epsilon(x, t)$. We first state the global positive upper and lower bounds, for which the proofs are postponed to Appendix A, and derive the Hölder regularity of ρ_ϵ and $\partial_x^2 n_\epsilon$ in some appropriately rescaled variables.

In Section 3, we apply the theory of normalized principal Floquet bundle for parabolic problems with Neumann boundary conditions, as developed in [8] and summarized in Appendix B, to define the effective Hamiltonian $H_\epsilon(z, t)$ and the corrector $\Phi_\epsilon(x, z, t)$, as functions of $\rho_\epsilon(x, t)$. We also state the uniqueness theorem, which allows us to define the approximate trajectory $\bar{z}_\epsilon(t)$ and phenotypic distribution $V_\epsilon(z, t)$ in terms of the effective Hamiltonian $H_\epsilon(z, t)$ and initial distribution $V_0(z)$.

In Section 4, we introduce the rate function $u_\epsilon(x, z, t) = -\epsilon \log n_\epsilon(x, z, t)$ and prove several technical estimates that enable us to obtain a lower bound $\delta_1 > 0$ of the step size in time with which we can continue the approximate trajectory.

In Section 5, we prove the main result Theorem 1.4.

The three appendices are devoted to the three main ingredients/tools that we develop for this singularly perturbed problem. In Appendix A, we prove the *a priori* L^∞ estimate of $\rho_\epsilon(x, t)$. In Appendix B, we state the existence and differentiability of the normalized principal Floquet bundle, which is used in the construction of effective Hamiltonian and corrector ($H_\epsilon(z, t), \Phi_\epsilon(x, z, t)$). In Appendix C, we prove the existence and uniqueness of solutions to the constrained Hamilton-Jacobi equations that we need.

2 A priori Estimates

We begin with the following result; see Proposition A.1 for the proof:

Proposition 2.1. *Let $\rho_\epsilon(x, t) = \int_I n_\epsilon(x, z, t) dz$, where n_ϵ is the solution to (1.6). Then there exists \widehat{C}_1 independent of $\epsilon > 0$ such that*

$$\frac{1}{\widehat{C}_1} \leq \rho_\epsilon(x, t) \leq \widehat{C}_1 \quad \text{for all } (x, t) \in D \times [0, \infty).$$

In the following, we extend $n_\epsilon(x, z, t)$ evenly and then periodically in z . Due to the Neumann boundary condition, the extended n_ϵ satisfies the same PDE with coefficients similarly extended to $D \times \mathbb{R} \times [0, \infty)$. Next, we rescale n_ϵ . Define, for $z_1 \in I$ and $t_1 \geq \epsilon$,

$$N_\epsilon(x, y, \tau) = N_\epsilon(x, y, \tau; z_1, t_1) := n_\epsilon(x, z_1 + \epsilon y, t_1 + \epsilon \tau),$$

and note that $N_\epsilon(x, y, \tau)$ satisfies a linear parabolic equation

$$\partial_\tau N_\epsilon - \alpha(z_1 + \epsilon y) \Delta_x N_\epsilon - \partial_y^2 N_\epsilon = N_\epsilon(m(x) - \rho_\epsilon(x, t_1 + \epsilon \tau)). \quad (2.1)$$

By Proposition 2.1, the above equation has L^∞ bounded coefficients, so we may apply local parabolic L^p estimates to obtain Hölder regularity of N_ϵ , which then allows us to use parabolic Schauder estimates to estimate $\partial_x^2 N$. Here $\partial_x^2 N$ denotes all second order partial derivatives in space $\partial_{x_i x_j}^2 N$.

Lemma 2.2. *For $\beta \in (0, 1)$, there exists $C > 0$ independent of ϵ such that*

$$\|\partial_x^2 N_\epsilon(x, y, \tau)\|_{C^{\beta, \beta, \beta/2}(\bar{D} \times [-\frac{1}{2}, \frac{1}{2}] \times [-\frac{1}{2}, 0])} \leq C \|N_\epsilon\|_{L^1(D \times (-1, 1) \times (-1, 0))}.$$

In particular, there exists a constant C independent of $t \geq \epsilon$ and $z \in I$ such that

$$\sup_{x \in D} |\partial_x^2 n_\epsilon(x, z, t)| \leq C \int_{t-\epsilon}^t \int_{-\epsilon}^\epsilon \int_D n_\epsilon(x', z + z', t') dx' dz' dt'. \quad (2.2)$$

Here we use the notation $\int_{s_1}^{s_2} = \frac{1}{s_2 - s_1} \int_{s_1}^{s_2}$, for any $s_1 < s_2$.

Proof. By Proposition 2.1, $\sup_{t \geq 0} \|\rho_\epsilon(\cdot, t)\|_{C(\bar{D})} \leq C$. Hence, the equation (2.1) has L^∞ bounded coefficients. So we may apply local parabolic L^p estimates to obtain Hölder regularity of N_ϵ . To this end, define $\Omega_{3/5} \subset \Omega_{4/5} \subset \Omega_1$ by

$$\Omega_R := D \times (-R, R) \times (-R, 0), \quad \text{for } R = \frac{3}{5}, \frac{4}{5} \text{ and } 1.$$

Then, recalling (2.1), we have

$$\|N_\epsilon\|_{C^{\beta, \beta, \beta/2}(\overline{\Omega_{3/5}})} \leq C \|N_\epsilon\|_{W_p^{2,2,1}(\Omega_{3/5})} \leq C \|N_\epsilon\|_{L^p(\Omega_{4/5})} \leq C \|N_\epsilon\|_{L^1(\Omega_1)}, \quad (2.3)$$

where the first, second and third inequality follows, respectively, from the Sobolev embedding, parabolic L^p estimate, and (A.5).

Integrating (2.3) over $z_1 \in I$, we have, for each $t_1 \geq \epsilon$,

$$\begin{aligned} \|\rho_\epsilon(\cdot, t_1 + \epsilon \cdot)\|_{W_p^{2,1}(D \times [-\frac{3}{5}, 0])} &\leq \int_I \|N_\epsilon(\cdot, 0, \cdot; t_1, z_1)\|_{W_p^{2,1}(D \times [-\frac{3}{5}, 0])} dz_1 \\ &\leq \int_I \|N_\epsilon(\cdot, 0, \cdot; t_1, z_1)\|_{L^1(D \times [-1, 0])} dz_1 \\ &\leq C \|\rho_\epsilon(\cdot, t_1 + \epsilon \cdot)\|_{L^1(D \times [-1, 0])} \leq C \end{aligned} \quad (2.4)$$

where we used the fact that $\rho_\epsilon(x, t_1 + \epsilon t) = \int_I N_\epsilon(x, 0, t; t_1, z_1) dz_1$ for the first and third inequalities, (2.3) in the second inequality, and Proposition A.1 for the last inequality. Similarly, for each $t_1 \geq \epsilon$ we have

$$\|\rho_\epsilon(\cdot, t_1 + \epsilon \cdot)\|_{C^{\beta, \beta/2}(\bar{D} \times [-\frac{3}{5}, 0])} \leq \int_I \|N_\epsilon(\cdot, 0, \cdot; t_1, z_1)\|_{C^{\beta, \beta/2}(\bar{D} \times [-\frac{3}{5}, 0])} dz_1 \leq C.$$

Now that $\rho_\epsilon(\cdot, t_1 + \epsilon \cdot)$ is Hölder continuous, we may apply parabolic Schauder estimates to (2.1) to get

$$\|\partial_x^2 N_\epsilon(x, y, \tau)\|_{C^{\beta, \beta, \beta/2}(\bar{D} \times [-\frac{1}{2}, -\frac{1}{2}] \times [-\frac{1}{2}, 0])} \leq C \|N_\epsilon\|_{C^{\beta, \beta, \beta/2}(\overline{\Omega_{3/5}})}. \quad (2.5)$$

The lemma follows from combining (2.3) and (2.5). \square

The following result follows from the proof of Lemma 2.2:

Corollary 2.3. (a) *For each $p > 1$, there exists some $\widehat{C}_p > 0$ independent of ϵ such that*

$$\sup_{\tau_0 \geq 1} \|\widetilde{\rho}_\epsilon(\cdot, \cdot)\|_{W_p^{2,1}(D \times [\tau_0, \tau_0 + 1])} \leq \widehat{C}_p, \quad \text{where } \widetilde{\rho}_\epsilon(x, \tau) := \rho_\epsilon(x, \epsilon \tau).$$

(b) *For each $\beta' \in (0, 1)$, there exists some $C_{\beta'} > 0$ independent of ϵ such that*

$$\|\widetilde{\rho}_\epsilon\|_{C^{\beta', \beta'/2}(\bar{D} \times [1, \infty))} \leq C_{\beta'}.$$

In particular, we have $\sup_{t \geq \epsilon} \|\rho_\epsilon(\cdot, t)\|_{C^{\beta'}(\bar{D})} \leq C_{\beta'}$.

This result requires an initial delay of order ϵ so as to take into account the possible initial layer on ρ_ϵ . This is responsible for the technical issues on the initial data that we encounter in the next section.

3 Approximate Trajectory via the Normalized Principal Floquet Bundle

Our next and fundamental task is to define the corrector φ_ϵ , closely related to $\log \Phi_\epsilon$ and effective Hamiltonian H_ϵ in terms of the principal Floquet bundle of certain parabolic problem with potential $m - \rho_\epsilon$. (Recall that ρ_ϵ is given by (1.7) in terms of the solution n_ϵ of (1.6).) This, in turn, enables us to define an approximate trajectory $\bar{z}_\epsilon(t)$ of the dominant trait.

Proposition 3.1. *For each fixed $\epsilon > 0$ and $z \in I$, there exists a unique classical solution $(\varphi_\epsilon(x, z, t), H_\epsilon(z, t))$ to the following linear parabolic problem in $D \times (-\infty, \infty)$:*

$$\begin{cases} \epsilon \partial_t \varphi_\epsilon - \alpha(z) \Delta_x \varphi_\epsilon + \alpha(z) |\nabla_x \varphi_\epsilon|^2 + m(x) - \rho_\epsilon(x, \max\{t, \epsilon\}) \\ \quad + H_\epsilon(t; z) = 0 & x \in D, t \in \mathbb{R}, \\ \partial_\nu \varphi_\epsilon = 0 & x \in \partial D, t \in \mathbb{R}, \\ \int_D e^{-\varphi_\epsilon(x, t; z)} dx = 1 & t \in \mathbb{R}. \end{cases} \quad (3.1)$$

Moreover, the quantities $(\varphi_\epsilon(x, t; z), H_\epsilon(z, t))$ depend smoothly on $z \in I$, i.e., for some constant \tilde{C}_0 independent of ϵ ,

$$\max_{i=0,1,2,3} \|\partial_z^i H_\epsilon(t; z)\|_\infty + \|\nabla_x \varphi_\epsilon(x, t; z)\|_\infty + \max_{i=0,1,2} \|\partial_z^i \varphi_\epsilon(x, t; z)\|_\infty \leq \tilde{C}_0, \quad (3.2)$$

where $\|\cdot\|_\infty$ denotes the L^∞ norm over $(x, z, t) \in D \times I \times \mathbb{R}$.

Proof. For $x \in D$ and $\tau \in \mathbb{R}$, define $c_\epsilon(x, \tau) := m(x) - \tilde{\rho}_\epsilon(x, \max\{\tau, 1\})$, where $\tilde{\rho}_\epsilon(x, \tau) = \rho_\epsilon(x, \epsilon\tau)$ as in Corollary 2.3, then $\|c_\epsilon\|_{C^{\beta, \beta/2}(\bar{D} \times \mathbb{R})}$ is uniformly bounded in ϵ thanks to Corollary 2.3(b). By Theorem B.2, we can define the corresponding normalized principal Floquet bundle

$$(\Phi_1(x, \tau; c_\epsilon, z), H_1(\tau; c_\epsilon, z)) \in C^{2+\beta, 1+\beta/2}(\bar{D} \times \mathbb{R}) \times C^{\beta/2}(\mathbb{R}),$$

which satisfies

$$\begin{cases} \partial_\tau \Phi_1 - \alpha(z) \Delta_x \Phi_1 = (m(x) - \rho_\epsilon(x, \epsilon \max\{\tau, 1\})) \Phi_1 + H_1(\tau; c_\epsilon, z) \Phi_1 & \text{for } x \in D, t \in \mathbb{R}, \\ \partial_\nu \Phi_1 = 0 & \text{for } x \in \partial D, t \in \mathbb{R}, \\ \Phi_1 > 0 \quad \text{for } x \in D, t \in \mathbb{R}, \quad \text{and} \quad \int_D \Phi_1 dx = 1 & \text{for } t \in \mathbb{R}. \end{cases}$$

Setting

$$\varphi_\epsilon(x, z, t) := -\log \Phi_1(x, t/\epsilon; c_\epsilon, z) \quad \text{and} \quad H_\epsilon(z, t) := H_1(t/\epsilon; c_\epsilon, z),$$

we obtain $(\varphi_\epsilon, H_\epsilon)$ satisfying (3.1). The smoothness follows from Proposition B.3. \square

We define the approximate trajectory by solving a constrained Hamilton-Jacobi equation by making use of uniqueness results under convexity assumption [30]. The proof is contained in Appendix C.

Proposition 3.2. *Suppose, for some $T > 0$, it holds that*

$$\liminf_{\epsilon \rightarrow 0} \left[\inf_{I \times [2\sqrt{\epsilon}, T]} \partial_z^2 H_\epsilon(z, t) \right] > 0. \quad (3.3)$$

Then for all $\epsilon > 0$ small, there exists a unique viscosity solution $(V_\epsilon(z, t), \bar{z}_\epsilon(t))$ to the following Hamilton-Jacobi equation with a constraint:

$$\begin{cases} \partial_t V_\epsilon + |\partial_z V_\epsilon|^2 - H_\epsilon(z, t) + H_\epsilon(\bar{z}_\epsilon(t), t) = 0 & \text{for } t \in [2\sqrt{\epsilon}, T], z \in I, \\ \partial_z V_\epsilon = 0 & \text{for } t \in [2\sqrt{\epsilon}, T], z \in \partial I, \\ \bar{z}_\epsilon(2\sqrt{\epsilon}) = \bar{z}_0, \quad \text{and} \quad V_\epsilon(z, 2\sqrt{\epsilon}) = V_0(z) & \text{for } z \in I, \\ \inf_{z \in I} V_\epsilon(z, t) = 0 & \text{for } t \in [2\sqrt{\epsilon}, T]. \end{cases} \quad (3.4)$$

Moreover, $\liminf_{\epsilon \rightarrow 0} \inf_{I \times [2\sqrt{\epsilon}, T]} \partial_{zz} V_\epsilon > 0$ (i.e. V_ϵ is strictly convex in z) and

$$V_\epsilon(\bar{z}_\epsilon(t), t) = 0, \quad \text{and} \quad V_\epsilon(z, t) > 0 \quad \text{for } z \neq \bar{z}_\epsilon(t),$$

and there exists a constant $C > 0$ independent of ϵ such that

$$\|\bar{z}_\epsilon(\cdot)\|_{C^{0,1}([2\sqrt{\epsilon}, T])} \leq C. \quad (3.5)$$

Remark 3.3. Thanks to (3.3) and the convexity of the initial data, V_ϵ and $\bar{z}_\epsilon(t)$ satisfy (3.4), except for the Neumann boundary condition, in the classical sense. The Neumann boundary condition, which has to be understood in the viscosity sense, is satisfied if $\partial_z V_\epsilon(\inf I, t) \leq 0 \leq \partial_z V_\epsilon(\sup I, t)$ for all t . This is indeed the case in our setting, where V_ϵ is convex in z such that the unique minimum point $\bar{z}_\epsilon(t)$ of $z \mapsto V_\epsilon(z, t)$ is attained in the interior for all time.

4 WKB Ansatz and Some Technical Lemmas

The rate function $u_\epsilon(x, z, t) = -\epsilon \log n_\epsilon(x, z, t)$ satisfies the equation

$$\begin{cases} \partial_t u_\epsilon - \frac{\alpha(z)}{\epsilon} \Delta_x u_\epsilon + \frac{\alpha(z)}{\epsilon^2} |\nabla_x u_\epsilon|^2 - \epsilon \partial_z^2 u_\epsilon + |\partial_z u_\epsilon|^2 + & m(x) - \rho_\epsilon(x, t) = 0 \\ & x \in D, z \in I, t > 0, \\ \partial_\nu u_\epsilon = 0 & x \in \partial D, z \in I, t > 0, \\ \partial_z u_\epsilon = 0 & x \in D, z \in \partial I, t > 0, \\ u_\epsilon(x, z, 0) = u_{\epsilon,0}(x, z) := -\epsilon \log n_{\epsilon,0}(x, z) & x \in D, z \in I. \end{cases} \quad (4.1)$$

Let $\varphi_\epsilon(x, z, t)$ be given in Proposition 3.1. Using the perturbed test function method, we define the function $v_\epsilon(x, z, t) := u_\epsilon(x, z, t) - \epsilon \varphi_\epsilon(x, z, t)$, which satisfies

$$\begin{cases} \partial_t v_\epsilon - \frac{\alpha(z)}{\epsilon} \Delta_x v_\epsilon + \frac{\alpha(z)}{\epsilon^2} |\nabla_x v_\epsilon|^2 + 2 \frac{\alpha(z)}{\epsilon} \nabla_x v_\epsilon \cdot \nabla_x \varphi_\epsilon - \epsilon \partial_z^2 v_\epsilon & + |\partial_z v_\epsilon|^2 + 2\epsilon \partial_z v_\epsilon \partial_z \varphi_\epsilon \\ = H_\epsilon(z, t) - \epsilon^2 (\partial_z^2 \varphi_\epsilon + |\partial_z \varphi_\epsilon|^2) + \rho_\epsilon(x, t) - \rho_\epsilon(x, \max\{t, \epsilon\}) & \text{for } x \in D, z \in I, t > 0, \\ \partial_\nu v_\epsilon = 0 & \text{for } x \in \partial D, z \in I, t > 0, \\ \partial_z v_\epsilon = -\epsilon \partial_z \varphi_\epsilon = O(\epsilon) & \text{for } x \in D, z \in \partial I, t > 0, \\ v_\epsilon(x, z, 0) = u_{\epsilon,0}(x, z) & \text{for } x \in D, z \in I. \end{cases} \quad (4.2)$$

Remark 4.1. The right hand side of (4.2) is essentially the Hamiltonian $H_\epsilon(z, t) + O(\epsilon)$, since the term $\rho_\epsilon(x, t) - \rho_\epsilon(x, \max\{t, \epsilon\})$ is identically zero except for $t \in [0, \epsilon]$.

Remark 4.2. By (3.2) we have

$$\|u_\epsilon - v_\epsilon\|_{L^\infty(D \times I \times (0, \infty))} = \epsilon \|\varphi_\epsilon\|_{L^\infty(D \times I \times (0, \infty))} = O(\epsilon),$$

thus upper and lower estimates of u_ϵ and v_ϵ are the same, up to an error of $O(\epsilon)$. We shall construct super- and sub-solutions of v_ϵ .

We are presented with a problem involving different time scales: n_ϵ solving equation (1.6) involves a fast time scale coupled with small mutation rate; the rate function u_ϵ satisfies in the limit a regular in time Hamilton-Jacobi equation of first order; the total population ρ_ϵ and the normalized principal Floquet bundle has a fast time scale which acts as a quasi-stationary limit. In this section, we prove some estimates that quantifies these interrelated facts.

Suppose $u_\epsilon(u, z, t)$ is close to a Dirac mass supported at $\{\hat{z}\}$ in the z variable at time \hat{t} , the following results give a quantitative estimate that says that it stays close in a short time.

Proposition 4.3. *Suppose that there exist $\hat{t} \geq \epsilon$, $\hat{z} \in I$, $V_1 \in C^2(\bar{I})$ and $\eta_1 > 0$, such that*

$$V_1(z) \geq 0, \quad V_1(z) = 0 \text{ iff } z = \hat{z}, \quad \partial_z^2 V_1(z) > \eta_1 > 0 \text{ for } \hat{z} - \eta_1 \leq z \leq \hat{z} + \eta_1 \quad (4.3)$$

and it holds that

$$\sup_{\hat{t}-\epsilon \leq t \leq \hat{t}} \|u_\epsilon(x, z, t) - V_1(z)\|_{C(\bar{D} \times \bar{I})} \rightarrow 0 \quad \text{as } \epsilon \rightarrow 0.$$

Then there exists $C > 0$ depending on η_1 , but independent of \hat{t} and \hat{z} and V_1 such that for each small $\delta > 0$,

$$\limsup_{\epsilon \rightarrow 0} \sup_{\hat{t} \leq t \leq \hat{t} + 2\delta} \left\| \int_I (z - \hat{z}) \partial_x^2 n_\epsilon(\cdot, z, t) dz \right\|_{C(\bar{D})} \leq C\delta^{1/3}. \quad (4.4)$$

Proof. Note that, similar to the argument in Step 3 of the proof of Proposition A.1, a lower solution for (4.1) can be constructed as:

$$\underline{U}(z, t) = V_1(z) - C_1(\mu_\epsilon + t - (\hat{t} - \epsilon)) - \frac{C}{\epsilon} \{ [z - (b - \sqrt{\epsilon})]_+^3 + [a + \sqrt{\epsilon} - z]_+^3 \}, \quad (4.5)$$

where $a = \inf I$, $b = \sup I$, and

$$\mu_\epsilon = \sup_{x \in D, z \in I} |u_\epsilon(x, z, \hat{t} - \epsilon) - V_1(z)|$$

is a constant tending to zero as $\epsilon \rightarrow 0$, and that $C_1 = C_1(\|m(x) - \rho_\epsilon\|_{C(\bar{D} \times [0, \infty))})$. By comparison in $D \times I \times [\hat{t} - \epsilon, +\infty)$, we have

$$u_\epsilon(x, z, t) \geq V_1(z) - C_1(\mu_\epsilon + |t - \hat{t} + \epsilon|) + O(\sqrt{\epsilon}) \quad \text{for } |z - \hat{z}| \geq \delta \quad \text{and } t > \hat{t} - \epsilon,$$

where

$$\inf_{|z - \hat{z}| \geq \delta} V_1 \geq \frac{\eta_1 \delta^2}{2} \quad \text{for } 0 < \delta \ll 1.$$

Therefore, for each fixed δ small, by choosing ϵ sufficiently small,

$$u_\epsilon(x, z, t) \geq \frac{\eta_1 \delta^2}{3} \quad \text{for } |z - \hat{z}| > \delta \quad \text{and } \hat{t} - \epsilon \leq t \leq \hat{t} + \delta^3,$$

which means

$$n_\epsilon(x, z, t) \leq \exp\left(-\frac{\eta_1 \delta^2}{3\epsilon}\right) \quad \text{for } |z - \hat{z}| > \delta \quad \text{and } \hat{t} - \epsilon \leq t \leq \hat{t} + \delta^3. \quad (4.6)$$

Now, for $t_1 \in [\hat{t}, \hat{t} + \delta^3]$,

$$\begin{aligned} & \left\| \int_I (z_1 - \hat{z}) \partial_x^2 n_\epsilon(\cdot, z_1, t_1) dz_1 \right\|_{C(\bar{D})} \leq \int_I |z_1 - \hat{z}| \|\partial_x^2 n_\epsilon(\cdot, z_1, t_1)\|_{C(\bar{D})} dz_1 \\ & \leq C \int_I |z_1 - \hat{z}| \left[\int_{t_1 - \epsilon}^{t_1} \int_{-\epsilon}^\epsilon \int_D n_\epsilon(x, z + z_1, t) dx dz dt \right] dz_1 \\ & \leq 2C\delta \int_{\hat{z} - 2\delta}^{\hat{z} + 2\delta} \left[\int_{t_1 - \epsilon}^{t_1} \int_{-\epsilon}^\epsilon \int_D n_\epsilon(x, z + z_1, t) dx dz dt \right] dz_1 \\ & \quad + C \int_{z_1: |z_1 - \hat{z}| > 2\delta} \left[\int_{t_1 - \epsilon}^{t_1} \int_{-\epsilon}^\epsilon \int_D n_\epsilon(x, z + z_1, t) dx dz dt \right] dz_1 \\ & \leq 2C\delta \int_{-\epsilon}^\epsilon \left[\int_{t_1 - \epsilon}^{t_1} \int_D \int_{\hat{z} - 2\delta}^{\hat{z} + 2\delta} n_\epsilon(x, z + z_1, t) dz_1 dx dt \right] dz + C' \exp\left(-\frac{\eta_1 \delta^2}{3\epsilon}\right) \\ & \leq 2C\delta \int_{-\epsilon}^\epsilon \left[\sup_{\hat{t} - \epsilon < t < \hat{t}} \|\rho_\epsilon(t, \cdot)\|_{L^1(D)} \right] dz + C' \exp\left(-\frac{\eta_1 \delta^2}{3\epsilon}\right), \end{aligned}$$

where we used (2.2) in Lemma 2.2 in the second inequality, and we switched the order of integration and used (4.6) to obtain the fourth inequality. By fixing δ to be small enough, we see that (4.4) holds for all ϵ sufficiently small. This completes the proof. \square

If we prescribe the diffusion rate to be slowly varying in time, and solve the diffusive logistic problem with a fast time scale, then the solution can be described via the corresponding quasi-steady-state approximation. The following is a quantitative result to this effect.

Lemma 4.4. *Let $F \in L^\infty([t_1, t_2]; L^p(D))$ be given and consider the equation*

$$\begin{cases} \epsilon \partial_t \rho - \alpha(z_1(t)) \Delta_x \rho = \rho(m(x) - \rho) + F(x, t) & \text{for } x \in D, t_1 \leq t \leq t_2, \\ \partial_\nu \rho = 0 & \text{for } x \in \partial D, t_1 \leq t \leq t_2. \end{cases}$$

Suppose, for some fixed constant $M > 1$,

$$z_1(t) \in I, \quad |z_1'(t)| \leq M \quad \text{for } t_1 \leq t \leq t_2.$$

Then for each $p > 1$, there exists $C, C_p > 1$ (independent of F) such that for $t \in [t_1, t_2]$,

$$\begin{aligned} & \|\rho(\cdot, t) - \theta_{z_1(t)}(\cdot)\|_{L^p(D)}^p \\ & \leq C \left[\|\rho(\cdot, t_1) - \theta_{z_1(t_1)}(\cdot)\|_{L^p(D)}^p e^{-(t-t_1)/(\epsilon C_p)} + \sup_{t_1 \leq t' \leq t} \|F(\cdot, t')\|_{L^p(D)}^p + \epsilon \right]. \end{aligned}$$

Proof. For each t , set $\alpha_1(t) := \alpha(z_1(t))$ and let $\theta_t(x) = \theta_{z_1(t)}(x)$ denote the unique positive solution of

$$\alpha_1(t) \Delta_x \theta + \theta(m - \theta) = 0 \quad \text{in } D, \quad \partial_\nu \theta = 0 \quad \text{on } \partial D.$$

Set $W(x, t) := \rho(x, t)/\theta_t(x) - 1$, it satisfies the Neumann boundary condition in x , i.e.,

$$\partial_\nu W(x, t) = 0 \quad \text{for } t \geq 0, x \in \partial D,$$

and, for $p > 1$, the equation

$$\begin{aligned} & \frac{\epsilon}{p} \partial_t (\theta_t^2 |W|^p) - \alpha_1(t) \nabla_x \cdot (\theta_t^2 |W|^{p-1} \nabla_x W) + \alpha_1(t) \frac{4(p-1)}{p^2} \theta_t^2 |\nabla_x W^{p/2}|^2 + \rho \theta_t^2 |W|^p \\ & = F \theta_t W |W|^{p-2} - \epsilon \theta_t \partial_t \theta_t W |W|^{p-2} - \epsilon \left(1 - \frac{2}{p}\right) \theta_t \partial_t \theta_t |W|^p, \end{aligned}$$

from which we derive the differential inequality

$$\begin{aligned} & \frac{\epsilon}{p} \partial_t (\theta_t^2 |W|^p) - \alpha_1(t) \nabla_x \cdot (\theta_t^2 |W|^{p-1} \nabla_x W) + \rho \theta_t^2 |W|^p \\ & \leq \theta_t |W|^{p-1} |F| + \hat{C}_0 \epsilon \theta_t |\partial_t \theta_t| (1 + |W|^p). \end{aligned} \quad (4.7)$$

Now, let \hat{C}_1 be the constant given in Proposition 2.1. We apply Young's inequality to get

$$\theta_t |W|^{p-1} |F| = \left[\left(\frac{q}{3\hat{C}_1} \right)^{\frac{1}{q}} \theta_t^{\frac{2}{q}} |W|^{p-1} \right] \left[\left(\frac{3\hat{C}_1}{q} \right)^{\frac{1}{q}} \theta_t^{1-\frac{2}{q}} |F| \right] \leq \frac{1}{3\hat{C}_1} \theta_t^2 |W|^p + C' |F|^p. \quad (4.8)$$

where $q = \frac{p}{p-1}$ and used that there is a constant C independent of ϵ such that

$$\frac{1}{C} \leq \theta_t(x) \leq C \quad \text{for } x \in D, \quad \text{and} \quad \sup_t \left[\|\partial_t \theta_t\|_{C^0(\bar{D})} + \|\theta_t\|_{C^2(\bar{D})} \right] \leq C, \quad (4.9)$$

where the estimates for θ_t follow from **(M)** and Schauder estimates, and the one for $\partial_t \theta_t$ follow from the implicit function theorem. By enlarging C' if necessary, we can also use (4.9) to deduce

$$\hat{C}_0 \epsilon \theta_t |\partial_t \theta_t| (1 + |W|^p) \leq \frac{1}{3\hat{C}_1} \theta_t^2 |W|^p + C' \epsilon. \quad (4.10)$$

Combining (4.7), (4.8), (4.10), and recalling that ρ is bounded from below by $\frac{1}{\hat{C}_1}$, we obtain

$$\begin{aligned} & \frac{\epsilon}{p} \partial_t (\theta_t^2 |W|^p) - \alpha_1(t) \nabla_x \cdot (\theta_t^2 |W|^{p-1} \nabla_x W) + \frac{1}{\hat{C}_1} \theta_t^2 |W|^p \\ & \leq \frac{1}{3\hat{C}_1} \theta_t^2 |W|^p + C' |F|^p + \frac{1}{3\hat{C}_1} \theta_t^2 |W|^p + C' \epsilon. \end{aligned}$$

Integrating in $x \in D$ and consolidating the terms with $\int_D \theta_t^2 |W|^p$, we obtain

$$\frac{\epsilon}{p} \frac{d}{dt} \int_D \theta_t^2 |W|^p dx + \frac{1}{3\hat{C}_1} \int_D \theta_t^2 |W|^p dx \leq C' \left(\int_D |F(x, t)|^p dx + \epsilon \right).$$

Let $M(t) := \int_D \theta_t^2 |W|^p dx$, we integrate the above in $t \in [t_1, t_2]$ to get

$$M(t_2) \leq M(t_1) e^{-(t_2-t_1)/(\epsilon C_p)} + 3\hat{C}_1 C' \left[\sup_{t_1 \leq t \leq t_2} \|F(\cdot, t)\|_{L^p(D)}^p + \epsilon \right],$$

where $C_p = 3\hat{C}_1/p$. The lemma follows since θ_t is uniformly bounded from above and below. \square

Suppose that the diffusion rate and the zero-th order coefficient are varying slowly in time, then the elliptic eigenvalue problem can be considered a quasi-steady-state approximation of the Floquet bundle equation (3.1), when ϵ is small. This is proved in the following result.

Proposition 4.5. *Given $T > 0$ and let $\|\bar{z}_\epsilon(t)\|_{C^{0,1}([0, T])}$ be uniformly bounded in $0 < \epsilon \ll 1$. For each $\eta > 0$, there exists $\nu = \nu(\eta) > 0$ such that if*

$$\limsup_{\epsilon \rightarrow 0} \sup_{[\sqrt{\epsilon}, T]} \|\rho_\epsilon(\cdot, t) - \theta_{\bar{z}_\epsilon(t)}(\cdot)\|_{L^2(D)} < \nu, \quad (4.11)$$

then

$$\limsup_{\epsilon \rightarrow 0} \sup_{[2\sqrt{\epsilon}, T-\sqrt{\epsilon}]} \|H_\epsilon(\cdot, t) - \lambda(\cdot, \bar{z}_\epsilon(t))\|_{C^2(\bar{I})} < \eta. \quad (4.12)$$

Proof. By Proposition B.3 in Appendix B, the principal Floquet bundle depends smoothly on the weight function and $z \in I$, i.e. the mapping

$$\begin{cases} H_1 : C^{\beta/2, \beta}(\mathbb{R} \times \bar{D}) \times \bar{I} \rightarrow C^{\beta/2}(\mathbb{R}) \\ (c(x, t), z) \mapsto H_1(\tau) \end{cases}$$

is smooth. Recall also the fact, when $c(x, t) = m(x) - \theta_{\hat{z}}(x)$

$$H_1(\tau) = H_1(\tau; m - \theta_{\hat{z}}, z) \equiv \lambda(z, \hat{z}) \quad \text{and is constant in } \tau \in \mathbb{R}.$$

Therefore, for each $\eta > 0$ there exists $\nu' > 0$ such that if

$$\|c(x, \tau) - (m(x) - \theta_{\hat{z}}(x))\|_{C^{\beta, \beta/2}(\bar{D} \times \mathbb{R})} < \nu' \quad \text{for some } \hat{z} \in \bar{I}, \quad (4.13)$$

then the corresponding principal Floquet bundle $H_1(\tau; c, z)$ satisfies

$$\|H_1(0; c, \cdot) - \lambda(\cdot, \hat{z})\|_{C^2(\bar{I})} \leq \sup_{\tau \in \mathbb{R}} \|H_1(\tau; c, \cdot) - \lambda(\cdot, \hat{z})\|_{C^2(\bar{I})} < \eta. \quad (4.14)$$

Now, fix $\beta' \in (\beta, 1)$, and choose by interpolation the constant $\nu > 0$ such that if

$$\begin{cases} \|c(x, \tau) - [m(x) - \theta_{\hat{z}}(x)]\|_{C^{\beta', \beta'/2}(\bar{D} \times \mathbb{R})} \leq \hat{C}_{\beta'} + \|\theta_{\hat{z}}(x) - m(x)\|_{C^{\beta'}(\bar{D})} \\ \sup_{\tau \in \mathbb{R}} \|c(x, \tau) - [m(x) - \theta_{\hat{z}}(x)]\|_{L^2(D)} < 2\nu, \end{cases} \quad (4.15)$$

then (4.13) and thus (4.14) holds. (Note that $\nu > 0$ depends on the uniform bound $\hat{C}_{\beta'}$ in Corollary 2.3 but is independent of ϵ, \hat{z} .)

We claim that if (4.11) holds for the constant ν we just specified, then (4.12) holds. Suppose not, then there exist sequences $\epsilon = \epsilon_j \rightarrow 0$ and $t_j \in [2\sqrt{\epsilon_j}, T - \sqrt{\epsilon_j}]$ such that $\bar{z}_{\epsilon_j}(t_j) \rightarrow \hat{z}$ and

$$\begin{cases} \lim_{\epsilon_j \rightarrow 0} \sup_{[\sqrt{\epsilon_j}, T]} \|\rho_{\epsilon_j}(\cdot, t) - \theta_{\bar{z}_{\epsilon_j}(t)}(\cdot)\|_{L^2(D)} \leq \nu, \\ \lim_{\epsilon_j \rightarrow 0} \|H_{\epsilon_j}(\cdot, t_j) - \lambda(\cdot, \bar{z}_{\epsilon_j}(t_j))\|_{C^2(\bar{I})} = \lim_{\epsilon_j \rightarrow 0} \|H_{\epsilon_j}(\cdot, t_j) - \lambda(\cdot, \hat{z})\|_{C^2(\bar{I})} \geq \eta. \end{cases} \quad (4.16)$$

Let $(\varphi_\epsilon(x, z, t), H_\epsilon(z, t))$ be the function given by Proposition 3.1, and define

$$\Phi_j(x, z, \tau) := e^{-\varphi_{\epsilon_j}(x, z, t_j + \epsilon_j \tau)}, \quad \text{and} \quad H_j(z, \tau) := H_{\epsilon_j}(z, t_j + \epsilon_j \tau).$$

Passing to a subsequence if necessary, we deduce that

$$\Phi_j(x, z, \tau) \rightarrow \hat{\Phi}(x, z, \tau) \quad \text{in } C_{loc}(\bar{D} \times \bar{I} \times \mathbb{R}), \quad \text{and} \quad H_j(z, \tau) \rightarrow \hat{H}(z, \tau) \quad \text{in } C_{loc}(\bar{I} \times \mathbb{R}),$$

where the limit $(\hat{\Phi}(x, z, \tau), \hat{H}(z, \tau))$ is a pair of the normalized principal Floquet bundle, i.e.

$$(\hat{\Phi}(x, z, \tau), \hat{H}(z, \tau)) = (\Phi_1(x, \tau; \hat{c}, z), H_1(x, \tau; \hat{c}, z)),$$

for some \hat{c} (which exists thanks to Corollary 2.3) given by

$$\hat{c}(x, \tau) = \lim_{\epsilon_j \rightarrow 0} (\rho_{\epsilon_j}(x, t_j + \epsilon_j \tau) - m(x)).$$

By the first statement of (4.16), we deduce that the weight \hat{c} satisfies (4.15), and hence the limit \hat{H} of H_{ϵ_j} (in the $C_{loc}(\mathbb{R})$ topology) satisfies (4.14). Moreover, recalling (3.2), we have

$$\sup_j \|H_j(\cdot, 0)\|_{C^3(\bar{I})} = \sup_j \|H_{\epsilon_j}(\cdot, t_j)\|_{C^3(\bar{I})} \leq C,$$

we deduce that $H_j(\cdot, 0) \rightarrow \hat{H}(\cdot, 0)$ in $C^2(\bar{I})$. Hence

$$\lim_{\epsilon_j \rightarrow 0} \|H_{\epsilon_j}(\cdot, t_j) - \lambda(\cdot, \hat{z})\|_{C^2(\bar{I})} = \|\hat{H}(\cdot, 0) - \lambda(\cdot, \hat{z})\|_{C^2(\bar{I})} < \eta,$$

where the last inequality holds since \hat{H} satisfies (4.14). This contradicts the second statement of (4.16). \square

Corollary 4.6. *Let V_1 satisfy the condition (4.3) of Proposition 4.3 for some given $\eta_1 > 0$, $\hat{z} \in I$, $\hat{t} \geq 3\sqrt{\epsilon}$. There exists $\delta_1 > 0$ depending on η_1 but independent of \hat{z}, \hat{t}, V_1 such that if*

$$\sup_{\hat{t} - 3\sqrt{\epsilon} \leq t \leq \hat{t} - 2\sqrt{\epsilon}} \|u_\epsilon(x, z, t) - V_1(z)\|_{C(\bar{D} \times \bar{I})} \rightarrow 0 \quad \text{as } \epsilon \rightarrow 0, \quad (4.17)$$

then

$$\liminf_{\epsilon \rightarrow 0} \inf_{I \times [\hat{t}, \hat{t} + \delta_1]} \partial_{zz}^2 H_\epsilon(z, t) \geq \delta_1 > 0. \quad (4.18)$$

Proof. Integrating (1.6) over z , we see that ρ_ϵ satisfies

$$\epsilon \partial_t \rho_\epsilon = \alpha(\hat{z}) \Delta_x \rho_\epsilon + \rho_\epsilon (m(x) - \rho_\epsilon(x, t)) + \int_I (\alpha(z) - \alpha(\hat{z})) \Delta_x n_\epsilon dz \quad (4.19)$$

for $(x, t) \in D \times [0, \infty)$ and the Neumann boundary condition on $\partial D \times [0, \infty)$. Next, taking

$$z_1(t) \equiv \hat{z}, \quad t_1 = \hat{t} - 2\sqrt{\epsilon}, \quad t_2 = \hat{t} + 2\delta_1, \quad F = \int_I (\alpha(z) - \alpha(\hat{z})) \Delta_x n_\epsilon dz,$$

and applying Lemma 4.4, we again obtain C' independent of ϵ and δ_1 such that for each $\delta_1 > 0$ small,

$$\begin{aligned} & \sup_{\hat{t}-\sqrt{\epsilon} \leq t \leq \hat{t}+2\delta_1} \|\rho_\epsilon(\cdot, t) - \theta_{\hat{z}}(\cdot)\|_{L^2(D)} \\ & \leq C' \left[\|\rho_\epsilon(\cdot, \hat{t} - 2\sqrt{\epsilon}) - \theta_{\hat{z}}(\cdot)\|_{L^2(D)} e^{-\sqrt{\epsilon}/(\epsilon C')} + \sup_{\hat{t}-2\sqrt{\epsilon} \leq t' \leq \hat{t}+2\delta_1} \|F(\cdot, t')\|_{L^2(D)} \right]. \end{aligned} \quad (4.20)$$

Next, apply Proposition 4.3, then (4.17) to deduce that

$$\limsup_{\epsilon \rightarrow 0} \sup_{t \in [\hat{t}-2\sqrt{\epsilon}, \hat{t}+2\delta_1]} \|F(x, t)\|_{L^2(D)} < C|\delta_1|^{1/3}. \quad (4.21)$$

Using the fact that $\|\rho_\epsilon\|_{L^\infty} + \|\theta_{\hat{z}}\|_{L^\infty}$ is uniformly bounded in ϵ and also (4.21), we can take limit superior in ϵ in (4.20) to obtain C'' independent of δ_1 such that for each $\delta_1 > 0$ small,

$$\limsup_{\epsilon \rightarrow 0} \sup_{\hat{t}-\sqrt{\epsilon} \leq t \leq \hat{t}+2\delta_1} \|\rho_\epsilon(\cdot, t) - \theta_{\hat{z}}(\cdot)\|_{L^2(D)} < C''|\delta_1|^{1/3}.$$

By choosing δ_1 small, Proposition 4.5 says that

$$\limsup_{\epsilon \rightarrow 0} \sup_{[\hat{t}, \hat{t}+\delta_1]} \|\partial_{z_1 z_1}^2 \lambda(\cdot, \hat{z}) - \partial_{z_1 z_1}^2 H_\epsilon(\cdot, t)\|_{C(\bar{U})} \leq \frac{1}{2} \inf_{I \times I} \lambda_{z_1 z_1},$$

since by **(H1)** the term on the right is a fixed positive constant. Hence, by choosing a suitably small $\delta_1 > 0$, (4.18) holds. \square

5 Proof of Theorem 1.4

Define

$$T^* := \sup\{T > 0 : \liminf_{\epsilon \rightarrow 0} \inf_{I \times [2\sqrt{\epsilon}, T]} \partial_{zz}^2 H_\epsilon(z, t) > 0\}. \quad (5.1)$$

Our purpose is to prove that $T^* = \infty$, which comes with the desired convergence results.

Applying Proposition 3.2, the two quantities $\bar{z}_\epsilon(t)$ and $V_\epsilon(z, t)$ are well-defined in the time interval $[2\sqrt{\epsilon}, T^*]$, for all sufficiently small ϵ . We will also define for convenience

$$V_\epsilon(z, t) = V_0(z), \quad \text{and} \quad \bar{z}_\epsilon(t) = \bar{z}_0 \quad \text{for } z \in I, t \in [0, 2\sqrt{\epsilon}].$$

Step 1. In this step, we show that $T^* \geq \delta_1$, where δ_1 is as given in Corollary 4.6.

By constructing super- and sub-solutions of the form

$$V_0(z) - \frac{1}{2}\epsilon |\log \epsilon| \pm C \left\{ (t + \epsilon) + \frac{1}{\epsilon} \left([z - (b - \sqrt{\epsilon})]_+^3 + [a + \sqrt{\epsilon} - z]_+^3 \right) \right\},$$

it follows by comparison using (4.2) that

$$\sup_{0 \leq t \leq 3\sqrt{\epsilon}} \|u_\epsilon(x, z, t) - V_0(z)\|_{C(\bar{D} \times \bar{I})} \rightarrow 0 \quad \text{as } \epsilon \rightarrow 0. \quad (5.2)$$

Based on (5.2), one can apply Corollary 4.6 to yield $T^* \geq \delta_1$.

Step 2. Next, we show

$$\sup_{D \times I \times [2\sqrt{\epsilon}, T^*]} \left| u_\epsilon(x, z, t) - V_\epsilon(z, t) - \int_{2\sqrt{\epsilon}}^t H_\epsilon(\bar{z}_\epsilon(s), s) ds \right| \rightarrow 0. \quad (5.3)$$

To this end, define $\mu_\epsilon = \|u_\epsilon(x, z, 2\sqrt{\epsilon}) - V_0(z)\|_{C(\bar{D} \times \bar{I})} = o(1)$ and

$$\begin{aligned}\bar{U}(z, t) &= V_\epsilon(z, t) + \int_{2\sqrt{\epsilon}}^t H_\epsilon(\bar{z}_\epsilon(s), s) ds + \mu_\epsilon + \frac{C}{\epsilon} \left\{ [(a + \sqrt{\epsilon}) - z]_+^3 + [z - (b - \sqrt{\epsilon})]_+^3 \right\}, \\ \underline{U}(z, t) &= V_\epsilon(z, t) + \int_{2\sqrt{\epsilon}}^t H_\epsilon(\bar{z}_\epsilon(s), s) ds - \mu_\epsilon - \frac{C}{\epsilon} \left\{ [(a + \sqrt{\epsilon}) - z]_+^3 - [z - (b - \sqrt{\epsilon})]_+^3 \right\}.\end{aligned}$$

Then (5.3) follows by comparison using (4.2).

Step 3. In this step, we show that as $\epsilon \rightarrow 0$,

$$\sup_{D \times I \times [0, T^*]} |u_\epsilon(x, z, t) - V_\epsilon(z, t)| \rightarrow 0. \quad (5.4)$$

We first claim that

$$\sup_{2\sqrt{\epsilon} \leq t \leq T^*} \left| \int_{2\sqrt{\epsilon}}^t H_\epsilon(\bar{z}_\epsilon(s), s) ds \right| = o(1). \quad (5.5)$$

Indeed, using the comparison functions in the previous step and that $\mu_\epsilon = o(1)$,

$$\begin{aligned}\int_D \rho_\epsilon(x, t) dx &= \iint_{D \times I} \exp\left(-\frac{u_\epsilon(x, z, t)}{\epsilon}\right) dz dx \\ &= \iint_{D \times I} \exp\left(-\frac{V_\epsilon(z, t)}{\epsilon} - \frac{\int_{2\sqrt{\epsilon}}^t H_\epsilon(\bar{z}_\epsilon(s), s) ds + o(1)}{\epsilon}\right) dz dx.\end{aligned} \quad (5.6)$$

Since $z \mapsto V_\epsilon(z, t)$ is strictly convex and attains a unique minimum at $\bar{z}_\epsilon(t)$, we can find constants $0 < c_* < c^*$ such that $-c^*(z - \bar{z}_\epsilon(t))^2 \leq V_\epsilon(z, t) \leq -c_*(z - \bar{z}_\epsilon(t))^2$. This gives

$$\sqrt{\frac{\pi}{c^*}} \leq \frac{1}{\sqrt{\epsilon}} \int \exp\left(-\frac{V_\epsilon(z, t)}{\epsilon}\right) dz \leq \sqrt{\frac{\pi}{c_*}}.$$

Using this $\frac{1}{C} \leq \int_D \rho_\epsilon(x, t) dx \leq C$ (from Proposition 2.1) in (5.6), we deduce

$$A_\epsilon = B_\epsilon \exp\left(-\frac{\int_{2\sqrt{\epsilon}}^t H_\epsilon(\bar{z}_\epsilon(s), s) ds + o(1)}{\epsilon}\right),$$

where $A_\epsilon \sim 1$ and $B_\epsilon \sim \sqrt{\epsilon}$. This proves (5.5). Combining (5.2), (5.3) and (5.5), we deduce (5.4).

Step 4. In this step, we show that

$$\sup_{\sqrt{\epsilon} \leq t \leq T^*} \|\rho_\epsilon(\cdot, t) - \theta_{\bar{z}_\epsilon(t)}\|_{C(\bar{D})} \rightarrow 0 \quad \text{as } \epsilon \rightarrow 0. \quad (5.7)$$

First, we deduce from (5.4) and Proposition 4.3 that

$$\sup_{[\epsilon, T^*]} \left\| \int_I (z - \bar{z}_\epsilon(t)) \partial_x^2 n_\epsilon(\cdot, z, t) dz \right\|_{C(\bar{D})} \rightarrow 0.$$

Note that Proposition 4.3 is applicable since the solution V_ϵ of (3.4) remains strictly convex in z , since H_ϵ is strictly convex in z up to time T^* thanks to (5.1). Then by taking $t_1 = \epsilon$ and $t_2 \in [\sqrt{\epsilon}, T^*]$ in Lemma 4.4, we deduce

$$\sup_{\sqrt{\epsilon} \leq t \leq T^*} \|\rho_\epsilon(\cdot, t) - \theta_{\bar{z}_\epsilon(t)}\|_{L^2(D)} \rightarrow 0. \quad (5.8)$$

The estimate (5.7) follows by interpolating (5.8) with the uniform estimate

$$\sup_{\sqrt{\epsilon} \leq t \leq T^*} \|\rho_\epsilon(\cdot, t) - \theta_{\bar{z}_\epsilon(t)}\|_{C^\beta(\bar{D})} \leq C,$$

which follows from Corollary 2.3(b) and that $\|\bar{z}_\epsilon(\cdot)\|_{C^{0,1}} \leq C$ (by Proposition 3.2).

Step 5. By (5.7) and Proposition 4.5, we have

$$\sup_{I \times [2\sqrt{\epsilon}, T^* - \sqrt{\epsilon}]} |H_\epsilon(z, t) - \lambda(z, \bar{z}_\epsilon(t))| \rightarrow 0. \quad (5.9)$$

Since $(V_\epsilon(z, t), \bar{z}_\epsilon(t))$ is the unique viscosity solution of (3.4), we may apply the stability theorem for viscosity solution, to deduce that, as $\epsilon \rightarrow 0$, $(V_\epsilon(z, t), \bar{z}_\epsilon(t))$ must converge to the unique viscosity solution $(V(z, t), \bar{z}(t))$ of (1.10), i.e.

$$\sup_{I \times [2\sqrt{\epsilon}, T^*]} |V_\epsilon(z, t) - V(z, t)| \rightarrow 0 \quad \text{and} \quad \sup_{[2\sqrt{\epsilon}, T^*]} |\bar{z}_\epsilon(t) - \bar{z}(t)| \rightarrow 0. \quad (5.10)$$

Step 6. We claim that $T^* = +\infty$. Suppose not, then $T^* < +\infty$. But then the above argument would allow us to extend the approximate trajectory by a fixed time-step $\delta_1 > 0$. This would yield a contradiction which proves that $T^* = +\infty$.

Step 7. To conclude, it remains to show (1.12) - (1.14). Indeed, (1.14) follows from $\rho_\epsilon(x, t) \rightarrow \theta_{\bar{z}_\epsilon(t)}(x)$ (by (5.7)) and $\bar{z}_\epsilon(t) \rightarrow \bar{z}(t)$ (by (5.10)). Next, (1.13) follows from

$$\sup_{D \times I \times [0, T^*]} |u_\epsilon(x, z, t) - V_\epsilon(z, t)| \rightarrow 0$$

(by (5.4)) and $V_\epsilon(z, t) \rightarrow V(z, t)$ (by (5.10)). Finally, Proposition 1.3 says for each t , $z \mapsto V(z, t)$ has a unique minimum point at $\bar{z}(t)$, so that

$$n_\epsilon(x, z, t) = \exp\left(-\frac{V_\epsilon(z, t) + o(1)}{\epsilon}\right) \approx \delta_0(z - \bar{z}(t))\rho_\epsilon(x, t), \quad (5.11)$$

where the function $\rho_\epsilon(x, t) = \int_I n_\epsilon(x, z, t) dz$ appeared since the above needs to be consistent upon integration over $z \in I$. Combining (1.14) and (5.11), we deduce (1.12).

6 Discussions and Generalizations

In Subsection 6.1, we discuss a special feature of the effective Hamiltonian λ that arises from evolution of random dispersal, as modeled by Laplacian in the spatial variable, which is the fact that $z_1 \mapsto \lambda(z_1, z_2)$ has the same monotonicity of α , regardless of choice of α and z_2 . Subsection 6.2 gives a concrete example of $\alpha(z)$ where **(H1)** can be verified. Subsection 6.3 discusses the generalization to multi-dimensional traits, as motivated by the evolution of conditional dispersal, where the optimal trait is not necessarily the slowest dispersal rate anymore.

6.1 Monotonicity of the effective Hamiltonian

The effective Hamiltonian can be written as

$$\lambda(z_1, z_2) = \Lambda(\alpha(z_1), \alpha(z_2)), \quad (6.1)$$

where $\Lambda(\alpha_1, \alpha_2)$ is the smallest eigenvalue of

$$\begin{cases} \alpha_1 \Delta \Phi + (m(x) - \Theta_{\alpha_2}(x))\Phi + \Lambda \Phi = 0 & \text{for } x \in D, \\ \partial_\nu \Phi = 0 & \text{for } x \in \partial D, \end{cases} \quad (6.2)$$

where $\Theta_\alpha(x)$ is the unique positive solution to

$$\alpha\Delta\Theta + (m(x) - \Theta)\Theta = 0 \quad \text{in } D, \quad \text{and } \partial_\nu\Theta = 0 \quad \text{on } \partial D. \quad (6.3)$$

Then it is well-known that Λ is smooth in $\mathbb{R}_+ \times \mathbb{R}_+$, and that $\partial_{\alpha_1}\Lambda(\alpha_1, \alpha_2) > 0$ [1, 15, 31]. We may conclude the following immediately.

$$\text{sgn}(\lambda_{z_1}(z_1, z_2)) = \text{sgn}(\alpha'(z_1)) \quad \text{for all } z_1, z_2. \quad (6.4)$$

Next, we observe that when $\alpha(z)$ satisfies **(H1)**, it is necessarily U-shaped and has a unique minimum point, which is where the dominant trait eventually converges.

Lemma 6.1. *Suppose $\alpha(z)$ is chosen such that **(H1)** holds, then there exists $z_{\min} \in \text{Int } I$ such that*

$$\alpha'(z) < 0 \quad \text{in } [a, z_{\min}), \quad \alpha'(z_{\min}) = 0 \quad \text{and} \quad \alpha'(z) > 0 \quad \text{in } (z_{\min}, b]; \quad (6.5)$$

and that $z_1 \mapsto \lambda(z_1, z_2)$ has the same minimum point z_{\min} for each $z_2 \in [a, b]$,

$$\partial_{z_1}\lambda(z, z_2) < 0 \quad \text{in } [a, z_{\min}), \quad \partial_{z_1}\lambda(z_{\min}, z_2) = 0 \quad \text{and} \quad \partial_{z_1}\lambda(z, z_2) > 0 \quad \text{in } (z_{\min}, b]. \quad (6.6)$$

Proof. Differentiating (6.1) with respect to z_1 , evaluating at $(z_1, z_2) = (a, a)$, and using **(H1)**, we have

$$0 > \partial_{z_1}\lambda(a, a) = \partial_{\alpha_1}\Lambda(\alpha(a), \alpha(a))\alpha'(a).$$

Using also that $\partial_{\alpha_1}\Lambda > 0$, we conclude that $\alpha'(a) < 0$ and similarly $\alpha'(b) > 0$. Thus α has at least one interior minimum point. Moreover, for each fixed z_2 , the mapping

$$z_1 \mapsto \lambda(z_1, z_2) = \Lambda(\alpha(z_1), \alpha(z_2))$$

is convex in z_1 , it has the desired monotonicity property in z_1 . This proves the second statement. Still because $\partial_{\alpha_1}\Lambda > 0$, we deduce that α also has the desired monotonicity as well. \square

6.2 An explicit example of $\alpha(z)$

We can construct explicitly some U-shaped dispersion $\alpha(z)$, so that the associated $\lambda(z_1, z_2)$ satisfies the convexity assumption **(H1)**. For any given interval $I_0 = [\alpha_0, \alpha_0 + L_0]$ with $\alpha_0, L_0 > 0$, define

$$\alpha(z) = \alpha_0 + \frac{1}{k_0} \int_0^z \tan z' dz' \quad \text{for } z \in [-z_M, z_M],$$

where $k_0 := \sup_{(\alpha_1, \alpha_2) \in I_0 \times I_0} \frac{|\partial_{\alpha_1}^2 \Lambda|}{\partial_{\alpha_1} \Lambda}$ and $z_M \in (0, \pi/2)$ is the unique number such that $\int_0^{z_M} \tan z dz = k_0 L_0$. With this choice of α , the associated $\lambda(z_1, z_2) = \Lambda(\alpha(z_1), \alpha(z_2))$ satisfies

$$\pm \partial_{z_1}\lambda(\pm z_M, z_2) = \pm \partial_{\alpha_1}\Lambda(\alpha(\pm z_M), \alpha(z_2))\alpha'(\pm z_M) > 0,$$

and, thanks to the definition of k_0 , we compute

$$\begin{aligned} \partial_{z_1}^2 \lambda(z_1, z_2) &\geq (\partial_{\alpha_1}\Lambda)\alpha''(z_1) - |\partial_{\alpha_1}^2 \Lambda|(\alpha'(z_1))^2 \\ &= \frac{\partial_{\alpha_1}\Lambda}{k_0} \left[(\tan z_1)' - \frac{1}{k_0} \frac{|\partial_{\alpha_1}^2 \Lambda|}{\partial_{\alpha_1}\Lambda} |\tan z_1|^2 \right] \\ &\geq \frac{\partial_{\alpha_1}\Lambda}{k_0} \left[(\tan z_1)' - |\tan z_1|^2 \right] \geq \frac{1}{k_0} \inf_{I_0 \times I_0} \partial_{\alpha_1}\Lambda > 0. \end{aligned}$$

6.3 Evolution of conditional dispersal

A limitation in our present study comes from the use of convexity to obtain various regularity results, in particular for the solutions of the constrained Hamilton-Jacobi equation. Here we present an example which motivates to look for more general methods. We have in mind the following model considering evolution of conditional dispersal, see [18] and the references therein.

$$\epsilon \partial_t n_\epsilon = \alpha(z) \Delta_x u - \beta(z) \nabla \cdot [n_\epsilon \nabla m(x)] + n_\epsilon(x)(m(x) - \rho_\epsilon(x, t)) + \epsilon^2 \partial_z^2 n_\epsilon \quad (6.7)$$

for $x \in D$, $z \in I$ and $t > 0$, with appropriate boundary conditions. Here α is the rate of unconditional dispersal, whereas β is the rate of the directed movement up the gradient of the prescribed function $m(x)$. One or both α, β can be dependent on the trait variable z .

The effective Hamiltonian is again given by the invasion exponent $\lambda(z_1, z_2)$ and can be similarly defined. Assuming $m(x) = x$, $\alpha(z) = z$ and $\beta = q$ for some small constant q , it can be shown that (6.7) possesses at least one positive equilibrium solution \tilde{u}_ϵ , which tends to a Dirac measure supported at two distinct points on the trait interval. In particular, the corresponding effective Hamiltonian is nonconvex. We conjecture that the time dependent problem supports moving Dirac-concentrations supported at two points $(\bar{z}_1(t), \bar{z}_2(t))$, which then converges to their equilibrium position.

Acknowledgment. K.-Y. L. is partially supported by National Science Foundation under grant DMS-1853561. K.-Y. L. and B.P. have received funding from the European Research Council (ERC) under the European Union's Horizon 2020 research and innovation programme (grant agreement No 740623). Part of this research is carried out while K.-Y. L. and B.P. are visiting the Institute for Mathematical Sciences at the National University of Singapore, and the Institut Henri Poincaré. The authors acknowledge support of the (UAR 839 CNRS-Sorbonne Université), and LabEx CARMIN (ANR-10-LABX-59-01) for the visit at IHP.

A A priori Estimates

Proposition A.1. *There exists $\hat{C}_1 > 1$, depending on $V_0(z), m(x), \underline{\alpha}, \bar{\alpha}, D$ but independent of ϵ , such that*

$$\frac{1}{\hat{C}_1} \leq \rho_\epsilon(x, t) \leq \hat{C}_1 \quad \text{for } x \in D, t \geq 0.$$

Proof. We prove the proposition in six steps.

Step 1. There exists C such that $\sup_{t \geq 0} \|\rho_\epsilon(\cdot, t)\|_{L^1(D)} \leq C$.

Integrate (1.6) over $(x, z) \in D \times I$, and use the Cauchy-Schwarz inequality, we obtain

$$\epsilon \frac{d}{dt} \int_D \rho_\epsilon(x, t) dx = \int_D \rho_\epsilon(x, t)(m(x) - \rho_\epsilon(x, t)) dx \leq \left(\int_D \rho_\epsilon dx \right) \left(m^* - \frac{1}{|D|} \int_D \rho_\epsilon dx \right) \quad (\text{A.1})$$

where $m^* = \sup_D m$. Hence we deduce from the differential inequality that

$$\int_D \rho_\epsilon(x, t) dx \leq \max \left\{ \int_D \rho_\epsilon(x, 0) dx, m^* |D| \right\} \quad \text{for all } t \geq 0. \quad (\text{A.2})$$

It remains to estimate the initial total population $\int_D \rho_\epsilon(x, 0) dx$ by **(H2)**:

$$\begin{aligned} \int_D \rho_\epsilon(x, 0) dx &= \int_D \int_I \exp\left(-\frac{u_\epsilon(x, z, 0)}{\epsilon}\right) dz dx \\ &\leq \frac{C}{\sqrt{\epsilon}} \int_I \exp\left(-\frac{V_0(z)}{\epsilon}\right) dz \leq \frac{C}{\sqrt{\epsilon}} \int_{\mathbb{R}} \exp\left(-\frac{K_1 |z - \bar{z}_0|^2}{\epsilon}\right) dz \leq C \sqrt{\frac{\pi}{K_1}}. \end{aligned}$$

Thus Step 1 is a direct consequence of (A.2).

Step 2. There exists C such that $\sup_{t \geq \epsilon} \|\rho_\epsilon(\cdot, t)\|_{C(\bar{D})} \leq C$.

It suffices to show the following assertion:

Claim A.2. *There exists C_1 such that for any $t_1 \geq \epsilon$,*

$$\sup_{x \in D} \rho_\epsilon(x, t_1) \leq C_1 \sup_{t \in [t_1 - \epsilon, t_1]} \int_D \rho_\epsilon(x, t) dx.$$

To prove the claim, we first extend $n_\epsilon(x, z, t)$ in the z variable by reflection across $z = b$, and then periodically in z to $D \times \mathbb{R} \times [0, \infty)$.

Consider, for each $(z_1, t_1) \in I \times [\epsilon, \infty)$, the rescaled function

$$N_\epsilon(x, y, \tau; z_1, t_1) := n_\epsilon(x, z_1 + \epsilon y, t_1 + \epsilon \tau), \quad (\text{A.3})$$

then N_ϵ satisfies, with $z = z_1 + \epsilon y$,

$$\begin{cases} \partial_t N_\epsilon - \alpha(z) \Delta_x N_\epsilon - \partial_y^2 N_\epsilon = N_\epsilon(m - \rho_\epsilon) \leq m N_\epsilon & \text{in } D \times \mathbb{R} \times [-1, \infty), \\ \partial_\nu N_\epsilon(x, y, \tau) = 0 & \text{on } \partial D \times \mathbb{R} \times [-1, \infty). \end{cases} \quad (\text{A.4})$$

Since N_ϵ is a subsolution of a linear, parabolic equation with L^∞ bounded coefficients⁵, we can apply the local maximum principle [26, Theorem 7.36] (see also [10, Section 6.2]), to obtain a constant C_1 independent of $\epsilon > 0$, $z_1 \in I$ and $t_1 \geq \epsilon$ such that

$$\|N_\epsilon(x, y, \tau)\|_{L^\infty(D \times (-4/5, 4/5) \times (-4/5, 0))} \leq C_1 \|N_\epsilon(x, y, \tau)\|_{L^1(D \times (-1, 1) \times (-1, 0))}. \quad (\text{A.5})$$

(Note that the spatial domain on both sides of the inequality can be taken to be the same, as a consequence of the Neumann boundary condition across ∂D .) Next, we write

$$\rho_\epsilon(x, t_1) = \int_I n_\epsilon(x, z_1, t_1) dz_1 = \int_I N_\epsilon(x, 0, 0; z_1, t_1) dz_1.$$

Taking supremum in $x \in D$, it follows that

$$\begin{aligned} \|\rho_\epsilon(\cdot, t_1)\|_{L^\infty(D)} &\leq \int_I \|N_\epsilon(\cdot, 0, 0; z_1, t_1)\|_{L^\infty(D)} dz_1 \\ &\leq \int_I \|N_\epsilon(x, y, \tau; z_1, t_1)\|_{L^\infty(D \times (-1/2, 1/2) \times (-1/2, 0))} dz_1 \\ &\leq C \int_I \|N_\epsilon(x, y, \tau; z_1, t_1)\|_{L^1(D \times (-1, 1) \times (-1, 0))} dz_1 \\ &\leq C \int_{-1}^0 \int_D \int_{(a-\epsilon, b+\epsilon)} n_\epsilon(x, z, t_1 + \epsilon \tau) dz dx d\tau \\ &\leq 2C \int_{-1}^0 \int_D \int_I n_\epsilon(x, z, t_1 + \epsilon \tau) dz dx d\tau \\ &\leq 2C \sup_{t \in [t_1 - \epsilon, t_1]} \int_D \rho_\epsilon(x, t) dx, \end{aligned}$$

where we used the periodicity of n_ϵ in the second to last inequalities. This proves Claim A.2.

Finally, we take supremum over $t_1 \geq \epsilon$ on both sides of the conclusion of Claim A.2, we deduce

$$\sup_{t \geq \epsilon} \|\rho_\epsilon(\cdot, t_1)\|_{L^\infty(D)} \leq C_1 \sup_{t_1 \geq \epsilon} \left[\sup_{t \in [t_1 - \epsilon, t_1]} \int_D \rho_\epsilon(x, t) dx \right] \leq C_1 \sup_{t \geq 0} \int_D \rho_\epsilon(\cdot, t) dx \leq C.$$

⁵Note that the term $-\rho_\epsilon$ is dropped.

This completes Step 2.

Step 3. There exists $C > 1$ such that $C^{-1} \leq \rho_\epsilon(x, t) \leq C$ for $x \in D$ and $t \in [0, \epsilon]$.

Based on **(H2)** we construct the following subsolution of (4.1):

$$\underline{U}(z, t) := V_0(z) - \frac{1}{2}\epsilon |\log \epsilon| - C_2(t + \epsilon) - \frac{C}{\epsilon} \{ [z - (b - \sqrt{\epsilon})]_+^3 + [a + \sqrt{\epsilon} - z]_+^3 \},$$

where C is chosen large such that $\underline{U}_z(b, t) \leq 0 \leq \underline{U}_z(a, t)$, and then C_2 is chosen⁶ large enough so that

$$\underline{U}(z, 0) \leq u_\epsilon(x, z, 0) \quad \text{for } x \in D, z \in I,$$

and the appropriate differential inequality (4.1) is satisfied in $D \times I \times (0, \infty)$. Hence for $0 \leq t \leq \epsilon$,

$$u_\epsilon(x, z, t) \geq \underline{U}(z, t) \geq \frac{1}{2}V_0(z) - \frac{1}{2}\epsilon |\log \epsilon| + O(\epsilon),$$

where the last inequality follows from the fact that $V_0(z) \geq 0$ and is bounded below by a positive constant near $z = a, b$, so that

$$\frac{1}{2}V_0(z) - \frac{C}{\epsilon} \{ [z - (b - \sqrt{\epsilon})]_+^3 + [a + \sqrt{\epsilon} - z]_+^3 \} \geq 0.$$

Integrating in z , we find

$$\rho_\epsilon(x, t) \leq \frac{C}{\sqrt{\epsilon}} \int_I \exp\left(-\frac{V_0(z) + O(\epsilon)}{\epsilon}\right) dz \leq C \quad \text{for } x \in D, t \in [0, \epsilon].$$

This proves the upper bound of Step 3. The lower bound can then be similarly proved, by using the upper bound and considering the upper solution⁷

$$\bar{U}(z, t) := V_0(z) - \frac{1}{2}\epsilon |\log \epsilon| + C_3(t + \epsilon).$$

Notice that, here, the term in curly bracket appearing in the definition of \underline{U} is not needed, as $V_0(z)$ has positive outer derivatives at $z \in \partial I$. This completes the proof of Step 3.

Step 4. There exists C_0 such that $\sup_{t \geq 0} \|\rho_\epsilon(\cdot, t)\|_{C(\bar{D})} \leq C_0$.

Step 4 is an immediate consequence of Steps 2 and 3.

Step 5. There exists C such that $\inf_{t \geq 0} \int_D \rho_\epsilon(x, t) dx \geq 1/C$.

We may assume that

$$\inf_{0 \leq t \leq \epsilon} \int_D \rho_\epsilon(x, t) dx > e^{-C_0} \frac{\inf_D m}{2C_1}$$

where C_0 is given in Step 4, and C_1 is given in Claim A.2. Indeed, by Step 3, such an inequality holds if we increase C_0 when necessary. Next, we assume to the contrary that there exist $t_1 > \epsilon$ such that

$$\int_D \rho_\epsilon(x, t_1) dx = e^{-C_0} \frac{\inf_D m}{2C_1} \quad \text{and} \quad \frac{d}{dt} \int_D \rho_\epsilon dx \Big|_{t=t_1} \leq 0. \quad (\text{A.6})$$

By Step 4 and (A.1), the function $A(t) := \int_D \rho_\epsilon(x, t) dx$ satisfies the differential inequality $\epsilon \frac{d}{dt} A(t) \geq -C_0 A(t)$, so that for $t \in [t_1 - \epsilon, t_1]$,

$$A(t) \leq e^{C_0(t_1-t)/\epsilon} A(t_1) \leq e^{C_0} e^{-C_0} \frac{\inf_D m}{2C_1} = \frac{\inf_D m}{2C_1}.$$

⁶Since the differential inequality obtained after dropping from (4.1) the $-\rho_\epsilon$ term, is independent of $-\rho_\epsilon$, the constant C_2 can be chosen independent of $\sup_{0 \leq t \leq \epsilon} \|\rho_\epsilon(\cdot, t)\|_{C(\bar{D})}$.

⁷Here the constant C_3 depends on the quantity $\sup_{0 \leq t \leq \epsilon} \|\rho_\epsilon(\cdot, t)\|_{C(\bar{D})}$, which has just been proved to be uniformly bounded.

By Claim A.2, we deduce that

$$\|\rho_\epsilon(\cdot, t_1)\|_{L^\infty(D)} \leq C_1 \sup_{t \in [t_1 - \epsilon, t_1]} A(t) \leq C_1 \frac{\inf_D m}{2C_1} = \frac{\inf_D m}{2}.$$

Hence, by (A.1),

$$\epsilon \frac{d}{dt} \int_D \rho_\epsilon dx \Big|_{t=t_1} = \int_D (m - \rho_\epsilon) \rho_\epsilon dx \Big|_{t=t_1} \geq \frac{\inf_D m}{2} A(t_1) > 0,$$

which is a contradiction to (A.6). This proves Step 5.

Step 6. There exists C such that $\rho_\epsilon(x, t) \geq 1/C$ for $x \in \bar{D}$ and $t \geq 0$.

For $0 \leq t \leq \epsilon$, the lower bound is proved in Step 3. For $t \geq \epsilon$, one first notices that $N_\epsilon(\tau, x, y; t_1, z_1)$ satisfies a parabolic equation with L^∞ bounded coefficients, and thus satisfies a weak Harnack inequality [26, Theorem 7.37] (see also [10, Section 6.5]). One then proves that $\rho_\epsilon(x, t)$ also satisfies such a weak Harnack inequality, so that for each $t_1 \geq \epsilon$,

$$\rho_\epsilon(x, t_1) \geq C \int_{t_1 - \epsilon}^{t_1 - \epsilon/2} \int_D \rho_\epsilon(x, t) dx dt,$$

where the last term is bounded from below, as proved in Step 5. This proves Step 6. Finally, the proposition follows from combining Steps 4 and 6. \square

B Differentiability of the Principal Floquet Bundle

The notion of a normalized principal Floquet bundle (see [35]) is a generalization to non-autonomous parabolic problems of the notion of principal eigenfunction of an elliptic, or periodic-parabolic operator. Its smooth dependence on parameters is recently established in [8].

B.1 The normalized principal Floquet bundle

Let $D \subset \mathbb{R}^N$ be a smooth bounded domain. Given $\alpha > 0$ and $c \in C^{\beta, \beta/2}(\bar{D} \times \mathbb{R})$, we say that the positive function $\phi_1(x, t)$ is the corresponding principal Floquet bundle if it satisfies

$$\begin{cases} \partial_t \phi_1 - \alpha \Delta \phi_1 - c(x, t) \phi_1 = H_1(t) \phi_1 & \text{for } x \in D, t \in \mathbb{R}, \\ \partial_\nu \phi_1(x, t) = 0 & \text{for } x \in \partial D, t \in \mathbb{R}, \\ \phi_1(x, t) > 0 & \text{for } x \in \bar{D}, t \in \mathbb{R}. \end{cases}$$

The existence and uniqueness is proved in [28], which is based on the abstract result of [35].

To formulate the smooth dependence on parameters, we need the notion of a normalized principal Floquet bundle.

Definition B.1. Given $\alpha > 0$ and $c \in C^{\beta, \beta/2}(\bar{D} \times \mathbb{R})$, we say that the pair $(\Phi_1(x, t), H_1(t))$ is the corresponding normalized principal Floquet bundle if it satisfies

$$\begin{cases} \partial_t \Phi_1 - \alpha \Delta \Phi_1 - c(x, t) \Phi_1 = H_1(t) \Phi_1 & \text{for } x \in D, t \in \mathbb{R}, \\ \partial_\nu \Phi_1(x, t) = 0 & \text{for } x \in \partial D, t \in \mathbb{R}, \\ \int_\Omega \Phi_1(x, t) dx \equiv 1 & \text{for } t \in \mathbb{R}, \\ \Phi_1(x, t) > 0 & \text{for } x \in \bar{D}, t \in \mathbb{R}. \end{cases} \quad (\text{B.1})$$

Theorem B.2. For each $\alpha > 0$ and $c \in C^{\beta, \beta/2}(\bar{D} \times \mathbb{R})$, there exists a unique pair

$$(\Phi_1(x, t), H_1(t)) \in C^{2+\beta, 1+\beta/2}(\bar{D} \times \mathbb{R}) \times C^{\beta/2}(\mathbb{R})$$

satisfying (B.1) in classical sense.

Proof. The existence and uniqueness of $(\Phi_1, H_1(t))$ follows from the existence of the principal Floquet bundle $\phi(x, t)$, by noting that $H_1(t)$ arises from the normalization $\int_{\Omega} \Phi_1(x, t) dx \equiv 1$; See [8, Theorem A.1] for detail. \square

We need the smooth dependence of the normalized principal Floquet bundle, which is recently established proved in [8].

Proposition B.3. *The normalized principal Floquet bundle, as a mapping*

$$\begin{aligned} (\alpha, c) &\mapsto (\Phi_1, H_1) \\ \mathbb{R}_+ \times C^{\beta, \beta/2}(\bar{\Omega} \times \mathbb{R}) &\rightarrow C^{2+\beta, 1+\beta/2}(\bar{\Omega} \times \mathbb{R}) \times C^{\beta/2}(\mathbb{R}) \end{aligned}$$

is smooth. In particular, there exists a constant $C = C(M)$ which is independent of $z \in [\frac{1}{M}, M]$ and $\|c\|_{C^{\beta, \beta/2}(\bar{D} \times \mathbb{R})} \leq M$ such that

$$\|\partial_x \Phi_1\|_{\infty} + \max_{i=0,1,2,3} \|\partial_z^i \Phi_1\|_{\infty} \leq C,$$

where $\|\cdot\|_{\infty}$ is the L^{∞} norm over $(x, z, t) \in D \times [\frac{1}{M}, M] \times \mathbb{R}$, and

$$\frac{1}{C} \leq \Phi_1(x, t) \leq C \quad \text{in } D \times \mathbb{R}.$$

Proof. The smooth dependence is proved in [8, Proposition A.4]. It remains to prove the positive upper and lower bounds on Φ_1 . For this purpose, we recall the uniform Harnack inequality [21, Theorem 2.5], which says that there exists some positive constant $C = C(M)$ such that

$$\sup_D \Phi_1(\cdot, t) \leq C \inf_D \Phi_1(\cdot, t) \quad \text{for } z \in \left[\frac{1}{M}, M\right], t \in \mathbb{R}.$$

Thanks to the normalization $\int_{\Omega} \Phi_1 dx \equiv 1$, we obtain

$$\frac{1}{C|\Omega|} \leq \frac{1}{C} \sup_D \Phi_1(\cdot, t) \leq \Phi_1(x, t) \leq C \inf_D \Phi_1(\cdot, t) \leq \frac{C}{|\Omega|}.$$

This completes the proof. \square

C Uniqueness for the Constrained Hamilton Jacobi Equation

We now establish the uniqueness of solutions to a constrained Hamilton-Jacobi equation in an open, bounded one-dimensional interval I under some monotonicity assumption. We begin with a proposition that does not assume convexity of the Hamiltonian and initial data in the trait variable z .

Proposition C.1. *For $i = 1, 2$, let $(V_i, \bar{z}_i) \in W^{1, \infty}(I' \times [0, T]) \times BV([0, T])$ be a solution to*

$$\begin{cases} \partial_t V + |\partial_z V|^2 + R(z, \bar{z}(t), t) = 0 & \text{for } z \in I', t \in [0, T], \\ \partial_z V(z, t) = 0 & \text{for } z \in \partial I', t \in [0, T] \end{cases} \quad (\text{C.1})$$

in the viscosity sense, and which verifies the initial data and the constraint

$$\begin{cases} V(z, 0) = V_0(z) & \text{for } z \in I', \\ \inf_{z \in I'} V(z, t) = 0 & \text{for } t \in [0, T] \end{cases} \quad (\text{C.2})$$

in the classical sense. Suppose that R is C^2 in all variables, and

$$\bar{z}_i(t) \in \text{Int } I' \quad \text{for } i = 1, 2, t \in [0, T], \quad (\text{C.3})$$

$$R(z_1, z_2, t) = 0 \quad \text{if and only if} \quad (z_1, z_2, t) \in \Gamma \quad (\text{C.4})$$

and

$$\partial_{z_1} R(z, z, t) > 0 \quad \text{in } \Gamma \quad \text{or} \quad \partial_{z_1} R(z, z, t) < 0 \quad \text{in } \Gamma, \quad (\text{C.5})$$

where $\Gamma := \{(z_1, z_2, t) \in I' \times I' \times [0, T] : z_1 = z_2\}$. Then

$$(V_1(z, t), \bar{z}_1(t)) = (V_2(z, t), \bar{z}_2(t)) \quad \text{for } z \in I' \text{ and } t \in [0, T].$$

We postpone the proof of this proposition and conclude the proofs of Propositions 1.3 and 3.2.

C.1 Proofs of Propositions 1.3 and 3.2

Proof of Proposition 1.3. Since assertions (ii) and (iii), and the existence part of assertion (i) are proved in [27, 30], we prove the uniqueness part of assertion (i) in the following.

Let $(V_i(z, t), \bar{z}_i(t))$ be two solutions such that the common initial data $V_0(z)$ is convex, smooth, and attains a unique minimum at some $\bar{z}_0 \in \text{Int } I$.

Step 1. Using (6.4) and the differential equation (1.11), we see that

$$\text{sgn} \left(\frac{d}{dt} \bar{z}_i(t) \right) = -\text{sgn} (\alpha'(\bar{z}_i(t))).$$

Since $\alpha'(\inf I) < 0 < \alpha'(\sup I)$ (thanks to **(H1)**), it follows that $\frac{d}{dt} \bar{z}_i(t)$ retains the same sign as $-\alpha'(\bar{z}_i(0))$ and that sign does not depend on i . Therefore, exactly one of the following cases holds:

$$(i) \quad \bar{z}_1(t) \equiv \bar{z}_2(t) \equiv \bar{z}_0, \quad (ii) \quad \left(\frac{d}{dt} \bar{z}_1(t) \right) \left(\frac{d}{dt} \bar{z}_2(t) \right) > 0 \quad \text{for all } t \in [0, T].$$

In case (i), the conclusion follows from standard uniqueness of viscosity solution to Neumann problem. We henceforth consider case (ii). In fact, by (1.11) we can assume without loss of generality that

$$\frac{d}{dt} \bar{z}_i(t) > 0 \quad \text{and} \quad \partial_{z_1} \lambda(\bar{z}_i(t), \bar{z}_i(t)) < 0 \quad \text{for } t \in [0, T] \text{ and } i = 1, 2. \quad (\text{C.6})$$

Step 2. We choose I' to be slightly larger than $[\bar{z}_0, \max_{i=1,2} \bar{z}_i(T)] = \cup_{i=1}^2 \bar{z}_i([0, T])$, then (C.3) and (C.4) hold.

Step 3. $(V_1(z, t), \bar{z}_1(t))$ and $(V_2(z, t), \bar{z}_2(t))$ are viscosity solutions to the same constrained Hamilton-Jacobi equation (with Neumann boundary conditions) on the restricted domain $I' \times [0, T]$:

$$\begin{cases} \partial_t V + |\partial_z V|^2 - \lambda(z, \bar{z}(t)) = 0 & \text{for } z \in I', t \in [0, T], \\ \partial_z V(z, t) = 0 & \text{for } z \in \partial I', t \in [0, T], \\ V(z, 0) = V_0(z) & \text{for } z \in I', \\ \inf_{z \in I'} V(z, t) = 0 & \text{for } t \in [0, T]. \end{cases} \quad (\text{C.7})$$

This step is valid since $z \mapsto V_i(z, t)$ is convex, and the unique minimum point $\bar{z}_i(t) \in \text{Int } I'$ for all t . Since the Hamiltonian $H(t, z, p) = |z|^2 - \lambda(z, \bar{z}(t))$ is convex in p and the solutions V_i are strictly convex in z with the minimum attained in the interior, we may extend H and V_i by reflection across the boundary $\partial I'$ in z with V_i remain viscosity solution of $\partial_t V_i + H(t, z, \partial_z V_i) = 0$ in the extended domain. Hence, they satisfy the Neumann boundary condition in $\partial I'$ in viscosity sense. The initial data and constraint are satisfied in the classical sense.

Step 4. We can now apply Proposition C.1 to conclude that $\bar{z}_1(t) = \bar{z}_2(t)$ a.e. in $[0, T]$. Then we can use the variational characterization (see (C.9)) to deduce that $V_1 \equiv V_2$ in the original domain $I \times [0, T]$ (not just in the smaller domain $I' \times [0, T]$). \square

Proof of Proposition 3.2. Fix $\epsilon > 0$ and let $R(z_1, z_2, t) := -H_\epsilon(z_1, t) + H_\epsilon(z_2, t)$. Once again, the existence of a viscosity solution $(V_\epsilon(z, t), \bar{z}_\epsilon(t))$ holds. Since $V_0(z)$ and $-R(z, \bar{z}, t)$ are convex in z , it follows from [30] or similarly to Proposition 1.3 that $V_\epsilon(z, t)$ is strictly convex in z for each t , and satisfies the differential equation

$$\frac{d}{dt} \bar{z}_\epsilon(t) = \frac{1}{\partial_{zz} V_\epsilon(\bar{z}_\epsilon(t), t)} \partial_{z_1} R(\bar{z}_\epsilon(t), \bar{z}_\epsilon(t), t). \quad (\text{C.8})$$

To show uniqueness, it suffices to repeat the proof of Proposition 1.3. We omit the details. \square

C.2 Proof of Proposition C.1

Suppose two sets of solutions (V_i, \bar{z}_i) , $i = 1, 2$, are given. First, extend the problem by reflection to the domain $[2 \inf I' - \sup I', \sup I'] \times [0, T]$ and then extend it periodically so that it is defined in $\mathbb{R} \times [0, T]$. We use the variational characterization:

$$V_i(z, t) = \inf_{\gamma(t)=z} \left\{ \int_0^t \left[\frac{|\dot{\gamma}(s)|^2}{4} - R(\gamma(s), \bar{z}_i(s), s) \right] ds + V_0(\gamma(0)) \right\} \quad (\text{C.9})$$

with the understanding that $V_0(z)$ is also being extended evenly and then periodically so that it is defined for all $z \in \mathbb{R}$.

Lemma C.2. *Assume (C.3), (C.4) and the first alternative of (C.5). Let $(V(z, t), \bar{z}(t)) \in W^{1,\infty}(\mathbb{R} \times [0, T]) \times \text{BV}([0, T])$ be a solution of (C.1) and (C.2). Then, at possible discontinuity points of $\bar{z}(\cdot)$ we have*

$$\bar{z}(t-) \leq \bar{z}(t+) \quad \text{and} \quad \mathcal{N}(t) \subset [\bar{z}(t-), \bar{z}(t+)] \quad (\text{C.10})$$

where $\mathcal{N}(t) = \{z \in \bar{I}' : V(z, t) = 0\}$.

Proof. By (C.4) and the first alternative of (C.5), we have

$$\text{sgn}(R(z_1, z_2, t)) = \text{sgn}(z_1 - z_2). \quad (\text{C.11})$$

Fix (z, t) such that $V(z, t) = 0$, it suffices to show that $\bar{z}(t-) \leq z \leq \bar{z}(t+)$. To show the first inequality, choose a minizing curve $\gamma(t)$ for (C.9) such that $\gamma(t) = z$ and

$$0 = V(z, t) = \int_0^t \left[\frac{|\dot{\gamma}(s)|^2}{4} - R(\gamma(s), \bar{z}(s), s) \right] ds + V_0(\gamma(0)). \quad (\text{C.12})$$

In fact, for any $h \in (0, t)$, the dynamic programming principle says that

$$0 \leq V(\gamma(t-h), t-h) = \int_0^{t-h} \left[\frac{|\dot{\gamma}(s)|^2}{4} - R(\gamma(s), \bar{z}(s), s) \right] ds + V_0(\gamma(0)).$$

Subtracting, we have

$$0 \leq - \int_{t-h}^t \left[\frac{|\dot{\gamma}(s)|^2}{4} - R(\gamma(s), \bar{z}(s), s) \right] ds \leq \int_{t-h}^t R(\gamma(s), \bar{z}(s), s) ds \quad \text{for all } h \in (0, t).$$

Dividing by h and letting $h \searrow 0$, we deduce that

$$R(z, \bar{z}(t-), t) = R(\gamma(t), \bar{z}(t-), t) \geq 0. \quad (\text{C.13})$$

By (C.11), we have $\bar{z}(t-) \leq z$.

Next, we fix as above $z, t, \gamma(\cdot)$, and define $\gamma_1 : [0, t+1] \rightarrow \mathbb{R}$ by

$$\gamma_1(s) = \begin{cases} \gamma(s) & \text{for } 0 \leq s \leq t, \\ z & \text{for } s > t. \end{cases}$$

Then by (C.9), we have for $0 < h < 1$,

$$0 \leq V(t+h, \gamma_1(t+h)) \leq \int_0^{t+h} \frac{|\dot{\gamma}_1(s)|^2}{4} - R(\gamma_1(s), \bar{z}(s), s) ds + V_0(\gamma(0)).$$

Subtracting (C.12) from the above, we have

$$0 \leq - \int_t^{t+h} R(z, \bar{z}(s), s) ds. \quad (\text{C.14})$$

Divide by h , and let $h \rightarrow 0$, then $R(z, \bar{z}(t+), s) \leq 0$. By (C.11), we have $\bar{z}(t+) \geq z$. \square

Remark C.3. From the above proof, for each $t_0 > 0$ and $\bar{z}_0 \in \mathcal{N}(t_0)$, it follows from (C.14) that

$$\int_{t_0}^{t_0+h} R(\bar{z}_0, \bar{z}(s), s) ds \leq 0 \quad \text{for all sufficiently small } h > 0.$$

Lemma C.4. *Assume (C.3), (C.4) and the first alternative of (C.5). Let $(V(z, t), \bar{z}(t)) \in W^{1,\infty}(\mathbb{R} \times [0, T]) \times \text{BV}([0, T])$ be a solution of (C.1) and (C.2), then \bar{z} is non-decreasing. Furthermore, we have*

$$\lim_{t \rightarrow 0^+} [\sup \mathcal{N}(t)] = \sup \mathcal{N}(0), \quad (\text{C.15})$$

and, still with $\mathcal{N}(t) = \{z \in \bar{I}' : V(z, t) = 0\}$, we have

$$\bar{z}(t+) = \sup \mathcal{N}(t) \quad \text{for all } t \quad (\text{C.16})$$

Proof. Without loss of generality, we choose the right-continuous representative of \bar{z} . Assume to the contrary that \bar{z} is not non-decreasing. i.e. $\bar{z}(t_1) > \bar{z}(t_2)$ for some $t_1 < t_2$. Since \bar{z} is right-continuous, we have $\bar{z}(t_1) > \bar{z}(t_2+)$, i.e. there exists $t_3 > t_2$ such that $\bar{z}(t_1) > \bar{z}(t)$ for all $t \in [t_2, t_3]$. Let $t_0 = \sup\{t \in [t_1, t_3] : \bar{z}(t) \geq \bar{z}(t_1)\}$. Then $t_0 \leq t_2 < t_3$, and

$$\bar{z}(t) < \bar{z}(t_1) \leq \bar{z}(t_0-) \quad \text{for } t \in (t_0, t_3). \quad (\text{C.17})$$

In particular, we deduce that $\bar{z}(t_0+) \leq \bar{z}(t_0-)$. By (C.10), it follows that $\bar{z}(t_0+) = \bar{z}(t_0-)$, and that $\mathcal{N}(t_0)$ consists of a single element, which we denote by \bar{z}_0 .

By the first alternative of (C.5), there exists a small $\delta_1 > 0$ such that

$$R(\bar{z}_0, z, t) > 0 \quad \text{for } z \in (\bar{z}_0 - \delta_1, \bar{z}_0), t \in [t_0 - \delta_1, t_0 + \delta_1]. \quad (\text{C.18})$$

Then choose $h \in (0, \delta_1)$ small enough so that $\bar{z}(s) \in (\bar{z}_0 - \delta_1, \bar{z}_0)$ for all $s \in (t_0, t_0 + h)$ (which is guaranteed by $\bar{z}(t_0+) = \bar{z}(t_0-) = \bar{z}_0$ and (C.17)). We deduce from (C.18) that

$$\int_{t_0}^{t_0+h} R(\bar{z}_0, \bar{z}(s), s) ds > 0,$$

where the last inequality follows from $\bar{z}(s) < \bar{z}_0$ (by (C.17)). We obtain a contradiction to Remark C.3. Thus \bar{z} is non-decreasing.

Next, we observe that (C.15) follows directly from the upper-semicontinuity of \mathcal{N} , i.e.

$$\limsup_{t' \rightarrow t^+} \mathcal{N}(t') \subset \mathcal{N}(t). \quad (\text{C.19})$$

It remains to show (C.16). It follows from (C.10) (proved in Lemma C.2) that $\bar{z}(t+) \geq \sup \mathcal{N}(t)$. Moreover, (C.10) and the monotonicity of \bar{z} imply that

$$\bar{z}(t+) \leq \bar{z}(t'-) \leq \inf \mathcal{N}(t') \leq \sup \mathcal{N}(t') \quad \text{for each } t' > t.$$

Using (C.19), we may let $t' \rightarrow t+$ to deduce $\bar{z}(t+) \leq \sup \mathcal{N}(t)$. This proves (C.16). \square

Proof of Proposition C.1. For $i = 1, 2$, let $(V_i(z, t), \bar{z}_i(t)) \in W^{1,\infty}(\mathbb{R} \times [0, T]) \times BV[0, T]$ be two solutions of (C.1) and (C.2). It suffices to show that $\bar{z}_1(t) = \bar{z}_2(t)$ a.e. in $[0, T]$. Without loss of generality, one can reduce to the case that for each $t > 0$, $\bar{z}_1 \neq \bar{z}_2$ in a set of positive measure in $(0, t)$. Furthermore, by considering $z' = -z$ if necessary, we can assume the first alternative of (C.5) to hold.

To apply [7, Section 3, Remark 2], it remains to verify the conditions **(U1)**-**(U3)** therein. Now, observe that the Hamiltonian function in (C.1) is smooth and satisfies $H(\bar{z}, t, z, p) = |p|^2 + R(z, \bar{z}, t)$, so that $L(\bar{z}, z, t, v) = |v|^2/4 - R(z, \bar{z}, t)$. This verifies **(U1)**. The condition **(U2)** also holds, as $V_i(z, t)$ admits the variational characterization (C.9).

It suffices to check **(U3)**. Define

$$z_i^t = \sup \mathcal{N}_i(t), \quad \text{where} \quad \mathcal{N}_i(t) = \{z \in I : V_i(z, t) = 0\},$$

and let $\gamma_i^t \in AC[0, t]$ be the minimizing path corresponding to the value $V_i(z_i^t, t) = 0$. We need to verify the following three conditions:

- (i) $\lim_{t \rightarrow 0+} z_i^t = \bar{z}_0 := \sup\{z \in I : V_0(z) = 0\}$,
- (ii) $\inf_{0 < \theta < 1} \partial_{z_2} R(\lim_{t \rightarrow 0+} [(1 - \theta)\bar{z}_1(t) + \theta\bar{z}_2(t)], \bar{z}_0, 0) < 0$,
- (iii) $\limsup_{t \rightarrow 0+} \|\dot{\gamma}_i^t\|_{L^\infty(0,t)} < +\infty$.

To verify condition (i), we apply (C.15) and (C.16) to get

$$\lim_{t \rightarrow 0+} \bar{z}_1(t) = \lim_{t \rightarrow 0+} \bar{z}_2(t) = \lim_{t \rightarrow 0+} z_i^t = \bar{z}_0. \quad (\text{C.20})$$

To verify condition (ii), observe that (C.4) and the first alternative of (C.5) imply that $\partial_{z_2} R(\bar{z}_0, \bar{z}_0, 0) = -\partial_{z_1} R(\bar{z}_0, \bar{z}_0, 0) < 0$. Using (C.20), we can then compute

$$\sup_{0 < \theta < 1} \partial_{z_2} R(\lim_{t \rightarrow 0+} [(1 - \theta)\bar{z}_1(t) + \theta\bar{z}_2(t)], \bar{z}_0, 0) = \partial_{z_2} R(\bar{z}_0, \bar{z}_0, 0) < 0.$$

For condition (iii), we observe that the initial data g and the Lagrangian function $\frac{|v|^2}{4} - R(I, t, x)$ are periodic in x , so the minimizing paths γ_i^t (corresponding to (z_i^t, t)) exists and is uniformly bounded in L^∞ . One can derive the regularity of $\dot{\gamma}_i^t$ by repeating the arguments in [7, Section 2].

Having verified **(U1)**-**(U3)**, one can then invoke [7, Section 3, Remark 2] to yield a contradiction. This proves that $\bar{z}_1(t) \equiv \bar{z}_2(t)$ a.e. in $[0, T]$. That $V_1(z, t) \equiv V_2(z, t)$ follows from the standard uniqueness results. \square

References

- [1] L. ALTENBERG, *Resolvent positive linear operators exhibit the reduction phenomenon*, Proc. Natl. Acad. Sci. USA, 109 (2012), pp. 3705–3710.
- [2] G. BARLES, S. MIRRAHIMI, AND B. PERTHAME, *Concentration in Lotka-Volterra parabolic or integral equations: a general convergence result*, Methods Appl. Anal., 16 (2009), pp. 321–340.
- [3] E. BOUIN AND C. HENDERSON, *Super-linear spreading in local bistable cane toads equations*, Nonlinearity, 30 (2017), pp. 1356–1375.
- [4] E. BOUIN, C. HENDERSON, AND L. RYZHIK, *The Bramson logarithmic delay in the cane toads equations*, Quart. Appl. Math., 75 (2017), pp. 599–634.

- [5] ———, *Super-linear spreading in local and non-local cane toads equations*, J. Math. Pures Appl. (9), 108 (2017), pp. 724–750.
- [6] E. BOUIN AND S. MIRRAHIMI, *A Hamilton-Jacobi approach for a model of population structured by space and trait*, Commun. Math. Sci., 13 (2015), pp. 1431–1452.
- [7] V. CALVEZ AND K.-Y. LAM, *Uniqueness of the viscosity solution of a constrained Hamilton-Jacobi equation*, Calc. Var. Partial Differential Equations, 59 (2020), pp. Paper No. 163, 22.
- [8] R. S. CANTRELL AND K.-Y. LAM, *On the evolution of slow dispersal in multispecies communities*, SIAM J. Math. Anal., 53 (2021), pp. 4933–4964.
- [9] N. CHAMPAGNAT, R. FERRIÈRE, AND S. MÉLÉARD, *Unifying evolutionary dynamics: From individual stochastic processes to macroscopic models*, Theoretical Population Biology, 69 (2006), pp. 297–321. ESS Theory Now.
- [10] Y.-Z. CHEN, *Second Order Parabolic Equations (in Chinese)*, Beijing University Mathematics series, Beijing University Press, Beijing, 2003.
- [11] L. DESVILLETES, P.-E. JABIN, S. MISCHLER, AND G. RAOUL, *On selection dynamics for continuous structured populations*, Commun. Math. Sci., 6 (2008), pp. 729–747.
- [12] U. DIECKMANN AND R. LAW, *The dynamical theory of coevolution: a derivation from stochastic ecological processes*, J. Math. Biol., 34 (1996), pp. 579–612.
- [13] O. DIEKMANN, *A beginner’s guide to adaptive dynamics*, in Mathematical modelling of population dynamics, vol. 63 of Banach Center Publ., Polish Acad. Sci. Inst. Math., Warsaw, 2004, pp. 47–86.
- [14] O. DIEKMANN, P.-E. JABIN, S. MISCHLER, AND B. PERTHAME, *The dynamics of adaptation: An illuminating example and a hamilton–jacobi approach*, Theoretical Population Biology, 67 (2005), pp. 257–271.
- [15] J. DOCKERY, V. HUTSON, K. MISCHAIKOW, AND M. PERNAROWSKI, *The evolution of slow dispersal rates: a reaction diffusion model*, J. Math. Biol., 37 (1998), pp. 61–83.
- [16] L. C. EVANS, *The perturbed test function method for viscosity solutions of nonlinear PDE*, Proc. Roy. Soc. Edinburgh Sect. A, 111 (1989), pp. 359–375.
- [17] S. GERITZ AND M. GYLLENBERG, *The mathematical theory of adaptive dynamics*, Cambridge University Press, Cambridge, (2008).
- [18] W. HAO, K.-Y. LAM, AND Y. LOU, *Concentration phenomena in an integro-PDE model for evolution of conditional dispersal*, Indiana Univ. Math. J., 68 (2019), pp. 881–923.
- [19] A. HASTINGS, *Can spatial variation alone lead to selection for dispersal?*, Theoretical Population Biology, 24 (1983), pp. 244–251.
- [20] C. HENDERSON, B. PERTHAME, AND P. E. SOUGANIDIS, *Super-linear propagation for a general, local cane toads model*, Interfaces Free Bound., 20 (2018), pp. 483–509.
- [21] J. HÚSKA, *Harnack inequality and exponential separation for oblique derivative problems on Lipschitz domains*, J. Differential Equations, 226 (2006), pp. 541–557.
- [22] P.-E. JABIN AND G. RAOUL, *On selection dynamics for competitive interactions*, J. Math. Biol., 63 (2011), pp. 493–517.

- [23] P.-E. JABIN AND R. S. SCHRAM, *Selection-mutation dynamics with spatial dependence*, 2016, preprint.
- [24] K.-Y. LAM, *Stability of Dirac concentrations in an integro-PDE model for evolution of dispersal*, Calc. Var. Partial Differential Equations, 56 (2017), pp. Paper No. 79, 32.
- [25] K.-Y. LAM AND Y. LOU, *An integro-PDE model for evolution of random dispersal*, J. Funct. Anal., 272 (2017), pp. 1755–1790.
- [26] G. M. LIEBERMAN, *Second order parabolic differential equations*, World Scientific Publishing Co., Inc., River Edge, NJ, 1996.
- [27] A. LORZ, S. MIRRAHIMI, AND B. PERTHAME, *Dirac mass dynamics in multidimensional nonlocal parabolic equations*, Comm. Partial Differential Equations, 36 (2011), pp. 1071–1098.
- [28] J. MIERCZYŃSKI, *Globally positive solutions of linear parabolic PDEs of second order with Robin boundary conditions*, J. Math. Anal. Appl., 209 (1997), pp. 47–59.
- [29] S. MIRRAHIMI AND B. PERTHAME, *Asymptotic analysis of a selection model with space*, J. Math. Pures Appl. (9), 104 (2015), pp. 1108–1118.
- [30] S. MIRRAHIMI AND J.-M. ROQUEJOFFRE, *A class of Hamilton-Jacobi equations with constraint: uniqueness and constructive approach*, J. Differential Equations, 260 (2016), pp. 4717–4738.
- [31] W.-M. NI, *The Mathematics of Diffusion*, vol. 82 of CBMS-NSF Regional Conference Series in Applied Mathematics, Society for Industrial and Applied Mathematics (SIAM), Philadelphia, PA, 2011.
- [32] S. NORDMANN AND B. PERTHAME, *Dynamics of concentration in a population structured by age and a phenotypic trait with mutations. Convergence of the corrector*, J. Differential Equations, 290 (2021), pp. 223–261.
- [33] B. PERTHAME AND G. BARLES, *Dirac concentrations in Lotka-Volterra parabolic PDEs*, Indiana Univ. Math. J., 57 (2008), pp. 3275–3301.
- [34] B. PERTHAME AND P. E. SOUGANIDIS, *Rare mutations limit of a steady state dispersal evolution model*, Math. Model. Nat. Phenom., 11 (2016), pp. 154–166.
- [35] P. POLÁČIK AND I. TEREŠČÁK, *Exponential separation and invariant bundles for maps in ordered Banach spaces with applications to parabolic equations*, J. Dynam. Differential Equations, 5 (1993), pp. 279–303.