



HAL
open science

Research progress on polysaccharide components of *Cistanche deserticola* as potential pharmaceutical agents

Na Cheng, Hao Wang, Huifang Hao, Faiz-Ur Rahman, Yongmin Zhang

► To cite this version:

Na Cheng, Hao Wang, Huifang Hao, Faiz-Ur Rahman, Yongmin Zhang. Research progress on polysaccharide components of *Cistanche deserticola* as potential pharmaceutical agents. *European Journal of Medicinal Chemistry*, 2023, 245 (Part 1), pp.114892. 10.1016/j.ejmech.2022.114892 . hal-03842136

HAL Id: hal-03842136

<https://hal.sorbonne-universite.fr/hal-03842136>

Submitted on 7 Nov 2022

HAL is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers.

L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.

Research progress on polysaccharide components of *Cistanche deserticola* as potential pharmaceutical agents

Na Cheng^a, Hao Wang^a, Huifang Hao^{a, b}, Faiz-Ur Rahman^a, Yongmin Zhang^{a, c, d, *}

^a Inner Mongolia University Research Center for Glycochemistry of Characteristic Medicinal Resources, School of Chemistry and Chemical Engineering, Inner Mongolia University, Hohhot, 010021, China.

^b School of Life Sciences, Inner Mongolia University, Hohhot, 010021, China.

^c Sorbonne Université, CNRS, Institut Parisien de Chimie Moléculaire, UMR 8232, 4 Place Jussieu, 75005 Paris, France.

^d Key Laboratory of Tropical Medicinal Resource Chemistry of Ministry of Education, College of Chemistry and Chemical Engineering, Hainan Normal University, Haikou, 571158, China.

Abstract

Cistanche deserticola is a traditional and precious Chinese herbal medicine, known as "desert ginseng", with anti-inflammatory, anti-oxidant, improving immunity, nourishing the kidneys and other pharmacological effects. Its chemical components mainly include phenylethanol glycosides, iridoids, polysaccharides and volatile components, among which polysaccharides have received extensive attention due to their biological activities such as regulating immune activity, anti-aging, anti-spleen deficiency and antitumor. In recent years, a large number of research have been carried out on the extraction and isolation, chemical structure analysis and biological activity of *Cistanche deserticola* polysaccharides. The methods of polysaccharide extraction mainly include traditional extraction method, ultrasonic assisted method, microwave assisted method and enzyme assisted method, etc. The extracted polysaccharides were analyzed by chemical methods including methylation, acid hydrolysis and Smith degradation and spectroscopy methods such as NMR and IR. A variety of polysaccharides with new structures were obtained, and some polysaccharides with known structures were also investigated for their biological activities and their structure-activity relationships. However, the relationship between polysaccharides structure and their biological activities is still unclear due to the large number of polysaccharide components, their complex structures and the lack of systematic research and analysis on them. It is expected that the subsequent study of polysaccharide structure and active conformational relationship will be highly valuable for the application of *Cistanche deserticola* in pharmaceutical sciences and health food.

Keywords: *Cistanche Deserticola*; Polysaccharides; Extraction and Isolation; Chemical Structure; Biological Activity; Research Progress; Pharmaceutical agents.

1. Introduction

Cistanche Hoffmg. Et Link is a perennial herbaceous parasitic plant of the Orobanchaceae family. There are about 27 species in the world, mainly distributed in the arid lands and deserts in the northern hemisphere and mainly concentrated in Inner Mongolia, Ningxia, Gansu, Xinjiang and Qinghai provinces in China ^[1]. A further study confirmed that four species and one variation of *Cistanche* distributed in China ^[2], i.e. *Cistanche deserticola* Y.C. Ma, *Cistanche tubulosa* (Schenk) R. Wight, *Cistanche salsa* (C.A. Mey.) G. Beck, *Cistanche sinensis* G. Beck and *Cistanche salsa* (C.A. Mey.) G. Beck var. *albiflora* P.F. Tu et Z.C. Lou (one variation of *Cistanche*).

The traditional Chinese medicine *Cistanche* consists of the dried succulent stems of *Cistanche* species of this genus, also known as Cunyun, Dayun, Goblin, etc. It is attached underground to the roots of the dicotyledonous plant *Haloxylon ammodendron* Bunge and develops by obtaining nutrients from the host plants that it parasitize. *Cistanche deserticola* is a rare and valuable Chinese herb in China ^[3] and honored as “Desert ginseng” attributing to its promising medical functions and nourishing value. It is proved that *Cistanche deserticola* Y.C. Ma which is mainly produced in the western part of Inner Mongolia are the authentic Chinese medicine *Cistanche* and documented in Chinese Pharmacopoeia ^[4]. Modern pharmacological research have disclosed that *Cistanche* possesses wide-reaching biological activities including anti-inflammatory, anti-oxidative, immunity-enhancing, anti-aging, nerve protection, memory-improving, prevention and treatment of osteoporosis, and anti-fatigue, etc ^[5-6]. Modern research has shown that the chemical components of *Cistanche deserticola* mainly include phenylethanoid glycosides, iridoid glycosides, lignans, polysaccharides, amino acid and volatile compounds, etc ^[7]. The active component that attracted more attention is reported to be phenylethanoid glycosides, which have significant effect in treating kidney deficiency ^[8] and protecting muscle cells ^[9] and so on.

In recent years, polysaccharide which obtained from natural herbs has been proved to possess multiple biological activities and lower cytotoxicity. Thus, polysaccharide as one of the main active ingredients of *Cistanche deserticola* has also received extensive attention, and it has been studied that polysaccharide has various pharmacological effects such as lowering blood sugar, blood lipids ^[10], regulating immunity ^[11] and anti-aging ^[12]. The biological activities of polysaccharides are closely related to their physicochemical properties, chemical composition and content of polysaccharides, etc. However, the relationship between the chemical structure of polysaccharides and their biological activities of *Cistanche deserticola* is unclear due to the complexity of the composition and structure of polysaccharides. Therefore, it is of great significance to study the structure-activity relationship of polysaccharides for the safety and efficacy of the use of *Cistanche deserticola*. The extraction and isolation process of polysaccharides is the first prerequisite for its in-depth study as it may affect their yield, chemical structure, and biological activities. Although these aspects have been reported in the literature ^[13], it is not sufficiently detailed and comprehensive, especially in terms of extraction, chemical structure and pharmaceutical application. The aim of this review is to introduce recent developments in the study of the extraction and isolation, chemical structure and biological activities of polysaccharides from *Cistanche deserticola*.

2. Extraction and Isolation

In the past, people mainly relied on traditional water extraction methods to obtain polysaccharides from *Cistanche deserticola*, which is not only time-consuming and labor-intensive, but the yield and purity of polysaccharides are not very advanced. With the increasing research interest in the polysaccharides of *Cistanche deserticola*, the extraction process of polysaccharides has been greatly developed. Various extraction methods including ultrasonic-assisted, microwave-assisted and enzyme-assisted have been continuously developed based on aqueous extraction process, with the aim of achieving high yield, energy-saving and less time consuming. Each extraction method has advantages and disadvantages, and the impact of these methods on the chemical structure and biological activity of polysaccharides should be considered comprehensively. These methods can be summarized in Table 1.

Table 1 *Cistanche deserticola* polysaccharide extraction method

Extraction Methods	Principle	Advantages	Disadvantages
Traditional Water Extraction	"Water extraction and alcohol precipitation"	Not require special operations, low cost	Time-consuming, labor-intensive, and there are more factors affecting the yield
Ultrasonic-Assisted Extraction	Ultrasonic waves generate strong cavitation and stirring effects, making polysaccharides easily diffuse and dissolve	Less solvent usage, shorter working time of extraction and lower extraction temperature	Prolonged ultrasonication may lead to the destruction of the polysaccharides structure
Microwave-Assisted Extraction	The microwave can generate strong pressure to break through cell membranes and cell walls, and promote the dissolution of polysaccharides	High efficiency, rapidity and energy saving	Use of microwave instruments is dangerous
Enzyme-Assisted Extraction	Biological enzymes effectively dissolve polysaccharides through the destruction of plant cell walls	The extraction conditions are mild and simple, less destructive to polysaccharide structure, and the extraction effect is significantly enhanced	Enzyme activity is unstable, high cost
Ultrasonic-Microwave Synergistic Extraction	-	Shorten the extraction time, obtain high-yield polysaccharides	Multiple methods are complicated to operate
Ultrasonic-Enzyme-Assisted Extraction	-	A fast and inexpensive method	Multiple methods are complicated to operate

2.1 Traditional Water Extraction Methods

The traditional extraction method utilizes the characteristics of traditional Chinese medicine

polysaccharides that are easily soluble in water but insoluble in organic solvents such as ethanol, petroleum ether, ether, etc., which is often referred to as "water extraction and alcohol precipitation" method. This method does not require special operations, however, it is time-consuming and labor-intensive, and there are more factors affecting the yield, such as extraction temperature, time, and solvent dosage. Naran and co-workers reported the subterranean parts of the holoparasite, *Cistanche deserticola*, was extracted with methanol and subsequently methanol-insoluble extracts was extracted using in succession cold water, hot water, 0.5 M NaOH and 0.01 M EDTA ^[14]. Yield analysis of the isolated polysaccharide fractions revealed that the yield of 0.5 M NaOH and 0.01 M EDTA extract was superior to that of cold and hot water. Naran and co-workers then described starch fractions have been isolated from the underground part of *Cistanche deserticola* Y. C. Ma using hot water and alkali treatments, respectively ^[15]. Chemical, enzymic and spectroscopic analyses revealed that the hot water and alkali extracts contained different proportions of amylose and amylopectin components. The yield of polysaccharides from the hot water extract was 1.3 %, while the yield of polysaccharides from the alkali extract was 2.9 %.

2.2 Ultrasonic-Assisted Extraction Methods

Ultrasonic-assisted extraction methods (UEM) utilizes ultrasonic waves to generate strong cavitation and stirring effects, making polysaccharides easily diffuse and dissolve, thereby increasing its extraction efficiency. But prolonged ultrasonication may lead to the destruction of the polysaccharides structure. With less solvent usage, shorter working time of extraction and lower extraction temperature, UEM is a fast and inexpensive method compared to traditional methods ^[16]. Lian and co-workers reported extraction of *Cistanche herba* polysaccharides by ultrasonic-assisted extraction and studied the main factors influencing the yield of polysaccharides by single factor test method ^[17]. The optimal extraction process is solid-liquid ratio was 1:25 with 75 min ultrasonic-assisted extraction under 60 °C, the yield of polysaccharides from *Cistanche herba* was 18.90 %.

2.3 Microwave-Assisted Extraction Methods

Microwave-assisted extraction method is based on the difference in the ability of substances to absorb microwaves, certain compounds are extracted by selective heating. The microwave has strong penetrating ability, high selectivity, and rapid heating. It can generate strong pressure to break through cell membranes and cell walls, and promote the dissolution of polysaccharides. The advantages of this method being widely used are high efficiency, rapidity and energy saving ^[18]. Teng and co-workers described four different methods for extracting polysaccharides, including microwave-assisted extraction, ultrasound-microwave synergistic extraction, ultrasonic-assisted extraction, etc ^[19]. The results demonstrated that the finest extraction method was ultrasound-microwave synergistic extraction, followed by microwave-assisted extraction. It can be observed that the penetrating and heating ability of microwaves play essential roles in the extraction process of polysaccharides.

2.4 Enzyme-Assisted Extraction Methods

The enzyme-assisted extraction method is to effectively dissolve polysaccharides through the destruction of plant cell walls by biological enzymes. The extraction conditions are mild and simple, less destructive to polysaccharide structure, and the extraction effect is significantly enhanced^[20]. Gao and co-workers reported employed enzyme-assisted extraction to extract the polysaccharides^[21]. Taking the enzyme dosage, enzymolysis temperature and enzymolysis time as the reference factors, the polysaccharides extraction yield was the indicator, and the optimal process of enzyme-assisted extraction was determined through orthogonal experiments. The results showed that the actual yield of polysaccharides increased from 4.61 % to 8.35 % when the enzyme dosage was 0.2 %, enzymatic hydrolysis time was 1.5 h, and hydrolysis temperature was 50 °C.

In addition, scientists have developed a combination of multiple extraction methods to achieve the effect of maximizing the strengths and avoiding weaknesses of a single extraction method, such as ultrasonic-microwave synergistic extraction, ultrasonic-enzyme-assisted extraction, etc. These methods not only shorten the extraction time, simplify the extraction process and reduce the operational difficulty, but obtain high-yield polysaccharides. Teng and co-workers firstly employed ultrasonic-microwave synergistic extraction method to extract polysaccharides from *Cistanche tubulosa*, which have the best extraction effect as it can exert stronger and more consistent force on the sample, and accelerate the process of dissolving polysaccharides into the solvent^[18]. Zhang and co-workers reported the effective extraction of *Cistanche tubulosa* polysaccharides utilizing ultrasound-cellulase-assisted^[22]. Single factoris was employed to further optimize ultrasound-cellulase-assisted extraction conditions and the maximum yield of polysaccharides was 22 % under the condition of pH of 5.2 for 31.5 min at 54.1 °C.

2.5 Other Extraction Methods

As the pharmacological effects of traditional Chinese medicine polysaccharides become more and more significant, their extraction process has also received extensive attention as the primary work of research. For the extraction research of *Cistanche deserticola* polysaccharides, scholars have tried other different extraction methods. Zhang and co-workers firstly used flash extraction method to extract polysaccharides from *Cistanche deserticola*, and studied the effects of solid-liquid ratio, voltage and extraction time on the yield of polysaccharides^[23]. Under the optimal extraction conditions, the polysaccharides yield was 12.35 %. This method has the advantages of time-saving, efficient and easy operation in the extraction of polysaccharides.

3. Chemical Structure

Cistanche deserticola is rich in chemical components, and scholars at home and abroad have isolated and structurally identified more than one hundred compounds, including phenylethanol glycosides, lignans, iridoids, polysaccharides, amino acids, etc^[24-25]. Among them, phenylethanol glycosides are considered to be the main active ingredients for the medicinal effects of *Cistanche deserticola*. However, as the research on the pharmacological effects of *Cistanche deserticola* has

intensified, it has been found that the polysaccharides also plays a crucial role in the pharmacological activity. The chemical structure of polysaccharides is the material basis for their biological activities and polysaccharides with different chemical structures have different pharmacological effect. Therefore, more and more attention has been paid to the structure research of polysaccharides from *Cistanche deserticola*. In the 90s, studies on the polysaccharides of *Cistanche deserticola* focused on the isolation , purification and compositional analysis of monosaccharides ^[26-27], while the precise structure of polysaccharides was not elucidated until 1997 ^[28]. Recent developments in modern chromatography and spectroscopy have also facilitated the characterization of the chemical structures of polysaccharides.

Ebringerová A and co-workers employed dilute alkali to extract three pectic polysaccharides (P1, P2 and P3) from *Cistanche deserticola* and structurally characterized them by chemistry, enzymology and NMR spectroscopy ^[28]. The results indicated that the main relative molecular mass peak of P1, P2 and P3 at 1.87×10^5 , 1.25×10^5 , 0.8×10^4 , respectively, as well as arabinose and galactose are prevailing neutral sugar fractions. The mainly differences between these pectins are proportions of homogalacturonan and rhamnogalacturonan RG-I sequences, contents of galacturonic acid, degree of methyl esterification and acetylation, etc. All pectins showed remarkable immunomodulating activities in both mitogenic and comitogenic tests. In addition the authors suggested that immunomodulatory properties of *Cistanche* pectins may be related to rhamnogalacturonan and α -3,5-arabinan moieties ^[29]. The acidic pectic polysaccharide, bupleuran 2IIb, from *Bupleurum falcatum* L. showed a potent enhancing immune activity ^[30]. The experimental result suggested that the ramified region which consisted of a rhamnogalacturonan core with neutral carbohydrate chains is important for this enhancing activity. These validated the relationship between the chemical structure of rhamnogalacturonan and immunomodulatory activity. This article not only examined chemical structure of the pectic polysaccharides in more detail, but specifically studied the structure in relation to its biological activity.

Subsequently Ebringerová A et al reported the crude polysaccharide extracted by cold water was purified by column chromatography to yield Cistan A with relative mean molecular mass was 2.01×10^5 ^[31]. Analysis of sugar composition revealed that Cistan A consisted of L-arabinose, D-galactose, L-rhamnose and D-galacturonic acid in a molar ratio of 6.3:10.0:1.0:0.8, in addition to very low amounts of D-xylose and D-glucose. NMR spectroscopy and methylation analysis demonstrated that Cistan A contains mainly a complex of pectic arabino-3,6-galactan type II with lowly-branched 3,5- α -L-arabinan. In immunological activities test, Cistan A showed the highest activities compared to commercial immunomodulator Zymosan and other extracted polysaccharides. Moreover the ramified regions of the galacturonan core and both the neutral side chains consisting of 3,6- β -D-galactan and 3,5- α -L-arabinan were considered to be engaged in the bioactive expression ^[29].

Tu and co-workers reported the structural characteristics of polysaccharide named CDP-4 isolated from the immunologically active polysaccharides of *Cistanche deserticola* ^[32]. Chemical and spectroscopic analyses measured a relative molecular mass of 1.4×10^4 , and TLC analysis of

development and industrial application of *Cistanche deserticola*.

4. Biological Activities

Polysaccharide, as one of the main active ingredients of *Cistanche deserticola*, has many pharmacological effects such as regulating immune activity, anti-aging, anti-liver damage, anti-virus, anti-tumor, and affecting intestinal flora, and is widely used in clinical and health food applications.

4.1 Regulation of immune activity

Cistanche deserticola polysaccharide (CDPS) is an important active component of *Cistanche deserticola* that exerts immunomodulatory effects. It promotes the proliferation of lymphocytes and improves immune function. And also has an activating effect on macrophages and enhances phagocytic activity, thus increasing the synthesis and secretion of cytokines that regulate immune response and playing an immunomodulatory role.

Zeng et al investigated immunomodulatory effects on mouse T cells of CDPS^[43]. CDPS was found to not only increase the proliferative response of mouse spleen and thymus lymphocytes alone, but also stimulate the proliferation of mouse thymus lymphocytes in concert with ConA and PHA. In addition, the polysaccharide significantly increased secretion of interleukin 2 by lymphocytes, so it is speculated that the promotional effect of CDPS on IL-2 secretion may be one of the mechanisms by which *Cistanche deserticola* can enhance immune function and anti-aging. Subsequently Zeng et al also investigated the mechanism of the immunomodulatory effects of CDPS^[44]. They determined the immunomodulatory function of CDPS on the proliferation of mouse thymic lymphocytes in vitro and studied the cell cycle and its intracellular Ca²⁺ concentration. Finally, it is speculated that the mechanism may be that CDPS promotes the release of calcium from mouse thymic lymphocytes, and the elevated calcium ion concentration promotes IL-2 gene expression, leading to T cell proliferation. Wang et al observed the effect of CDPS on lymphocyte proliferation in vivo and in vitro, and also found that this effect is closely related to CDPS promoting the release of IL-2 from lymphocytes^[45].

Wang et al. determined the activation effect of CDPS on macrophages^[46]. The results showed that CDPS significantly enhanced the phagocytosis of macrophages and the release of NO, TNF- α and IL-1 from RAW264.7 cells, a mouse mononuclear macrophage commonly used to test immune-related experiments, as well as the release of NO from immunosuppressed RAW264.7 cells, which demonstrated that the immune enhancing effect of CDPS was correlated with its activation of macrophages. Zhang et al also discovered that CDPS may exert immunomodulatory functions by enhancing phagocytosis of THP-1 cells and promoting the release of cytokines^[47].

4.2 Anti-aging

Research has demonstrated that CDPS has significant anti-aging activity, mainly through anti-oxidation, anti-fatigue, improved learning and memory, and enhanced telomerase activity.

Sun et al used D-galactose to cause experimental aging model in mice and observed the antioxidant function of CDPS on aging mice ^[48]. The results showed that CDPS inhibited lung collagen increase and elastin decrease and altered the imbalance of oxidative/antioxidative function of the body. It is hypothesized that it may eliminate the increase in free radical production caused by D-galactose and avoid lung tissue damage due to depletion of antioxidant substances, thus achieving a slowing effect on aging. Subsequently the group studied the effect of ozone-injured experimental lung aging mice administered with CDPS ^[49]. It was found that after injection of CDPS, the hypoxia tolerance of aging mice was significantly improved, the activity of superoxide dismutase for scavenging free radicals was enhanced, and the content of lipid peroxidation metabolites was reduced, indicating that CDPS can enhance the antioxidant function of aging mice.

Reduced fatigue resistance is one of the main features of aging. Yan et al examined the effect and mechanism of CDPS on anti-fatigue in D-galactose-induced aging mice ^[50]. It was revealed that both high and low doses (400 mg/kg, 100 mg/kg) of CDPS had anti-fatigue effects on aging mice. The mechanism may be to reduce serum urea nitrogen and lactate levels, increase content of liver glycogen and muscle glycogen, and enhance the ability of mice to fight free radical damage and antioxidant.

Recently one of the causes of Alzheimer's disease is thought to be a decrease in the number of neurons in brain tissue caused by apoptosis, and CDPS has anti-aging effects, so scholars have studied their related effects. Yin et al observed the effect of CDPS on learning and memory ability of model rats with Alzheimer's disease ^[51]. It was found that CDPS inhibited NO production and release through the inhibition of ROS activity, and improved the decrease of learning and memory ability in rats. The mechanism may be to reduce the damage of oxygen free radicals and accelerating the scavenging of free radicals in vivo. Ma et al discovered that CDPS was effective in improving learning and memory ability in D-galactose-induced ageing mice ^[52]. It is speculated that CDPS enhances learning memory function by upregulating cAMP response element binding protein expression in neurons, promoting neuronal survival and neuronal morphological and functional recovery, and reducing apoptosis. Wu et al examined the effect of CDPS on the improvement of learning memory ability in aging mice and revealed the mechanism was that CDPS modestly increased the release of excitatory neurotransmitter through upregulation of the cAMP/PKA/CREB/BDNF signaling pathway ^[53].

Zhang et al employed a D-galactose aging mouse model to observe the effect of CDPS on it ^[54]. It was observed that CDPS significantly reduced malondialdehyde content in tissues and increased telomerase activity in heart and brain tissues, lymphocyte proliferative response and

phagocytosis of macrophages. Therefore, it is inferred that CDPS has good effects in antagonizing free radical damage, enhancing telomerase activity and immune function, and delaying the effect on aging.

4.3 Gut Protection and laxative effect

Jia et al revealed that polysaccharide-rich aqueous extracts of *Cistanche deserticola* can effectively prevent colorectal cancer and intestinal inflammation in a mouse model ^[55]. The mechanism may be to stimulate the immune active system, reducing inflammatory mucosal hyperplasia and intestinal helicobacter infection. This study demonstrated that CDPS may have clinical potential in the prevention and treatment of colorectal cancer. Fu et al investigated the effect of CDPS on gut microbiota and discovered that all species of polysaccharides could regulate the diversity of intestinal microbiota, increase beneficial bacteria, promote the absorption of echinacoside, and improve the disordered gut microbiota ^[56]. In addition, low molecular weight polysaccharides may treat diseases related to the gut microbiota. Gao et al investigated the effects of CDPS on mice with cognitive decline and dysbiosis of gut microflora due to D-galactose ^[57]. The results indicate that CDPS improving low cognitive performance in mice by restoring gut microbial homeostasis.

Liu et al demonstrated the regulatory effects of CDPS on the constipation of aged rats and CDPS markedly increased the content of beneficial bacteria while dramatically decreasing the content of harmful bacteria ^[58]. In addition, CDPS enhances immunity and prevents pathogenic bacterial infections. The mechanism of the laxative effect of CDPS mainly included metabolic energy and amino acid synthesis.

4.4 Anti-spleen deficiency and anti-liver damage

Xu et al investigated the regulatory effect of CDPS on CCl₄-induced liver qi stagnation and spleen deficiency in mice ^[59]. The results revealed that CDPS alleviated weight loss in mice with hepatic depression and spleen deficiency, and also modulated immune deficiency. Wang et al investigated the protective effects of CDPS on CCl₄-induced acute liver injury in mice and showed that CDPS could reduce malondialdehyde content and increase superoxide dismutase levels in mouse liver ^[60]. CDPS significantly improved the hepatic redox status in mice, and the mechanism may be that CDPS counteracts lipid peroxide damage to hepatocytes, improves their antioxidant capacity, and performs normal defense and compensatory functions. Guo et al extracted three polysaccharides from *Cistanche deserticola*, of which the polysaccharide named CDP-C had higher anti-liver damage and antioxidant activity ^[61]. Bioactivity experiments suggest that CDP-C may exert protective effects against alcohol-induced chronic liver injury by reducing malondialdehyde and triglyceride levels and modulating the activity of related enzymes. The

biological activity correlates with the fact that CDP-C contains a higher proportion of galacturonic acid.

4.5 Prevent and treat osteoporosis

Song et al investigated the function and mechanism of CDPS on osteoporosis. The result showed CDPS inhibited osteoclastogenesis and hydroxyapatite absorption, also restrained the expression of osteoclast marker genes. Consequently, CDPS is expected to be applied clinically for the treatment of osteoporosis ^[62]. Wang et al explored the types of active ingredients and mechanisms involved in the treatment of osteoporosis in *Cistanche deserticola*. Senescence accelerated mice were administrated *Cistanche deserticola* extract total glycosides and polysaccharides. The result demonstrated that these extracts can promote osteoblastogenic bone formation and improve bone microstructure injury by modulating Wnt/ β -catenin signaling pathway ^[63]. Xiao et al investigated the effect of extracted and purified CDPS on the generation and function of osteoclasts and the treatment of osteoporosis. The results exhibited that CDPS treatment prevented OVX-induced osteoporosis and improved bone loss by inhibiting osteoclast activity and function ^[64].

4.6 Other biological activities

In terms of biological activities, CDPS not only exhibits common effects such as regulation of immune function, anti-aging, anti-splenic deficiency, anti-hepatic injury and intestinal protection. But many scholars have also studied the pharmacological activities in other aspects, including anti-tumor, wound healing, promotion of bone marrow hematopoietic function and improvement of menopause, in order to provide reference for *Cistanche deserticola* in clinical applications and health care drugs.

Zhang et al revealed that CDPS significantly inhibited Lewis lung tumor and solid tumor S₁₈₀ in mice and prompted natural killer cell and interleukin-2 activity ^[65]. Therefore, it is speculated that the tumor suppressive effect of CDPS is related to its ability to increase the activity of natural killer cells and interleukin-2, which alleviates the depressed immune function of mice. These effects increase with increasing doses of CDPS, which may be used clinically as an adjuvant to tumor treatment. Gao et al investigated the effects of various concentrations of CDPS on cultured human fibroblasts in vitro and found that the drug significantly promoted the growth of fibroblasts in vitro ^[66]. The concentration of CDPS has an impact on its effect, and the appropriate concentration should be adopted to promote the healing of wounds, providing a theoretical basis for its clinical application. Chen et al observed the effect of CDPS on hematopoietic function in myelosuppressed anemic mice and confirmed that CDPS may promote the recovery of bone marrow hematopoietic function through promoting the transformation of the bone marrow cell cycle in myelosuppressed anemic mice ^[67]. Zhang et al explored the effect of CDPS on the regulation of serum endocrine function in rats with menopausal hypertension and established an animal model by removing the ovaries of rats and giving them dry chow ^[68]. CDPS was identified to enhance HPOA function to achieve regulation of neurotransmitters and endocrine hormones, improve neurological function, stabilize the body environment, and accelerate metabolism, thus

improving menopausal symptoms.

5. Conclusion

Cistanche deserticola has been used in China for more than 2,000 years. Its taste is sweet and slightly salty, and it has many health functions including nourishing the kidneys, laxative effect, delaying aging and enhancing memory. Polysaccharide is one of the important active ingredients extracted from *Cistanche deserticola*, which has received wide attention for its pharmacological activities such as immunomodulation, anti-aging, anti-tumor and anti-liver damage. Therefore, the extraction and isolation of polysaccharides, chemical structure analysis and the study of its biological activity have important research significance. At present, the extraction methods of *Cistanche deserticola* polysaccharides mainly include traditional extraction method, ultrasonic-assisted method, microwave-assisted method and enzyme-assisted method, etc. Analysis of the chemical structure of polysaccharides is the primary prerequisite for studying its pharmacological activity and efficiently utilizing the transformation. The extracted polysaccharides were analyzed by chemical methods such as methylation, periodate oxidation, acid hydrolysis and Smith degradation, and various spectroscopy methods including NMR, IR and HPLC to obtain a variety of polysaccharides with new structures. The relationship between the chemical structure and biological activity of polysaccharides has been studied by a small number of scholars, but the relationship remains unclear due to the large number of components, complex structure and lack of systematic studies on polysaccharides. In addition, there are few studies on the effects of *Cistanche deserticola* polysaccharide extraction methods on its chemical structure and biological activity. It is expected that the subsequent research on polysaccharide structure and active conformational relationship will be of high value to realize the precise transformation of big health products, with a view to achieving the modernization of *Cistanche deserticola* in Chinese medicine and making major breakthroughs and innovations in pharmaceutical sciences, clinical medicine, food and health products, functional cosmetics, etc.

Declaration of competing interest

The authors declare that they have no known competing financial interests or personal relationships that could have appeared to influence the work reported in this paper.

Acknowledgements

The authors acknowledge the “JUN-MA” High-level Talents Program of Inner Mongolia University (No. 21300-5195112, No. 21300-5205107), and the funding from the Science & Technology Department of Inner Mongolia Autonomous Region (No. 2021CG0029, No. 21300-5213402).

References

[1] C.-M. Gu, X.-Y. Yang, L.-F. Huang, *Cistanches Herba*: A neuropharmacology review, *Front.*

Pharmacol. 7 (2016) 289.

- [2] P.-F. Tu, Y.-P. He, Z.-C. Lou, Survey and protection of medicinal resources of desertying Cistanche(*Cistanche deserticola*), *Chin. Tradit. Herb. Drugs*. 24 (1994) 205-208.
- [3] T. Wang, X.-Y. Zhang, W.-Y. Xie, *Cistanche deserticola* Y. C. Ma, "desert ginseng": a review, *Am. J. Chin. Med.* 40 (2012) 1123-1141.
- [4] The State of Pharmacopoeia Commission of the People's Republic of China. Pharmacopoeia of the People's Republic of China(2005). China Medical Science and Technology Press, Beijing, 2005.
- [5] Z.-F. Fu, X. Fan, X.-Y. Wang, X.-M. Gao, *Cistanches Herba*: an overview of its chemistry, pharmacology, and pharmacokinetics property, *J. Ethnopharmacol.* 219 (2018) 233-247.
- [6] Z.-M. Li, H.-N. Lin, L. Gu, J.-W. Gao, C.-M. Tzeng, *Herba Cistanche*(Rou Cong-rong): One of the best pharmaceutical gifts of traditional Chinese medicine, *Front. Pharmacol.* 7 (2016) 41.
- [7] L. Lei, Z.-H. Song, P.-F. Tu, Advances in research of chemical constituents in plants of *Cistanche* Hoffing. et Link, *Chin. Tradi. Herb. Drugs*. 34 (2003) 473-476.
- [8] J. Li, D. Huang, L.-Q. He, Effect of Roucongrong (*Herba Cistanches Deserticolae*) on reproductive toxicity in mice induced by glycoside of Leigongteng (*Radix et Rhizoma Tripterygii*), *J. Tradi. Chin. Med.* 34 (2014) 324-328.
- [9] E. Wat, C.F. Ng, C.M. Koon, C. Zhang, S. Gao, B. Tomlinson, C. B. S. Lau, The adjuvant value of *Herba Cistanches* when used in combination with statin in murine models, *Sci. Rep-UK*. 7 (2017) 1-17.
- [10] W.-T. Xiong, L. Gu, C. Wang, H.-X. Sun, X. Liu, Anti-hyperglycemic and hypolipidemic effects of *Cistanche tubulosa* in type 2 diabetic *db/db* mice, *J. Ethnopharmacol.* 150 (2013) 935-945.
- [11] X.-X. Gao, J. Chen, Y.-L. Peng, Research Situation of Pharmacological Effects of *Cistanche deserticol* Polysaccharides, *Food and Drug*. 17 (2015) 136-139.
- [12] Z.-F. Sui, T.-M. Gu, B. Liu, S.-W. Peng, Z.-L. Zhao, L. Li, D.-F. Shi, R.-Y. Yang, Water-soluble carbohydrate compound from the bodies of *Herba Cistanches*: Isolation and its scavenging effect on free radical in skin, *Carbohydr. Polym.* 85 (2011) 75-79.
- [13] N. Deng, Y.-J. Shen, H. Ding, X.-B. Song, L.-J. Zhang, Research progress on pharmacological action of polysaccharides in *Cistanche Deserticola*, *Journal of Liaoning University of TCM*. 22 (2020) 67-71.
- [14] R. Naran, A. Ebringerová, Z. Hromádková, V. Patoprstý, Carbohydrate polymers from underground parts of *Cistanche deserticola*, *Phytochemistry*. 40 (1995) 709-715.
- [15] R. Naran, A. Ebringerová, Z. Hromádková, K. Babor, Bratislava, The starch component of the wood parasite *Cistanche deserticola* Y. C. Ma, *Starch- Stärke*. 48 (1996) 255-259.
- [16] I.P. Claver, H.-H. Zhang, Q. Li, Z. Kexue, H.-M. Zhou, Optimization of ultrasonic extraction of polysaccharides from Chinese Malted Sorghum using response surface methodology, *Pak. J. Nutr.* 9 (2010) 336-342.
- [17] L. Lian, M.-B. Gao, X.-Y. Yin, X.-Q. Zhao, X.-H. Hu, B. Zhang, Optimization of Ultrasonic-assisted Extraction of Polysaccharides from *Cistanche herba*, *Guangzhou Chemical*

Industry. 46 (2018) 62-63.

- [18] A.M. Cikoš, S. Jokić, D. Šubarić, I. Jerković, Overview on the application of modern methods for the extraction of bioactive compounds from *Marine Macroalgae*, *Mar. Drugs*. 16 (2018) 348-368.
- [19] L.-P. Teng, L. Mu, F. Luo, Z.-B. Zhou, Extraction of Crude Polysaccharides from *Cistanche tubulosa* and its Effect of Scavenging DPPH Radicals, *Guizhou Agricultural Sciences*. 39 (2011) 66-68.
- [20] Y. Chen, F.-K. Yao, K. Ming, D.-Y. Wang, Y.-L. Hu, J.-G. Liu, Polysaccharides from traditional Chinese medicines: Extraction, purification, modification, and biological activity, *Molecules*. 21 (2016) 1705-1728.
- [21] J.-D. Gao, X. Liu, X.-Y. Zhu, R. Wu, T.-T. Shi, T. Ma, Optimization of enzymatic extraction process for *Cistanche deserticola* polysaccharides and its kinetic analysis, *J. Gansu University of Chinese Medicine*. 34 (2017) 26-30.
- [22] W.-J. Zhang, J. Huang, W. Wang, Q. Li, Y. Chen, W.-W. Feng, D.-H. Zheng, T. Zhao, G.-H. Mao, L.-Q. Yang, X.-Y. Wu, Extraction, purification, characterization and antioxidant activities of polysaccharides from *Cistanche tubulosa*, *Int. J. Biol. Macromol.* 93 (2016) 448-458.
- [23] X.-F. Zhang, M.-M. Deng, G.-H. Luo, S.-H. Yang, D.-X. Wang, Optimization of flash extraction of polysaccharides from *Cistanche deserticola* and its antiglycosylation activity, *Journal of Gansu Agricultural University*. 54 (2019) 193-198.
- [24] Y.-L. Song, K.-W. Zeng, Y. Jiang, P.-F. Tu, *Cistanches Herba*, from an endangered species to a big brand of Chinese medicine, *Med. Res. Rev.* 41 (2021) 1539-1577.
- [25] Y. Jiang, P.-F. Tu, Analysis of chemical constituents in *Cistanche* species, *J. Chromatogr. A*. 1216 (2009) 1970-1979.
- [26] R. Naran, A. Ebringerová, D. Badgaa, Carbohydrate components of the holoparasite *Cistanche deserticola* general characteristics of the underground part, *Chem. Pap.* 49 (1995) 35- 38.
- [27] B. Kosiková, A. Ebringerová, R. Naran, Characterization of lignin- carbohydrate fractions isolated from the wood parasite *Cistance deserticola* YC Ma, *Holzforschung*. 53 (1999) 33- 38.
- [28] A. Ebringerová, Z. Hromádková, E. Machová, R. Naran, V. Hřibalová, Isolation and Characterization of Mitogenic Pectic Polysaccharides from *Cistanche deserticola* YC Ma, *Chem. Pap.* 51 (1997) 289- 294.
- [29] J.-F. Zhao, H. Kiyohara, H. Yamada, N. Takemoto, H. Kawamura, Heterogeneity and characterisation of mitogenic and anticomplementary pectic polysaccharides from the roots of *Glycyrrhiza uralensis* Fisch et D.C., *Carbohyd. Res.* 219 (1991) 149-172.
- [30] T. Matsumoto, J.-C. Cyong, H. Kiyohara, H. Matsui, A. Abe, M. Hirano, H. Danbara, H. Yamada, The Pectic polysaccharide from *Bupleurum falcatum* L. enhances immune-complexes binding to peritoneal macrophages through Fc receptor expression, *Int. J. Immunopharmac.* 15 (1993) 683-693.
- [31] A. Ebringerová, Z. Hromádková, V. Hřibalová, J. Hirsch, An Immunomodulating Pectic Arabinogalactan from Roots of *Cistanche deserticola*, *Chem. Pap.* 56 (2002) 320- 325.
- [32] X.-M. Wu, P.-F. Tu, Studies on the chemical structure of polysaccharide CDP-4 isolated from *Cistanche deserticola*, *Journal of Peking University*. 36 (2004) 24-26.
- [33] X.-M. Wu, P.-F. Tu, Chemical properties of a mannoglucan from *Cistanche deserticola*,

Pharmazie. 59 (2004) 815- 816.

- [34] X.-M. Wu, X.-M. Gao, K.W.K. Tsim, P.-F. Tu, An arabinogalactan isolated from the stems of *Cistanche deserticola* induces the proliferation of cultured lymphocytes, *Int. J. Biol. Macromol.* 37 (2005) 278- 282.
- [35] X.-S. Wang, Y. Zheng, J.-P. Zuo, J.-N. Fang. Structural features of an immunoactive acidic arabinogalactan from *Centella asiatica*, *Carbohydr. polym.* 59 (2005) 281-288.
- [36] B.P. da Silva, J.P. Parente, Chemical properties and biological activity of a polysaccharide from *Melocactus depressus*, *Planta Med.* 68 (2002) 74-76.
- [37] X.-M. Wu, P.-F. Tu, Isolation and characterization of α - (1 \rightarrow 6)- glucans from *Cistanche deserticola*, *J. Asian Nat. Prod. Res.* 7 (2005) 823- 828.
- [38] N. Radna, E. Anna, A. Juraj, The Cell Wall Material of the Wood Parasite *Cistanche deserticola* Y.C.Ma, *Halzforschung.* 49 (1995) 293-299.
- [39] W. Zhao, H. Yan, Z.-Y. Liang, Y.-J. Zhang, X.-Q. Jiao, Structural Analysis of Water-soluble Polysaccharide SPA Isolated from the Stem of the *Cistanche Deserticola* Ma, *Chem. J. Chinese U.* 26 (2005) 461-463.
- [40] W. Zhao, H. Yan, Z.-Y. Liang, Y.-J. Zhang, X. Huo, Purification and characterization on water soluble polysaccharides SPA isolated from the stem of *Cistanche deserticola* Ma, *Journal of northeast normal university.* 36 (2004) 111-115.
- [41] Q. Dong, J. Yao, J.- N. Fang, K. Ding, Structural characterization and immunological activity of two cold- water extractable polysaccharides from *Cistanche deserticola* Y. C. Ma, *Carbohydr. Res.* 342 (2007) 1343- 1349.
- [42] H. Zeng, L.-L. Huang, L.-S. Zhou, P.-P. Wang, X. Chen, K. Ding, A galactoglucan isolated from of *Cistanche deserticola* Y. C. Ma. and its bioactivity on intestinal bacteria strains, *Carbohydr. Polym.* 223 (2019) 115038.
- [43] Q.-L. Zeng, J.-H. Mao, Z.-L. Lv, Purification of Polysaccharide of *Cistanche Deserticola* Y C Ma and its Immunomodulatory Effects on T Cell Function, *Journal of Zhejiang Medical University.* 27 (1998) 108-111.
- [44] Q.-L. Zeng, Y.-F. Zheng, Z.-L. Lv, Immunomodulatory effects of polysaccharide of *Cistanche Deserticola* Y Ma, *Journal of Zhejiang University (Medical Sciences).* 31 (2002) 284-287.
- [45] X.-Y. Wang, Y. Qi, R.-L. Cai, X.-H. Li, M.-H. Yang, Y. Shi, Enhancing Effect of Polysaccharides of *Cistanche deserticola* Y C Ma on Lymphocyte Proliferation, *Acta Laboratorium Animalis Scientia Sinica.* 17 (2009) 424-427.
- [46] X.-Y. Wang, Y. Qi, R.-L. Cai, X.-H. Li, M.-H. Yang, Y. Shi, The effect of *Cistanche deserticola* polysaccharides (CDPS) on macrophages activation, *Chin. Pharmacol. Bull.* 25 (2009) 787-790.
- [47] T. Zhang, W.-S. Xu, Y.-B. Jia, G.-R. Bai, J. Gao, L.-X. Cui, L.-H. Lang, Y.-J. Zhang, The Effects and Mechanism of *Cistanche deserticola* Polysaccharide on the THP-1 Cell's Phagocytosis, *Life Science Research.* 17 (2013) 148-151.

- [48] Y. Sun, D.-J. Wang, J. Zhu, H.-Q. Zhang, Effects of *cistanche desertica* polysaccharides on the constitution of protein and anti-oxidative capacity of lung in aging mice, *Chin. Pharmacol. Bull.* 17 (2001) 101-103.
- [49] Y. Sun, D.-J. Wang, X.-M. Liu, L. Shi, H.-Q. Zhang, The change of functions and hypermicro-structure in lung in experimental aging mice and effect of *cistanche desertica* polysaccharides on the change, *Chin. Pharmacol. Bull.* 18 (2002) 84-87.
- [50] L. Yan, J.-P. Hu, X.-D. Sun, K.-R. Cai, Study on the anti-fatigue effect and mechanism of *cistanche deserticola* polysaccharide on D-galactose-induced aging mice, *Hebei J. TCM.* 41 (2019) 96-100.
- [51] G. Yin, D.-K. Gong, B.-H. Liu, C.-J. Yao, Effect of Polysaccharides of *Cistanche Deserticola* on learning and memory ability and oxygen stress of model rats with Alzheimer's disease, *J Apoplexy and Nervous Diseases.* 30 (2013) 504-507.
- [52] H. Ma, R.-X. Yin, M. Guo, Y. Bao, Z.-H. Cui, G. Li, Effect of *Cistanche deserticola* Polysaccharides on Expression of CREB in D-galactose Induced Aging Model Mice, *Chinese Journal of Experimental Traditional Medical Formulae.* 20 (2014) 137-141.
- [53] Y. Wu, H. Zhang, R. Bu, H. Ma, M. Su, G. Li, *In vivo* and *in vitro* studies of protective effect of CDPS on acute aging mouse model induced by D-galactose, *Chin. Pharmacol. Bull.* 33 (2017) 927-933.
- [54] H.-Q. Zhang, Y. Li, Y.-Y. Song. Effect of Polysaccharides of *Cistanche deserticola* on Immune Cells and Telomerase Activity in Aging Mice, *Chin. Pharm. J.* 46 (2011) 1081-1083.
- [55] Y.-M. Jia, Q.-N. Guan, Y.-H. Guo, C.-G. Du, Reduction of Inflammatory Hyperplasia in the Intestine in Colon Cancer-prone Mice by Water-extract of *Cistanche deserticola*, *Phytother. Res.* 26 (2012) 812-819.
- [56] Z.-F. Fu, L.-F. Han, Z. Peng, H.-P. Mao, H. Zhang, Y.-F. Wang, X.-M. Gao, E. Liu, *Cistanche* polysaccharides enhance echinacoside absorption *in vivo* and affect the gut microbiota, *Int. J. Biol. Macromol.* 149 (2020) 732-740.
- [57] Y. Gao, B. Li, H. Liu, Y.-J. Tian, C. Gu, X.-L. Du, R. Bu, J. Gao, Y. Liu, G. Li, *Cistanche deserticola* polysaccharides alleviate cognitive decline in aging model mice by restoring the gut microbiota-brain axis, *Aging-US.* 13 (2021) 15320-15335.
- [58] X.-J. Liu, C. Jian, M.-Y. Li, F.-X. Wei, H.-L. Liu, X.-M. Qin, Microbiome-metabolomics deciphers the effects of *Cistanche deserticola* polysaccharides on aged constipated rats, *Food Funct.* 13 (2022) 3993-4008.
- [59] Z.-X. Xu, Y. Sun, A.-P. Lin, H.-Q. Zhang, The regulation effects of *Herba Cistanches* polysaccharides on immunological functions of liver qi stagnation and spleen deficiency induced by CCl₄ in mice, *Chin. J. Integr. Med.* 17 (1997) 261-263.
- [60] Y.-F. Wang, J.-J. Zhao, M.-H. Li, H.-Y. Zhang, S. Jiang, The Protective Effect of *Cistanche Deserticola* polysaccharide on Acute Liver Injury Induced by Carbon Tetrachloride in Mice, *Journal of Baotou Medical College.* 30 (2014) 3-5.
- [61] Y.-H. Guo, L.-L. Cao, Q.-S. Zhao, L.-J. Zhang, J.-J. Chen, B.-Y. Liu, B.-Zhao, Preliminary characterizations, antioxidant and hepatoprotective activity of polysaccharide from *Cistanche*

deserticola, Int. J. Biol. Macromol. 09 (2016) 039.

[62] D.-Z. Song, Z. Cao, Z.-B. Liu, J. Tickner, H. Qiu, C. Wang, K. Chen, Z.-Y. Wang, S.-W. Dong, J.-K. Xu, *Cistanche deserticola* polysaccharide attenuates osteoclastogenesis and bone resorption via inhibiting RANKL signaling and reactive oxygen species production, J. Cell. Physiol. 233 (2018) 9674-9684.

[63] F.-J. Wang, P.-F. Tu, K.-W. Zeng, Y. Jiang, Total glycosides and polysaccharides of *Cistanche deserticola* prevent osteoporosis by activating Wnt/ β -catenin signaling pathway in SAMP6 mice, J. Ethnopharmacol. 271 (2021) 113899.

[64] W. Xiao, Y.-Y. Wei, F. Yang, X.-Y. Lu, S.-W. Liu, Y.-L. Long, *Cistanche deserticola* polysaccharide inhibits OVX-induced bone loss in mice and RANKL-induced osteoclastogenesis, J. Funct. Foods. 81 (2021) 104464.

[65] H.-Q. Zhang, A.-H. Xu, D.-D. Li, Pharmacological Action of *Cistanche Deserticola* Polysaccharides on Tumor Bearing Mice, Chin. J. Integr. Med. 17 (1997) 155-156.

[66] H. Gao, H. Yan, H.-Q. Zhang, K. Tong, The proliferative effects of *Cistanche* polysaccharide on human fibroblasts, Chin. Tradit. Patent Med. 20 (1998) 43-44.

[67] Z.-W. Chen, B.-D. Zhu, H.-Y. Xu, Experiment Study of Desertliving *Cistanche* Polysaccharides on Regulated Haematogenesis in Bone Marrow Depressed Anemia Mice, Chin. Arch. Tradit. Chin. Med. 25 (2007) 1473-1474.

[68] N.-N. Zhang, Effects of Polysaccharides of *Cistanche Deserticola* on Endocrine Regulation in Rats with Menopausal Hypertension of Dryness Syndrome, Acta. Chin. Med. Pharmacol. 46 (2018) 65-69.