



HAL
open science

Effectiveness of Influenza Vaccine against Influenza A in Europe in Seasons of Different A(H1N1)Pdm09 and the Same A(H3N2) Vaccine Components (2016–17 and 2017–18)

Esther Kissling, Francisco Pozo, Silke Buda, Ana-Maria Vilcu, Caterina Rizzo, Alin Gherasim, Judit Krisztina Horváth, Mia Brytting, Lisa Domegan, Adam Meijer, et al.

► To cite this version:

Esther Kissling, Francisco Pozo, Silke Buda, Ana-Maria Vilcu, Caterina Rizzo, et al.. Effectiveness of Influenza Vaccine against Influenza A in Europe in Seasons of Different A(H1N1)Pdm09 and the Same A(H3N2) Vaccine Components (2016–17 and 2017–18). *Vaccine*: X, 2019, 3, pp.100042. 10.1016/j.jvacx.2019.100042 . hal-03846533

HAL Id: hal-03846533

<https://hal.sorbonne-universite.fr/hal-03846533>

Submitted on 21 Apr 2023

HAL is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers.

L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.



Distributed under a Creative Commons Attribution - NonCommercial - NoDerivatives 4.0 International License



Effectiveness of influenza vaccine against influenza A in Europe in seasons of different A(H1N1)pdm09 and the same A(H3N2) vaccine components (2016–17 and 2017–18)

Esther Kissling^{a,*}, Francisco Pozo^b, Silke Buda^c, Ana-Maria Vilcu^d, Caterina Rizzo^{e,f}, Alin Gherasim^{g,h}, Judit Krisztina Horváthⁱ, Mia Brytting^j, Lisa Domegan^k, Adam Meijer^l, Iwona Paradowska-Stankiewicz^m, Ausenda Machadoⁿ, Vesna Višekruna Vučina^o, Mihaela Lazar^p, Kari Johansen^q, Ralf Dürwald^r, Sylvie van der Werf^{s,t}, Antonino Bella^e, Amparo Larrauri^{g,h}, Annamária Ferencziⁱ, Katherina Zakikhany^j, Joan O'Donnell^k, Frederika Dijkstra^l, Joanna Bogusz^m, Raquel Guiomar^u, Sanja Kurečić Filipović^o, Daniela Pitigoi^v, Pasi Penttinen^q, Marta Valenciano^a, I-MOVE/I-MOVE+ study team¹

^a Epidemiology Department, Epiconcept, 47 rue de Charenton, 75012 Paris, France

^b National Centre for Microbiology, National Influenza Reference Laboratory, WHO-National Influenza Centre, Institute of Health Carlos III, Madrid, Spain

^c Robert Koch Institute, Department of Infectious Disease Epidemiology, Respiratory Infections Unit, Seestrass 10, 13353 Berlin, Germany

^d Sorbonne Université, INSERM, Institut Pierre Louis d'épidémiologie et de Santé Publique (IPLESP UMRs 1136), F-75012, Paris, France

^e Department of Infectious Diseases, National Institute of Health, Rome, Italy

^f Bambino Gesù Children's Hospital, Rome, Italy

^g National Epidemiology Centre, Institute of Health Carlos III, Madrid, Spain

^h CIBER de Epidemiología y Salud Pública (CIBERESP), Institute of Health Carlos III, Madrid Spain

ⁱ Office of the Chief Medical Officer, Budapest, Hungary

^j Public Health Agency of Sweden, Stockholm, Sweden

^k Health Service Executive-Health Protection Surveillance Centre, 25-27 Middle Gardiner Street, Dublin 1 D01 A4A3, Ireland

^l National Institute for Public Health and the Environment (RIVM), Bilthoven, the Netherlands

^m National Institute of Public Health-National Institute of Hygiene, Warsaw, Poland

ⁿ Departamento de Epidemiologia, Instituto Nacional de Saúde Dr. Ricardo Jorge, Lisbon, Portugal

^o Croatian Institute of Public Health, Zagreb, Croatia

^p "Cantacuzino" National Medico-Military Institute for Research and Development, Bucharest, Romania

^q European Centre for Disease Prevention and Control (ECDC), Gustav III:s boulevard 40, 169 73 Solna, Sweden

^r Robert Koch Institute, National Reference Center for Influenza, Seestrass 10, 13353 Berlin, Germany

^s Unité de Génétique Moléculaire des Virus à ARN, Institut Pasteur, CNRS UMR3569, Université Paris Diderot SPC, France

^t CNR des Virus des Infections Respiratoires, WHO National Influenza Center, Institut Pasteur, France

^u Departamento de Doenças Infecciosas, Instituto Nacional de Saúde Dr. Ricardo Jorge, Portugal

^v Carol Davila University of Medicine and Pharmacy, Bucharest, Romania

* Corresponding author at: Esther Kissling, Epiconcept, 47 rue de Charenton, 75012 Paris, France.

E-mail address: e.kissling@epiconcept.fr (E. Kissling).

¹ I-MOVE/I-MOVE+ study team: list of co-authors: **Croatia:** Goranka Petrović, Croatian Institute of Public Health, Zagreb, Croatia; Zvezdana Lovrić Makarić, Croatian Institute of Public Health, Zagreb, Croatia; Iva Pem Novosel, Croatian Institute of Public Health, Zagreb, Croatia; Martina Zajec, Croatian Institute of Public Health, Zagreb, Croatia; Bernard Kaić, Croatian Institute of Public Health, Zagreb, Croatia. **France:** Sylvie Behillil, Unité de Génétique Moléculaire des Virus à ARN, UMR 3569 CNRS, Université Paris Diderot SPC, Institut Pasteur, Paris, France; CNR des virus des infections respiratoires, Institut Pasteur, Paris, France; Vincent Enouf, Unité de Génétique Moléculaire des Virus à ARN, UMR 3569 CNRS, Université Paris Diderot SPC, Institut Pasteur, Paris, France; CNR des virus des infections respiratoires, Institut Pasteur, Paris, France; Bruno Lina, Laboratoire de Virologie des HCL, Institut des Agents Infectieux, CNR des virus à transmission respiratoire, Groupement Hospitalier Nord, Lyon, France; Laboratoire Virpath, Centre International de Recherche en Infectiologie (CIRI), Université de Lyon, Inserm U1111, CNRS UMR 5308, ENS de Lyon, UCBL, Lyon, France; Martine Valette, Laboratoire de Virologie des HCL, Institut des Agents Infectieux, CNR des virus à transmission respiratoire, Groupement Hospitalier Nord, Lyon, France; Laboratoire Virpath, Centre International de Recherche en Infectiologie (CIRI), Université de Lyon, Inserm U1111, CNRS UMR 5308, ENS de Lyon, UCBL, Lyon, France. **Germany:** Ute Preuß, Robert Koch Institute, Department of Infectious Diseases, Berlin, Germany; Annicka Reuß, Robert Koch Institute, Department of Infectious Diseases, Berlin, Germany; Barbara Biere, Robert Koch Institute, National Reference Center for Influenza, Berlin, Germany; Maria Smallfield, Robert Koch Institute, National Reference Center for Influenza, Berlin, Germany; Brunhilde Schweiger, Robert Koch Institute, National Reference Center for Influenza, Berlin, Germany; Marianne Wedde, Robert Koch Institute, National Reference Center for Influenza, Berlin, Germany. **Hungary:** Beatrix Orszó, National Center for Epidemiology, Budapest, Hungary; Zita Vizler, Office of the Chief Medical Officer, Budapest, Hungary; Éva Hercegh, National Center for Epidemiology, Influenza Virus Laboratory, Budapest, Hungary; Bálint Szalai, National Center for Epidemiology, Influenza Virus Laboratory, Budapest, Hungary. **Ireland:** Laura Coughlan, Health Service Executive, Health Protection Surveillance Centre, Dublin, Ireland; Linda Dunford, National Virus Reference Laboratory, University College Dublin, Ireland; Joanne Moran, National Virus Reference Laboratory, University College Dublin, Ireland; Jeff Connell, National Virus Reference Laboratory, University College Dublin, Ireland; Cillian de Gascun, National Virus Reference Laboratory, University College Dublin, Ireland; Grainne Tuite, National Virus Reference Laboratory, University College Dublin, Ireland; Claire Collins, Irish College of General Practitioners, Dublin, Ireland; Michael Joyce, Irish College of General Practitioners, Dublin, Ireland; Olga Levis, Irish College of General Practitioners, Dublin, Ireland.

<https://doi.org/10.1016/j.jvacx.2019.100042>

2590-1362/© 2019 The Authors. Published by Elsevier Ltd.

This is an open access article under the CC BY-NC-ND license (<http://creativecommons.org/licenses/by-nc-nd/4.0/>).

ARTICLE INFO

Article history:

Received 16 May 2019

Received in revised form 13 September 2019

Accepted 15 September 2019

Available online 17 September 2019

Keywords:

Influenza

Influenza vaccine

Vaccine effectiveness

Multicentre study

Case-control study

Europe

ABSTRACT

Introduction: Influenza A(H3N2) viruses predominated in Europe in 2016–17. In 2017–18 A(H3N2) and A(H1N1)pdm09 viruses co-circulated. The A(H3N2) vaccine component was the same in both seasons; while the A(H1N1)pdm09 component changed in 2017–18. In both seasons, vaccine seed A(H3N2) viruses developed adaptations/alterations during propagation in eggs, impacting antigenicity.

Methods: We used the test-negative design in a multicentre primary care case-control study in 12 European countries to measure 2016–17 and 2017–18 influenza vaccine effectiveness (VE) against laboratory-confirmed influenza A(H1N1)pdm09 and A(H3N2) overall and by age group.

Results: During the 2017–18 season, the overall VE against influenza A(H1N1)pdm09 was 59% (95% CI: 47–69). Among those aged 0–14, 15–64 and ≥ 65 years, VE against A(H1N1)pdm09 was 64% (95% CI: 37–79), 50% (95% CI: 28–66) and 66% (95% CI: 42–80), respectively. Overall VE against influenza A(H3N2) was 28% (95% CI: 17–38) in 2016–17 and 13% (95% CI: –15 to 34) in 2017–18. Among 0–14-year-olds VE against A(H3N2) was 28% (95% CI: –10 to 53) and 29% (95% CI: –87 to 73), among 15–64-year-olds 34% (95% CI: 18–46) and 33% (95% CI: –3 to 56) and among those aged ≥ 65 years 15% (95% CI: –10 to 34) and –9% (95% CI: –74 to 32) in 2016–17 and 2017–18, respectively.

Conclusions: Our study suggests the new A(H1N1)pdm09 vaccine component conferred good protection against circulating strains, while VE against A(H3N2) was $< 35\%$ in 2016–17 and 2017–18. The egg propagation derived antigenic mismatch of the vaccine seed virus with circulating strains may have contributed to this low effectiveness. A(H3N2) seed viruses for vaccines in subsequent seasons may be subject to the same adaptations; in years with lower than expected VE, recommendations of preventive measures other than vaccination should be given in a timely manner.

© 2019 The Authors. Published by Elsevier Ltd. This is an open access article under the CC BY-NC-ND license (<http://creativecommons.org/licenses/by-nc-nd/4.0/>).

1. Introduction

WHO recommended the same influenza A(H3N2) Northern Hemisphere vaccine component for the 2016–17 and 2017–18 influenza seasons: A/Hong Kong/4801/2014 (H3N2)-like virus. An A/California/7/2009 (H1N1)pdm09-like virus vaccine component was recommended for the seventh consecutive year in the 2016–17 season. This was replaced with an A/Michigan/45/2015 (H1N1)pdm09-like virus vaccine component in the 2017–18 season.

Influenza A(H3N2) was the predominant circulating influenza virus in Europe in the 2016–17 season with very little A(H1N1)pdm09 and B circulating [1,2]. While overall in Europe in the 2017–18 season influenza B/Yamagata virus lineage-mismatched to the trivalent vaccine was the main circulating strain, both A(H1N1)pdm09 and A(H3N2) circulated in varying patterns across

countries [2]. Excess all-cause mortality was seen in both seasons, particularly among the elderly, in 2016–17 coinciding with a pre-dominance of A(H3N2) [3,4].

Since 2008–9, the I-MOVE/I-MOVE+ (Influenza Monitoring Vaccine Effectiveness in Europe) primary care multicentre case control study (MCCS) has provided vaccine effectiveness (VE) estimates by influenza virus (sub)type, age group, and target population. Since 2012–13, VE has also been estimated by vaccine type, and since 2015–16, by virus genetic clade [5–7].

We present the I-MOVE/I-MOVE+MCCS VE estimates against influenza A by subtype for the 2016–17 and 2017–18 seasons. We estimate VE by age group (including birth cohorts who may be more susceptible for infection [8]), target population, previous vaccination, vaccine type, time within the season and also estimate VE against genetic clade. The 2017–18 VE against influenza B

Italy: Maria Rita Castrucci, National Influenza Center National Institute of Health, Rome Italy; Simona Puzelli, National Influenza Center National Institute of Health, Rome Italy; Valeria Alfonsi, Department of Infectious Diseases, National Institute of Health, Rome Italy; Eva Appelgreen, Department of Infectious Diseases, National Institute of Health, Rome Italy. **The Netherlands:** Marit de Lange, National Institute for Public Health and the Environment (RIVM), Bilthoven, Netherlands; Sharon van den Brink, National Institute for Public Health and the Environment (RIVM), Bilthoven, Netherlands; Pieter Overduin, National Institute for Public Health and the Environment (RIVM), Bilthoven, Netherlands; Anne Teirlinck, National Institute for Public Health and the Environment (RIVM), Bilthoven, Netherlands; Daphne Reukers, National Institute for Public Health and the Environment (RIVM), Bilthoven, Netherlands; Wim van der Hoek, National Institute for Public Health and the Environment (RIVM), Bilthoven, Netherlands; Gé Donker, Netherlands Institute for Health Services Research (Nivel), Utrecht, the Netherlands; Mariette Hooiveld, Netherlands Institute for Health Services Research (Nivel), Utrecht, the Netherlands. **Portugal:** Verónica Gomez, Departamento de Epidemiologia, Instituto Nacional de Saúde Dr. Ricardo Jorge, Lisbon, Portugal; Irina Kislaya, Departamento de Epidemiologia, Instituto Nacional de Saúde Dr. Ricardo Jorge, Lisbon, Portugal; Baltazar Nunes, Departamento de Epidemiologia, Instituto Nacional de Saúde Dr. Ricardo Jorge, Lisbon, Portugal; Ana Paula Rodrigues, Departamento de Epidemiologia, Instituto Nacional de Saúde Dr. Ricardo Jorge, Lisbon, Portugal; Patrícia Conde, Departamento de Doenças Infeciosas, Instituto Nacional de Saúde Dr. Ricardo Jorge, Portugal; Inês Costa, Departamento de Doenças Infeciosas, Instituto Nacional de Saúde Dr. Ricardo Jorge, Portugal; Paula Cristóvão, Departamento de Doenças Infeciosas, Instituto Nacional de Saúde Dr. Ricardo Jorge, Portugal; Pedro Pechirra, Departamento de Doenças Infeciosas, Instituto Nacional de Saúde Dr. Ricardo Jorge, Portugal. **Romania:** Emilia Lupulescu, “Cantacuzino” National Medico-Military Institute for Research and Development, Bucharest; Ivanciuc Alina, “Cantacuzino” National Medico-Military Institute for Research and Development, Bucharest; Mihai Maria Elena, “Cantacuzino” National Medico-Military Institute for Research and Development, Bucharest; Carmen Maria Cherciu “Cantacuzino” National Medico-Military Institute for Research and Development, Bucharest; Elena Stoian, “Cantacuzino” National Medico-Military Institute for Research and Development, Bucharest. **Spain:** Clara Mazagatos, National Epidemiology Centre, Institute of Health Carlos III, Madrid Spain; CIBER de Epidemiología y Salud Pública (CIBERESP), Institute of Health Carlos III, Madrid Spain; Inmaculada Casas, National Centre for Microbiology, National Influenza Reference Laboratory, WHO-National Influenza Centre, Institute of Health Carlos III, Madrid, Spain; Luis García Comas, Dirección General de Salud Pública, Comunidad de Madrid, Madrid, Spain; María Esther Insua Marisquerena, Dirección General de Salud Pública, Comunidad de Madrid, Madrid, Spain; Juan Carlos Galán, Laboratorio Hospital Ramón y Cajal, CIBER de Epidemiología y Salud Pública (CIBERESP), Madrid, Spain; Fernando Gonzalez Carril, Departamento de Salud, Gobierno del País Vasco, Spain; Rosa Sancho Martínez, Departamento de Salud, Gobierno del País Vasco, Spain; Gustavo Cilla, Laboratorio Hospital Donostia, CIBERESP, Spain; Jesús Castilla, Navarra Institute for Health Research (IdiSNA), Pamplona, CIBERESP, Spain; Manuel García Cenoz, Navarra Institute for Health Research (IdiSNA), Pamplona, CIBERESP, Spain; Ana Navascués, Complejo Hospitalario de Navarra, Spain; Carmen Quiñones Rubio, Dirección General de Salud Pública y Consumo de La Rioja, Spain; Eva Martínez Ochoa, Dirección General de Salud Pública y Consumo de La Rioja, Spain; Miriam Blasco, Laboratorio Hospital San Pedro de Logroño, Spain; Jaume Gimenez Duran Servicio de Epidemiología, Dirección General de Salud Pública, Mallorca, Spain; Juana Maria Vanrell, Servicio de Epidemiología, Dirección General de Salud Pública, Mallorca, Spain; Jordi Reina, Laboratorio del Hospital Son Espases, Mallorca, Spain; Daniel Castrillejo, Servicio de Epidemiología, DGSC, Consejería de Bienestar Social y Sanidad, Ciudad Autónoma de Melilla, Spain. **Sweden:** Åsa Wiman, Public Health Agency of Sweden, Stockholm, Sweden; AnnaSara Carnahan, Public Health Agency of Sweden, Stockholm, Sweden; **Epiconcept:** Alain Moren, Epidemiology Department, Epiconcept, Paris, France.

lineage-mismatched to the trivalent vaccine will be addressed elsewhere in context of VE against influenza B in post-pandemic seasons.

2. Methods

Eleven European study sites (in Croatia, France, Germany, Ireland, Italy, Poland, Portugal, Romania, Spain, Sweden and the Netherlands) participated in our 2016–17 and 2017–18 I-MOVE/I-MOVE+MCCS, while one (in Hungary) participated in the in the 2016–17 MCCS only. Each study site used the test-negative design using the European Centre for Disease Prevention and Control (ECDC) generic case-control study protocol and the I-MOVE+protocol [9,10]. The methods are described in detail elsewhere [5,11].

Briefly, for each season, participating practitioners interviewed and collected nasopharyngeal or combined naso- and oropharyngeal specimens from a systematic sample of consenting patients seeking medical attention for influenza-like illness (ILI). Practitioners collected information including symptoms, date of onset and swabbing, current seasonal influenza vaccination status, date of influenza vaccination and vaccine product, seasonal influenza vaccination status from the previous season, sex, age and presence of chronic medical conditions in the past 12 months.

In the pooled analysis we included patients meeting the European Union ILI case definition [12], swabbed within 7 days of symptom onset, and who had not received antivirals in the 14 days prior to swabbing. Study sites with fewer than 10 influenza-positive cases by influenza subtype or with fewer than 10 vaccinated patients were excluded from the pooled analysis.

A case of confirmed influenza was an ILI patient who had been swabbed and whose test result was positive for influenza A virus using real-time, reverse-transcription polymerase chain reaction (RT-PCR). Controls were ILI patients who tested negative for any influenza virus using RT-PCR.

We defined a person as vaccinated if they had received at least one dose of current seasonal influenza vaccine more than 14 days before ILI symptom onset. Those vaccinated fewer than 15 days before ILI onset were excluded. All other patients were classified as unvaccinated.

2.1. Statistical methods

We conducted a complete case analysis, in which patients with missing values for any of the variables in the model measuring adjusted VE are excluded. We computed the pooled VE for each influenza A subtype as $(1-OR) * 100$ using a one-stage logistic regression model with study site as a fixed effect.

We calculated VE adjusting for a priori confounding factors: symptom onset date, age, sex, and presence of at least one chronic disease or other risk conditions such as pregnancy and obesity (where available). The continuous variables symptom onset date and age in years were modelled as restricted cubic splines with 3, 4 or 5 knots, or as categorical variables (age group, onset month or onset week), the choice of which was determined by sample size and the Akaike information criterion. We used the “one in ten” rule of covariate degrees of freedom to events to determine if we were overfitting the model [13]. If the number of events/parameters was less than 10, we conducted a sensitivity analysis using Firth’s method of penalised logistic regression [14].

To study the effect of prior seasonal influenza vaccination on the current season VE, we conducted an indicator analysis using four categories: individuals unvaccinated in both seasons (reference group), vaccinated in the previous season only, vaccinated in the current season only, and those vaccinated in both seasons.

We did not measure effect of previous vaccination among children aged <9 years, due to their multi-seasonal vaccination schedule.

We measured VE by age group (0–14, 15–64 and 65 years and older) and in the target group for vaccination, which includes persons aged 60 or 65 years and older (depending on study site), persons with underlying medical conditions and other risk groups [15]. For the A(H1N1)pdm09 age-specific analysis among adults, we also stratified VE by the Linderman birth cohort of heightened susceptibility (1965–1979) due to influenza pre-exposure history, resulting in the age groups of 15–37, 38–52, and 53 years and over [8]. We measured VE by type of vaccine (inactivated subunit, inactivated split virion and MF59 adjuvanted), including only study sites in the analysis where vaccines of that type were available.

We measured VE by calendar time, estimating influenza A subtype-specific VE from October to December, in January and in February–April, in order to obtain early, peak and end of season estimates, respectively.

Data management and statistical analyses were carried out using Stata 15.1 [16].

2.2. Laboratory methods

Nine study sites in 2016–17 and eight in 2017–18 selected either all influenza virus-positive specimens or a random proportion of specimens for sequencing the haemagglutinin gene segment (HA) for each influenza A subtype. HA consensus sequences were uploaded by each site to GISAID and downloaded for centralised phylogenetic and amino acid substitution analysis of the HA1 coding portion in MEGA6 to determine clade distribution at the National Influenza Centre, Madrid. Amino acid composition analysis was done relative to the egg-adapted high growth reassortant X-263B vaccine strain of A/Hong Kong/4801/2014 H3N2 (GISAID accession number EPI731469). For H1N1pdm09 viruses the egg-adapted high growth reassortant X-179A vaccine strain of A/California/07/2009 (GISAID accession number EPI257201) and X-275A vaccine strain of A/Michigan/45/2015 (GISAID accession number EPI830230) were used as a reference for seasons 2016–17 and 2017–18, respectively.

3. Results

In the 2016–17 season we included 4909 influenza A(H3N2) cases and 6098 test-negative controls. Due to low numbers of influenza A(H1N1)pdm09 (N = 118) and B (N = 190) cases, no VE analysis against these was attempted (Fig. 1).

In the 2017–18 season, we included 1958 influenza A(H1N1)pdm09 cases, 756 influenza A(H3N2) cases and 5887 test-negative controls in the I-MOVE/I-MOVE+MCCS VE analyses (Fig. 1).

3.1. Participant profile and virological description

3.1.1. Influenza A(H3N2): 2016–17 and 2017–18

The median age of A(H3N2) cases was 32 years in 2016–17 and 41 years in 2017–18, compared to 31 and 32 years in controls in the same seasons, respectively (Supplementary Table 1). The vaccine coverage among cases was 11% in 2016–17 and 16% in 2017–18, compared to 12% in controls in both seasons.

In the 2016–17 and 2017–18 influenza seasons we included 1050 and 312 sequenced A(H3N2) viruses, respectively in the genetic analysis (Table 1). In 2016–17, 256 (24%) belonged to the 3C.2a clade represented by A/HongKong/4801/2014, 781 (74%) belonged to the 3C.2a1 clade represented by A/Singapore/INFIM H-16-0019/2016 (in 2016–17 the representative virus for this clade was A/Bolzano/7/2016, however we use the 2017–18




Fig. 1. Number of ILI patients by case status (test negative controls and cases by influenza (sub)type) and week of symptom onset, I-MOVE/I-MOVE+ primary care multicenter case control study, Europe, influenza seasons 2016–17 and 2017–18. ILI: Influenza-like illness; ISO: International Organization for Standardization.

representative virus for easier comparison between seasons) and 13 (1%) belonged to the 3C.3a clade represented by A/Switzerland/9715293/2013. In 2017–18, 173 (55%) belonged to the 3C.2a clade, 134 (43%) belonged to the 3C.2a1 clade and 5 (2%) belonged to the 3C.3a clade.

In 2016–17 there was a variety of amino acid substitutions in antigenic sites within the 3C.2a and 3C.2a1 clades in most studies, the proportion of which varied between countries (Table 1).

Conversely, in 2017–18, 167/173 (97%) of the viruses of the A/HongKong/4801/2014 3C.2a clade belonged to the 3C.2a2 subclade with the T131K, R142K and R261Q amino acid substitutions relative to the vaccine strain A/HongKong/4801/2014. Among the 134 viruses in 2017–18 belonging to the A/Singapore/INFIMH-16-0019/2016 3C.2a1 clade, 132 (99%) belonged to the 3C.2a1b subclade with the N171K, N121K, K92R and H311Q amino acid substitutions relative to the vaccine strain A/HongKong/4801/2014.

3.1.2. Influenza A(H1N1)pdm09: 2017–18

The median age of influenza A(H1N1)pdm09 cases was 19 years, compared to 32 years for influenza test-negative controls (Supplementary Table 1). Among A(H1N1)pdm09 cases, 47% belonged to the 0–14 years age group, compared to 33% of controls. The vaccination coverage was 4% among A(H1N1)pdm09 cases, compared to 12% among controls.

All A(H1N1)pdm09 sequenced viruses belonged to the clade represented by the A/Michigan/45/2015 vaccine virus (clade 6B.1) (Table 1).

3.1.3. Vaccines used in 2016–17 and 2017–18

In both seasons, all vaccines used were egg-propagated. Among vaccinated controls with known vaccine brand in 2016–17 (572/685, 84%), 100 (17%) received adjuvanted vaccine, 52 with aluminium phosphate gel and 48 with MF59 adjuvant, and nine (2%) received a live attenuated influenza vaccine (LAIV) (Supplementary table 1). Among vaccinated controls with known vaccine brand in 2017–18 (556/687, 81%), 30 (5%) received adjuvanted vaccine, all with MF59 adjuvant, and eight vaccinated participants (1%) received LAIV. In 2016–17 and 2017–18, 79% and 84% of vaccines with known brand were inactivated non-adjuvanted trivalent vaccines.

3.2. 2016–17 and 2017–18 vaccine effectiveness estimates against A (H3N2)

3.2.1. Overall

The VE against influenza A(H3N2) among all ages in 2016–17 was 28% (95% CI: 17–38) and in 2017–18 13% (95% CI: –15 to 34) (Table 2). Among all ages in the target group for vaccination the VE was 22% (95% CI: 7–34) in 2016–17 and 13% (95% CI: –21 to 38) in 2017–18

3.2.2. By age group

The VE against A(H3N2) in 2016–17 was 28% (95% CI: –10 to 53), 34% (95% CI: 18–46) and 15% (95% CI –10 to 34) among those aged 0–14, 15–64 and ≥ 65 years, respectively.

The VE against influenza A(H3N2) in 2017–18 was 29% (95% CI: –87 to 73), 33% (95% CI: –3 to 56) and –9% (–74 to 32) among those aged 0–14, 15–64 and ≥ 65 years, respectively.

3.2.3. VE by clade/subclade

The VE in 2016–17 against all variants of 3C.2a excluding 3C.3a1 was 31% (95% CI: –12 to 57) and the VE against the 3C.2a3 subclade was 37% (95% CI: –9 to 63) (Table 3). In 2016–17 the VE against all 3C.2a1 variants was 25% (95% CI: –1 to 44). The VE against 3C.2a1 with no mutations in A-E antigenic sites except for N171K was 56% (95% CI: 20–76). The VE against the 3C.2a1b subclade was 7% (95% CI: –56 to 45).

In 2017–18 the VE against all variants of 3C.2a excluding 3C.3a1 was –29% (–109 to 21). The vast majority (160/166) of 3C.2a viruses belonged to the subclade 3C.2a2, against which the VE was –19 (95% CI: –95 to 27). The VE against 3C.2a1b was 43% (95% CI: –16 to 72).

3.2.4. By vaccine group

In 2016–17, among all ages, the VE of the trivalent subunit vaccine against A(H3N2) was 31% (95% CI: 13–46) and of the trivalent split virion vaccine was 23% (95% CI: 4–39). The VE of the MF59 adjuvanted trivalent subunit vaccine, used in three study sites, against A(H3N2) was 49% (95% CI: 12–71).

In 2017–18, the VE of the trivalent subunit vaccine against A (H3N2) was –3% (95% CI: –50 to 30) and the VE of the trivalent split virion vaccine against A(H3N2) was 19 (95% CI: –29 to 49).

Table 1

Genetic group distribution among nine study sites participating in the random sequencing of influenza virus positive specimens, I-MOVE/I-MOVE+ primary care multicenter case control study, Europe, influenza season 2016–17 and 2017–18.

	Characterised viruses ¹	Clade/subclade	DE		ES		FR		HU		IE		NL		PT		RO		SE		Total	
			N	%	N	%	N	%	N	%	N	%	N	%	N	%	N	%	N	%	N	%
2016–17	A(H3N2)		207		346		87		51		63		80		124		25		67		1050	
	A/HongKong/4801/2014-like*	3C.2a	47	23	59	17	28	32	15	29	14	22	16	20	42	34	10	40	25	37	256	24
			1	2	0	0	0	0	0	0	0	0	2	13	2	5	0	0	0	0	5	2
	N121K+S144K+(N122D+262N)	3C.2a3	30	64	54	92	27	96	14	93	11	79	8	50	38	90	10	100	16	64	208	81
	T131K+R142K+R261Q	3C.2a2	16	34	5	8	1	4	1	7	3	21	6	38	2	5	0	0	9	36	43	17
	A/Bolzano/7/2016-like ²	3C.2a1	159	77	280	81	59	68	36	71	45	71	63	79	82	66	15	60	42	63	781	74
	N171K		14	9	137	49	40	68	1	3	1	2	9	14	1	1	6	40	7	17	216	28
	N171K+N121K+H140M	–	32	20	58	21	7	12	2	5	6	13	14	22	6	7	7	47	8	19	140	18
	N171K+N121K+K92R+H311Q	3C.2a1b	43	27	59	21	1	2	4	11	5	11	18	29	29	35	1	7	1	2	161	21
	N171K+N121K+T135K	3C.2a1a	54	34	8	3	2	3	28	78	1	2	11	17	0	0	1	7	22	52	128	16
	N171K+R142G	–	16	10	18	6	9	15	1	3	32	71	11	17	46	56	0	0	4	10	137	18
	A/Switzerland/9715293/2013-like	3C.3a	1	0	7	2	0	0	0	0	4	6	1	1	0	0	0	0	0	0	13	1
	Characterised viruses	Clade/subclade	DE		ES		FR		HU		IE		NL		PT		RO³		SE		Total	
			N	%	N	%	N	%	N	%	N	%	N	%	N	%	N	%	N	%	N	%
2017–18	A(H1N1)pdm09		43		10		64		–		13		31		22		13		9		205	
	A/Michigan/45/2015-like	6B.1	43	100	10	100	64	100			13	100	31	100	22	100	13	100	9	100	205	100
	A(H3N2)		25		141		35		–		48		26		13		–		24		312	
	74A/HongKong/4801/2014-like*	3C.2a	23	92	44	31	25	71	–		41	85	18	69	7	54	–		15	63	173	55
			0	0	0	0	0	0	–		0	0	0	0	0	0	–		0	0	0	0
	N121K+S144K+(N122D+262N)	3C.2a3	2	9	0	0	1	4	–		3	7	0	0	0	0	–		0	0	6	3
	T131K+R142K+R261Q	3C.2a2	21	91	44	100	24	96	–		38	93	18	100	7	100	–		15	100	167	97
	A/Singapore/INFIMH-16-0019/2016-like ²	3C.2a1	2	8	97	69	8	23	–		6	13	8	0	6	46	–		7	29	134	43
	N171K+N121K		0	0	0	0	0	0	–		0	0	0	0	0	0	–		0	0	0	0
	N171K+N121K+H140M	–	0	0	0	0	0	0	–		0	0	0	0	0	0	–		0	0	0	0
	N171K+N121K+K92R+H311Q	3C.2a1b	1	50	96	99	8	100	–		6	100	8	100	6	100	–		7	100	132	99
	N171K+N121K+T135K	3C.2a1a	1	50	1	1	0	0	–		0	0	0	0	0	0	–		0	0	2	1
	N171K+R142G	–	0	0	0	0	0	0	–		0	0	0	0	0	0	–		0	0	0	0
	A/Switzerland/9715293/2013-like	3C.3a	0	0	0	0	2	6	–		1	2	0	0	0	0	–		2	8	5	2
	Total 2017–18		68		151		89		–		61		57		35		13		33		517	

* Viruses with no amino acid substitutions or substitutions outside of the A–E antigenic sites.

¹ We use for both seasons the clade/subclade denomination created in mid-season in 2017–18 (report prepared by the Francis Crick Institute for the WHO annual consultation on the composition of influenza vaccine for the Northern Hemisphere 2018–2019). Therefore some clades/subclades circulating in 2016–17, but not in 2017–18, have no specific clade/subclade denomination. All amino acid substitutions shown for the A(H3N2) overarching clade 3C.2a viruses are relative to the vaccine virus A/HongKong/4801/2014.

² A/Bolzano/7/2016 and A/Singapore/INFIMH-16-0019/2016 are two representative viruses of the same clade 3C.2a1, carrying the N171K substitution in the HA gene compared to the vaccine virus A/HongKong/4801/2014. In 2016–17, A/Bolzano/7/2016 was used as the representative virus and in 2017–18 A/Singapore/INFIMH-16-0019/2016 was used being representative for clade 3C.2a1 viruses carrying N121K in addition to N171K and chosen as vaccine strain for the 2018 Southern Hemisphere and 2018/2019 Northern Hemisphere vaccines.

³ Romania sequenced 7 A(H3N2) viruses, but they are excluded from the pooled analysis as they had fewer than 10 cases.

Table 2
Pooled adjusted seasonal vaccine effectiveness against influenza A(H1N1)pdm09 and A(H3N2), overall, by age groups and by vaccine type, previous vaccination and time within the season. I-MOVE/I-MOVE+ primary care multicenter study, Europe, influenza seasons 2016–17 and 2017–18.

Influenza A(H3N2) 2016–17					
Age group	Population	N ^a	Cases; vacc /Controls; vacc ^b	Adjusted VE (%)	95% CI (%)
All ages		10,591	4733;516/5858;680	28	17–38
0–14 years		3452	1494;46/1958;66	28	–10 to 53
15–64 years		5840	2606;183/3234;286	34	18–46
≥65 years		1298	633;287/665;328	15	–10 to 34
All ages	Target group for vaccination	3181	1440;431/1741;543	22	7–34
All ages	Subunit vaccine ^a	8304	3832;179/4472;238	31	13–46
	Split virion vaccine ^a	9303	4209;188/5094;220	23	4–39
	Adjuvanted vaccine ^a	4678	2027;32/2651;48	49	12–71
≥9 years	Unvaccinated	7822	3168/3343 ^b	Ref	
	2016–17 vaccine only		69/132 ^b	50	32–63
	2015–16 vaccine only		87/144 ^b	29	5–47
	2015–16 and 2016–17 vaccines		412/467 ^b	20	5–32
All ages	Oct-Dec	3541	1442;141/2099;206	38	16–54
	Jan	4379	2389;260/1990;289	34	18–47
	Feb-April	2671	902;115/1769;185	1	–34 to 27
Influenza A(H3N2) 2017–18					
Age group	Population	N	Cases; vacc /Controls; vacc	Adjusted VE (%)	CI (%)
All ages		5607	731;119/4876;623	13	–15 to 34
0–14 years		1772	160;6/1612;74	29	–87 to 73
15–64 years		3128	444;33/2684;245	33	–3 to 56
≥65 years		673	127;80/546;283	–9	–74 to 32
All ages	Target group	1763	251;106/1512;512	13	–21 to 38
All ages	Subunit vaccine ^a	4556	471;56/3739;290	–3	–50 to 30
	Split virion vaccine ^a	4637	555;36/3906;140	19	–29 to 49
≥9 years	Unvaccinated	4051	491/2783 ^b		
	17–18 vaccine only		11/105 ^b	49	1–74
	16–17 vaccine only		17/119 ^b	16	–47 to 52
	16–17 and 17–18 vaccines		95/430 ^b	7	–29 to 33
All ages	Oct-Dec	1413	138;19/1275;133	11	–72 to 54
	Jan	1849	230;42/1619;234	5	–54 to 41
	Feb-April	2345	363;58/1982;256	18	–24 to 45
Influenza A(H1N1)pdm09 2017–18					
Age group	Population	N	Cases; vacc /Controls; vacc	Adjusted VE (%)	CI (%)
All ages		7175	1875;81/5300;643	59	47–69
0–14 years		2680	888;18/1792;80	64	37–79
15–64 years		3806	893;37/2913;256	50	28–66
≥65 years		684	94;26/590;307	66	42–80
15–37 years		1609	339;4/1270;54	71	18 to 90
38–52 years		1311	369;18/942;70	23	–38 to 57
53+years		1572	279;41/1293;439	64	47 to 76
All ages	Target group	2075	424;65/1651;526	56	40–68
All ages	Subunit vaccine ^a	6268	1745;39/4192;292	59	40–71
	Split virion vaccine ^a	6184	1745;11/4288;140	71	45–85
≥9 years	Unvaccinated	4784	981/3023 ^b		
	17–18 vaccine only		6/110 ^b	79	51–91
	16–17 vaccine only		25/137 ^b	24	–22 to 52
	16–17 and 17–18 vaccines		61/441 ^b	46	26–61
All ages	Oct-Dec	2009	326;4/1683;152	87	62–95
	Jan	2277	673;33/1604;230	53	27–69
	Feb-April	2826	856;44/1970;257	53	32–67

CI: confidence interval; vacc: vaccinated; VE: vaccine effectiveness.

^a Among study sites where this vaccine type was used. Subunit and split virion vaccine refers to egg-propagated inactivated trivalent subunit and split virion vaccines, respectively. For adjuvanted vaccine only study sites using the MF59 adjuvant included.

^b Cases/controls.

3.2.5. By previous vaccination

In 2016–17, using those unvaccinated in current and prior season as a reference group, the VE among those aged ≥9 years was 50% (95% CI: 32–63) in those vaccinated in the current season only, 20% (95% CI: 5–32) in those vaccinated in both seasons, and 29% (95% CI: 5–47) in those vaccinated in previous season only.

In 2017–18, using those unvaccinated in current and prior season as a reference group, the VE among those aged ≥9 years was

49% (95% CI: 1–74) in those vaccinated in the current season only, 7% (95% CI: –29 to 33) in those vaccinated in both seasons and 16% (95% CI: –47 to 52) in those vaccinated in previous season only.

3.2.6. By calendar time

In 2016–17 the VE was 38% (95% CI: 16–54) in October to December, 34% (95% CI: 18–47) in January and 1% (95% CI: –34 to 27) in February to April among all ages.

Table 3

Pooled adjusted seasonal vaccine effectiveness against influenza A(H3N2) clades and subclades, I-MOVE/I-MOVE+ primary care multicenter study, Europe, influenza seasons 2016–17 and 2017–18.

Season	Clade/subclade	N	Cases; vacc /Controls; vacc	Adjusted VE (%)	CI (%)
2016–17	3C.2a (all variants excluding 3C.2a1)	4492	252;28/4240;496	31	–12 to 57
	3C.2a3	4445	205;20/4240;496	37	–9 to 63
	N121K+S144K+(N122D+262 N)				
	3C.2a1 (all variants)	5007	767;99/4240;496	25	–1 to 44
	3C.2a1 (no mutations in A-E antigenic sites other than N171K)	4454	214;19/4240;496	56	20 to 76
	3C.2a1b	4397	157;30/4240;496	7	–56 to 45
2017–18	N171K+N121K+K92R+H311Q				
	3C.2a (all variants)	3732	166;40/3566;445	–29	–109 to 21
	3C.2a2	3726	160;38/3566;445	–19	–95 to 27
	T131K+R142K+R261Q				
	3C.2a1 (all variants)	3698	132;17/3566;445	41	–19 to 71
	3C.2a1b	3697	131;17/3566;445	42	–18 to 71
	N171K+N121K+K92R+H311Q				

CI: confidence interval; vacc: vaccinated; VE: vaccine effectiveness.

In 2017–18 the VE was 11% (95% CI: –72 to 54), 5% (95% CI: –54 to 41) and 18% (95% CI: –24 to 45) in October to December, January and in February to April among all ages, respectively.

3.3. 2017–18 vaccine effectiveness estimates against A(H1N1)pdm09

3.3.1. Overall and by age group

In the 2017–18, 1875 A(H1N1)pdm09 cases and 5300 test-negative controls were included in the complete case VE analysis. The overall VE was 59% (95% CI: 47–69). Among the target group for vaccination (N = 2075) the VE was 56% (95% CI: 42–68) (Table 2).

The VE was 64% (95% CI: 37–79), 50% (95% CI: 28–66%) and 66% (95% CI 42–80) among patients aged 0–14, 15–64 and ≥64 years, respectively.

The VE among those aged 15–37 years was 71% (95% CI: 18–90), among those aged 38–52 years (corresponding to the Linderman birth cohort of susceptibility of 1965–79) 23% (95% CI: –38 to 57) and among those aged 53 and over 64% (95% CI: 47–76).

3.3.2. By vaccine type

VE was 59% (95%CI: 40–71) for the trivalent inactivated subunit vaccine and 71% (95% CI: 45–85) for the trivalent inactivated split virion vaccine. Numbers were too low to estimate VE for the MF59 adjuvanted vaccine.

3.3.3. By previous vaccination

In the ≥9 years olds (N = 4784), compared to those receiving no vaccine in current or prior season, the VE was 79% (95% CI: 51–91) among those receiving the 2017–18 vaccine only and 46% (95% CI: 26–61) among those receiving both 2016–17 and 2017–18 vaccine.

3.3.4. By calendar time

VE was 87% (95% CI: 62–95) in October to December, 53% (95% CI: 27–69) in January and 53% (95% CI: 32–67%) in February to April (see Table 2).

4. Discussion

In 2017–18 the influenza VE against primary care medically attended ILI due to influenza A(H1N1)pdm09 in the I-MOVE/I-MOVE+MCCS in Europe was 56 and 59% in the total and target population, respectively. The VE against A(H3N2) in 2016–17 was 22% and 28% in the total and target populations, respectively, and in 2017–18 it was 13% in both populations.

4.1. Influenza A(H1N1)pdm09

The 2017–18 season was the first season of the new A/Michigan/45/2015-like virus vaccine component after seven consecutive seasons of the A/California/7/2009 (H1N1)-like virus vaccine component. The A/California/7/2009 (H1N1)-like virus was maintained from 2010–11 to 2016–17 in the Northern Hemisphere vaccine. Despite evolution of the A(H1N1)pdm09 viruses into distinct genetic clades there was no significant antigenic drift, making a vaccine strain switch unnecessary until recently. Clades 6B and 6C started to circulate in Europe in 2012–13. From 2014 to 15 onwards, all genetically characterised viruses reported to ECDC belonged to genetic subgroup 6B, characterised by a K163Q substitution in HA1 [17,18]. In 2015–16, 90% (652/723) of all systematically selected A(H1N1)pdm09 viruses characterised in the I-MOVE/I-MOVE+MCCS belonged to the 6B.1 subgroup, defined by the S84N, I216T and S162N substitutions. This latter substitution may result in a potential new glycosylation site [6]. These recent genetic clades were antigenically distinct when using human antisera following vaccination [1,18]. This was also reflected in the 2015–16 VE against A(H1N1)pdm09 of 33% (95% CI: –4 to 57) among all ages in the I-MOVE/I-MOVE+MCCS, which was lower than previous seasons. This decline in VE across seasons was also seen in Canada, but not in the UK [6,7,15,19–21]. The I-MOVE/I-MOVE+MCCS 2017–18 season VE of 59% (95% CI: 47–69) indicates that the vaccine component change was successful, underlining the importance of complementation of antigenic characterisation of viruses using naïve ferret antisera with antisera derived from vaccinated humans from different age groups [22].

The process of “original antigenic sin” is where human antibody responses cross-react to influenza strains experienced in early childhood when challenged by a newer viral strain [23]. In 2014, Linderman et al. proposed that this process explained a heightened susceptibility to the circulating A(H1N1)pdm09 6B clade in some birth cohorts. These people may have experienced childhood exposure to A(H1N1) viruses with the K163 epitope, not the case in younger adults whose childhood exposure was to A(H1N1) viruses with this epitope masked [8]. Linderman et al. reported that particular birth cohorts 1965 to 1979 had K163-specific antibodies in their sera. They therefore recommended a change in vaccine from the A/California/7/2009 (H1N1)-like virus, whilst indicating nevertheless the uncertainty that a vaccine change will break the “original antigenic sin”. The VE against A(H1N1)pdm09 among the 1965–1979 birth cohort (38–52 years) in the 2017–18 I-MOVE/I-MOVE+MCCS was 22%, lower than among other age groups. This VE against A(H1N1)pdm09 is similar to the VE reported in 2015–16 in the US among similar birth cohorts (VE = 23%) [24]. The dip

in VE was seen in slightly older age groups in Canada and was not apparent in the 2015–16 I-MOVE/I-MOVE+MCCS, where VE was low overall [6,20]. The age-specific VE we observe in the our study in the 2017–18 season may be due to random variation, or due to a more complex interplay of influenza virus exposure and vaccinations [20]. Monitoring age-specific effects is warranted to better understand susceptible populations in future influenza seasons.

Our interim VE results against A(H1N1)pdm09 were higher at 68%, with data from the start of the 2017–18 season up to the 26th of January 2018 than these end of season results (VE = 59%) [25]. Indeed, the VE in October to December (87%) was higher than the VE in January and February to April (54% and 53%). A decline in VE against A(H1N1)pdm09 by time since vaccination was also seen in the I-MOVE/I-MOVE+MCCS in 2015–16 and a decline in VE against A(H1N1)pdm09 by calendar time was seen in Canada in 2015–16 [20], but not in the I-MOVE MCCS between 2010 and 11 and 2014–15 [26]. While monitoring declining vaccine effect against the 6B.1 clade warrants further research, the decline by calendar time could possibly be a statistical artefact of the study. However, the decline was not seen in VE against A(H3N2) in 2017–18, indicating that this may be at least in part a real rather than an artefactual phenomenon for influenza A(H1N1)pdm09.

Vaccine coverage was low and numbers were small for estimates of vaccine type specific VE and indicated no real difference between VE of inactivated trivalent subunit and split virion vaccine among all ages, similar to the I-MOVE/I-MOVE+MCCS results from the 2013–14, 2014–15 and 2015–16 seasons [6,15,19]. Numbers for quadrivalent and adjuvanted vaccine types were too low to estimate vaccine type-specific VE.

VE among those receiving both current and prior season influenza vaccine was lower than among those receiving current season only vaccination (46% vs.79%), compared to a reference group of those not receiving vaccine in either season. A similar pattern for A(H1N1)pdm09, albeit with low precision, was seen in the 2015–16 season in the I-MOVE/I-MOVE+MCCS study and also in Canada, but not in the US [6,20,27]. The findings in the 2017–18 season may be explained by negative interference of the prior vaccine strain, but should be seen also in context of different profiles among unvaccinated, single season vaccinees, vaccinated and repeat vaccinated participants, as well as in the context of the influence of different antibody landscapes among individuals.

4.2. Influenza A(H3N2)

Similar clades of A(H3N2) were circulating in the 2016–17 and 2017–18 influenza seasons, albeit in different proportions and with some differing subclades, with the same vaccine component in both seasons. VE was approximately 30% among those aged <65 years in both seasons, and lower among those aged 65 and older.

A(H3N2) vaccine seed viruses can develop adaptations/alterations during propagation in eggs, impacting antigenicity, which may negatively affect VE [28–30]. Egg-propagated A/HongKong/4801/2014 (H3N2)-like seed vaccine viruses have the egg adaptation induced T160K substitution, whereas 160T is present in the original A/HongKong/4801/2014 virus and current circulating 3C.2a viruses. Viruses with 160T grow poorly in eggs, and the T160K egg adaptation results in the loss of a putative N-glycosylation site in the HA antigenic site B [31]. In the I-MOVE/I-MOVE+MCCS 98–99% of all systematically sequenced A(H3N2) viruses had 160T. These changes due to egg-adaptation may contribute to the low VE observed in the 2016–17 and 2017–18 seasons in our study. The A/Singapore/INFIMH-16-0019/2016 (H3N2)-like vaccine component for the Southern Hemisphere 2018 and the Northern Hemisphere 2018–19 influenza season, is also subject to the same problems during egg adaptation in the vaccine manufacturing process in the same critical antigenic site:

vaccine seed strains having 160K instead of 160T in the original virus. If similar genetic variants of A(H3N2) with 160T are circulating in the future we may expect VE to be similar to the 2016–17 and 2017–18 seasons, making recommendations for other preventive measures already at start of the season vital. This may be particularly crucial for older age groups, where A(H3N2) can also cause severe disease, as younger age groups had a higher VE compared to those 65 years of age and older. That said, WHO reported that A(H3N2) strains circulating in the 2017–18 season were more similar in HA assays to the egg-propagated A/Singapore/INFIMH-16-0019/2016 (H3N2)-like virus (the recommended 2018–19 Northern hemisphere A(H3N2) vaccine virus component) than the 2017–18 egg-propagated A/HongKong/4801/2014 (H3N2)-like virus vaccine strain [32]; however early results from Europe indicate that 2018–19 season VE against A(H3N2) may be low [33].

Low VE against influenza A(H3N2) may also be explained by the genetic variation among circulating strains. In 2016–17 the VE against 3C.2a variants, excluding 3C.2a1, was similar to the VE against 3C.2a1 (31% vs 25%). In 2017–18, the VE against 3C.2a, excluding 3C.2a1, of which almost all belonged to the 3C.2a2 subclade, was lower than the VE against 3C.2a1 (–29 vs 41%). The same pattern of difference in 2017–18 between 3C.2a (with 3C.2a1 excluded) and 3C.2a1 VE was seen in other study sites [34] [R Pebody, PHE, UK, Personal communication]. Sample size was low, however, in particular in the 2017–18 season, and it was not possible to disentangle age-specific and strain-specific effects on VE. Increased sequencing of A(H3N2) viruses in seasons with A(H3N2) circulating would help with interpretation of VE in light of genetic changes.

There was no statistical difference between VE of inactivated trivalent subunit and split virion vaccine in any of the age groups in 2016–17 or in 2017–18, as in the previous A(H3N2) season in 2014–15 [19].

The 2015–16 influenza A(H3N2) vaccine component was A/Switzerland/9715293/2013, which has 160K in the original and egg-adapted strains. In 2016–17 the VE among those receiving both current and prior vaccine was lower than the VE among those receiving current vaccine and no prior season vaccine. The pattern was the same in 2017–18 when those receiving both current and prior vaccine had a VE of 7% compared to the VE of 49% among those receiving current vaccine only. The results suggest a lower VE with combined use of current and prior vaccination, as per the antigenic distance hypothesis [35,36]. It is possible that this is a true phenomenon in the study, as due to passaging-related adaptations in the vaccine virus, the antigenic distance between vaccine and circulating strains may be large and the vaccine viruses are either identical (2017–18) or similar (2016–17), compared to previous seasons. Other potential explanations include confounding due to different participant profiles of repeat and single season vaccinees, recall bias and random variation. Prospective longitudinal studies are needed to better understand the phenomenon of effects of previous vaccination.

The interim VE against A(H3N2) in 2016–17 was 38% (95% CI: 21–51)[37], higher than the end-of-season estimate of 28% (95% CI: 17–38). When grouped into calendar months (Oct-Dec, January and Feb-April), the 2016–17 season VE declined along the season resulting in no effectiveness in February to April. This could be due to virological changes in the A(H3N2) virus across the season, waning of immunity within the individual against the vaccine, be a statistical artefact, or a combination of all. In 2016–17, 6% (14/216) of clade 3C.2a1 viruses with no mutations in A-E antigenic sites other than the N171K occurred in February to April, compared to 18% (99/565) of clade 3C.2a1 viruses with mutations in the same period (data not shown).

In 2017–18, the interim VE against A(H3N2) was –16 (95% CI: –96 to 13) compared to the 13% (–15 to 34) end-of-season estimate

[25]. Differences in point estimates were likely due to random variation due to sample size and there was no evidence of change in VE by calendar time. However, there was also less diversity of subclades, so that within clade 3C.2a (excluding clade 3C.2a1) and clade 3C.2a1, respectively, there was no consistent change in variants over time. Additionally, we may not expect any measurable waning of immunity, with such a low overall VE (13%).

This study uses the test-negative design that has its strengths in controlling for health care-seeking behaviour and ease of study implementation [38]. However as with all observational studies, we cannot rule out bias due to unmeasured confounding and selection bias. We believe these are minimised by the study design, as the practitioners are blinded to the case status of their patients, and to the adherence to the protocols, in which practitioners are asked to systematically or exhaustively select patients to swab and interview [10].

5. Conclusions

Our study indicates that the new vaccine component A/Michigan/45/2015 conferred greater protection to the circulating 6B.1 clade than the A/California/7/2009 vaccine strain. Receiving influenza vaccine in the 2017–18 season reduced the risk of being an A(H1N1)pdm09 case by more than half among all ages in our study. All viruses sequenced in our study belonged to the 6B.1 clade, so we expect a similar protection of the Northern Hemisphere vaccine, which contains the same A(H1N1)pdm09 strain, against A(H1N1)pdm09, should this virus circulate in the future, without additional common amino acid changes. To monitor this, representative genetic and amino acid composition characterisation of circulating A(H1N1)pdm09 should complement VE studies in coming seasons.

The VE of the A/HongKong/4801/2014 vaccine component against A(H3N2) was <30% in the 2016–17 and 2017–18 seasons. The circulating subclades in both seasons may have contributed towards the low VE, as well as the amino acid substitutions within the vaccine virus during the egg-adaptation process of manufacturing. No vaccines based on cell-propagated viruses were used by participants of the 2016–17 and 2017–18 I-MOVE/I-MOVE+MCCS. The A(H3N2) component A/Singapore/INFIMH-16-0019/2016 recommended for the 2018–19 Northern Hemisphere influenza vaccine is subject to the same issues of amino acid substitutions during the egg-adaptation process. As influenza A(H3N2) and A(H1N1)pdm09 co-circulated strain in the 2018–19 season, it is possible that influenza VE against A(H3N2) will be low, particularly among older adults, based on the results of our study. Excess influenza-associated mortality and morbidity has been reported among the elderly during previous seasons with A(H3N2) circulation, when the H3N2 vaccine component showed poor or moderate VE. Therefore, national authorities need to be prepared in such a situation for rapid actions in reallocation of health care resources, ensuring surge capacity and ensuring timely prescriptions and administration of influenza antivirals.

All authors attest they meet the ICMJE criteria for authorship

Declaration of Competing Interest

The authors declare that they have no known competing financial interests or personal relationships that could have appeared to influence the work reported in this paper.

Acknowledgements

The I-MOVE/I-MOVE+ study team is very grateful to all patients, general practitioners, paediatricians, hospital teams, laboratory

teams, regional epidemiologists who have contributed to the study.

We acknowledge the authors, originating and submitting laboratories of the sequences from GISAID's EpiFlu Database used for this study. All submitters of data may be contacted directly via the GISAID website www.gisaid.org.

Funding

This project has received funding from the European Union's Horizon 2020 research and innovation programme under grant agreement no. 634446 to conduct the study in individuals aged 65 years or more.

ECDC has contributed funds for the coordination and some study sites under the Framework contract no. ECDC/2014/026 for the individuals aged less than 65 years.

The WHO Regional office for Europe has contributed funds for the Romanian study site.

Appendix A. Supplementary material

Supplementary data to this article can be found online at <https://doi.org/10.1016/j.jvax.2019.100042>.

References

- [1] ECDC, 'Influenza virus characterisation - Summary Europe, June 2016', Jul. 2016.
- [2] C. Adlhoch, R. Snacken, A. Melidou, S. Ionescu, P. Penttinen, and The European Influenza Surveillance Network, 'Dominant influenza A(H3N2) and B/Yamagata virus circulation in EU/EEA, 2016/17 and 2017/18 seasons, respectively', *Euro Surveill. Bull. Eur. Sur Mal. Transm. Eur. Commun. Dis. Bull.*, vol. 23, no. 13, 2018.
- [3] Vestergaard LS et al. Excess all-cause and influenza-attributable mortality in Europe, December 2016 to February 2017. *Euro. Surveill. Bull. Eur. Sur Mal. Transm. Eur. Commun. Dis. Bull.* 2017;22(14). Apr.
- [4] EuroMOMO, 'European monitoring of excess mortality for public health action - EuroMOMO'. [Online]. Available: [Accessed: 14-Aug-2018]. <http://www.euromomo.eu/index.html>.
- [5] Valenciano M, Ciancio B, I-MOVE study team. I-MOVE: a European network to measure the effectiveness of influenza vaccines. *Euro Surveill. Bull. Eur. Sur Mal. Transm. Eur. Commun. Dis. Bull.* 2012;17(39). Sep.
- [6] Kissling E et al. 2015/16 I-MOVE/I-MOVE+ multicentre case-control study in Europe: moderate vaccine effectiveness estimates against influenza A(H1N1)pdm09 and low estimates against lineage-mismatched influenza B among children. *Influenza Other Respir. Viruses* 2018;12(4):423–37. Jul.
- [7] Kissling E et al. 'Influenza vaccine effectiveness estimates in Europe in a season with three influenza type/subtypes circulating: the I-MOVE multicentre case-control study, influenza season 2012/13. *Euro Surveill Bull. Eur. Sur Mal. Transm. Eur. Commun. Dis. Bull.* 2014;19(6). Feb.
- [8] Linderman SL et al. Potential antigenic explanation for atypical H1N1 infections among middle-aged adults during the 2013–2014 influenza season. *Proc. Natl. Acad. Sci. U. S. A.* 2014;111(44):15798–803. Nov.
- [9] ECDC, 'European Centre for Disease Prevention and Control (ECDC). Protocol for case control studies to measure pandemic and seasonal vaccine effectiveness in the European Union and European Economic Area', European Centre for Disease Prevention and Control, Stockholm, Sweden, 2010.
- [10] 'Generic protocol for the test negative design case control studies to measure pandemic and seasonal influenza vaccine effectiveness in the European Union and European Economic Area Member States', European Union, Jul. 2015.
- [11] Valenciano M et al. Estimates of pandemic influenza vaccine effectiveness in Europe, 2009–2010: results of Influenza Monitoring Vaccine Effectiveness in Europe (I-MOVE) multicentre case-control study. *PLoS Med.* 2011;8(1):. [doi:10.1371/journal.pmed.1000388](https://doi.org/10.1371/journal.pmed.1000388).
- [12] ECDC, 'European Commission. Commission Decision 2009/363/EC of 30 April 2009 amending Decision 2002/253/EC laying down case definitions for reporting communicable diseases to the Community network under Decision No 2119/98/EC of the European Parliament and of the Council.', Jan. 2009.
- [13] Peduzzi P, Concato J, Kemper E, Holford TR, Feinstein AR. A simulation study of the number of events per variable in logistic regression analysis. *J Clin Epidemiol* 1996;49(12):1373–9. Dec.
- [14] Firth D. Bias reduction of maximum likelihood estimates. *Biometrika* 1993;80(1):27–38. Mar.
- [15] Valenciano M et al. The European I-MOVE multicentre 2013–2014 case-control study. homogeneous moderate influenza vaccine effectiveness against A(H1N1)pdm09 and heterogeneous results by country against A(H3N2). *Vaccine* 2015;33(24):2813–22. Jun.
- [16] StataCorp, Stata statistical software. College Station, TX: StataCorp LLC.

- [17] European Centre for Disease Prevention and Control, 'Influenza virus characterisation, Summary Europe, June 2015', European Centre for Disease Prevention and Control, Stockholm, Sweden.
- [18] WHO, 'Recommended composition of influenza virus vaccines for use in the 2017 southern hemisphere influenza season', WHO, Sep. 2016.
- [19] Valenciano M et al. Vaccine effectiveness in preventing laboratory-confirmed influenza in primary care patients in a season of co-circulation of influenza A (H1N1)pdm09, B and drifted A(H3N2), I-MOVE Multicentre Case-Control Study, Europe 2014/15. *Euro Surveill. Bull. Eur. Sur Mal. Transm. Eur. Commun. Dis. Bull.* 2016;21(7). p. pii=30139.
- [20] Skowronski DM et al. Beyond antigenic match: possible agent-host and immuno-epidemiological influences on influenza vaccine effectiveness during the 2015–2016 season in Canada. *J. Infect. Dis.* 2017;216(12):1487–500. Dec.
- [21] Pebody R et al. Effectiveness of seasonal influenza vaccine for adults and children in preventing laboratory-confirmed influenza in primary care in the United Kingdom: 2015/16 end-of-season results. *Euro Surveill. Bull. Eur. Sur Mal. Transm. Eur. Commun. Dis. Bull.* Sep. 2016;21(38).
- [22] Hensley SE. Challenges of selecting seasonal influenza vaccine strains for humans with diverse pre-exposure histories. *Curr. Opin. Virol.* 2014;8:85–9. Oct.
- [23] Davenport FM, Hennessy AV, Francis T. Epidemiologic and immunologic significance of age distribution of antibody to antigenic variants of influenza virus. *J. Exp. Med.* 1953;98(6):641–56. Dec.
- [24] Flannery B et al. Influence of Birth Cohort on Effectiveness of 2015–2016 Influenza Vaccine Against Medically Attended Illness Due to 2009 Pandemic Influenza A(H1N1) Virus in the United States. *J. Infect. Dis.* 2018;218(2):189–96. Jun.
- [25] Rondo M. Interim 2017/18 influenza seasonal vaccine effectiveness: combined results from five European studies. *Euro Surveill. Bull. Eur. Sur Mal. Transm. Eur. Commun. Dis. Bull.* 2018;23(9).
- [26] Kissling E et al. I-MOVE multicentre case-control study 2010/11 to 2014/15: Is there within-season waning of influenza type/subtype vaccine effectiveness with increasing time since vaccination? *Euro Surveill Bull. Eur. Sur Mal. Transm. Eur. Commun. Dis. Bull.* 2016;21(16). Apr.
- [27] Jackson ML et al. Influenza Vaccine Effectiveness in the United States during the 2015–2016 Season. *N. Engl. J. Med.* 2017;377(6):534–43. 10.
- [28] Chen Z, Zhou H, Jin H. The impact of key amino acid substitutions in the hemagglutinin of influenza A (H3N2) viruses on vaccine production and antibody response. *Vaccine* May 2010;28(24):4079–85.
- [29] Skowronski DM, De Serres G. Role of egg-adaptation mutations in low influenza A(H3N2) vaccine effectiveness during the 2012–13 season. *Clin. Infect. Dis. Off. Publ. Infect. Dis. Soc. Am.* 2018. Apr.
- [30] Parker L et al. Effects of egg-adaptation on receptor-binding and antigenic properties of recent influenza A (H3N2) vaccine viruses. *J. Gen. Virol.* 2016;97(6):1333–44.
- [31] Zost SJ. Contemporary H3N2 influenza viruses have a glycosylation site that alters binding of antibodies elicited by egg-adapted vaccine strains. *Proc. Natl. Acad. Sci. U. S. A.* 2017;114(47):12578–83. 21.
- [32] Worldwide Influenza Centre, 'Report prepared for the WHO annual consultation on the composition of influenza vaccine for the Northern Hemisphere 2018–19', The Francis Crick Institute, Feb. 2018
- [33] Kissling E et al. Interim 2018/19 influenza vaccine effectiveness: six European studies, October 2018 to January 2019. *Eurosurveillance* 2019;24(8). Feb.
- [34] Skowronski, Danuta M., 'Viral genomic variation and vaccine effectiveness across consecutive influenza A(H3N2) epidemics in Canada, 2016–17 and 2017–18', presented at the Canadian Immunization Conference, Ottawa, Canada, 04–Dec-2018.
- [35] Skowronski DM et al. Serial vaccination and the antigenic distance hypothesis: effects on influenza vaccine effectiveness during A(H3N2) epidemics in Canada, 2010–2011 to 2014–2015. *J. Infect. Dis.* 2017;215(7):1059–99. Apr.
- [36] Smith DJ, Forrest S, Ackley DH, Perelson AS. Variable efficacy of repeated annual influenza vaccination. *Proc. Natl. Acad. Sci. U. S. A.* 1999;96(24):14001–6. Nov.
- [37] Kissling E, Rondo M, I-MOVE/I-MOVE+ study team. Early 2016/17 vaccine effectiveness estimates against influenza A(H3N2): I-MOVE multicentre case control studies at primary care and hospital levels in Europe. *Euro Surveill. Bull. Eur. Sur Mal. Transm. Eur. Commun. Dis. Bull.* 2017;22(7). 16.
- [38] Fukushima W, Hirota Y. Basic principles of test-negative design in evaluating influenza vaccine effectiveness. *Vaccine* 2017;35(36):4796–800. 24.