



HAL
open science

Comparative Effectiveness of Improvement in Pain and Physical Function for Baricitinib versus Adalimumab, Tocilizumab and Tofacitinib Monotherapies in Rheumatoid Arthritis Patients Who Are Naïve to Treatment with Biologic or Conventional Synthetic Disease-Modifying Antirheumatic Drugs: A Matching-Adjusted Indirect Comparison

B Fautrel, B Zhu, P C Taylor, M van De Laar, P Emery, F de Leonardis, C L Kannowski, C Nicolay, Z Kadziola, I de la Torre, et al.

► To cite this version:

B Fautrel, B Zhu, P C Taylor, M van De Laar, P Emery, et al.. Comparative Effectiveness of Improvement in Pain and Physical Function for Baricitinib versus Adalimumab, Tocilizumab and Tofacitinib Monotherapies in Rheumatoid Arthritis Patients Who Are Naïve to Treatment with Biologic or Conventional Synthetic Disease-Modifying Antirheumatic Drugs: A Matching-Adjusted Indirect Comparison. *RMD Open: Rheumatic & Musculoskeletal Diseases*, 2020, 6 (1), pp.e001131. 10.1136/rmdopen-2019-001131 . hal-03849936

HAL Id: hal-03849936

<https://hal.sorbonne-universite.fr/hal-03849936v1>

Submitted on 15 Nov 2022

HAL is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers.

L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.

ORIGINAL RESEARCH


Comparative effectiveness of improvement in pain and physical function for baricitinib versus adalimumab, tocilizumab and tofacitinib monotherapies in rheumatoid arthritis patients who are naïve to treatment with biologic or conventional synthetic disease-modifying antirheumatic drugs: a matching-adjusted indirect comparison

B Fautrel,¹ B Zhu,² P C Taylor,³ M van de Laar,⁴ P Emery,⁵ F De Leonardis,² C L Kannowski,² C Nicolay,² Z Kadziola,² I De La Torre,² R Fleischmann⁶

To cite: Fautrel B, Zhu B, Taylor PC, et al. Comparative effectiveness of improvement in pain and physical function for baricitinib versus adalimumab, tocilizumab and tofacitinib monotherapies in rheumatoid arthritis patients who are naïve to treatment with biologic or conventional synthetic disease-modifying antirheumatic drugs: a matching-adjusted indirect comparison. *RMD Open* 2020;**6**:e001131. doi:10.1136/rmdopen-2019-001131

► Additional material is published online only. To view please visit the journal online (<http://dx.doi.org/10.1136/rmdopen-2019-001131>).

Received 24 October 2019
Revised 29 February 2020
Accepted 2 March 2020



© Author(s) (or their employer(s)) 2020. Re-use permitted under CC BY-NC. No commercial re-use. See rights and permissions. Published by BMJ.

For numbered affiliations see end of article.

Correspondence to

Dr Bruno Fautrel;
bruno.fautrel@aphp.fr

ABSTRACT

Objective To compare improvement in pain and physical function for patients treated with baricitinib, adalimumab, tocilizumab and tofacitinib monotherapy from randomised, methotrexate (MTX)-controlled trials in conventional synthetic disease-modifying antirheumatic drugs (csDMARDs)/biologic (bDMARD)-naïve RA patients using matching-adjusted indirect comparisons (MAICs).

Methods Data were from Phase III trials on patients receiving monotherapy baricitinib, tocilizumab, adalimumab, tofacitinib or MTX. Pain was assessed using a visual analogue scale (0–100 mm) and physical function using the Health Assessment Questionnaire-Disability Index (HAQ-DI). An MAIC based on treatment-arm matching, an MAIC with study-level matching and Bucher's method without matching compared change in outcomes between therapies. Matching variables included age, gender, baseline disease activity and baseline value of outcome measure.

Results With all methods, greater improvements were observed in pain and HAQ-DI at 6 months for baricitinib compared with adalimumab and tocilizumab ($p < 0.05$). Differences in treatment effects (TEs) favouring baricitinib for pain VAS for treatment-arm matching, study-level matching and Bucher's method, respectively, were –12, –12 and –12 for baricitinib versus adalimumab and –7, –7 and –9 for baricitinib versus tocilizumab; the difference in TEs for HAQ-DI was –0.28, –0.28 and –0.30 for adalimumab and –0.23, –0.23 and –0.26 for tocilizumab. For baricitinib versus tofacitinib, no statistically significant differences for pain improvement were observed except with one of the three methods (Bucher method) and none for HAQ-DI.

Conclusions Results suggest greater pain reduction and improved physical function for baricitinib monotherapy compared with tocilizumab and adalimumab monotherapy. No statistically significant differences in pain reduction and improved physical function were observed between baricitinib and tofacitinib with the MAIC analyses.

Key messages

What is already known about this subject?

► Large, randomised clinical trials have demonstrated the efficacy of baricitinib, adalimumab, tocilizumab and tofacitinib monotherapy in pain reduction and HAQ-DI improvement compared with methotrexate monotherapy, but there are no head-to-head trials between these treatments in patients with RA who are naïve to treatment with conventional synthetic or biologic disease-modifying antirheumatic drugs.

What does this study add?

► The results from this study add evidence, through indirect comparison, that suggest greater pain reduction and improved physical function for baricitinib monotherapy compared with tocilizumab and adalimumab monotherapy.

How might this impact on clinical practice or future developments?

► The findings from this study will help clinicians evaluate different therapies to reduce pain and improve physical function in the treatment of RA patients.

INTRODUCTION

Despite substantial improvements over the last two decades in the management of patients with rheumatoid arthritis (RA), the treat-to-target approach has led rheumatologists to focus on inflammatory disease activity, whereas patients generally consider the reduction of pain and fatigue and improvement of physical

function to be more important.¹⁻³ Their assessment, in addition to healthcare provider (HCP)-reported disease activity measures, should help physicians determine the best treatment management for the patient. In the RA-BEAM randomised controlled trial (RCT), with concomitant methotrexate (MTX), baricitinib 4 mg one time per day demonstrated greater improvements in pain and physical function compared with adalimumab 40 mg every other week in a population of patients who had had an insufficient response to MTX.⁴ There is an absence, however, of prospective, head-to-head trials between different biologic or targeted synthetic disease-modifying antirheumatic drugs (b/tsDMARDs) in MTX-naïve RA patients, a population that could be considered more sensitive to change in PROs because they had not yet experienced the irreversible consequences of the longstanding disease.

In the absence of data from RCT, indirect comparison methodologies, such as Network Meta-Analysis (NMA) and, in more recent years, Matching-Adjusted Indirect Comparison (MAIC), have been proposed to compare the efficacy of different therapies based on aggregate data from different RCTs, and they are commonly used for the purposes of health technology appraisal.⁵⁻⁷ Compared with an NMA, which is based on the assumption that treatment effects (TEs) are only relative to a common comparator (eg, placebo) with no additional difference between the trials in the distribution of effect-modifying variables,^{7 8} MAIC builds upon the indirect comparison through additional adjustment of effect-modifying variables.

An MAIC analysis uses patient-level data of a drug to match with published data from comparators. Specifically, individual patient data from one or more studies for one treatment are reweighted to match with the baseline characteristics, which are known to be TE modifiers, from a published study of another treatment. To have an appropriate analysis, the study with patient-level data and the study with published data must have a common reference arm for matching. After the matching with the individual patient data, the weighted difference in mean values of an outcome measure between the active arm and the reference arm of one study is calculated and compared with the difference from the other published study.⁵

The objective of this analysis was to compare improvement in pain and physical function between baricitinib, adalimumab, tocilizumab and tofacitinib monotherapy with an MAIC using data from randomised, MTX-controlled trials in conventional synthetic DMARD (csDMARD)/bDMARD-naïve RA patients.

METHODS

Study eligibility

The studies included in this analysis were derived from a prior systematic literature review (SLR) that was designed for a NMA conducted by Eli Lilly. The SLR synthesised the evidence of treatments on measures of treatment response and effectiveness, disease activity, physical function, radiographic outcomes,

safety and other key measures for adult patients with moderate-to-severe RA among studies conducted from 1999 to 2016. The criteria for selection in the SLR and a flow chart describing the screening for inclusion are in online supplementary figure 1. For the purposes of the current analysis, we focused on the population of patients with limited or no treatment with csDMARDs in the SLR; 27 studies met this criterion. Of these studies, 12 included monotherapy and an MTX treatment arm, which constitutes the common comparator; and of these 12 studies, 5 reported on pain and physical function, as measured by the Health Assessment Questionnaire-Disability Index (HAQ-DI), at 6 months or 24±2 weeks, depending upon the time points reported in the studies. These five studies were included in the current analysis (table 1). The study designs and inclusion and exclusion criteria for the studies have been previously reported.⁶⁻¹¹ The doses of the medications included in the analysis were oral 4 mg of baricitinib daily,⁹ subcutaneous 40 mg of adalimumab every other week,^{6 7} intravenous 8 mg/kg of tocilizumab every 4 weeks^{8 10} and oral 5 mg tofacitinib two times per day.¹¹

Outcome measures

Pain was measured with the patient's assessment of pain, a visual analogue scale (VAS), ranging from 0 to 100 mm. Physical function was measured with the HAQ-DI.^{12 13} The HAQ-DI consists of 24 questions referring to eight domains: dressing/grooming, arising, eating, walking, hygiene, reach, grip and activities. The score for the HAQ-DI ranges from 0 to 3, with lower scores reflecting better physical function and thus, less disability.

Matching-adjusted indirect comparisons (MAICs) and sensitivity analyses

The primary MAIC in this analysis was based on the Signorovitch method with weights applied to treatment arms.^{5 14} Specifically, data from the baricitinib 4 mg treatment arm from the RA-BEGIN trial⁹ were weighted to match the baseline characteristics that are TE modifiers (age, gender, Disease Activity Score-28 erythrocyte sedimentation rate [DAS28-ESR], pain VAS and HAQ-DI) from the adalimumab arm from PREMIER,^{6 7} tofacitinib 5 mg twice a day arm from ORAL-START twice a day¹¹ and tocilizumab 8 mg/kg arm from FUNCTION.⁸ For reference, the MTX monotherapy arms were also matched between the trials. Analyses were conducted on patients from RA-BEGIN who met the inclusion and exclusion criteria of the respective comparator trials. Sensitivity analyses were conducted with the inclusion of disease duration as an additional matching variable. Two other approaches, an MAIC based on the Signorovitch method with study-level matching (matching on the entire study, rather than by treatment arm)⁵ and Bucher's method without matching adjustment,¹⁵ were also conducted as sensitivity analyses to determine the consistency of the findings. Because of the prior experience patients in the AMBITION study had with MTX, we also conducted

Table 1 Study design and characteristics

Study	Study design and dosage information	Inclusion criteria	Key exclusion criteria
Baricitinib (RA-BEGIN, NCT01711359) ⁹	Patients were randomised 4:3:4 to oral MTX one time per week (N=210), baricitinib 4 mg (monotherapy) one time per day (N=159), or the combination of baricitinib+MTX (N=215)	<ul style="list-style-type: none"> ▶ Patients were ≥18 years ▶ Moderately-to-severely active RA ▶ Patients had active disease (TJC ≥6 and SJC ≥6) ▶ Serum CRP level ≥3.6 mg/L ▶ Seropositive for RF or ACPA ▶ No prior csDMARD therapy and no prior bDMARD 	<ul style="list-style-type: none"> ▶ Recent clinically significant infection and select laboratory abnormalities
Tocilizumab (AMBITION, NCT00109408) ¹⁰	Patients were randomised to tocilizumab (TCZ) 8 mg/kg intravenously every 4 weeks (N=286), or to MTX oral capsules, weekly together with folate (>5 mg/week) (N=284)	<ul style="list-style-type: none"> ▶ Patients were ≥18 years ▶ ≥3 months of moderately-to-severely active RA ▶ SJC of ≥6, TJC ≥8, CRP level ≥1.0 mg/dL or ESR ≥28 mm/h at baseline ▶ Oral glucocorticoids and NSAIDs were permitted if stable >6 weeks 	<ul style="list-style-type: none"> ▶ Clinically unstable concurrent illnesses ▶ Active or untreated latent TB ▶ Unsuccessful treatment with TNFi ▶ Received MTX within 6 months of randomisation or discontinued MTX previously
Tocilizumab (FUNCTION, NCT01007435) ⁸	Patients were randomised to 4 mg/kg TCZ+MTX (N=288), 8 mg/kg TCZ+MTX (N=290), 8 mg/kg TCZ+placebo (N=292) or placebo+MTX (N=287); TCZ or placebo were administered intravenously every 4 weeks	<ul style="list-style-type: none"> ▶ Patients were ≥18 years ▶ ≤2 years of moderate-to-severe RA ▶ SJC of ≥4, TJC of ≥6, CRP level ≥1.0 mg/dL or ESR ≥28 mm/h at baseline ▶ Positive RF or ACPA or ≥1 erosion of hands, wrists or feet attributable to RA based on a central radiographic reading 	<ul style="list-style-type: none"> ▶ Clinically unstable concurrent illnesses and screened according to local standards ▶ Active or untreated latent TB ▶ Had been unsuccessfully treated with TNFi ▶ Had received MTX 6 months prior to randomisation or had discontinued MTX
Tofacitinib (ORAL-START, NCT01039688) ¹¹	Patients were randomised to tofacitinib 5 mg two times per day (BID, N=373) or tofacitinib 10 mg BID (N=397) or MTX (N=186)	<ul style="list-style-type: none"> ▶ Patients were ≥18 years ▶ ≥3 months of moderately-to-severely active RA ▶ SJC of ≥6, TJC of ≥6, CRP level >7.0 mg/L or ESR >28 mm/h at baseline ▶ ≥3 distinct joint erosions on radiographs, positive test for IgM RF or ACPA 	<ul style="list-style-type: none"> ▶ Prior treatment with lymphocyte-depleting or alkylating agents ▶ Select lab abnormalities ▶ History of: another autoimmune rheumatic disease except Sjögren's syndrome ▶ Serious infection ▶ Lymphoproliferative disorder ▶ Malignancy except adequately treated non-metastatic basal/squamous cell cancer of the skin or cervical carcinoma in situ ▶ Evidence of active, latent or inadequately treated Mycobacterium TB infection
Adalimumab (PREMIER, NCT00195663) ^{6,7}	Patients were randomised to adalimumab 40 mg subcutaneously every other week + weekly oral MTX (N=268); adalimumab 40 mg subcutaneously every other week (adalimumab + placebo; N=274); or weekly oral MTX (N=257)	<ul style="list-style-type: none"> ▶ Patients were ≥18 years ▶ <3 years of RA ▶ SJC of ≥8, TJC of ≥10, CRP level ≥1.5 mg/dL or ESR ≥28 mm/h at baseline 	<p>Patients who had received treatment with MTX, cyclophosphamide, cyclosporine, azathioprine or >2 other DMARDs</p>

ACPA, anti-citrullinated protein antibodies; bDMARD, biologic disease-modifying antirheumatic drugs; CRP, C reactive protein; ESR, erythrocyte sedimentation rate; IgM, immunoglobulin M; MTX, methotrexate; RA, rheumatoid arthritis; RF, rheumatoid factor; SJC, swollen joint count; TB, tuberculosis; TCZ, tocilizumab; TJC, tender joint count; TNFi, tumour necrosis factor inhibitor.

separate MAICs between baricitinib and tocilizumab, one with data from AMBITION alone and the second with data from AMBITION and FUNCTION combined.^{8–10}

Statistical analyses

Differences between weighted TE in mean change in pain VAS and HAQ-DI from baseline to 6 months for baricitinib and the reported TE for adalimumab, tocilizumab or tofacitinib were compared. For adalimumab, mean changes in pain and HAQ-DI were based on the mean pain and HAQ-DI values reported at Week 26.⁷ The variance of the weighted TE was estimated with the bootstrap method with 1000 iterations.^{16 17} The differences and their associated 95% CIs are presented and a $p < 0.05$ was considered statistically significant. Analyses were not adjusted for multiplicity, and they were conducted with SAS version 9.4 (Cary, NC) and R (version 3.3.3).

RESULTS

Baseline characteristics

Baseline characteristics are presented in [table 2](#). For the MTX arms across trials, the mean baseline pain VAS ranged from 59 to 65 mm and the 6-month mean change in pain ranged from -28.3 to -33.5 mm. Likewise, for the MTX arm, the mean baseline HAQ-DI values ranged from 1.5 to 1.7, and the 6-month mean change in HAQ-DI ranged from -0.5 to -0.74 ([table 3](#)). The similarity of the baseline pain and HAQ-DI scores and the similar change in pain and HAQ-DI from the MTX control arm across studies suggest comparability between the trials.

The baseline values of the variables used in matching for all the trials are shown in [table 4](#), which includes the baseline variables for RA-BEGIN after the matching on those variables. Because of the matching, the baseline values for baricitinib are the same as those from the published data for the respective comparator drugs. The effective sample sizes from RA-BEGIN, with individual patient-level data, are reduced as the consequence of weighting and matching.

Pain

For the primary MAIC analysis, baricitinib-treated patients showed greater improvement in pain at 6 months compared with adalimumab (treatment difference: -12.3 , 95% CI -17.9 to -6.6) and tocilizumab (treatment difference: -7.3 , 95% CI -14.2 to -0.38) ([figure 1](#)). Consistent results were observed with the other indirect comparison methods. There were numerical, but no statistically significant, differences in pain improvement between baricitinib and tofacitinib with the primary analysis. With the sensitivity analyses, no statistically significant differences were observed using the MAIC with study-level matching; there were, however, significant differences with Bucher method (treatment difference: -7.1 ; -13.5 to -0.65). ([figure 1](#))

Physical function

For the primary MAIC analysis ([figure 1](#)), baricitinib-treated patients were shown to have greater improvement in physical function at 6 months compared with adalimumab (treatment difference: -0.28 , 95% CI -0.44 to -0.13) and tocilizumab (treatment difference: -0.23 , 95% CI -0.39 to -0.07). Similar results were observed with the other indirect comparisons. There were no differences between baricitinib and tofacitinib with all methods ([figure 1](#)).

Sensitivity analyses

To confirm the robustness of these results, we conducted sensitivity analyses in which disease duration was an additional matching variable, and when data from AMBITION and FUNCTION were analysed together. These sensitivity analyses were generally consistent with the direction and magnitude of the primary results, except for the comparison with the AMBITION data ([figures 2a,b](#)). The different patient characteristics from AMBITION from those in the FUNCTION and RA-BEGIN studies may have contributed to the differences, as described in the discussion below.

DISCUSSION

The gold standard for assessing the relative effectiveness of one medication compared with another is a properly powered head-to-head study using an appropriate metric as the primary endpoint. There has not been a study conducted comparing one JAK inhibitor with another, or with a bDMARD, in MTX-naïve patients, as monotherapy. In the absence of a head-to-head RCT, we applied an MAIC to compare improvement in pain and physical function for patients treated with baricitinib, adalimumab, tocilizumab and tofacitinib monotherapy from randomised, MTX-controlled trials in RA patients who were naïve to csDMARDs and bDMARDs. The MAIC enables greater flexibility to adjust for patient characteristics and TE modifiers and should provide a more robust indirect comparison than other traditional indirect comparison methods, such as a network meta-analysis.¹⁸ This MAIC analysis has been used in the indirect comparison of efficacy in other rheumatic diseases, such as psoriatic arthritis.^{19 20} The results of the current analysis suggest greater pain reduction with improved physical function for baricitinib monotherapy compared with tocilizumab and adalimumab monotherapy with the primary MAIC and sensitivity analyses. For comparisons between baricitinib and tofacitinib monotherapy, greater pain reduction with baricitinib was not consistently observed across the MAIC analyses, which did not allow for a robust conclusion on a difference between the two molecules. There were no differences observed between the two JAK inhibitors for HAQ-DI. Similar observations were also observed with models in which disease duration was an additional matching variable and with models in which the AMBITION and FUNCTION data were analysed together.

Table 2 Baseline characteristics from trials in the indirect comparisons

Characteristics	RA-BEGIN ⁹		AMBITION ¹⁰		FUNCTION ⁸		PREMIER ^{6,7}		ORAL-START ¹¹	
	MTX (N=210)	Baricitinib 4 mg (N=159)	MTX (N=284)	Tocilizumab 8 mg/kg (N=286)	MTX (N=287)	Tocilizumab 8 mg/kg (N=292)	MTX (N=257)	Adalimumab 40 mg (N=274)	MTX (N=186)	Tofacitinib 5 mg (N=373)
Mean duration of RA, years	1.3	1.9	6.2	6.4	0.4	0.5	0.8	0.7	2.7	2.9
SJC, 66 joints	16.4	16.1	19.2	19.1	16.2	16.5	22.1	21.8	16.8	16.3
TJC, 68 joints	27	26	31.1	31.8	27.4	28.7	32.3	31.8	25.4	25.7
CRP, mg/L	22	24	31	30	23	25	40	41	26	23
DAS28-ESR	6.6	6.6	6.8	6.8	6.6	6.7	6.3	6.4	6.6	6.6
Patient's Global Assessment of Disease, 0–100 mm VAS	66	65	66	64	64	68	63	68	58	60
MTX dosing information	MTX initiated at 10 mg/week and, if tolerated, increased to 20 mg/week by Week 8; if clinically indicated, an initial dosage of 7.5 mg one time per week and maximum dosage of 12.5 mg once weekly									
Mean MTX dosing achieved, mg/week	17.7	15.5	N/A (81% achieved 20 mg/week)		16.9	16.9	MTX initiated at 10 mg/week for the first 4 weeks. If tolerated, dosage was increased to 15 mg/week during Weeks 4–8, and to 20 mg/week at Week 9. Dosage could be reduced to as low as 7.5 mg/week			

CRP, C reactive protein; DAS28, Disease Activity Score for 28 joints; ESR, erythrocyte sedimentation rate; MTX, methotrexate; RA, rheumatoid arthritis; SJC, swollen joint count; TJC, tender joint count; VAS, visual analogue scale.

Table 3 Pain and HAQ-DI for studies included in the MAIC

Endpoint values at baseline and either change from baseline or mean values at 6 months	RA-BEGIN ⁹		AMBITION ¹⁰		FUNCTION ⁸		PREMIER ^{6,7}		ORAL-START ¹¹	
	MTX (N=210)	Baricitinib 4 mg (N=159)	MTX (N=284)	Tocilizumab 8 mg/kg (N=286)	MTX (N=287)	Tocilizumab 8 mg/kg (N=292)	MTX (N=257)	Adalimumab 40 mg (N=274)	MTX (N=186)	Tofacitinib 5 mg (N=373)
Patient's assessment of pain, 0–100 mm VAS										
Baseline	65	64	62	59	60	63	60	65	59	59
6 months	-30	-41	-31	-35	-34	-36	Adjusted absolute mean*: 29	Adjusted absolute mean*: 31	-28	-32
HAQ-DI, 0-3										
Baseline	1.7	1.6	1.5	1.6	1.5	1.6	1.5	1.6	1.5	1.5
6 months	-0.7	-1.0	-0.5	-0.7	-0.4	-0.04(vs MTX)	Adjusted absolute mean*: 0.9	Adjusted absolute mean*: 0.9	-0.6	-0.8

*Adjusted mean scores, rather than change from baseline, were reported in Strand *et al.*⁷ HAQ-DI, Health Assessment Questionnaire-Disability Index; MTX, methotrexate; VAS, visual analogue scale. Pain and HAQ-DI were collected at 6 months or 24±2 weeks.

Table 4 Baseline variables used in the matching across all trials and baseline variables from RA-BEGIN after matching

Study	Baseline variables from different studies before matching					Baseline variables from RA-BEGIN after matching*						
	Sample size	Age, mean (years)	Gender (%)	DAS28-ESR	Pain VAS	HAQ-DI	MTX sample size, pain/HAQ-DI	Age, mean (years)	Gender (%)	DAS28-ESR	Pain VAS	HAQ-DI
RA-BEGIN⁹												
MTX	210	51	70%	6.6	65	1.7						
Baricitinib 4 mg	159	51	76%	6.6	64	1.6						
AMBITION/ FUNCTION^{6 10}												
MTX	571	50	80%	6.7	61	1.49	179.4/160.9	50	80%	6.7	61	1.49
Tocilizumab 8 mg/kg	578	50	79%	6.8	61	1.59	142.7/147.6	50	79%	6.8	61	1.59
PREMIER^{6 7}												
MTX	257	52	74%	6.3	60	1.5	181.4/180.1	52	74%	6.3	60	1.5
Adalimumab 40 mg	274	52	77%	6.4	65	1.6	151.5/153.4	52	77%	6.4	65	1.6
ORAL-START¹¹												
MTX	186	49	78%	6.6	59	1.5	177.8/172.8	49	78%	6.6	59	1.5
Tofacitinib 5 mg	373	50	77%	6.6	59	1.5	146.5/148.9	50	77%	6.6	59	1.5

*Effective sample size and values for baricitinib or MTX from RA-BEGIN decreased after matching with different studies.

DAS28, Disease Activity Score for 28 joints; ESR, erythrocyte sedimentation rate; HAQ-DI, Health Assessment Questionnaire-Disability Index; MTX, methotrexate; VAS, visual analogue scale.




Figure 1 Treatment differences from indirect comparisons with matching by treatment arm (primary analysis), matching by study and without matching. HAQ-DI, Health Assessment Questionnaire-Disability Index; MTX, methotrexate; VAS, visual analogue scale.


To our knowledge, this is the first systematic analysis that compares patients receiving different csDMARD and bDMARD monotherapies in MTX-naïve patients with RA. This analytic approach offers many advantages over other conventional pairwise meta-analyses.²¹ Of note, the inclusion of active comparators provides more clinically relevant information compared with meta-analyses with only placebo. We also included data from well-designed RCTs that included large enough sample sizes to allow for more reliable estimations of differences between treatments.

There are, however, limitations intrinsic to the MAIC approach and, for this reason, our results should be interpreted with caution. The MAIC analysis matches based on observed TE modifiers, but it is not possible to control for variables that are unobserved. Additionally, the application of the MAIC reduces the effective sample size for the study with patient-level data, RA-BEGIN in the current analysis, which subsequently results in reduced power and capability to detect differences between medications. Importantly, the RCTs in the current analysis were conducted at different time periods during which more aggressive therapy was being introduced, with different patients and investigators in different regions of the world. These factors have the potential to increase the variations in baseline characteristics among the included studies, with consequent challenges in matching patients accurately. Of note, the AMBITION trial included 33% of patients who had been previously treated with MTX, but who had stopped MTX >6 months. Patients from AMBITION

had a longer duration of disease, higher tender and swollen joint counts and higher CRP levels than the other studies included in this MAIC analysis; whereas, patients from FUNCTION tended to show more similar characteristics to those in RA-BEGIN; the other studies included patients who were naïve to treatment. Because of this, we included the AMBITION trial only as a sensitivity analysis. Also, parameters, such as race and geographic location, were not included in the analysis, because these parameters were not widely reported in the original trial publications. Additionally, these variables and the inclusion of geographic location have rarely been explored in indirect comparisons. Lastly, the baricitinib RCT in the MAIC was conducted before the drug and dosage for baricitinib received regulatory approvals; a monotherapy study with a 2 mg dose of baricitinib was not conducted.

In conclusion, this MAIC suggests that among RA patients who are naïve to csDMARDs and bDMARDs, baricitinib 4 mg provides statistically significant greater pain reduction and improvement of physical function compared with adalimumab 40 mg and tocilizumab 8 mg/kg. No difference in pain reduction was observed between baricitinib and two times per day tofacitinib 5 mg with two of the three analyses employed, and no difference was observed in improving physical function. Well-designed, properly powered, head-to-head clinical trials are needed to confirm whether there is a class effect for JAK inhibitors over bDMARDs.

a) Model with duration of disease



b) Data from FUNCTION and AMBITION presented in combination and separately




Figure 2 Sensitivity analyses with (a) disease duration included in the model and (b) with data from AMBITION and FUNCTION analysed separately. HAQ-DI, Health Assessment Questionnaire-Disability Index; MTX, methotrexate; VAS, visual analogue scale.

Author affiliations

¹Sorbonne University, Pierre Louis Institute for Epidemiology and Public Health; Assistance Publique Hopitaux de Paris, Pitie-Salpetriere University Hospital, Rheumatology Dept, Paris, France
²Eli Lilly and Company, Indianapolis, Indiana, USA
³Botnar Research Centre, Univ of Oxford, Headington, UK
⁴University of Twente and Arthritis Center Twente, Enschede, Netherlands

⁵Leeds MSK Biomed/Chapel Allerton Hosp, Leeds, UK
⁶University of Texas Southwestern Med Ctr, Dallas, Texas, USA

Correction notice This article has been corrected since it was published online. The open access licence of the article has been reinstated.

Acknowledgements The authors would like to thank Molly E. Tomlin, MS, of Eli Lilly and Company for her assistance with manuscript preparation and process

support and Julie A Sherman of Eli Lilly and Company for her assistance in creating the figures.

Contributors All authors participated in the interpretation of data, provided critical comments and input and reviewed and approved the final manuscript. B. Zhu, C. Nicolay and K. Kadziola additionally conducted the analyses.

Funding This study was funded by Eli Lilly and Company and Incyte Corporation.

Competing interests B. Fautrel: Grant/research support from: AbbVie, Lilly, MSD, Pfizer; Consultant and consultancy fees from: AbbVie, Biogen, BMS, Celgene, Janssen, Lilly, Medac, MSD, NORDIC Pharma, Novartis, Pfizer, Roche, Sanofi-Aventis, SOBI, UCB; P. C. Taylor: Research grants from Celgene, Galapagos, Janssen, Lilly. Consultation fees from AbbVie, Biogen, Galapagos, Gilead, GlaxoSmithKline, Janssen, Lilly, Novartis, Pfizer, Roche, Sanofi, Nordic Pharma, Fresenius and UCB. M. van de Laar: Grant/research support from: AbbVie, Eli Lilly and Company, Sanofi-Genzyme, Pfizer; Janssen-Cilag. Consultant and consulting fees for: Eli Lilly and Company, Sanofi Genzyme, Abbvie. P. Emery: Consultant and consulting fees for: Pfizer, MSD, Abbvie, BMS, UCB, Roche, Novartis, Samsung, Sandoz, Eli Lilly and Company. R. Fleischmann: Grant/research support from: AbbVie, Amgen, AstraZeneca, Bristol-Myers Squibb, Celgene, Centrexion, Genetech, GlaxoSmithKline, Janssen, Eli Lilly and Company, Merck, Pfizer, Regeneron, Roche, Sanofi, Aventis, UCB; Consultant and consulting fees for: AbbVie, Amgen, Bristol-Myers Squibb, Celgene, Celltrion, GSK, Janssen, Eli Lilly and Company, Novartis, Pfizer, Samsung, Sanofi-Aventis. B. Zhu, F. De Leonardi, C.L. Kannowski, C. Nicolay, Z. Kadziola, and I. De La Torre: employees and shareholders of Eli Lilly and Company.

Patient consent Not required.

Ethics approval Not applicable.

Data sharing statement Lilly provides access to relevant anonymised patient-level data from studies on approved medicines and indications as defined by the sponsor-specific information on clinicalstudydatarequest.com. For details on submitting a request, see the instructions provided at clinicalstudydatarequest.com.

Provenance and peer review Not commissioned; externally peer reviewed.

Open access This is an open access article distributed in accordance with the Creative Commons Attribution Non Commercial (CC BY-NC 4.0) license, which permits others to distribute, remix, adapt, build upon this work non-commercially, and license their derivative works on different terms, provided the original work is properly cited, appropriate credit is given, any changes made indicated, and the use is non-commercial. See: <http://creativecommons.org/licenses/by-nc/4.0/>.

REFERENCES

- Fautrel B, Alten R, Kirkham B, *et al*. Call for action: how to improve use of patient-reported outcomes to guide clinical decision making in rheumatoid arthritis. *Rheumatol Int* 2018;38:935–47.
- Taylor PC, Moore A, Vasilescu R, *et al*. A structured literature review of the burden of illness and unmet needs in patients with rheumatoid arthritis: a current perspective. *Rheumatol Int* 2016;36:685–95.
- Oude Voshaar M, Das Gupta Z, Van de Laar MA, *et al*. Op0329-hpr outcomes that matter to people living with inflammatory arthritis; a global standard set, developed by the international consortium for health outcome measurement (ichom) working group for inflammatory arthritis. *Ann Rheum Dis* 2018;77:211.
- Taylor PC, Keystone EC, van der Heijde D, *et al*. Baricitinib versus placebo or adalimumab in rheumatoid arthritis. *N Engl J Med* 2017;376:652–62.
- Signorovitch JE, Wu EQ, Yu AP, *et al*. Comparative effectiveness without head-to-head trials: a method for matching-adjusted indirect comparisons applied to psoriasis treatment with adalimumab or etanercept. *Pharmacoeconomics* 2010;28:935–45.
- Breedveld FC, Weisman MH, Kavanaugh AF, *et al*. The premier study: a multicenter, randomized, double-blind clinical trial of combination therapy with adalimumab plus methotrexate versus methotrexate alone or adalimumab alone in patients with early, aggressive rheumatoid arthritis who had not had previous methotrexate treatment. *Arthritis Rheum* 2006;54:26–37.
- Strand V, Rentz AM, Cifaldi MA, *et al*. Health-related quality of life outcomes of adalimumab for patients with early rheumatoid arthritis: results from a randomized multicenter study. *J Rheumatol* 2012;39:63–72.
- Burmester GR, Rigby WF, van Vollenhoven RF, *et al*. Tocilizumab in early progressive rheumatoid arthritis: function, a randomised controlled trial. *Ann Rheum Dis* 2016;75:1081–91.
- Fleischmann R, Schiff M, van der Heijde D, *et al*. Baricitinib, methotrexate, or combination in patients with rheumatoid arthritis and no or limited prior disease-modifying antirheumatic drug treatment. *Arthritis Rheumatol* 2017;69:506–17.
- Jones G, Sebba A, Gu J, *et al*. Comparison of tocilizumab monotherapy versus methotrexate monotherapy in patients with moderate to severe rheumatoid arthritis: the ambition study. *Ann Rheum Dis* 2010;69:88–96.
- Lee EB, Fleischmann R, Hall S, *et al*. Tofacitinib versus methotrexate in rheumatoid arthritis. *N Engl J Med* 2014;370:2377–86.
- Bruce B, Fries JF. The health assessment questionnaire (HAQ). *Clin Exp Rheumatol* 2005;23:S14–8.
- Ramey D, Fries J, Singh G. The health assessment questionnaire 1995-status and review. In: Spilker B, ed. *Pharmacoeconomics and quality of life in clinical trials*. Philadelphia, Lippincott-Raven, 1996: 227–37.
- Petto H, Kadziola Z, Brnabic A, *et al*. Alternative weighting approaches for anchored matching-adjusted indirect comparisons via a common comparator. *Value Health* 2019;22:85–91.
- Bucher HC, Guyatt GH, Griffith LE, *et al*. The results of direct and indirect treatment comparisons in meta-analysis of randomized controlled trials. *J Clin Epidemiol* 1997;50:683–91.
- Erder MH, Xie J, Signorovitch JE, *et al*. Cost effectiveness of guanfacine extended-release versus atomoxetine for the treatment of attention-deficit/hyperactivity disorder: application of a matching-adjusted indirect comparison. *Appl Health Econ Health Policy* 2012;10:381–95.
- Sikirica V, Findling RL, Signorovitch J, *et al*. Comparative efficacy of guanfacine extended release versus atomoxetine for the treatment of attention-deficit/hyperactivity disorder in children and adolescents: applying matching-adjusted indirect comparison methodology. *CNS Drugs* 2013;27:943–53.
- Strand V, McInnes I, Mease P, *et al*. Matching-adjusted indirect comparison: secukinumab versus infliximab in biologic-naive patients with psoriatic arthritis. *J Comp Eff Res* 2019;8:497–510.
- Nash P, McInnes IB, Mease PJ, *et al*. Secukinumab versus adalimumab for psoriatic arthritis: comparative effectiveness up to 48 weeks using a matching-adjusted indirect comparison. *Rheumatol Ther* 2018;5:99–122.
- Strand V, Betts KA, Mittal M, *et al*. Comparative effectiveness of adalimumab versus secukinumab for the treatment of psoriatic arthritis: a matching-adjusted indirect comparison. *Rheumatol Ther* 2017;4:349–62.
- Mills EJ, Thorlund K, Ioannidis JP. Demystifying trial networks and network meta-analysis. *BMJ* 2013;346:f2914.