



Modifiable factors associated with postoperative atrial fibrillation in older patients with hip fracture in an orthogeriatric care pathway: a nested case–control study

Pierre Gay, Bastien Genet, Audrey Rouet, Rana Alkouri, Judith Cohen-Bittan, Jacques Boddaert, Lorène Zerah

► To cite this version:

Pierre Gay, Bastien Genet, Audrey Rouet, Rana Alkouri, Judith Cohen-Bittan, et al.. Modifiable factors associated with postoperative atrial fibrillation in older patients with hip fracture in an orthogeriatric care pathway: a nested case–control study. *BMC Geriatrics*, 2022, 22 (1), pp.845. 10.1186/s12877-022-03556-9 . hal-03856720

HAL Id: hal-03856720

<https://hal.sorbonne-universite.fr/hal-03856720>

Submitted on 31 Jan 2023

HAL is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers.

L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.

**BLOOD TRANSFUSION AND DELAYED SURGERY ASSOCIATED WITH
POSTOPERATIVE ATRIAL FIBRILLATION IN OLDER PATIENTS WITH HIP
FRACTURE IN A DEDICATED ORTHOGERIATRIC CARE PATHWAY: A
NESTED CASE–CONTROL STUDY**

**Pierre Gay¹, Bastien Genet¹, Audrey Rouet¹, Rana Alkouri², Judith Cohen-Bittan¹,
Boddaert Jacques^{1,3}, Lorène Zerah^{1,4}**

1. Assistance Publique-Hôpitaux de Paris (AP-HP), Hôpital Pitié-Salpêtrière,
Département de Gériatrie, Paris, France
2. Assistance Publique-Hôpitaux de Paris (AP-HP), Hôpital Pitié-Salpêtrière,
Département de Biochimie, Paris, France
3. Sorbonne Université, INSERM, Centre d'Immunologie et des Maladies
Infectieuses (Cimi-Paris), Paris, France
4. Sorbonne Université, INSERM, Institut Pierre Louis d'Épidémiologie et de Santé
Publique, Paris, France

Corresponding author:

Dr Lorène Zerah, Department of Geriatric Medicine, Hôpital la Pitié-Salpêtrière, 47-83 Bd de
l'Hôpital 75013 Paris, France. Email: lorene.zerah@aphp.fr

Tel: +33 1 84 82 81 95

Fax: +33 1 42 16 06 26

ORCID no: 0000-0002-3474-2200

Word count: N = 3074, Tables and figures: N = 4, ref: N = 37

ABSTRACT (249 words)

Background Few data are available regarding post-operative atrial fibrillation (POAF) in non-cardiothoracic surgery, particularly orthopedic surgery. Hence, given the frequent incidence of POAF after surgery and its marked impact, we need to identify modifiable risk factors associated with POAF after hip fracture surgery in older patients.

Methods We conducted a nested case–control study in the unit for perioperative geriatric care of an academic hospital in Paris from July 1, 2009 to December 31, 2019, enrolling all consecutive patients aged ≥ 70 years with hip fracture surgery and no history of permanent atrial fibrillation before admission. Patients with and without POAF were matched 1:5 on 5 baseline characteristics (age, hypertension, diabetes, coronary artery disease, cardiac failure).

Results Of the 757 patients included, 384 were matched, and 64 had POAF. The incidence of POAF was 8.5%. The mean age was 86 ± 6 years, 298 (78%) patients were female, and the median Charlson Comorbidity Index was 6 (interquartile range IQR 4–8). The median time between surgery and the occurrence of POAF was 2 days (1–4). On multivariable logistic regression analysis (matched cohort), the modifiable factors present at admission associated with POAF were time to surgery greater than 48 hr (odds ratio OR= 1.66, 95% Confidence interval CI [1.01–2.81]) and more than 2 transfusions of packed red blood cells (OR = 3.94, [1.50–10.03]).

Conclusions This study provides new information about POAF in older patients with hip fracture surgery, a surgical emergency whose complexity requires multidisciplinary care.

Key words: arrhythmia, orthopedic surgery, geriatrics

INTRODUCTION

Hip fracture, with the total number of cases increasing in line with the ageing population, is a major public health concern associated with poor vital and functional outcomes.(1–3) In parallel, atrial fibrillation (AF), the most common sustained cardiac arrhythmia in adults, is particularly common in older patients,(4) reaching 10% in patients over 80 years of age.(5) An increase in the number of surgical populations among older people over the last decade has increased the overall prevalence of postoperative AF (POAF), defined as new-onset AF in the immediate postoperative period: 20% to 50% of patients after cardiac surgery and 10% to 30% after non-cardiac thoracic surgery.(4)

POAF, whose usual peak of incidence is between day 2 and 4 postoperatively,(6) is associated with numerous complications such as cardioembolic stroke, acute coronary syndrome, and acute heart failure, with an impact on in-hospital mortality, length of hospital stay and medical health costs.(4,7–9) Several studies, largely focused on POAF after cardiothoracic surgery, identified triggering factors including baseline characteristics (age, male sex, hypertension, chronic heart failure, ischemic heart disease, chronic kidney disease), and perioperative and postoperative complications (hypovolemia, electrolyte disorders, hypoxemia).(4,6,9–12) However, few data are available regarding POAF after non-cardiothoracic surgery, particularly emergent orthopedic surgery.(4,13)

Hence, given the frequent incidence of POAF after surgery and its marked impact, we need to identify risk factors, particularly modifiable risk factors, associated with POAF after orthopedic surgery. We previously demonstrated that optimizing patient care by improving the management of peri-operative factors and thus decreasing postoperative complications could reduce by a maximum of one quarter the 6-month mortality rate after hip fracture.(14) Identifying the modifiable factors associated with the occurrence of POAF in a population with

multimorbidity and polypharmacy would be an essential step that could indicate the future directions for care improvement.

Our primary objective was to identify the modifiable acute risk factors (triggers) or protective factors associated with the occurrence of POAF in older patients with hip fracture surgery in a dedicated orthogeriatric pathway.

MATERIAL AND METHODS

The database was declared to the French National Commission on Computing and Liberty (CNIL) of the Assistance Publique-Hôpitaux de Paris (APHP) for this study (no. 20190822165316). This report follows the STROBE recommendations (eMethods 1 in the Supplement).(15)

Study design, study setting and eligibility criteria

A nested case–control study was conducted in the unit for perioperative geriatric care of an academic hospital. The unit for perioperative geriatric care is part of a dedicated orthogeriatric care pathway including coordination between the department of emergency medicine and surgery, department of anaesthesiology and critical care, department of orthopaedic surgery and department of rehabilitation. This dedicated orthogeriatric care pathway is defined as 1) an early alert from the emergency department (ED), 2) considering hip fracture as requiring surgery as soon as feasible (i.e., 24 hr/day), 3) rapid transfer to the unit for perioperative geriatric care after surgery, and 4) rapid transfer of stable patients to a dedicated rehabilitation unit.(16,17)

The management strategy in this unit for perioperative geriatric care, previously described(16), focused on early mobilization with the aim of chair-sitting and walking within 24 and 48 hr after arrival, respectively; pain management; the provision of air-filled mattresses

for patients with pressure sores or at high risk of pressure sores; swallowing disorders detected using a systematic medical survey; detection of stool impaction and urinary retention using bedside ultrasonography; correction of anemia with transfusion of packed red blood cells (usually when haemoglobin level is $<8 \text{ g.L}^{-1}$); and detection of delirium, AF and malnutrition.

From July 1, 2009 to December 31, 2019, all consecutive patients with hip fracture admitted to the unit for perioperative geriatric care were evaluated for eligibility. Patients were included if they were ≥ 70 years old and their primary presentation was hip fracture (first hospitalization after surgery in the unit for perioperative geriatric care). We excluded patients with multiple or metastatic or periprosthetic fractures, a history of permanent AF before admission in the unit for perioperative geriatric care, post-operative supraventricular arrhythmia other than AF; and missing data (missing anesthesia records, missing data from the ED, no electrocardiogram before admission in the unit for perioperative geriatric care). Patients were followed until death or the end of hospitalization in the unit for perioperative geriatric care. Some patients had been included in previous studies.(14,16–25)

Outcomes

Our main outcome measure was the occurrence of POAF defined as new-onset AF in the immediate period after surgery (until the end of hospitalization in the unit for perioperative geriatric care). The occurrence of POAF was retrospectively adjudicated by 2 geriatricians (AR, PG) who independently reviewed medical charts ($\kappa = 0.948$, 95% confidence interval [CI] 0.87–0.99). In case of disagreement, consensus was reached with a third independent senior expert (JB).

Data collection methods

Since the opening of the unit for perioperative geriatric care in 2009, we have created a

dedicated research database that is prospectively implemented by 3 senior geriatricians (JB, JCB, LZ), experts in orthogeriatrics, and that integrates all the data from the orthogeriatric care pathway for each patient.

The following variables were collected prospectively by interviewing patients, their family members or their physicians and pharmacists during the hospital stay and were defined as baseline characteristics before hip fracture: age, sex, home or nursing home living conditions, walking ability, previous medical history including cardiovascular and neurologic diseases, chronic medications, and type of fracture (radiological definition by an orthopedic surgeon).

Co-morbidity severity was assessed with the Charlson Comorbidity Index(26) because all comorbidity scores are equivalent in predicting mortality in this population(20). Functional status was evaluated with an activities of daily living scale(27). Repeated falls was defined as 2 or more falls in the previous year, chronic renal failure as Cockcroft creatinine clearance $< 30 \text{ ml} \cdot \text{min}^{-1}$, hypokalemia as potassium level $< 3.5 \text{ mmol/L}$ and anemia as haemoglobin level $< 12 \text{ g} \cdot \text{dL}^{-1}$ for women and $13 \text{ g} \cdot \text{dL}^{-1}$ for men.

During the perioperative period, we prospectively recorded the surgical treatment, the delay and duration of surgery, the anesthetic drugs used and all drugs and transfusions administered from the ED to the unit for perioperative geriatric care. Cardiovascular drugs taken at baseline and their suspension were recorded retrospectively.

After surgery, delays to first sitting and first walking, destination (home or rehabilitation) at discharge from the unit for perioperative geriatric care and length of stay in acute care were recorded. All postoperative complications during the acute care period were prospectively recorded.

Statistical analysis

Because the database was prospectively supplemented, all authors were “blinded” to the research question at the time of data collection. The study is based on data for all available patients during the study period, and thus no a priori power calculation was conducted. Data are presented as mean \pm SD or median (interquartile range) for continuous variables and number (percentage) for categorical variables. Comparison of quantitative variables involved unpaired Student *t* test or Mann-Whitney test depending on the normal distribution of data. Normality was assessed by graphical representation of the data distribution. Comparison of categorical variables involved chi-squared or Fisher’s exact test, as appropriate.

Patients with (cases) and without POAF (controls) were matched 1:5 on 5 baseline characteristics (age, hypertension, diabetes, coronary artery disease, cardiac failure). We selected those 5 factors from a literature review(4) and deliberations of a panel of 9 independent experts. This panel included 3 geriatricians, 4 cardiologists and 2 anesthesiologists who were all blinded to the research question at this time and to the other experts’ answers. We asked each expert the following standardized question: “According to you, what are the 9-10 main baseline predisposing factors of POAF among patients of 70 and more undergoing hip fracture surgery?” The 4 most frequently given answers were age (89%), diabetes (89%), hypertension (89%) and coronary artery disease (78%) (eMethods 2 in the Supplement). Then, 4 other factors were mentioned with equal frequency: history of paroxysmic AF, chronic heart failure, chronic kidney disease, and valvular disease. We chose to keep chronic heart failure because it was most homogeneously proposed by the 3 represented medical specialties and was previously described as a strong predisposing factor in this specific context (28–30).

Then, using the matched dataset, we performed a logistic conditional regression analysis to assess independent modifiable variables present at admission in the unit for perioperative geriatric care that were associated with POAF; adjusted odds ratios (ORs) and their 95% CIs

were calculated. To avoid overestimation, a conservative approach was used(31,32): all variables with $P < .20$ on univariate analyses and all clinically relevant variables were included.

Statistical analyses were performed with R v 1.4.1717 (package MatchIt). All p-values were two-tailed and $p < .05$ was considered statistically significant.

RESULTS

Demographic data, patient baseline characteristics, surgery and biology

We included 757 patients, and 384 were matched; 64 had POAF (**Figure 1**). Excluded patients ($n = 607$, 45%) had similar characteristics as patients included with one exception: fewer patients excluded had haemoglobin level $< 10 \text{ g.dL}^{-1}$ before surgery (68% vs 74%, $p = 0.03$) (eMethods 3 in the Supplement).

Baseline characteristics are reported in **Table 1** (matched cohort). The mean age was 86 ± 6 years; 298 (78%) patients were female; the median Charlson Comorbidity Index was 6 (IQR 4–8); and 17% were living in a nursing home before the hip fracture.

The incidence of POAF was 8.5% in our cohort (**Figure 1**), with a median time between hospitalization in our unit and the occurrence of POAF of 2 days (1–4). On univariate analysis (matched cohort), patients with and without POAF did not differ in comorbidities (except for history of paroxysmic AF), autonomy, type of hip fracture or surgery (**Table 1**). The median time to surgery was 26 hr (20–47), and for more patients with than without POAF, the time to surgery was longer than 48 hr: 23 (36%) versus 62 (19%) ($p = 0.004$). The groups did not differ in biologic factors on the first day of hospitalization (haemoglobin and potassium levels) (**Table 1**).

Perioperative therapeutics

On univariate analysis (matched cohort), patients with and without POAF did not differ

in use of anesthetic drugs or techniques (**Table 2**). Patients with POAF more frequently received more than 2 transfusions of packed red blood cells in the emergency room or recovery room than patients without POAF: 8 (13%) versus 9 (3%) ($p = 0.003$) (**Table 2**). The groups did not differ in cardiovascular drug use (treatment at baseline and suspension of treatment before surgery), with the exception of flecainide, used more at baseline in the POAF than non-POAF group (eMethods 4 in the Supplement).

Main outcome

On multivariable logistic regression analysis (matched cohort; all variables detailed in **Table 3**), the modifiable factors associated with POAF, present at admission in the unit for perioperative geriatric care, were time to surgery greater than 48 hr (OR = 1.66 [1.01–2.81]) and more than 2 transfusions of packed red blood cells (OR = 3.94 [1.50–10.03]).

Post-operative complications

On univariate analysis (matched cohort), stroke, acute heart failure and infection were more frequent in patients with than without POAF (**Table 2**). In 5 patients with acute heart failure, the heart failure occurred before the discovery of POAF. Also, in 3 patients with acute coronary syndrome, the acute coronary syndrome occurred before the discovery of POAF.

DISCUSSION

This study aimed to identify modifiable risk factors associated with POAF after hip fracture surgery in a geriatric population receiving treatment in a dedicated orthogeriatric care pathway. The incidence of POAF was 8.5% in our cohort. The occurrence of POAF was associated with time to surgery greater than 48 hr and more than 2 transfusions of packed red blood cells before hospitalization in the unit for perioperative geriatric care.

The reported incidence of POAF after non-thoracic non-cardiac surgery varies widely, from 0.5% to 15%, which reflects the heterogeneity in definitions (inclusion or not of patients with a history of AF), methods used to identify POAF, study populations (geriatric or not) and settings (type of surgeries).(4,6,13) We chose to include in this cohort patients with a history of paroxysmic AF, as for previous cohorts of patients with hip fracture,(10,11) to minimize classification bias. In studies by Leibowitz et al.(11) (n = 410, mean age 80 ± 7.8 years) and Rostagno et al. (10) (n = 2922, mean age 83.7 ± 8.2 years), the incidence of newly diagnosed AF among older comorbid patients undergoing hip fracture surgery was 3.7% and 3.6%, respectively. Because the definition of POAF and the populations were similar to our cohort,(10,11) the higher incidence we found in our cohort may have been related to differences in monitoring and screening POAF in this highly specialized environment.(14,16)

We found delayed surgery associated with the occurrence of POAF, which agrees with the study of Rostagno et al., finding a time to surgery less than 48 hr for 80% of patients in the control group versus 49% in the AF group (p = 0.0001).(10) Once again, surgery for a hip fracture should be considered an emergency. A meta-analysis including 191 873 patients found that early hip fracture surgery (cut-off between 24 and 48 hr) was associated with significantly reduced risk of death (OR = 0.74 [0.67–0.81]).(33) Of note, reducing the time to surgery is not included in the European Society of Cardiology recommendations to prevent POAF.(4)

We also found the occurrence of POAF associated with more than 2 transfusions of packed red blood cells before hospitalization in the unit for perioperative geriatric care, which may reflect complications of anemia or the transfusion itself.(34,35) Perioperative anemia is a known risk factor of POAF. Anemia may lead to relative ischemia of atrial cells and myocardial conduction tissue, thus altering the cell electric properties and leading to arrhythmias.(12,36) In addition, anemia, especially acute anemia, produces an intense adrenergic activation that may trigger POAF in predisposed patients.(12,36) However, in

cardiothoracic surgery, there is evidence that packed red blood cell transfusion increases the incidence of POAF.(37) We previously found in our unit that a restrictive transfusion strategy was associated with fewer in-hospital cardiovascular complications (including AF, coronary syndrome, heart failure and stroke) with no significant difference in long-term mortality.(19)

We found no association between hypokalemia and POAF in our cohort,(36) nor with cardiovascular drug use (beta-blocker interruption),(4,6) possibly owing to lack of power. In addition, we collected the serum potassium level only on the first day of hospitalization, which could explain why we did not find an association of hypokalemia with POAF occurring later during the hospital stay.

Our study has several limitations. First, it was an observational study, and causality cannot be demonstrated. Second, it was a retrospective study, and bias due to other procedural changes, which may have influenced the occurrence of POAF during the same time cannot be excluded. Third, many patients had to be excluded because of missing preoperative electrocardiogram results (including missing anesthesia or emergency records) to fully meet the definition (exclusion of permanent AF). However, these patients had the same characteristics as included patients with the exception that they had less anaemia before surgery (potential trigger for POAF). Fourth, patients were not continuously monitored, which may have resulted in an under-diagnosis of the incidence of paroxysmal POAF and misclassification bias. Fifth, because our study was conducted in a highly specialized environment that is associated with reduced mortality and postoperative complications as compared with patients admitted to an orthopaedic department,(14,16) our results may not be extrapolated to other wards.

This study may have significant implications for the care of perioperative patients. The complexity of POAF, in particular in a population with multimorbidity and polypharmacy, requires a multifaceted, holistic, and multidisciplinary approach to the management of AF

patients, with active involvement in partnership with geriatricians, surgeons, anesthesiologists and emergency physicians.(4)

CONCLUSIONS

Among older patients hospitalized in a dedicated orthogeriatric care pathway for hip fracture surgery, the occurrence of POAF was associated with 2 modifiable factors: time to surgery greater than 48 hr and more than 2 transfusions of packed of red blood cells before the hospitalization in the unit for perioperative geriatric care. Further studies are needed to demonstrate a possible causal link between these factors and POAF.

NOTES

Conflict of interest: All authors have completed and submitted the ICMJE Form for Disclosure of Potential Conflicts of Interest. Pr Boddaert reports personal fees for lectures from VIFOR Pharma and Baxter companies outside the submitted work. No other disclosures are reported.

Authors' contributions:

Study concept and design: Boddaert, Zerah, Rouet, Genet, Gay

B Genet had full access to all of the data in the study and take responsibility for the integrity of the data and the accuracy of the data analysis.

Acquisition, analysis, or interpretation of data: All authors

Drafting of the manuscript: Gay, Zerah

Critical revision of the manuscript for important intellectual content: All authors

Statistical analysis: Genet, Zerah

Study supervision: Zerah, Boddaert, Rouet

English editing: Laura Smales (BioMedEditing, Toronto, Canada).

Funding/support: This work was supported only by institutional sources

Transparency statement: The lead author affirms that this manuscript is an honest, accurate and transparent account of the study being reported; that no important aspects of the study have been omitted; and that any discrepancies from the study as planned have been explained.

Ethics approval: The database was declared to the French National Commission on Computing and Liberty (CNIL) of the Assistance Publique-Hôpitaux de Paris (APHP) for this study (no. 20190822165316)

Data sharing: Data can be obtained on request to Pr Jacques Boddaert (Jacques.boddaert@aphp.fr).

Competing interests: No conflict of interest to declare

Acknowledgments: We thank Dr Guillaume Coutance, Dr Sabine Drevet, Dr Thomas Gilbert, Dr Jules Greze, Dr Romuald Henry, Dr Léopold Oliver, Dr Cyril Sellier, Dr Francesca Santi, Dr Arnaud Savoure for their expertise (9 independent experts for matching).

REFERENCES

1. Berry SD, Kiel DP, Colón-Emeric C. Hip fractures in older adults in 2019. *JAMA*. 2019;321(22):2231-2232. doi:10.1001/jama.2019.5453.
2. Brauer CA, Coca-Perraillon M, Cutler DM, Rosen AB. Incidence and mortality of hip fractures in the United States. *JAMA*. 2009;302(14):1573-1579. doi:10.1001/jama.2009.1462.
3. Hannan EL, Magaziner J, Wang JJ, et al. Mortality and locomotion 6 months after hospitalization for hip fracture: risk factors and risk-adjusted hospital outcomes. *JAMA*. 2001;285(21):2736-2742. doi:10.1001/jama.285.21.2736.
4. Hindricks G, Potpara T, Dagres N, et al. 2020 ESC Guidelines for the diagnosis and management of atrial fibrillation developed in collaboration with the European Association for Cardio-Thoracic Surgery (EACTS): The Task Force for the diagnosis and management of atrial fibrillation of the European Society of Cardiology (ESC) Developed with the special contribution of the European Heart Rhythm Association (EHRA) of the ESC. *Eur Heart J*. 2021;42(5):373-498. doi:10.1093/eurheartj/ehaa612.
5. Go AS, Hylek EM, Phillips KA, et al. Prevalence of diagnosed atrial fibrillation in adults: national implications for rhythm management and stroke prevention: the AnTicoagulation and Risk Factors in Atrial Fibrillation (ATRIA) Study. *JAMA*. 2001;285(18):2370-2375. doi:10.1001/jama.285.18.2370.
6. Dobrev D, Aguilar M, Heijman J, Guichard J-B, Nattel S. Postoperative atrial fibrillation: mechanisms, manifestations and management. *Nat Rev Cardiol*. 2019;16(7):417-436. doi:10.1038/s41569-019-0166-5.

7. Subramaniam B, Shankar P, Shaefi S, et al. Effect of Intravenous Acetaminophen vs Placebo Combined With Propofol or Dexmedetomidine on Postoperative Delirium Among Older Patients Following Cardiac Surgery: The DEXACET Randomized Clinical Trial. *JAMA*. 2019;321(7):686-696. doi:10.1001/jama.2019.0234.
8. Lin M-H, Kamel H, Singer DE, Wu Y-L, Lee M, Ovbiagele B. Perioperative/Postoperative Atrial Fibrillation and Risk of Subsequent Stroke and/or Mortality. *Stroke*. 2019;50(6):1364-1371. doi:10.1161/STROKEAHA.118.023921.
9. Bhavé PD, Goldman LE, Vittinghoff E, Maselli J, Auerbach A. Incidence, predictors, and outcomes associated with postoperative atrial fibrillation after major noncardiac surgery. *Am Heart J*. 2012;164(6):918-924. doi:10.1016/j.ahj.2012.09.004.
10. Rostagno C, Cartei A, Rubbieri G, et al. Postoperative atrial fibrillation is related to a worse outcome in patients undergoing surgery for hip fracture. *Intern Emerg Med*. 2021;16(2):333-338. doi:10.1007/s11739-020-02372-6.
11. Leibowitz D, Abitbol C, Alcalai R, Rivkin G, Kandel L. Perioperative atrial fibrillation is associated with increased one-year mortality in elderly patients after repair of hip fracture. *Int J Cardiol*. 2017;227:58-60. doi:10.1016/j.ijcard.2016.11.067.
12. Bessissow A, Khan J, Devereaux PJ, Alvarez-Garcia J, Alonso-Coello P. Postoperative atrial fibrillation in non-cardiac and cardiac surgery: an overview. *J Thromb Haemost JTH*. 2015;13 Suppl 1:S304-312. doi:10.1111/jth.12974.
13. Subramani Y, El Tohamy O, Jalali D, Nagappa M, Yang H, Fayad A. Incidence, Risk Factors, and Outcomes of Perioperative Atrial Fibrillation following Noncardiothoracic Surgery: A Systematic Review and Meta-Regression Analysis of Observational Studies. *Anesthesiol Res Pract*. 2021;2021:5527199. doi:10.1155/2021/5527199.

14. Zerah L, Hajage D, Raux M, et al. Attributable Mortality of Hip Fracture in Older Patients: A Retrospective Observational Study. *J Clin Med*. 2020;9(8). doi:10.3390/jcm9082370.
15. von Elm E, Altman DG, Egger M, et al. The Strengthening the Reporting of Observational Studies in Epidemiology (STROBE) statement: guidelines for reporting observational studies. *PLoS Med*. 2007;4(10):e296. doi:10.1371/journal.pmed.0040296.
16. Boddaert J, Cohen-Bittan J, Khiami F, et al. Postoperative admission to a dedicated geriatric unit decreases mortality in elderly patients with hip fracture. *PloS One*. 2014;9(1):e83795. doi:10.1371/journal.pone.0083795.
17. Boddaert J, Raux M, Khiami F, Riou B. Perioperative management of elderly patients with hip fracture. *Anesthesiology*. 2014;121(6):1336-1341. doi:10.1097/ALN.0000000000000478.
18. Vallet H, Chenevier-Gobeaux C, Villain C, et al. Prognostic Value of Serum Procalcitonin After Orthopedic Surgery in the Elderly Population. *J Gerontol A Biol Sci Med Sci*. 2017;72(3):438-443. doi:10.1093/gerona/glw097.
19. Zerah L, Dourthe L, Cohen-Bittan J, et al. Retrospective evaluation of a restrictive transfusion strategy in older adults with hip fracture. *J Am Geriatr Soc*. 2018;66(6):1151-1157. doi:10.1111/jgs.15371.
20. Boddaert J, Na N, Le Manach Y, et al. Prediction of postoperative mortality in elderly patients with hip fracture: are specific and geriatric scores better than general scores? *Br J Anaesth*. 2017;118(6):952-954. doi:10.1093/bja/aex148.

21. Vallet H, Breining A, Le Manach Y, et al. Isolated cardiac troponin rise does not modify the prognosis in elderly patients with hip fracture. *Medicine (Baltimore)*. 2017;96(7):e6169. doi:10.1097/MD.00000000000006169.
22. Magny E, Vallet H, Cohen-Bittan J, et al. Pressure ulcers are associated with 6-month mortality in elderly patients with hip fracture managed in orthogeriatric care pathway. *Arch Osteoporos*. 2017;12(1):77. doi:10.1007/s11657-017-0365-9.
23. Zerah L, Cohen-Bittan J, Raux M, et al. Association between Cognitive Status before Surgery and Outcomes in Elderly Patients with Hip Fracture in a Dedicated Orthogeriatric Care Pathway. *J Alzheimers Dis JAD*. 2017;56(1):145-156. doi:10.3233/JAD-160655.
24. Genet B, Lamy T, Cohen-Bittan J, et al. Lack of Association Between Perioperative Medication and Postoperative Delirium in Hip Fracture Patients in an Orthogeriatric Care Pathway. *J Am Med Dir Assoc*. October 2021:S1525-8610(21)00857-4. doi:10.1016/j.jamda.2021.09.022.
25. Thietart S, Baque M, Cohen-Bittan J, et al. Short-term administration of nonsteroidal anti-inflammatory drugs in older patients with hip fracture: A cohort study. *Eur J Anaesthesiol*. 2021;38(9):1003-1005. doi:10.1097/EJA.0000000000001482.
26. Charlson ME, Pompei P, Ales KL, MacKenzie CR. A new method of classifying prognostic comorbidity in longitudinal studies: development and validation. *J Chronic Dis*. 1987;40(5):373-383. doi:10.1016/0021-9681(87)90171-8.
27. Katz S, Akpom CA. 12. Index of ADL. *Med Care*. 1976;14(5 Suppl):116-118. doi:10.1097/00005650-197605001-00018.

28. Gupta BP, Huddleston JM, Kirkland LL, et al. Clinical presentation and outcome of perioperative myocardial infarction in the very elderly following hip fracture surgery. *J Hosp Med*. 2012;7(9):713-716. doi:10.1002/jhm.1967.
29. Leibowitz D, Abitbol C, Alcalai R, Rivkin G, Kandel L. Perioperative atrial fibrillation is associated with increased one-year mortality in elderly patients after repair of hip fracture. *Int J Cardiol*. 2017;227:58-60. doi:10.1016/j.ijcard.2016.11.067.
30. Khormae S, Do HT, Mayr Y, et al. Risk of Ischemic Stroke After Perioperative Atrial Fibrillation in Total Knee and Hip Arthroplasty Patients. *J Arthroplasty*. 2018;33(9):3016-3019. doi:10.1016/j.arth.2018.04.009.
31. Sun GW, Shook TL, Kay GL. Inappropriate use of bivariable analysis to screen risk factors for use in multivariable analysis. *J Clin Epidemiol*. 1996;49(8):907-916. doi:10.1016/0895-4356(96)00025-x.
32. Harrell FE, Lee KL, Matchar DB, Reichert TA. Regression models for prognostic prediction: advantages, problems, and suggested solutions. *Cancer Treat Rep*. 1985;69(10):1071-1077.
33. Moja L, Piatti A, Pecoraro V, et al. Timing matters in hip fracture surgery: patients operated within 48 hours have better outcomes. A meta-analysis and meta-regression of over 190,000 patients. *PloS One*. 2012;7(10):e46175. doi:10.1371/journal.pone.0046175.
34. Carson JL, Terrin ML, Noveck H, et al. Liberal or restrictive transfusion in high-risk patients after hip surgery. *N Engl J Med*. 2011;365(26):2453-2462. doi:10.1056/NEJMoa1012452.

35. Willett LR, Carson JL. Management of postoperative complications: anemia. *Clin Geriatr Med*. 2014;30(2):279-284. doi:10.1016/j.cger.2014.01.006.
36. Chelazzi C, Villa G, De Gaudio AR. Postoperative atrial fibrillation. *ISRN Cardiol*. 2011;2011:203179. doi:10.5402/2011/203179.
37. Sood N, Coleman CI, Kluger J, White CM, Padala A, Baker WL. The association among blood transfusions, white blood cell count, and the frequency of post-cardiothoracic surgery atrial fibrillation: a nested cohort study from the Atrial Fibrillation Suppression Trials I, II, and III. *J Cardiothorac Vasc Anesth*. 2009;23(1):22-27. doi:10.1053/j.jvca.2008.06.009.

Tables Legends

Table 1: Demographic data and baseline characteristics of older patients with and without postoperative atrial fibrillation (POAF) (matched population)

Table 2: Management, complications and in-hospital outcomes of older patients with and without POAF (matched population)

Table 3: Logistic regression analysis of factors associated with occurrence of POAF (matched population)

Figure Legends

Figure 1: Flow chart of participants in the study.

Table 1: Demographic data and baseline characteristics of older patients with and without postoperative atrial fibrillation (POAF) (matched population)

	All patients N = 384	Without POAF N = 320	With POAF N = 64	P value
Age (years)*	86 ± 6	86 ± 6	86 ± 6	0.99
<85	141 (37)	116 (36)	25 (39)	0.67
85 to 90	145 (38)	122 (38)	23 (36)	0.74
>90	98 (26)	82 (26)	16 (25)	0.92
Male sex	86 (22)	71 (22)	15 (23)	0.83
Medical history				
CCI	6 (4-8)	6 (4-8)	7 (4-7)	0.71
Paroxysmic atrial fibrillation	77 (20)	53 (17)	24 (38)	<0.001
Hypertension*	264 (69)	220 (69)	44 (69)	0.99
Diabetes*	24 (6)	20 (6)	4 (6)	0.99
Coronary artery disease*	66 (17)	55 (17)	11 (17)	0.99
Cardiac failure*	48 (13)	40 (13)	8 (13)	0.99
Valvular heart disease	30 (8)	22 (7)	8 (13)	0.13
COPD	30 (8)	24 (8)	6 (9)	0.61
Chronic renal failure†	35 (9)	29 (9)	6 (9)	0.99
Autonomy before surgery				
ADL	5.5 (4-6)	5.5 (3.5-6)	5.5 (5-6)	0.31
iADL	2 (0-4)	2 (0-4)	2 (1-3)	0.94
Living in institution	65 (17)	64 (20)	1 (2)	<0.001
Walking before surgery	373 (97)	309 (97)	64 (100)	0.13
Walking with assistance	227 (59)	186 (58)	41 (64)	0.38
Fracture				
Intertrochanteric fracture	206 (54)	169 (53)	37 (58)	0.48
Femoral neck fracture	156 (41)	133 (42)	23 (36)	0.39
Surgery				
Time to surgery (hr)	26 (20-47)	26 (19-46)	33 (20-54)	0.11
Time to surgery > 48 hr	85 (22)	62 (19)	23 (36)	0.004
Duration of surgery (min)	143 ± 48	141 ± 43	151 ± 68	0.13
Missing values	4 (1)	4 (1)	0 (0)	-
Type of surgery				
Gamma nail	196 (51)	162 (51)	34 (53)	0.79
Dynamic hip screw	22 (6)	17 (5)	5 (8)	0.39
Unipolar prosthesis	138 (36)	117 (37)	21 (33)	0.52
Bipolar prosthesis	12 (3)	10 (3)	2 (3)	0.99
Missing values	4 (1)	4 (1)	0 (0)	-
Biologic factors				
Haemoglobin < 10 g.dL ⁻¹ before surgery	292 (76)	245 (77)	47 (73)	0.56
Missing values	1 (0)	1 (0)	0 (0)	-
Kalemia < 3.5 mmol/L at admission	42 (11)	34 (11)	8 (13)	0.66
Missing values	32 (8)	27 (8)	5 (8)	-

Data are mean ± SD, median (interquartile range), or number (percentage). Missing values are specified only if present.

* Matching on these 5 characteristics

Abbreviations: CCI = Charlson Comorbidity; COPD = chronic obstructive pulmonary disease; chronic renal failure = creatinine clearance <60 ml/min; ADL = activities of daily living, iADL = index of activities of daily living

Table 2: Management, complications and in-hospital outcomes of older patients with and without postoperative atrial fibrillation (POAF) (matched population)

	All patients N = 384	Without POAF N = 320	With POAF N = 64	P value
Perioperative therapeutics				
General anesthesia	342 (89)	285 (89)	57 (89)	0.89
Missing values	31 (8)	26 (8)	5 (8)	-
Femoral block	154 (40)	12 (4)	31 (48)	0.17
Missing values	37 (10)	32 (10)	5 (8)	-
Morphinics	338 (88)	280 (88)	58 (91)	0.73
Missing values	38 (10)	33 (10)	5 (8)	-
Hypnotics	333 (87)	275 (86)	58 (91)	0.36
Missing values	38 (10)	33 (10)	5 (8)	-
Fluranes	200 (52)	167 (52)	33 (52)	0.42
Missing values	80 (21)	70 (22)	10 (16)	-
Catecholamines	301 (78)	252 (79)	49 (77)	0.32
Missing values	39 (10)	34 (11)	5 (8)	-
Others				
Prostigmin	14 (4)	13 (4)	1 (2)	0.48
Missing values	40 (10)	35 (11)	5 (8)	-
Atropine	25 (7)	21 (7)	4 (6)	0.99
Missing values	39 (10)	34 (11)	5 (8)	-
Transfusion in ER or RR	111 (29)	88 (28)	23 (36)	0.16
> 2 transfusions of packed RBCs	17 (4)	9 (3)	8 (13)	0.003
Missing values	3 (1)	3 (1)	0 (0)	-
In-hospital complications				
Infection	68 (18)	51 (16)	17 (27)	0.04
Acute heart failure	36 (9)	23 (7)	13 (20)	0.001
Acute coronary syndrome	27 (7)	20 (6)	7 (11)	0.18
Thromboembolic disease	27 (7)	23 (7)	4 (6)	0.99
Stroke	4 (1)	0 (0)	4 (6)	<0.001
Discharge				
In-hospital mortality	7 (2)	5 (2)	2 (3)	0.33
Back home at discharge	229 (60)	190 (59)	39 (61)	0.82
Length of stay (days)	11 ± 6.1	11 ± 5.5	13 ± 8.5	0.007

Data are mean ± SD, median (interquartile range), or number (percentage). Missing values are specified only if present.

Abbreviations: ER = emergency room; RR = recovery room; RBC = red blood cell

Table 3: Logistic regression analysis of factors associated with occurrence of postoperative atrial fibrillation (POAF) (matched population)

	Univariate analysis OR (95% CI)	P value	Multivariable analysis OR (95% CI)	P value
Baseline				
CCI (for 1-point increase)	0.94 (0.86–1.02)	0.15	0.96 (0.85–1.09)	0.54
Male sex	1.08 (0.74–1.59)	0.70	1.19 (0.71–2.01)	0.52
History of paroxysmic atrial fibrillation	2.45 (1.70–3.54)	<0.001	2.94 (1.75–4.92)	0.001
Time to surgery > 48 hr	2.36 (1.62–3.43)	<0.001	1.66 (1.01–2.81)	0.05
Duration of surgery (min)	1.01 (1.00–1.02)	0.03	0.99 (0.99–1.01)	0.18
Hypokalemia	1.16 (0.68–1.98)	0.59	1.68 (0.84–3.35)	0.14
Therapeutics				
Femoral block	1.54 (1.10–2.15)	0.01	0.86 (0.56–1.33)	0.50
> 2 transfusions of packed RBCs	4.88 (2.28–10.4)	<0.001	3.94 (1.50–10.3)	0.005

N = 321 (cases: 54, controls: 267), Concordance = 0.69

Hypokalemia = potassium level <3.5 mmol.l⁻¹ at admission in unit for perioperative geriatric care

Abbreviations: OR = odds ratio, CI: confidence interval; CCI = Charlson Comorbidity Index; RBC = red blood cell

Figure 1: Flow chart of participants in the study

