



**HAL**  
open science

## Emerging Opportunities for Life Course Research on Neighbourhoods and Mental Health

Milagros Ruiz, Basile Chaix

► **To cite this version:**

Milagros Ruiz, Basile Chaix. Emerging Opportunities for Life Course Research on Neighbourhoods and Mental Health. *Journal of Epidemiology and Community Health*, 2019, 73 (8), pp.689–690. 10.1136/jech-2019-212419 . hal-03889757

**HAL Id: hal-03889757**

**<https://hal.sorbonne-universite.fr/hal-03889757>**

Submitted on 21 Feb 2023

**HAL** is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers.

L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.

**Title:** Emerging opportunities for life course research on neighbourhoods and mental health

**List of authors:** Milagros Ruiz<sup>1</sup> and Basile Chaix<sup>2</sup>

**Author affiliations:** <sup>1</sup>Research Department of Epidemiology and Public Health, University College London, London, United Kingdom; <sup>2</sup>Sorbonne Université, INSERM, Institut Pierre Louis d'Épidémiologie et de Santé Publique, Nemesis research team, Paris, France

**Corresponding author:** Milagros Ruiz, Research Department of Epidemiology and Public Health, University College London, 1-19 Torrington Place, London WC1E 6BT, UK, m.a.ruiz@ucl.ac.uk, +44 (0)20 7679 8252

**Word count:** 1,200 (1,200 max.)

**Keywords:** adolescents, depression, depressive symptoms, life course, mental health, neighbourhood effects, US, young adults

**Funding:** MR and BC are supported by a European Commission Horizon 2020 Grant, 667661, as part of the Promoting Mental Wellbeing in the Ageing Population: Determinants, Policies and Interventions in European Cities (MINDMAP) research project.

**License for Publication:** The Corresponding Author has the right to grant on behalf of all authors and does grant on behalf of all authors, an exclusive license (or non exclusive for government employees) on a worldwide basis to the BMH Publishing Group Ltd to permit this article (if accepted) to be published in JECH and any other BMJPG products and sublicenses such use and exploit all subsidiary rights, as set out in our license (<http://group.bmj.com/products/journals/instructions-for-authors/licence-forms>).

**Competing interests:** None declared.

Social capital, referring to the quality and quantity of social relationships in a community, and the socio-economic characteristics of neighbourhoods are considered influential determinants for the mental wellbeing of residents,<sup>1-3</sup> but most of the empirical evidence on these relationships is limited in several ways. Cross-sectional studies cannot distinguish between social causation and social selection/drift mechanisms. For example, poor mental health may constrain individuals to migrate into, or remain in, neighbourhoods with worse conditions.<sup>1 2</sup> A further, and equally critical, limitation of the cross-sectional design for this research area is the absence of the life course approach.

While common mental disorders can occur at any age, the first onset of major depression most likely occurs during young adulthood. Consequently, depression onset in young adulthood is a powerful predictor of chronic depression or recurrent depressive episodes in later life.<sup>4</sup> Therefore, understanding the links between the neighbourhood environment, as intertwined with and influencing other life circumstances (job market integration, social integration), and depression during young adulthood is crucial to assess how these conditions may increase the risk of developing this disorder. Given limited and conflicting reports for emerging adults, it is unclear whether life circumstances and more proximate risk factors play a greater role than neighbourhood factors at this life stage, or whether neighbourhoods are important through their impact on these determinants.<sup>5</sup>

Goldstein and colleagues add much-needed life course evidence to this literature by investigating the associations between social fragmentation (a composite index based on the share of female-headed households, residents living in the area <5 years, foreign-born residents and renters), income inequality (GINI index) and economic disadvantage (household income) with depressive symptoms in the US NEXT Generation Health Study.<sup>5</sup> Using annual assessments on depressive symptoms, the study tracked a nationally representative sample of US adolescents from 15-16 to 21-22 years of age, thereby capturing their transition into adulthood. Markers of the neighbourhood environment were derived using objective Census-tract level data geocoded using residential addresses at four time points over the follow-up. Contrary to most studies in this literature that relied on measurements at a single occasion,<sup>1-3</sup> repeated exposure data allows the study to account for changes in neighbourhood conditions.

Goldstein et al employed a multilevel prospective design to investigate depressive symptom differences by the neighbourhood environmental attributes independent of individual- and family-level factors. None of the nested models showed any evidence of symptom differences by social fragmentation, income inequality or economic disadvantage over the six-year period. Their lack of findings echo previous studies whereby neighbourhood socio-economic conditions did not increase the risk of depressive symptoms among young adults over a 10- and 14-year study in Finland and the US, respectively.<sup>5</sup>

The authors express methodological concern regarding the use of US census tracts to identify neighbourhoods, which group an average of 4,000 residents according to administrative boundaries. As census tracts do not accurately capture egocentric definitions of neighbourhoods that are more likely to reflect the daily experienced neighbourhood, this spatial unit may have underestimated true associations.<sup>6</sup> Although assessing social fragmentation, income inequality and economic disadvantage using egocentrically defined neighbourhoods is one strategy to improve studies like the one conducted by Goldstein et al,

these markers remain an indirect sociodemographic proxy of the social environment that are unlikely to efficiently capture influential dimensions such as social cohesion, collective efficacy, and social incivilities and delinquency.<sup>7</sup>

Notably, the abovementioned US study<sup>5</sup> found that depressive symptoms did not increase by neighbourhood disadvantage, using similar data as for Goldstein et al's<sup>5</sup> marker of social fragmentation; but perceived neighbourhood safety and physical neglect strongly predicted higher symptom increases from adolescence to adulthood.<sup>5</sup> A dominant criticism of evidence relating self-reported neighbourhood characteristics, such as cognitive (perceived) social capital, with mental health is the likelihood of cognitive distortion bias or same-source bias.<sup>13</sup> Reverse causation, however, is less probable among young adults because depression rarely occurs prior to this life stage.<sup>4</sup> Ultimately however, neighbourhood perceptions become informative when they can be compared to objective neighbourhood data of the same dimension. Despite the tendency to prioritize objective over subjective measurements, both are vital for a nuanced exploration of the relationships between neighbourhood social and physical environments and mental health. As little work has employed methodological techniques that can refine the measurement of neighbourhoods using subjective appraisals,<sup>8,9</sup> we encourage future studies to implement econometric or multilevel factor analytic methods to capture the abovementioned neighbourhood characteristics related to the social interactions of residents.

Goldstein et al caution against using their null findings to dismiss the importance of neighbourhood environments on the mental wellbeing of emerging adults.<sup>5</sup> For instance, their study focussed on the residential neighbourhood, while young adults are particularly mobile and likely to spend a substantial fraction of their time out of their residential environment. Daily dynamics of individuals over space are essential because they reflect key social and behavioural processes and expose them to distinct geographic environments. Recent advances and novel methods in sensor-based tracking, paired with smartphone ecological momentary assessment, can help generate knowledge on how individual and neighbourhood determinants experienced over daily activities coincide to influence health-related behaviours, and physical and mental health.<sup>10</sup> Applying these tools to studies of late adolescents and young adults would be particularly valuable as many social and health trajectories take shape during this period.<sup>4</sup>

Moreover, the full breadth of the neighbourhood context entails not only the social environment, but also the physical and built environment and access to health and social services. These additional dimensions portray other harmful forms of neighbourhood disadvantage, including poor aesthetic qualities of place, low street integration and dwindling infrastructure to sparsity of service providers and poor quality of care.<sup>7</sup> Furthermore, in addition to the neighbourhood local social and physical environment, it is relevant to recognise the wider regulatory and policy context at the meso- and macro-scales. It is obviously no easy task to gather data on this wide set of environmental determinants and the MINDMAP project bears significance on this research area. By linking individual-level data from multiple cohort studies of ageing across Europe and North America with not only area-level information on the social and physical environment, but also with policy indicators, MINDMAP provides a unique opportunity to understand how neighbourhoods and the wider social, physical and political environment influences mental wellbeing and cognition in the

later life course.<sup>11</sup> Data integration of this kind can enrich other life course studies interested in neighbourhood effects.

Life course environmental epidemiology provides a framework to enhance knowledge on the relationship of the neighbourhood context with mental wellbeing. For example, certain exposures may be important through sensitive period, accumulation of risk or chain of risk processes across the life course.<sup>4</sup> As maternal depression is a grave risk factor for offspring depression,<sup>4</sup> evidence relating neighbourhood disadvantage with poor mental health among mothers of infants and young children<sup>3</sup> also highlights the importance of intergenerational processes. Applying these models to understand the causal effects of neighbourhoods on mental health net of a large set of confounders requires investigators to grapple with a complex hierarchy of individual- and area-level factors thoughtfully organized within a causal diagram (e.g., a directed acyclic graph).<sup>12</sup> Multilevel longitudinal investigations should draw on the emerging opportunities discussed here in order to promote knowledge on neighbourhoods and mental health across the life course.

## References

1. Ehsan AM, De Silva MJ. Social capital and common mental disorder: a systematic review. *J Epidemiol Community Health* 2015;69(10):1021-8. doi: 10.1136/jech-2015-205868
2. Richardson R, Westley T, Garipey G, et al. Neighborhood socioeconomic conditions and depression: a systematic review and meta-analysis. *Soc Psychiatry Psychiatr Epidemiol* 2015;50(11):1641-56. doi: 10.1007/s00127-015-1092-4
3. Blair A, Ross NA, Garipey G, et al. How do neighborhoods affect depression outcomes? A realist review and a call for the examination of causal pathways. *Soc Psychiatry Psychiatr Epidemiol* 2014;49(6):873-87. doi: 10.1007/s00127-013-0810-z
4. Rudenstine S. Applying a life course perspective to depression. In: Koenen K, Rudenstine S, Susser E, et al., eds. *A Life Course Approach to Mental Disorders*. First edition ed. Oxford, UK: Oxford University Press 2014:88-96.
5. Goldstein RB, Lee AK, Haynie DL, et al. Neighbourhood disadvantage and depressive symptoms among adolescents followed into emerging adulthood. *J Epidemiol Community Health* 2019 doi: 10.1136/jech-2018-212004
6. Duncan DT, Kawachi I, Subramanian SV, et al. Examination of how neighborhood definition influences measurements of youths' access to tobacco retailers: a methodological note on spatial misclassification. *Am J Epidemiol* 2014;179(3):373-81. doi: 10.1093/aje/kwt251
7. Galea S, Vlahov D. Urban Health: Evidence, Challenges, and Directions. *Annu Rev Public Health* 2004;26(1):341-65. doi: 10.1146/annurev.publhealth.26.021304.144708
8. Chaix B, Lindström M, Rosvall M, et al. Neighbourhood social interactions and risk of acute myocardial infarction. *J Epidemiol Community Health* 2007;62:62-68.
9. Dunn EC, Masyn KE, Johnston WR, et al. Modeling contextual effects using individual-level data and without aggregation: an illustration of multilevel factor analysis (MLFA) with collective efficacy. *Popul Health Metr* 2015;13(1):12. doi: 10.1186/s12963-015-0045-1
10. Chaix B. Mobile Sensing in Environmental Health and Neighborhood Research. *Annu Rev Public Health* 2018;39(1):367-84. doi: 10.1146/annurev-publhealth-040617-013731
11. Beenackers MA, Doiron D, Fortier I, et al. MINDMAP: establishing an integrated database infrastructure for research in ageing, mental well-being, and the urban environment. *BMC Public Health* 2018;18(1):158. doi: 10.1186/s12889-018-5031-7
12. Nandi A, Welsh L. Social context and mental health over the life course. In: Koenen K, Rudenstine S, Susser E, et al., eds. *A Life Course Approach to Mental Disorders*. First edition ed. Oxford, UK: Oxford University Press 2014:227-37.