



HAL
open science

Quadrupolar $^{23}\text{Na}^+$ NMR relaxation as a probe of subpicosecond collective dynamics in aqueous electrolyte solutions

Iurii Chubak, Leeor Alon, Emilia V Silletta, Guillaume Madelin, Alexej Jerschow, Benjamin Rotenberg

► To cite this version:

Iurii Chubak, Leeor Alon, Emilia V Silletta, Guillaume Madelin, Alexej Jerschow, et al.. Quadrupolar $^{23}\text{Na}^+$ NMR relaxation as a probe of subpicosecond collective dynamics in aqueous electrolyte solutions. Nature Communications, 2023, 14 (1), pp.84. 10.1038/s41467-022-35695-3 . hal-03928032

HAL Id: hal-03928032

<https://hal.sorbonne-universite.fr/hal-03928032>

Submitted on 6 Jan 2023

HAL is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers.

L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.

Quadrupolar $^{23}\text{Na}^+$ NMR Relaxation as a Probe of Ultrafast Collective Dynamics in Aqueous Electrolyte Solutions

Iurii Chubak,^{1,*} Leeor Alon,^{2,3,*} Emilia V. Silletta,^{4,5}

Guillaume Madelin,^{2,3} Alexej Jerschow,^{6,†} and Benjamin Rotenberg^{1,‡}

¹*Sorbonne Université CNRS, Physico-Chimie des électrolytes et Nanosystèmes Interfaciaux, F-75005 Paris, France*

²*New York University School of Medicine, Department of Radiology,*

Center for Biomedical Imaging, 660 First Avenue, New York, NY 10016, USA

³*Center for Advanced Imaging Innovation and Research, Department of Radiology, New York University Grossman School of Medicine, New York, NY, 10016, USA*

⁴*Universidad Nacional de Córdoba, Facultad de Matemática, Astronomía, Física y Computación, Medina Allende s/n, X5000HUA, Córdoba, Argentina.*

⁵*Instituto de Física Enrique Gaviola, CONICET, Medina Allende s/n, X5000HUA, Córdoba, Argentina.*

⁶*New York University, Department of Chemistry, 100 Washington Square E, New York, NY 10003, USA*

Nuclear magnetic resonance relaxometry represents a powerful tool for extracting dynamic information. Yet, obtaining links to molecular motion is challenging for many ions that relax through the quadrupolar mechanism, which is mediated by electric field gradient fluctuations and lacks a detailed microscopic description. For sodium ions in dilute and concentrated aqueous electrolytes, we show that combining *ab initio* calculations and classical molecular dynamics to account for electron cloud effects and long-time sampling of fluctuations enables reaching a good agreement between the computed and experimental relaxation rates over a broad range of temperatures. While the commonly-assumed rotational models fail to rationalize the measured effects, we find the subpicosecond timescale of electric field gradient relaxation to be comparable to that of solution structural rearrangements. This work thus shows that quadrupolar sodium relaxometry can sense the fast collective dynamics around the solute, thereby elevating its role in exploring electrolyte or bioelectrolytic systems.

INTRODUCTION

The proper characterization and modeling of the solvation structure of alkaline cations (e.g., Li^+ , Na^+ , and K^+) in aqueous solution is of considerable interest both in physiological systems [1–8] and electrolytes used for electrochemical devices [9–13]. Nuclear magnetic resonance (NMR) spectroscopy provides an excellent source of dynamic and structural information for a number of nuclear species, including ^{23}Na with a nuclear spin $3/2$ and close to 100% natural abundance that produces the second strongest NMR signal after protons in biological tissues [14]. The NMR sensitivity of sodium is 9.2% of that of proton, while its typical concentration can be three, or more, orders of magnitude lower. Thus, in biological systems the sodium signal-to-noise ratio is 3,000–12,000 times lower than that of ^1H [4]. Nonetheless, the longitudinal relaxation time T_1 of ^{23}Na (typically 40 ms and below) is short compared to that of ^1H (on the order of seconds) [4], allowing for rapid averaging of the signals such that quantitative analysis is made possible within reasonable time scales [15].

The shortness of the ^{23}Na NMR relaxation times is due to a fluctuating quadrupole interaction related to the changes in the solvation shell and the proximity of other ions [16]. The relaxation rate is determined from a combination of the strength of the electric field gradient (EFG) at the nucleus quantified by means of the quadrupolar

coupling constant (QCC) C_Q and the characteristic correlation time τ_c with which the memory of fluctuations is lost. While the knowledge of C_Q and τ_c can potentially provide information about the hydration sphere structure [17, 18] and useful dynamic properties (e.g., diffusion coefficients, viscosity, or conductivity [19–21]), respectively, their unambiguous determination from the experimentally-measured rates in solution has remained essentially impossible [22].

Different models have been suggested to rationalize quadrupolar relaxation using dielectric description [23, 24], mode-coupling analysis [25], definite molecular processes (e.g., water reorientation [26–28] and collective symmetry-breaking fluctuations [29]), or Brownian rotational diffusion [19–21]. *Ab initio* [30–34] and classical [29, 35–42] molecular dynamics (MD) simulations have become indispensable tools in assessing predictions of such theories, and invalidated the isotropic monoexponential character of the quadrupolar relaxation that is often assumed under a continuous solvent description. A pronounced role of intermolecular cross-correlations on the relaxation was emphasized [17, 29, 38]. Contrary to classical MD that relies on approximations [39, 42–44], *ab initio* methods provide the best accuracy of the computed EFG at the ion position that incorporates the electron cloud contribution [30, 32, 45–47]. However, the associated high computational cost often impedes the long-time sampling of EFG fluctuations [30, 31] and, hence, the ac-

curacy of the correlation time estimates, even for aqueous ions at infinite dilution [30, 32]. Therefore, the fully first-principles approach may not be appropriate in concentrated electrolytes, for which the quadrupolar NMR relaxation slows down with increasing salt concentration [26–28, 48, 49].

Here we show that C_Q parametrized at the ab initio level combined with τ_c evaluated classically allows reaching good agreement between the calculated and experimentally-obtained quadrupolar rates of $^{23}\text{Na}^+$ in electrolyte solutions at multiple salt concentrations and temperatures. We find that the main effect of increased relaxivity is due to a lengthening of the correlation times, rather than a change of the average quadrupolar coupling constant. Contrary to the commonly-assumed picture, we conclude that rotational models based on the water dipole reorientation or Stokes-Einstein-Debye relation significantly overestimate the EFG correlation times. Rather, our results indicate that the EFG relaxation occurs over a time scale comparable to that of solution structural rearrangements, suggesting that the short-time collective dynamics of the liquid primarily drives the quadrupolar relaxation at the sodium ion position.

RESULTS

Electron cloud contribution to electric field gradients

We perform density functional theory (DFT) calculations to determine quantum EFGs at the Na^+ position in aqueous sodium chloride (NaCl) solutions at varying salt concentrations $c = 1\text{--}5$ molal (denoted with $\text{mol}\cdot\text{kg}^{-1}$ or m) at $T = 25^\circ\text{C}$ (see Methods for details). The projector-augmented wave (PAW) method [45, 47, 50] is used to reconstruct the all-electron charge density at the nucleus. A configuration of a NaCl solution at $5\text{ mol}\cdot\text{kg}^{-1}$ with converged charge densities is highlighted in Fig. 1a.

In classical MD, the electron cloud contribution to the EFG can be incorporated by means of the Sternheimer approximation [43, 44], in which the full EFG at the nucleus \mathbf{V} is proportional to the EFG created by the external charge distribution \mathbf{V}_{ext} : $\mathbf{V} \simeq (1 + \gamma)\mathbf{V}_{\text{ext}}$. Here, the electron cloud polarization is included via the simple rescaling factor $1 + \gamma$, with the Sternheimer (anti)shielding factor γ being typically large $\gamma \gg 1$ [44]. To validate the Sternheimer approximation for present systems, we have compared ab initio, \mathbf{V}_{AI} , against classical, \mathbf{V}_{ext} , EFGs at the Na^+ position, as determined on the same set of classically-generated solution configurations (see Methods). Consistently with aqueous ions at infinite dilution [39, 42], we find a strong correlation between \mathbf{V}_{AI} and \mathbf{V}_{ext} for all $c = 1\text{--}5\text{ mol}\cdot\text{kg}^{-1}$, as seen in Fig. 1b for the two extreme cases. The latter allows us to define effective Sternheimer factors γ_{eff} through the linear fit $V_{\alpha\beta}^{\text{AI}} = (1 + \gamma_{\text{eff}})V_{\alpha\beta}^{\text{ext}}$. As seen in Fig. 1c, the resulting




FIG. 1. Electron cloud contribution to the EFG at the Na^+ position. **a** Representative system configuration of a NaCl solution at 5 m. The gray opaque regions around water molecules show charge densities obtained with DFT PAW calculations (see Methods). **b** Component-wise comparison of ab initio EFGs, $V_{\alpha\beta}^{\text{AI}}$, against classical EFGs, $V_{\alpha\beta}^{\text{ext}}$, at the position of Na^+ ions on the same set of configurations for different salt concentrations c . The solid lines indicate the fit for an effective Sternheimer factor γ_{eff} : $V_{\alpha\beta}^{\text{AI}} = (1 + \gamma_{\text{eff}})V_{\alpha\beta}^{\text{ext}}$. **c** Effective Sternheimer factors for Na^+ obtained from the linear fit (yellow squares) or from the ratio $(1 + \gamma'_{\text{eff}})^2 = \langle V_{\text{AI}}^2 \rangle / \langle V_{\text{ext}}^2 \rangle$ (blue circles) at different c . **d** EFG variance at the Na^+ position for different c as obtained directly with ab initio calculations (green triangles), or using the value of γ_{eff} (yellow squares) and γ'_{eff} (blue circles) at infinite dilution. The error bars in (c) and (d) were calculated using bootstrapping.

γ_{eff} feature a small increase with c (less than 5% compared to the infinite dilution value $\gamma_{\text{eff},0} = 10.54 \pm 0.11$ [42]) associated with the modifications of the ion's solvation sphere (see Supplementary Note IIB and IIC).

Despite the small changes of γ_{eff} with increasing c , the Sternheimer approximation for the EFG variance, $(1 + \gamma_{\text{eff}})^2 \langle V_{\text{ext}}^2 \rangle$, that is necessary for the NMR rate computation (Eq. (1) in Methods) underestimates the ab initio value $\langle V_{\text{AI}}^2 \rangle$ by more than 20% (highlighted in Fig. 1d using $\gamma_{\text{eff},0}$). This again underlines the deficiencies of the Sternheimer approximation [42] that does not take into account non-electrostatic electron cloud polarization effects, such as short-range repulsion [18, 51, 52]. To improve upon the predictions, we formally define the Sternheimer factor γ'_{eff} as $(1 + \gamma'_{\text{eff}})^2 = \langle V_{\text{AI}}^2 \rangle / \langle V_{\text{ext}}^2 \rangle$ with state-dependent values of $\langle V_{\text{AI}}^2 \rangle$ and $\langle V_{\text{ext}}^2 \rangle$. Sim-

ilarly, γ'_{eff} slowly grows with c , yet starting from a markedly enhanced value of $\gamma'_{\text{eff},0} = 12.09 \pm 0.14$ at infinite dilution (Fig. 1c). The EFG variance prediction $(1 + \gamma'_{\text{eff},0})^2 \langle \mathbf{V}_{\text{ext}}^2 \rangle$ using $\gamma'_{\text{eff},0}$ at infinite dilution is within 5% accuracy of $\langle \mathbf{V}_{\text{AI}}^2 \rangle$ within the considered concentration range, a much better estimate in comparison to the simple Sternheimer approximation (Fig. 1d). While not capturing all condensed-phase effects that arise with increasing c , the estimate $(1 + \gamma'_{\text{eff},0})^2 \langle \mathbf{V}_{\text{ext}}^2 \rangle$ provides a fair accuracy, reproduces the trend of $\langle \mathbf{V}_{\text{AI}}^2 \rangle$ to decrease with the salt concentration (see Fig. 1d), and permits to avoid computationally expensive DFT calculations at multiple system state points of interest. As discussed below, in combination with the EFG relaxation dynamics captured at the classical level, this approach provides a good description of the quadrupolar $^{23}\text{Na}^+$ NMR rates in aqueous solutions.

Relaxation of electric field gradient fluctuations

We perform classical MD simulations employing the Madrid-2019 force field (FF) [53, 54] to facilitate the long-time sampling of EFG fluctuations and to investigate the mechanisms behind the concentration and temperature behavior of the quadrupolar $^{23}\text{Na}^+$ NMR rate in aqueous sodium chloride, bromide (NaBr), and fluoride (NaF) solutions (see Methods for simulation details). Two facts give confidence in this approach: (i) a very strong correlation between the full and classical (external) EFGs (Fig. 1b), indicating that the dynamics of the former should be largely determined by that of the latter; (ii) while classical MD with rigid water molecules do not quantitatively reproduce the librational or hydrogen-bond stretching water dynamics that occur at very short times below ~ 50 fs [55], it is expected that these high frequency motions do not significantly affect the dominating long-time (~ 1 ps) EFG relaxation mode (e.g., see Ref. [40] and below).

Increasing salt concentration c or decreasing temperature T causes a profound slow-down of the EFG fluctuations at the ion position (Fig. 2). Due to a qualitative similarity of the EFG relaxation in the solutions considered, here we will focus on the case of NaCl; see Supplementary Information (SI) for NaBr and NaF. Fig. 2a shows the ACFs of the classical EFG at the Na^+ position, $C_{\text{EFG}}(t) \equiv \langle \mathbf{V}_{\text{ext}}(0) \cdot \mathbf{V}_{\text{ext}}(t) \rangle$, as a function of c at $T = 25^\circ\text{C}$ (see Supplementary Fig. S8 for other T). Similarly to a single Na^+ in water [32, 38, 39, 42], $C_{\text{EFG}}(t)$ relaxes in two steps: (i) a rapid initial decay happening at $t \lesssim 0.2$ ps that corresponds to $\approx 70\%$ of the EFG decorrelation. This is in good agreement with the ab initio MD results for Na^+ at infinite dilution [32], highlighting the validity of the classical approach; (ii) a much slower secondary decay occurring in the picosecond regime. As seen in Fig. 2a, the increase in c leaves the initial fast de-




FIG. 2. Relaxation of EFG fluctuations. **a** Normalized autocorrelation functions $C_{\text{EFG}}(t)/C_{\text{EFG}}(0)$ of the EFG at the position of a Na^+ ion obtained using classical MD simulations for different salt concentrations c at $T = 25^\circ\text{C}$ in aqueous NaCl solutions. Qualitatively similar trends are found for other concentrations and temperatures (see Supplementary Fig. S8). Insets in (a) highlight the short-time behavior of the ACFs for $t < 1$ ps and a typical system configuration at $c = 4$ m (Na^+ and Cl^- ions are blue and green, respectively). **b** Temperature behavior of $C_{\text{EFG}}(t)$ at $c = 4$ m. **c** Long-time behavior of $C_{\text{EFG}}(t)$ plotted on a double logarithmic scale at $c = 4$ m for different temperatures (the legend shown in (b)). The black solid and dashed lines highlight a power-law scaling $\sim t^\alpha$ with $\alpha = -5/2$ and $\alpha = -3/2$, respectively. Shaded regions in (a), (b), and (c) indicate standard errors from multiple independent simulation runs.

cay practically unchanged, while causing a pronounced slow-down of the second decay mode. The latter is highlighted in the inset of Fig. 2a showing the EFG ACFs for $t < 1$ ps for different c at $T = 25^\circ\text{C}$ (see also Supplementary Fig. S9). A qualitatively similar trend is found with decreasing temperature, as we show in Fig. 2b at $c = 4$ m and in Supplementary Fig. S8 for other c .

The form of the EFG ACF decay in Fig. 2 suggests a collective pathway behind the relaxation. After the initial fast decay that can be described with an exponential $\sim e^{-t/\tau_f}$ with $\tau_f \approx 62$ fs, we find a development of a much slower relaxation mode that profoundly depends on c and T . Compared to earlier results [40], our long-

time sampling reveals that the slow part of the EFG ACF is not exponential, as clearly seen from the behavior of $C_{\text{EFG}}(t)$ on a semi-logarithmic scale in Figs. 2a-b and as we show with explicit fits in Supplementary Note IIE. Except at very long times, we find that the slow decay can be modeled either with a two-exponential (Supplementary Fig. S10) or a stretched exponential fit $\sim e^{-(t/\tau_s)^\beta}$ with $\beta = 0.67 \pm 0.05$ (Supplementary Fig. S11 and S12), which suggests a broad distribution of contributing relaxation modes (Supplementary Fig. S13). Although observed over a limited time range (up to a decade), we find that the long-time tail of the EFG ACFs is consistent with a power law $\sim t^{-5/2}$, as shown with $C_{\text{EFG}}(t)$ on a log-log scale for $c = 4$ m in Fig. 2c. Such a hydrodynamic tail was predicted by a mode-coupling theory of Bosse *et al.* for the EFG ACF in molten salts [25]. It originates from the coupling between the ion motion and shear excitations in the liquid, a mechanism causing the well-known $\sim t^{-3/2}$ tail of the velocity ACF [56]. While sampling of the EFG fluctuations at even longer time scales is necessary to decisively confirm to presence of $\sim t^{-5/2}$ regime, our results for Na^+ in Fig. 2c suggest that its relative contribution may be marginal because the apparent onset of the algebraic decay occurs at times when the ACF has decayed considerably.

Quadrupolar relaxation rates

The combination of EFG fluctuations captured at the classical level and a consistent inclusion of the electron cloud contribution to the EFG enables reaching a good quantitative agreement between the calculated and experimentally-measured quadrupolar NMR rate for $^{23}\text{Na}^+$ in aqueous NaCl, as we compare in Fig. 3 with filled and open symbols, respectively. As seen in Eq. (1) in Methods, the quadrupolar rate is proportional to the product of the effective correlation time of EFG fluctuations, $\tau_c = C_{\text{EFG}}^{-1}(0) \int_0^\infty dt C_{\text{EFG}}(t)$, and the EFG variance, which we approximate as $\langle \mathbf{V}^2 \rangle = (1 + \gamma'_{\text{eff},0})^2 \langle \mathbf{V}_{\text{ext}}^2 \rangle$ with $\gamma'_{\text{eff},0} = 12.09$ and $\langle \mathbf{V}_{\text{ext}}^2 \rangle = C_{\text{EFG}}(0)$. The integration of $C_{\text{EFG}}(t)$ over tens of picoseconds is necessary to obtain well-converged correlation times τ_c (Supplementary Fig. S14), notably at high salt concentrations and low temperatures (Fig. 2). Finally, based on estimates in Supplementary Note IC, here we assume that the $^{23}\text{Na}^+$ rate $1/T_1$ is almost entirely determined by the quadrupolar interaction, whereas possible contributions from the dipole-dipole coupling between the spin of the ^{23}Na nucleus and that of ^1H , ^{23}Na , ^{35}Cl are marginal.

The NMR rate $1/T_1$ grows with increasing salt concentration c and with reducing temperature T (Fig. 3), suggesting that the slowing down of EFG fluctuations determines the rate behavior (Fig. 2). Experimentally, $1/T_1$ rises by about 50% within the considered range of concentrations $c = 0.17$ – 5.1 m for temperatures $T = 20$ –




FIG. 3. **Concentration and temperature dependence of the quadrupolar relaxation rate.** T_1^{-1} of $^{23}\text{Na}^+$ in aqueous NaCl as a function of the salt concentration c as obtained in experiments (E, solid lines and open symbols) and simulations (S, filled symbols) for different temperatures. The error bars for simulation results are associated with the approximation for incorporating the electron cloud contribution to the EFG.

50 °C, in line with the previous results [22, 28, 48]. At $T = 30$ °C, $1/T_1$ increases from around 15.9 s $^{-1}$ at $c = 0.17$ m to 25.2 s $^{-1}$ at $c = 5.1$ m. With increasing T from 25 to 50 °C, $1/T_1$ reduces by more than 25% for considered salt concentrations. In general, our computational results for $1/T_1$ of $^{23}\text{Na}^+$ agree well with the experimental data, especially at lower salt concentrations $c \lesssim 2.5$ m, reproducing both the concentration and temperature behavior. For higher salt concentrations, we find that $1/T_1$ grows systematically faster with increasing c as compared to the experiments, yet the relative error remains less than 15% over the considered range of conditions. The latter difference is likely caused by the shortcomings of the employed FF in capturing dynamic properties of the solution for $c \gtrsim 2$ m [54].

Microscopic parameters of the relaxation

We find that the slowing down of EFG fluctuations at the Na^+ position primarily causes a marked increase in the quadrupolar NMR relaxation rate with increasing c and decreasing T (Fig. 3). In Fig. 4, we quantify the role of dynamic and static effects that are reflected in the changes of τ_c and $\langle \mathbf{V}^2 \rangle$, respectively, with varying salt concentration and temperature, as obtained in MD simulations of aqueous NaCl (see Supplementary Fig. S18 for other electrolyte solutions). While τ_c increases by a factor of ~ 1.5 – 2.5 with increasing c and decreasing T within the considered range of parameters (Fig. 4a), the value of

$\langle \mathbf{V}^2 \rangle$ reduces concurrently by up to 10% (Fig. 4c), indicating that the augmented correlation times are mainly responsible for the rate behavior.

For considered c and T , τ_c of Na^+ is quite short and below 1 ps (Fig. 4a), a feature already pointed out in previous classical [35, 38, 39, 42] and ab initio [30, 32] MD studies at infinite dilution. At $T = 25^\circ\text{C}$, we find that τ_c increases from 0.41 ps at $c \approx 0.06$ m to 0.65 ps at $c = 4$ m. Despite the rapid decorrelation of EFG ACFs for $t \lesssim 0.2$ ps (Fig. 2), we find that the contribution of the slow relaxation process to τ_c yields more than 85% of its overall value and also grows with increasing c and decreasing T (Fig. 4b). The dominance of the slow non-exponential decay of EFG ACFs over the τ_c behavior again exemplifies the governing role of collective processes behind the quadrupolar Na^+ relaxation.

While the EFG variance at the Na^+ position is largely determined by the first solvation shell contribution (Fig. 4d), a quantitative understanding of the QCC is only achieved if we take into account point charges within a radius of $r \gtrsim 8$ Å around the central ion, approximately the length scale of pronounced ion-ion and ion-solvent correlations (Supplementary Figs. S3 and S4). We find that $\langle \mathbf{V}^2 \rangle$ is reduced in bipyramidal complexes with octahedral symmetry, coordinated by six water molecules, yet only by 10% compared to the ensemble average (Supplementary Note IID). Contrary to the full EFG variance, the contribution of the first solvation shell features an increase with c (Fig. 4d), correlated with the fact that the six-coordinated state becomes less likely with increasing the salt concentration (Supplementary Fig. S7). Our consistently calculated QCC for $^{23}\text{Na}^+$ in aqueous NaCl is in the range between $19 \cdot 10^6$ and $20.6 \cdot 10^6$ $\text{rad} \cdot \text{s}^{-1}$ for considered c and T (Supplementary Fig. S15), a value approximately 3–4 times larger than previous estimates based on the assumption that the EFG primarily decorrelates by translational and reorientational water dynamics with $\tau_c \approx 3$ –7 ps [19–22, 57]. We thus conclude that the aforementioned modes of motion provide only a minor contribution to the observed relaxation.

Assessment of the relaxation models

We utilize information available in experiments and molecular simulations in Fig. 5 to shed light on the mechanisms behind the quadrupolar relaxation. First, we focus on the possibility to model the EFG correlation time τ_c using the commonly-used Stokes-Einstein-Debye (SED) relation $\tau_c^{\text{SED}} = 4\pi\eta r_0^3 / 3k_B T$, where η is the dynamic viscosity of the solution, r_0 is the sodium's hydrodynamic (Stokes) radius, and k_B is the Boltzmann constant. Within the SED picture, the EFG relaxation at the Na^+ position is governed by the Brownian rotational diffusion, likely to be related with collective reorientations of ion-water solvation complexes [48]. While the




FIG. 4. **Microscopic parameters of the EFG relaxation.** **a** Effective correlation time τ_c of the EFG fluctuations at the Na^+ position as a function of salt concentration c for different temperatures T . **b** Relative contribution of the slow EFG relaxation mode to τ_c for different c and T . The contribution was estimated using the stretched exponential fit of the normalized EFG ACFs (Supplementary Note II E). **c** Variance of the total EFG at the ion position $\langle \mathbf{V}^2 \rangle = (1 + \gamma'_{\text{eff},0})^2 \langle \mathbf{V}_{\text{ext}}^2 \rangle$ as a function of c for different temperatures. The legend in (c) is the same as in (a). **d** EFG variance, $\langle \mathbf{V}^2 \rangle$, evaluated from water molecules and ions located within a different number of solvation shells around the central Na^+ ion as a function of c at $T = 25^\circ\text{C}$. The error bars from multiple independent simulation runs are either explicitly shown or do not exceed the symbol size.

SED model assumptions are not expected to hold down to the molecular scale [59, 60], we systematically explore τ_c^{SED} in relation to τ_c , as it is often exploited to rationalize quadrupolar relaxation dynamics of $^{23}\text{Na}^+$ [19–21].

We use the translational Stokes-Einstein relation $D = k_B T / 6\pi\eta r_0$ to determine the concentration- and temperature-dependent values of the Stokes radius from the experimental Na^+ diffusion coefficients (Fig. 5a) and highly accurate NaCl viscosity values provided by Kestin *et al.* [58] (Fig. 5b). D and η calculated in our MD simulations (see Methods) are in good agreement with the experiments, especially for $c \lesssim 2$ m, capturing both the concentration and temperature behavior (compare filled and open symbols in Figs. 5a–b). The viscosity η in MD was obtained via the Green-Kubo formula using the stress ACFs (Fig. 5c), as detailed in Eqs. (3) and (4) in




FIG. 5. Assessing models of the quadrupolar relaxation. **a** Na^+ diffusion coefficient as a function of salt concentration c for different temperatures T in experiments (solid lines with open symbols, legend shown in (a)) and simulations (filled symbols, legend shown in (b)). **b** Dynamic solution viscosity as a function of c for different T in experiments (solid lines with open symbols, legend shown in (a)) and simulations (filled symbols, legend shown in (b)). Experimental viscosities were taken from Ref. [58]. **c** Stress tensor ACFs $C_{\text{stress}}(t)$ normalized by the system volume V and $k_B T$ for increasing C at $T = 25^\circ\text{C}$. The black dotted line shows the stretched exponential fit of the long-time decay at $c = 4$ m. Inset: the time scale of solution structural relaxation τ_{struct} , as extracted from the long-time decay of $C_{\text{stress}}(t)$ (see main text), as a function of c at multiple temperatures. T decreases from top to bottom, the legend is shown in (f) and (c). **d**, Stokes-Einstein-Debye time plotted versus the EFG correlation time τ_c^{exp} , as extracted from experimental data for different c and T . **e** τ_c^{exp} as a function of $\eta/k_B T$ for different temperatures. r_0^{eff} is the effective hydrodynamic radius of a Na^+ ion extracted under assumption that τ_c^{exp} can be modeled by a SED relation (the gray line shows the best fit). **d** and **e** share the same legend. **f** Mean water dipole reorientation time τ_{dip} plotted versus the EFG correlation time τ_c , as extracted in simulations for different c and T . **g** τ_{struct} plotted versus τ_c for different c and T . The gray line indicates the linear dependence, $\tau_{\text{struct}} = \tau_c$.

Methods. In Fig. 5d, we compare τ_c^{SED} calculated from state-dependent Stokes radii $r_0(c, T)$ against the effective EFG correlation time τ_c^{exp} obtained from the experimental NMR rates and the sodium's QCC from simulations (Supplementary Fig. S15) rather than those from previous estimates [19–22, 57]. r_0 assumes values between 1.5 and 2.0 Å for considered parameters (Supplementary Fig. S17) While both τ_c^{SED} and τ_c^{exp} generally lengthen with increasing c and decreasing T , τ_c^{SED} exceeds τ_c^{exp} by a factor of 8–17 (Fig. 5d). Similar results are obtained in our simulations (Supplementary Fig. S17). At $T = 25^\circ\text{C}$, τ_c^{SED} increases from around 7.5 to 8.2 ps, larger by more than one order of magnitude than τ_c^{exp} that grows from 0.44 to 0.66 ps for increasing c from 0.17 to 5.1 m. Thus, we conclude that the EFG correlation times cannot be understood on the basis of the SED relation parameterized using the translational hydrodynamic

radius of sodium ions $r_0 = k_B T / 6\pi\eta D$.

This is further illustrated in Fig. 5e showing τ_c^{exp} plotted against $\eta/k_B T$ for various temperatures. While a Stokes-Einstein-like relation holds for τ_c^{exp} , that is a strong correlation $\tau_c^{\text{exp}} \propto \eta/k_B T$ exists for the considered range of parameters, the effective Stokes radius $r_0^{\text{eff}} = 0.69$ Å that would correspond to the EFG correlation time τ_c^{exp} within the SED model is clearly unphysical and smaller than the ionic radius 1.02 Å. r_0^{eff} was obtained from the fit $\tau_c^{\text{exp}} = 4\pi\eta [r_0^{\text{eff}}]^3 / 3k_B T + \tau_0^{\text{eff}}$ with an additional intercept $\tau_0^{\text{eff}} = 0.11$ ps needed for the best data representation [60]. We obtain a similar value of $r_0^{\text{eff}} \approx 0.68$ Å from our MD simulations (Supplementary Fig. S17). Therefore, the validity of the relation $\tau_c^{\text{exp}} \propto \eta/k_B T$ explains the correlation between τ_c^{exp} and D^{-1} reported in Refs. [19–21], rather than simplified assumptions of the rotational Brownian diffusion that yield

much larger estimates of τ_c^{exp} (Fig. 5d).

We now return to microscopic time scales of molecular motion in relation to that of EFG fluctuations. The average water dipole reorientation time $\tau_{\text{dip}} = \int_0^\infty dt \langle P_1[\mathbf{u}(t) \cdot \mathbf{u}(0)] \rangle$ assumed to drive the quadrupolar relaxation within the Hertz model [17, 26, 27, 38] is 11-14 times larger compared to τ_c , as extracted in our simulations (Fig. 5f and Supplementary Fig. S16). Above, \mathbf{u} is a unit vector pointing along the HOH bisector of a water molecule and $P_1(x) = x$ is the first Legendre polynomial. This indicates that the single molecule reorientation with neglected intermolecule cross-correlations cannot explain the EFG relaxation dynamics. Yet, as seen in Fig. 5f, both τ_{dip} and τ_c increase similarly with increasing c and decreasing T , suggesting that the overall deceleration of the electrolyte dynamics, marked by an enhanced viscosity, impacts in a similar way both the motions that drive water reorientation as well as those that cause the EFG relaxation at the ion position.

To illustrate the relationship between these effects, in our MD simulations we extract a typical time scale of solution structural relaxation τ_{struct} using the stress tensor ACFs (Fig. 5c). While the short-time behavior of $C_{\text{stress}}(t)$ corresponding to elastic, vibrational contributions features little changes with varying c and T [61], its long-time tail slows down with increasing c and decreasing T , indicating an overall deceleration of the viscous dynamics of the liquid. We find that the long-time tail can be modeled well using a stretched exponential decay, $\sim e^{-(t/\tau_K)^{\beta_K}}$, with $\beta_K \approx 0.61 \pm 0.04$ (consistent with earlier simulations of pure water [61] and time-resolved spectroscopy experiments [62]). τ_K is in the range between 0.13 and 0.70 ps for considered parameters. The mean structural relaxation time $\tau_{\text{struct}} = \tau_K \beta_K^{-1} \Gamma(\beta_K^{-1})$, defined through the integral of the stretched exponential expression, is strongly correlated and comparable to the subpicosecond EFG correlation time τ_c (Fig. 5g). While the stress and EFG tensors are not directly related to each other, both quantities are inherently collective, the relaxation of their fluctuations is mainly driven by many-body correlations and features a similar stretched decay for $t \gtrsim 0.4$ ps. All these observations suggest that the fast collective dynamics of the liquid that drive its structural rearrangements are also responsible for the quadrupolar NMR relaxation.

CONCLUSIONS AND PERSPECTIVES

We have shown that the multiscale methodology combining DFT PAW calculations to parameterize the QCC and classical MD simulations to sample long-time EFG fluctuations enables an accurate description of the concentration and temperature behavior of the quadrupolar NMR relaxation rates of $^{23}\text{Na}^+$ in aqueous electrolyte solutions. The resulting NMR rates are in very good

agreement with the experimental data, especially at low salt concentrations, as validated in aqueous NaCl at multiple concentrations and temperatures. We find that the growth of the relaxation rate T_1^{-1} with increasing c and decreasing T is primarily due to the slowing down of the EFG fluctuations reflected in the augmented EFG correlation time τ_c , while the concurrent changes in the QCC are rather small. We have demonstrated that the commonly-assumed rotational relaxation models based on either the water dipole reorientation [26–28] or on the Stokes-Einstein-Debye relation [19–21] overestimate the consistently-determined τ_c by at least an order of magnitude. This disagreement is understandable as these models restrict the relaxation description to one- or two-body correlations, oversimplifying the inherently collective character of the EFG at the ion position [17, 29, 38]. The quantitative interpretation of the EFG correlation times in terms of such simple isotropic models should therefore be used with caution. Instead, our results indicate that the Na^+ EFG relaxation occurs over a subpicosecond time scale comparable to that of solution structural rearrangements τ_{struct} . This again invalidates a continuous-solvent hydrodynamic description assuming that $\tau_c \gg \tau_{\text{struct}}$.

Our results suggest that the quadrupolar NMR relaxometry of $^{23}\text{Na}^+$ may be used as a complementary tool to analyze electrolyte dynamics in the THz domain. As the quadrupolar relaxation is largely determined by the processes in the immediate vicinity of the solute, it can provide supplementary information on the fast, collective, molecular motions in ionic solvation cages. The ability to capture the NMR rates by means of classical MD allows elucidating the quadrupolar phenomena that occur in multicomponent systems, such as concentrated aqueous solutions of multiple salts [63], mixtures of salts with glycerol [64], or polyelectrolytes [65], where the relaxation dynamics may be influenced by environment heterogeneity, interface formation, microphase separation, or ion binding to polyelectrolyte chains. Future work could also focus on developing mesoscopic approaches that would allow a quantitative description of quadrupolar relaxation in complex biological-type compartments, characterized either by slow-motion conditions with dynamics within intracellular and extracellular spaces in biological tissues that may include structural anisotropy with residual quadrupolar coupling and a distribution of correlation times (as, for example, in articular cartilage where sodium ions are surrounded by a collagen matrix [4]).

METHODS

Quadrupolar NMR relaxation rates

The quadrupolar mechanism dominates the relaxation of nuclei with spin $I > 1/2$ and is due to the coupling between their quadrupolar moment eQ with the EFG tensor \mathbf{V} at the nucleus position [16]. While the NMR relaxation of spin components is generally bi-exponential for ^{23}Na with $I = 3/2$ [22, 66], it is possible to define effective longitudinal and transverse quadrupolar relaxation rates, $1/T_1$ and $1/T_2$, respectively, provided that the ‘‘fast motion’’ or ‘‘extreme narrowing’’ regime holds [30, 67, 68]. In this case, the characteristic EFG correlation time τ_c is much smaller than the Larmor period ω_0^{-1} of the nucleus, $\omega_0\tau_c \ll 1$. The latter can be shown to be fulfilled for all cases considered in this work as the relevant correlation times for ^{23}Na in electrolyte solutions are below 100 ps, and the magnetic field used in the experiments is 11.7 T that corresponds to $\omega_0^{-1} \approx 7.6$ ns. As we show in more detail in the SI, the two quadrupolar relaxation rates become equal in the fast motion regime and, combined with the rotational invariance of the system, can be expressed as [38]

$$\frac{1}{T_1} = \frac{2I + 3}{20I^2(2I - 1)} \left(\frac{eQ}{\hbar} \right)^2 \langle \mathbf{V}^2 \rangle \tau_c \quad (1)$$

where \hbar is the reduced Planck constant, τ_c is an effective correlation time of EFG fluctuations

$$\tau_c = \langle \mathbf{V}^2 \rangle^{-1} \int_0^\infty dt \langle \mathbf{V}(0) : \mathbf{V}(t) \rangle, \quad (2)$$

where $\langle \mathbf{V}(0) : \mathbf{V}(t) \rangle = \sum_{\alpha, \beta} \langle V_{\alpha\beta}(0) V_{\alpha\beta}(t) \rangle$ with $\alpha, \beta = x, y, z$ and the brackets $\langle \dots \rangle$ denoting an ensemble average, and $\langle \mathbf{V}^2 \rangle \equiv \langle \mathbf{V}(0) : \mathbf{V}(0) \rangle$ is the EFG variance at the ion position. For a ^{23}Na nucleus with $I = 3/2$ and $Q = 104 \cdot 10^{-31} \text{ m}^2$ [69], the rate constant $1/T_1$ can be recast as $1/T_1 = C_Q^2 \tau_c / 10$ with the quadrupolar coupling constant (QCC) defined as $C_Q^2 = \frac{2}{3} \left(\frac{eQ}{\hbar} \right)^2 \langle \mathbf{V}^2 \rangle$ [16, 30]. Finally, Eq. (1), which follows from linear response theory, allows to calculate the quadrupolar spin-lattice relaxation rate $1/T_1$ from the EFG fluctuations in equilibrium MD simulations without an imposed magnetic field.

NMR experiments

Solution samples with 9 different NaCl concentrations were prepared by mixing x mg of NaCl in $(y-x)$ mg of deionized water in a beaker, with $x = 0.1, 0.2, 0.5, 0.8, 1.1, 1.4, 1.7, 2.0, 2.3$ mg and $y = 10$ mg, to make solutions of concentrations 0.173, 0.349, 0.901, 1.488, 2.115, 2.786, 3.505, 4.278, and 5.111 mol \cdot kg $^{-1}$ to 5 mm NMR tubes (sample volume = 0.5 mL). All mass measurements were

performed on a Mettler Toledo ME204E balance with a resolution of 0.1 mg. The solution at 26% weight corresponds to NaCl saturation in water at 20 °C [70].

NMR experiments were performed on an 11.7 T NMR Bruker Avance I spectrometer operating at 132.3 MHz for ^{23}Na , using a 5 mm double resonance broadband probe. The test tubes with the solutions were placed inside the spectrometer where the sample temperature could be controlled using gas flow and a temperature sensor providing a precise, stable and reliable temperature regulation. After each desired temperature reached steady state, a standard free induction decay was acquired followed by a longitudinal relaxation time T_1 mapping sequence, and a diffusion pulse sequence. At each temperature, the tuning and matching was checked. The duration of the 90° pulse was 9.6 μs , whereas that for the 180° pulse was 19.6 μs . A standard inversion-recovery pulse sequence was used to acquire T_1 with 32 logarithmically spaced steps. The delay was varied from 1 ms to 400 ms for ^{23}Na . Diffusion coefficients were measured using a Pulsed-Gradient-Spin-Echo in 32 steps with a maximum b -value of 2200 s \cdot mm $^{-2}$. The maximum diffusion gradient was 1 T \cdot m $^{-1}$ and the duration was 4 ms.

Molecular dynamics simulations

Aqueous sodium chloride (NaCl), bromide (NaBr), and fluoride (NaF) solutions were simulated using classical MD employing the Madrid-2019 FF [53] that is based on the TIP4P/2005 water model [71] and uses scaled charges of $+0.85e$ and $-0.85e$ (e is the fundamental unit of charge) for Na^+ cations and Cl^- , Br^- , and F^- anions, respectively. The FF parameters are listed in the SI. The scaled ionic charges aim at taking into account the electronic contribution to the dielectric constant at high frequencies in a mean-field fashion [72]. At a moderate computational cost in comparison to fully polarizable models, the EFG relaxation within the Madrid-2019 FF [53] has recently been shown to accurately describe the quadrupolar NMR relaxation rates of alkali metal ions at infinite dilution [42], in particular that of Na^+ . Solutions comprised of $N = 1000$ water molecules and N_p ion pairs were initialized at different salt concentrations c between 0.06 m ($N_p = 1$) and 4 m ($N_p = 72$) in a cubic box at the equilibrium solution density $\rho(c, T)$ obtained in NPT simulations at $P = 1$ bar. The densities are in excellent agreement with the experimental ones, as discussed in Supplementary Note IB.

The equilibrated electrolyte systems were then simulated in the NVT ensemble. Both NPT and NVT simulation runs were carried out in the open-source MetalWalls package on graphics processing units [73] with electrostatic interactions computed with Ewald summation [74] and a short-range cutoff of 1.24 nm. The constant temperature was maintained using the Nose-Hoover

chains thermostat with a time constant of 1 ps. System temperatures in range from 10 °C to 50 °C were considered. The equations of motion were integrated using the velocity Verlet algorithm and an integration time step of 1 fs. The effective rigidity of water molecules was imposed with the help of the RATTLE algorithm with a precision of 10^{-9} . For each (c, T) state point, at least five independent runs of length 5 ns were performed to measure the EFG at the ion positions (sampled every 50 fs). Full Ewald summation expressions [74] were used in the computation of the EFGs, as recently implemented in MetalWalls [42]. For the considered system parameters, the relaxation of EFG fluctuations was found not to be affected by the finite box size, as we discuss in Supplementary Note. II A.

Ab initio calculations

To determine EFGs with the electron cloud contribution, smaller systems containing 55 water molecules and $N_p = 1, 2, 3, 4,$ and 5 NaCl ions pairs, corresponding to the salt concentrations $c = 1, 2, 3, 4, 5$ mol·kg $^{-1}$, were simulated in the same way as the larger ones using the Madrid-2019 FF. In a single NVT simulation run at $T = 25$ °C, 2000 configurations were sampled with a period of 10 ps, and were later used in DFT-based EFG calculations with periodic boundary conditions in the Quantum Espresso (QE) package [75]. No additional geometry optimization of the configurations was performed in the DFT calculations. The pseudopotential-based projector-augmented wave (PAW) method [45, 47, 50] was used to reconstruct the all-electron charge density in the vicinity of the nucleus using the QE-GIPAW package [76]. The self-consistent electron densities were calculated using the PBE functional [77], a kinetic energy cutoff of 80 Ry, and norm-conserving pseudopotentials of the GIPAW package [78]. In the case of Na $^+$ ions, the EFGs obtained with the PBE functional were shown to be in good agreement [32] with those obtained with the hybrid PBE0 functional [79].

Dynamical properties of electrolyte solutions

The shear viscosity of aqueous electrolyte solutions was obtained using the Green-Kubo relation [80]:

$$\eta = \frac{V}{k_B T} \int_0^{+\infty} dt C_{\text{stress}}(t), \quad (3)$$

with V being the system volume and k_B standing for the Boltzmann constant. The stress tensor ACF $C_{\text{stress}}(t)$ was computed as [80]

$$C_{\text{stress}}(t) = \frac{1}{10} \sum_{\alpha, \beta} \langle P_{\alpha\beta}(t) P_{\alpha\beta}(0) \rangle, \quad (4)$$

where α, β run over the three Cartesian components and $P_{\alpha\beta}$ is the traceless symmetrized part of the stress tensor $\sigma_{\alpha\beta}$: $P_{\alpha\beta} = \frac{1}{2}(\sigma_{\alpha\beta} + \sigma_{\beta\alpha}) - \frac{1}{3}\delta_{\alpha\beta} \sum_{\gamma} \sigma_{\gamma\gamma}$. For each salt concentration, the viscosity was measured over more than 5 independent simulation runs of length 5 ns with the stress tensor sampled every integration time step (1 fs).

The Na $^+$ diffusion coefficients were extracted from the long time limit of the ion's mean-square displacement:

$$D = \lim_{t \rightarrow \infty} \frac{1}{6N_p t} \sum_{i=1}^{N_p} \langle [\mathbf{r}_i(t) - \mathbf{r}_i(0)]^2 \rangle, \quad (5)$$

where N_p is the number of sodium ions in the system, $\mathbf{r}_i(t)$ is the position of the i -th ion at time t , and the brackets $\langle \dots \rangle$ stand for ensemble averaging. The obtained diffusion coefficients were corrected for finite-size effects using the Yeh-Hummer relation [81]:

$$D_{\infty} = D + \frac{k_B T \xi}{6\pi\eta L} \quad (6)$$

with the diffusion coefficient D_{∞} corresponding to a macroscopic system, D being obtained in a cubic simulation box with side length L , and $\xi \approx 2.837297$. The calculated values of viscosity η in Eq. (3) were used for evaluating D_{∞} in Eq. (6). The finite-size correction term corresponded to 17–22 % of the measured value D .

ACKNOWLEDGEMENTS

We would like to thank Dr. Seena Dehkharghani for the thoughtful discussions on the influence of temperature on water dynamics and NMR relaxation parameters and Dr. Antoine Carof for useful discussions on the modeling of quadrupolar NMR relaxation. This work was supported in part by the National Institutes of Health (NIH): grant no. R01EB026456. This project received funding from the European Research Council under the European Union's Horizon 2020 research and innovation program (grant agreement no. 863473). The authors acknowledge access to HPC resources from GENCI (grant no. A0110912966) and AJ wishes to acknowledge the HPC resources of NYU. AJ acknowledges funding from the US National Science Foundation under award no. CHE2108205.

-
- * Equal contribution
 † alexej.jerschow@nyu.edu
 ‡ benjamin.rotenberg@sorbonne-universite.fr
- [1] Ohtaki, H. & Radnai, T. Structure and dynamics of hydrated ions. *Chem. Rev.* **93**, 1157–1204 (1993).
 - [2] Inglese, M. *et al.* Brain tissue sodium concentration in multiple sclerosis: A sodium imaging study at 3 Tesla. *Brain* **133**, 847–857 (2010).
 - [3] Ouwerkerk, R., Bleich, K. B., Gillen, J. S., Pomper, M. G. & Bottomley, P. A. Tissue sodium concentration in human brain tumors as measured with ^{23}Na MR imaging. *Radiology* **227**, 529–537 (2003).
 - [4] Madelin, G., Lee, J.-S., Regatte, R. R. & Jerschow, A. Sodium MRI: Methods and applications. *Prog. Nucl. Magn. Reson. Spectrosc.* **79**, 14–47 (2014).
 - [5] Guermazi, A. *et al.* Compositional MRI techniques for evaluation of cartilage degeneration in osteoarthritis. *Osteoarthr. Cartil.* **23**, 1639–1653 (2015).
 - [6] Silletta, E. V., Jerschow, A., Madelin, G. & Alon, L. Multinuclear absolute magnetic resonance thermometry. *Commun. Phys.* **2**, 152 (2019).
 - [7] Hu, R. *et al.* X-nuclei imaging: Current state, technical challenges, and future directions. *J. Magn. Reson. Imaging* **51**, 355–376 (2020).
 - [8] Madelin, G. *X-Nuclei Magnetic Resonance Imaging* (CRC Press, 2022).
 - [9] Slater, M. D., Kim, D., Lee, E. & Johnson, C. S. Sodium-ion batteries. *Adv. Funct. Mater.* **23**, 947–958 (2013).
 - [10] Yabuuchi, N., Kubota, K., Dahbi, M. & Komaba, S. Research development on sodium-ion batteries. *Chem. Rev.* **114**, 11636–11682 (2014).
 - [11] Hwang, J.-Y., Myung, S.-T. & Sun, Y.-K. Sodium-ion batteries: Present and future. *Chem. Soc. Rev.* **46**, 3529–3614 (2017).
 - [12] Chandrashekar, S. *et al.* ^7Li MRI of Li batteries reveals location of microstructural lithium. *Nature Mater.* **11**, 311–315 (2012).
 - [13] Pecher, O., Carretero-González, J., Griffith, K. J. & Grey, C. P. Materials’ methods: NMR in battery research. *Chem. Mater.* **29**, 213–242 (2017).
 - [14] Wertz, J. E. & Jardetzky, O. Nuclear spin resonance of aqueous sodium ion. *J. Chem. Phys.* **25**, 357–358 (1956).
 - [15] Headley, L. C. Nuclear magnetic resonance relaxation of ^{23}Na in porous media containing NaCl solution. *J. Appl. Phys.* **44**, 3118–3121 (1973).
 - [16] Abragam, A. *The Principles of Nuclear Magnetism* (Oxford University Press, 1961).
 - [17] Versmold, H. Interaction induced magnetic relaxation of quadrupolar ionic nuclei in electrolyte solutions. *Mol. Phys.* **57**, 201–216 (1986).
 - [18] Aidas, K., Ågren, H., Kongsted, J., Laaksonen, A. & Mocchi, F. A quantum mechanics/molecular dynamics study of electric field gradient fluctuations in the liquid phase. the case of Na^+ in aqueous solution. *Phys. Chem. Chem. Phys.* **15**, 1621–1631 (2013).
 - [19] Price, W. S., Chapman, B. E. & Kuchel, P. W. Correlation of viscosity and conductance with $^{23}\text{Na}^+$ NMR T_1 measurement. *Bull. Chem. Soc. Jpn.* **63**, 2961–2965 (1990).
 - [20] Mitchell, J. Can sodium NMR provide more than a tracer for brine in petrophysics? *J. Pet. Sci. Eng.* **146**, 360–368 (2016).
 - [21] D’Agostino, C., Davis, S. J. & Abbott, A. P. ^{23}Na NMR T_1 relaxation measurements as a probe for diffusion and dynamics of sodium ions in salt–glycerol mixtures. *J. Chem. Phys.* **154**, 224501 (2021).
 - [22] Woessner, D. E. NMR relaxation of spin-3/2 nuclei: Effects of structure, order, and dynamics in aqueous heterogeneous systems. *Concepts Magn. Reson. A: Bridg. Educ. Res.* **13**, 294–325 (2001).
 - [23] Hynes, J. T. & Wolynes, P. G. A continuum theory for quadrupole relaxation of ions in solution. *J. Chem. Phys.* **75**, 395–401 (1981).
 - [24] Perng, B.-C. & Ladanyi, B. M. A dielectric theory of spin-lattice relaxation for nuclei with electric quadrupole moments. *J. Chem. Phys.* **109**, 676–684 (1998).
 - [25] Bosse, J., Quitmann, D. & Wetzel, C. Mode-coupling theory of field-gradient correlation functions: The quadrupolar relaxation rate in liquids. *Phys. Rev. A* **28**, 2459–2473 (1983).
 - [26] Hertz, H. G. Magnetic relaxation by quadrupole interaction of ionic nuclei in electrolyte solutions part I: Limiting values for infinite dilution. *Ber. Bunsenges. Phys. Chem.* **77**, 531–540 (1973).
 - [27] Hertz, H. G. Magnetic relaxation by quadrupole interaction of ionic nuclei in electrolyte solutions part II: Relaxation at finite ion concentrations. *Ber. Bunsenges. Phys. Chem.* **77**, 688–697 (1973).
 - [28] Hertz, H. G., Holz, M., Keller, G., Versmold, H. & Yoon, C. Nuclear magnetic relaxation of alkali metal ions in aqueous solutions. *Ber. Bunsenges. Phys. Chem.* **78**, 493–509 (1974).
 - [29] Carof, A., Salanne, M., Charpentier, T. & Rotenberg, B. Collective water dynamics in the first solvation shell drive the NMR relaxation of aqueous quadrupolar cations. *J. Chem. Phys.* **145**, 124508 (2016).
 - [30] Badu, S., Truffandier, L. & Autschbach, J. Quadrupolar NMR spin relaxation calculated using ab initio molecular dynamics: Group 1 and group 17 ions in aqueous solution. *J. Chem. Theory Comput.* **9**, 4074–4086 (2013).
 - [31] Schmidt, J., Hutter, J., Spiess, H.-W. & Sebastiani, D. Beyond isotropic tumbling models: Nuclear spin relaxation in liquids from first principles. *ChemPhysChem* **9**, 2313–2316 (2008).
 - [32] Philips, A., Marchenko, A., Truffandier, L. A. & Autschbach, J. Quadrupolar NMR relaxation from ab initio molecular dynamics: Improved sampling and cluster models versus periodic calculations. *J. Chem. Theory Comput.* **13**, 4397–4409 (2017).
 - [33] Philips, A., Marchenko, A., Ducati, L. C. & Autschbach, J. Quadrupolar ^{14}N NMR relaxation from force-field and ab initio molecular dynamics in different solvents. *J. Chem. Theory Comput.* **15**, 509–519 (2019).
 - [34] Philips, A. & Autschbach, J. Quadrupolar NMR relaxation of aqueous $^{127}\text{I}^-$, $^{131}\text{Xe}^+$, and $^{133}\text{Cs}^+$: A first-principles approach from dynamics to properties. *J. Chem. Theory Comput.* **16**, 5835–5844 (2020).
 - [35] Engström, S., Jönsson, B. & Jönsson, B. A molecular approach to quadrupole relaxation. Monte Carlo simulations of dilute Li^+ , Na^+ , and Cl^- aqueous solutions. *J. Magn. Reson. (1969-1992)* **50**, 1–20 (1982).
 - [36] Engström, S., Jönsson, B. & Impey, R. W. Molecular dynamic simulation of quadrupole relaxation of atomic ions in aqueous solution. *J. Chem. Phys.* **80**, 5481–5486 (1984).

- [37] Linse, P. & Halle, B. Counterion N.M.R. in heterogeneous aqueous systems. *Mol. Phys.* **67**, 537–573 (1989).
- [38] Roberts, J. E. & Schnitker, J. Ionic quadrupolar relaxation in aqueous solution: Dynamics of the hydration sphere. *J. Phys. Chem.* **97**, 5410–5417 (1993).
- [39] Carof, A., Salanne, M., Charpentier, T. & Rotenberg, B. Accurate quadrupolar NMR relaxation rates of aqueous cations from classical molecular dynamics. *J. Phys. Chem. B* **118**, 13252–13257 (2014).
- [40] Carof, A., Salanne, M., Charpentier, T. & Rotenberg, B. On the microscopic fluctuations driving the NMR relaxation of quadrupolar ions in water. *J. Chem. Phys.* **143**, 194504 (2015).
- [41] Mohammadi, M., Benders, S. & Jerschow, A. Nuclear magnetic resonance spin-lattice relaxation of lithium ions in aqueous solution by NMR and molecular dynamics. *J. Chem. Phys.* **153**, 184502 (2020).
- [42] Chubak, I., Scalfi, L., Carof, A. & Rotenberg, B. NMR relaxation rates of quadrupolar aqueous ions from classical molecular dynamics using force-field specific Sternheimer factors. *J. Chem. Theor. Comput.* **17**, 6006–6017 (2021).
- [43] Sternheimer, R. On nuclear quadrupole moments. *Phys. Rev.* **80**, 102–103 (1950).
- [44] Sternheimer, R. M. Shielding and antishielding effects for various ions and atomic systems. *Phys. Rev.* **146**, 140–160 (1966).
- [45] Blöchl, P. E. Projector augmented-wave method. *Phys. Rev. B* **50**, 17953–17979 (1994).
- [46] Autschbach, J., Zheng, S. & Schurko, R. W. Analysis of electric field gradient tensors at quadrupolar nuclei in common structural motifs. *Concepts Magn. Reson. A* **36A**, 84–126 (2010).
- [47] Charpentier, T. The PAW/GIPAW approach for computing NMR parameters: A new dimension added to NMR study of solids. *Solid State Nucl. Magn. Reson.* **40**, 1–20 (2011).
- [48] Eisenstadt, M. & Friedman, H. L. Nuclear magnetic relaxation in ionic solution. I. Relaxation of ^{23}Na in aqueous solutions of NaCl and NaClO_4 . *J. Chem. Phys.* **44**, 1407–1415 (1966).
- [49] Eisenstadt, M. & Friedman, H. L. Nuclear magnetic relaxation in ionic solution. II. Relaxation of ^{23}Na in aqueous solutions of various diamagnetic salts. *J. Chem. Phys.* **46**, 2182–2193 (1967).
- [50] Petrilli, H. M., Blöchl, P. E., Blaha, P. & Schwarz, K. Electric-field-gradient calculations using the projector augmented wave method. *Phys. Rev. B* **57**, 14690–14697 (1998).
- [51] Calandra, P., Domene, C., Fowler, P. W. & Madden, P. A. Nuclear quadrupole coupling of ^{17}O and ^{33}S in ionic solids: Invalidation of the Sternheimer model by short-range corrections. *J. Phys. Chem. B* **106**, 10342–10348 (2002).
- [52] Gambuzzi, E., Charpentier, T., Menziani, M. C. & Pedone, A. Computational interpretation of ^{23}Na MQMAS NMR spectra: A comprehensive investigation of the Na environment in silicate glasses. *Chem. Phys. Lett.* **612**, 56–61 (2014).
- [53] Zeron, I. M., Abascal, J. L. F. & Vega, C. A force field of Li^+ , Na^+ , K^+ , Mg^{2+} , Ca^{2+} , Cl^- , and SO_4^{2-} in aqueous solution based on the TIP4P/2005 water model and scaled charges for the ions. *J. Chem. Phys.* **151**, 134504 (2019).
- [54] Blazquez, S., Conde, M. M., Abascal, J. L. F. & Vega, C. The Madrid-2019 force field for electrolytes in water using TIP4P/2005 and scaled charges: Extension to the ions F^- , Br^- , I^- , Rb^+ , and Cs^+ . *J. Chem. Phys.* **156**, 044505 (2022).
- [55] Carlson, S., Brüning, F. N., Loche, P., Bonthuis, D. J. & Netz, R. R. Exploring the absorption spectrum of simulated water from MHz to infrared. *J. Phys. Chem. A* **124**, 5599–5605 (2020).
- [56] Alder, B. J. & Wainwright, T. E. Decay of the velocity autocorrelation function. *Phys. Rev. A* **1**, 18–21 (1970).
- [57] Civan, M. M. & Shporer, M. *NMR of Sodium-23 and Potassium-39 in Biological Systems*, 1–32 (Springer US, Boston, MA, 1978).
- [58] Kestin, J., Khalifa, H. E. & Correia, R. J. Tables of the dynamic and kinematic viscosity of aqueous NaCl solutions in the temperature range 20–150 °C and the pressure range 0.1–35 MPa. *J. Phys. Chem. Ref. Data* **10**, 71–88 (1981).
- [59] Laage, D. & Hynes, J. T. A molecular jump mechanism of water reorientation. *Science* **311**, 832–835 (2006).
- [60] Turton, D. A. & Wynne, K. Stokes–Einstein–Debye failure in molecular orientational diffusion: Exception or rule? *J. Phys. Chem. B* **118**, 4600–4604 (2014).
- [61] Hansen, J. S., Kisliuk, A., Sokolov, A. P. & Gainaru, C. Identification of structural relaxation in the dielectric response of water. *Phys. Rev. Lett.* **116**, 237601 (2016).
- [62] Torre, R., Bartolini, P. & Righini, R. Structural relaxation in supercooled water by time-resolved spectroscopy. *Nature* **428**, 296–299 (2004).
- [63] Dubouis, N. *et al.* Chasing aqueous biphasic systems from simple salts by exploring the LiTFSI/LiCl/H₂O phase diagram. *ACS Cent. Sci.* **5**, 640–643 (2019).
- [64] Abbott, A. P., D’Agostino, C., Davis, S. J., Gladden, L. F. & Mantle, M. D. Do group 1 metal salts form deep eutectic solvents? *Phys. Chem. Chem. Phys.* **18**, 25528–25537 (2016).
- [65] Becher, M., Becker, S., Hecht, L. & Vogel, M. From local to diffusive dynamics in polymer electrolytes: NMR studies on coupling of polymer and ion dynamics across length and time scales. *Macromolecules* **52**, 9128–9139 (2019).
- [66] Hubbard, P. S. Nonexponential nuclear magnetic relaxation by quadrupole interactions. *J. Chem. Phys.* **53**, 985–987 (1970).
- [67] Spiess, H. W. Rotation of molecules and nuclear spin relaxation. In *Dynamic NMR Spectroscopy*, 55–214 (Springer Berlin Heidelberg, Berlin, Heidelberg, 1978).
- [68] Cowan, B. *Nuclear Magnetic Resonance and Relaxation* (Cambridge University Press, 1997).
- [69] Pyykkö, P. Year-2017 nuclear quadrupole moments. *Mol. Phys.* **116**, 1328–1338 (2018).
- [70] Lide, D. R. *CRC Handbook of Chemistry and Physics* (CRC Press, 2004), 85 edn.
- [71] Abascal, J. L. F. & Vega, C. A general purpose model for the condensed phases of water: TIP4P/2005. *J. Chem. Phys.* **123**, 234505 (2005).
- [72] Kirby, B. J. & Jungwirth, P. Charge scaling manifesto: A way of reconciling the inherently macroscopic and microscopic natures of molecular simulations. *J. Phys. Chem. Lett.* **10**, 7531–7536 (2019).
- [73] Marin-Lafèche, A. *et al.* Metalwalls: A classical molecular dynamics software dedicated to the simulation of electrochemical systems. *J. Open Source Softw.* **5**, 2373

- (2020).
- [74] Aguado, A. & Madden, P. A. Ewald summation of electrostatic multipole interactions up to the quadrupolar level. *J. Chem. Phys.* **119**, 7471–7483 (2003).
- [75] Giannozzi, P. *et al.* QUANTUM ESPRESSO: A modular and open-source software project for quantum simulations of materials. *J. Phys. Condens. Matter* **21**, 395502 (2009).
- [76] Varini, N., Ceresoli, D., Martin-Samos, L., Giroto, I. & Cavazzoni, C. Enhancement of DFT-calculations at petascale: Nuclear magnetic resonance, hybrid density functional theory and Car–Parrinello calculations. *Comput. Phys. Commun.* **184**, 1827–1833 (2013).
- [77] Perdew, J. P., Burke, K. & Ernzerhof, M. Generalized gradient approximation made simple. *Phys. Rev. Lett.* **77**, 3865–3868 (1996).
- [78] GIPAW Norm-Conserving Pseudopotentials. <https://sites.google.com/site/dceresoli/pseudopotentials>. Accessed: 2020-04-28.
- [79] Adamo, C. & Barone, V. Toward reliable density functional methods without adjustable parameters: The PBE0 model. *J. Chem. Phys.* **110**, 6158–6170 (1999).
- [80] Chen, T., Smit, B. & Bell, A. T. Are pressure fluctuation-based equilibrium methods really worse than nonequilibrium methods for calculating viscosities? *J. Chem. Phys.* **131**, 246101 (2009).
- [81] Yeh, I.-C. & Hummer, G. System-size dependence of diffusion coefficients and viscosities from molecular dynamics simulations with periodic boundary conditions. *J. Phys. Chem. B* **108**, 15873–15879 (2004).