



Investigation of the crystal field in rare earth titanate pyrochlores by resonant inelastic x-ray scattering

Octave Duros, Christian Brouder, Renaud Delaunay, Hebatalla Elnaggar,
Romain Jarrier, Amélie Juhin, Gheorghe S Chiuzbaian

► To cite this version:

Octave Duros, Christian Brouder, Renaud Delaunay, Hebatalla Elnaggar, Romain Jarrier, et al.. Investigation of the crystal field in rare earth titanate pyrochlores by resonant inelastic x-ray scattering. 17th SOLEIL User's Meeting, Jan 2023, Saint-Aubin, France. hal-03991376

HAL Id: hal-03991376

<https://hal.sorbonne-universite.fr/hal-03991376>

Submitted on 15 Feb 2023

HAL is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers.

L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.

Investigation of the crystal field in rare earth titanate pyrochlores by resonant inelastic x-ray scattering

Octave Duros^{1,2}, C. Brouder², R. Delaunay¹, H. Elnaggar², R. Jarrier¹, A. Juhin², G. S. Chiuzbăian¹

¹ Sorbonne Université, CNRS, Laboratoire de Chimie Physique – Matière et Rayonnement, LCPMR, F-75005 Paris, France

² Sorbonne Université, Muséum National d'Histoire Naturelle, CNRS, Institut de Minéralogie, de Physique des Matériaux et de Cosmochimie, IMPMC, F-75005 Paris, France

Context

- Rare earth titanate pyrochlores $R_2Ti_2O_7$ ($R = Sm, Tb, Dy, Er, Yb$) possess **fascinating magnetic states** [1-2] depending on the rare earth element R .
- The **structural argument** (e.g. different lattice parameters) **cannot solely explain** the variety of observed states: quantum spin liquids, spin ices or spin glasses.
- Characterization of the **crystal electric field** (CEF) acting on the R sites is a key ingredient for the understanding of those states.
- The **large variety** of CEF parametrizations that came over time led to many misconceptions in the litterature, thus needing a **clear definition** for further work.

1. CEF potential expansion

- CEF potential H_{CEF} perturbs the spherical symmetry of the free ion Hamiltonian H_F :

$$H = H_F + H_{\text{CEF}}$$

- One can express H_{CEF} as the sum of the potential V felt by every single electron i at a position r_i of the R metal:

$$H_{\text{CEF}} = -e \sum_{i=1}^n V(\mathbf{r}_i)$$

with V expressed with a time dependent charge $V(\mathbf{r}_i) = \int \frac{\rho(\mathbf{R})}{|\mathbf{R} - \mathbf{r}_i|} d\tau$ distribution $\rho(\mathbf{R})$


$$H_{\text{CEF}} = -e \sum_{i=1}^n \int \rho(\mathbf{R}) \sum_{k=0}^{\infty} \frac{r_i^k}{R^{k+1}} P_k(\cos \omega) d\tau$$

where $P_k(\cos \omega)$ are the *Legendre polynomials* [3]:

$$P_k(\cos \omega) = \frac{4\pi}{2k+1} \sum_{q=-k}^k Y_k^{q*}(\theta, \varphi) Y_k^q(\theta_i, \varphi_i)$$

| | |
|---|---|
| expansion coefficients | operators |
| $Y_k^{q*}(\theta, \varphi) = (-1)^q Y_k^{-q}$ | $Y_k^q(\theta_i, \varphi_i) = Y_k^q(i)$ |

3. The $R_2Ti_2O_7$ case



- R atoms and Ti atoms are situated in two different **apexes-sharing tetrahedrons sub-lattices** that interpenetrate into one another;
- Magnetic moments** are geometrically **frustrated** (3D kagomé lattices);
- D_{3d} point symmetry group, k and q are limited and CEF in $R_2Ti_2O_7$ compounds can be described by:

$$H_{\text{CEF}} = B_0^2 C_0^2 + B_0^4 C_0^4 + B_3^4 (C_{-3}^4 - C_3^4) + B_0^6 C_0^6 + B_3^6 (C_{-3}^6 - C_3^6) + B_6^6 (C_{-6}^6 + C_6^6)$$

5. Conclusion and outlook

- Choice of a clear definition based on spherical tensors, that can directly be used in Quanty code.
- First observation of the CEF effect in pyrochlore crystals at the $R N_{4,5}$ edges with RIXS.
- Precise extraction of CEF parameters in $R_2Ti_2O_7$ ongoing, with new experiments and machine learning.

2. CEF as a sum of parameters

$$P_k(\cos \omega) = \frac{4\pi}{2k+1} \left[Y_k^0 Y_k^0(i) + \sum_{q=-k}^k (-1)^q (Y_k^{-q} Y_k^q(i) + Y_k^q Y_k^{-q}(i)) \right]$$

We introduce the **tesseral harmonics** Z_{kq} to get rid of complex quantities:

$$Z_{k0}^c = Y_k^0, \quad Z_{kq}^c = \frac{1}{\sqrt{2}} (Y_k^{-q} + (-1)^q Y_k^q), \quad Z_{kq}^s = \frac{i}{\sqrt{2}} (Y_k^{-q} - (-1)^q Y_k^q)$$

as well as **tensor operators** which have the same transformation properties as Y_k^q :

$$C_q^k = \sqrt{\frac{4\pi}{2k+1}} Y_k^q(i)$$

It comes:


$$H_{\text{CEF}} = \sum_{k=0}^{\infty} \sqrt{\frac{4\pi}{2k+1}} \left[Z_{k0}^c C_0^k + \sum_{q=-k}^k \left(Z_{kq}^c \frac{1}{\sqrt{2}} (C_{-q}^k + (-1)^q C_q^k) \right. \right. \\ \left. \left. + Z_{kq}^s \frac{i}{\sqrt{2}} (C_{-q}^k - (-1)^q C_q^k) \right) \right] \times \int \rho(\mathbf{R}) \frac{r_i^k}{R^{k+1}} d\tau$$

We eventually introduce **CEF parameters** $B_q^k = \sqrt{\frac{4\pi}{2k+1}} Z_{k0}^c \int \rho(\mathbf{R}) \frac{r_i^k}{R^{k+1}} d\tau$ to simplify the former expression:

$$B_q^k = \sqrt{\frac{4\pi}{2k+1}} \frac{Z_{k0}^c}{\sqrt{2}} \int \rho(\mathbf{R}) \frac{r_i^k}{R^{k+1}} d\tau \quad B'_q^k = \sqrt{\frac{4\pi}{2k+1}} \frac{Z_{k0}^s}{\sqrt{2}} \int \rho(\mathbf{R}) \frac{r_i^k}{R^{k+1}} d\tau$$

$$H_{\text{CEF}} = \sum_{k=0}^{\infty} \left[B_0^k C_0^k + \sum_{q=-k}^k \left(B_q^k (C_{-q}^k + (-1)^q C_q^k) + B'_q^k (C_{-q}^k - (-1)^q C_q^k) \right) \right]$$

4. How to characterize the CEF



RIXS process representation and RIXS spectra of the CEF effect in $Yb_2Ti_2O_7$ measured on SEXTANTS beamline in SOLEIL, at the $Yb N_{4,5}$ edge.

- High-resolution RIXS** allows probing the CEF;
- AERHA spectrometer [4] on SEXTANTS is **worldwide** the only instrument offering the required **energy resolution**
- Simulations run through **Quanty** software package [5]: **Ligand Multiplet Theory** calculations with CEF contribution, expanded on **spherical tensors**.