



HAL
open science

Role of MuscleBlind-Like proteins in the regulation of expression of $CaV\beta 1$ isoforms in adult skeletal muscle

Amélie Vergnol, Alain Sureau, A. Traoré, X. Lornage, G. Gourdon, Denis Furling, France Piétri-rouxel, S. Falcone

► **To cite this version:**

Amélie Vergnol, Alain Sureau, A. Traoré, X. Lornage, G. Gourdon, et al.. Role of MuscleBlind-Like proteins in the regulation of expression of $CaV\beta 1$ isoforms in adult skeletal muscle. Myology 2022, Sep 2022, Nice (FRANCE), France. hal-03999540

HAL Id: hal-03999540

<https://hal.sorbonne-universite.fr/hal-03999540>

Submitted on 6 Mar 2023

HAL is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers.

L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.


Role of MuscleBlind-Like proteins in the regulation of expression of CaVβ1 isoforms in adult skeletal muscle

A. Vergnol¹, A. Sureau¹, M. Traoré¹, X. Lornage¹, G. Gourdon¹, D. Furling¹, F. Pietri-Rouxel^{1*} and S. Falcone^{1*}


¹ Sorbonne Université, Inserm, Institut De Myologie, Centre De Recherche En Myologie F-75013 – Paris (France), * Equally contributed to this work

INTRODUCTION


CaVβ1-E/GDF5 axis in muscle mass homeostasis



Cacnb1 isoforms in skeletal muscle



Implication of MBNLs in DM1 pathophysiology




Voltage-gated calcium channels (CaVs or VGCCs) are major regulators of calcium-related cellular functions. In skeletal muscle, though the essential component of the pore channel is the CaVα1 subunit, the CaVβ1 subunit is an essential subunit guaranteeing CaV fine-tuning activity.

CaVβ1-E and CaVβ1-D are two different isoforms of CaVβ1 protein in skeletal muscle, expressed during embryogenesis and in healthy innervated adult muscle, respectively. Importantly, our recent study demonstrated that the embryonic CaVβ1-E expression increases after a nerve damage in adult skeletal muscle and enables the expression of GDF5 (Growth Differentiation Factor 5) to counteract excessive muscle wasting (Traoré et al. 2019). However, the mechanisms leading to the increase in CaVβ1-E expression are unknown to date.


Our RNAseq data analysis in innervated versus denervated muscles revealed MuscleBlind-Like (MBNL) proteins as potential candidates regulating CaVβ1 expression in skeletal muscle. Interestingly, in a human model of Dystrophy Myotonic 1 (DM1), the sequestration of MBNLs in toxic nuclear aggregates is related to an impaired splicing of CaVβ1 transcript (CACNB1) (Arandel et al. 2017).

Here, we evaluate the effect of a modulation of MBNLs protein levels on the expression of CaVβ1 isoforms in both *in vitro* and *in vivo* systems as well as in pathological mouse models of DM1.

Correlation between MBNLs and Cacnb1 expressions




Mouse model of Mbn1 downregulation




MBNLs modulates the expression of CaVβ1 isoforms *in vivo*

Variations of Cacnb1 isoforms at transcriptional level




A downregulation of Mbn1 induces an increase of the expression of CaVβ1-E in mouse skeletal muscle through a direct or indirect mechanism.


CaVβ1E (70 kDa)




Variations of CaVβ1 isoforms at protein level



MBNLs impacts Cacnb1 mRNA stability



Cacnb1-E increases in DM1 pathological models



CONCLUSIONS & PERSPECTIVES

- Mbn1 negatively regulates Mbn3
- Downregulation of Mbn1 *in vivo* is associated with impaired muscle and neuromuscular functions
- A downregulation of Mbn1, associated with an increase of Mbn3, leads to increased CaVβ1-E and decreased CaVβ1D expression levels *in vivo*
- Cacnb1 transcripts stability is modulated by MBNLs *in vitro* through their 3'UTR
- Mouse models of DM1 are associated with an increased Cacnb1-E expression

- Characterization of the splicing events occurring at Cacnb1 Ex2-3 and Ex13-14
- Studying a potential cross-regulation of CaVβ1-D on CaVβ1-E expression
- Deciphering the role of CaVβ1-E in DM1 pathophysiology