



HAL
open science

Neighborhood built environment typologies and adiposity in children and adolescents

Tracie A Barnett, Adrian E Ghenadenik, Andraea van Hulst, Gisele Contreras, Yan Kestens, Basile Chaix, Marie-Soleil Cloutier, Melanie Henderson

► **To cite this version:**

Tracie A Barnett, Adrian E Ghenadenik, Andraea van Hulst, Gisele Contreras, Yan Kestens, et al.. Neighborhood built environment typologies and adiposity in children and adolescents. *International Journal of Obesity*, 2022, 46 (3), pp.588-596. 10.1038/s41366-021-01010-1 . hal-04016825

HAL Id: hal-04016825

<https://hal.sorbonne-universite.fr/hal-04016825v1>

Submitted on 6 Mar 2023

HAL is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers.

L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.

Neighborhood built environment typologies and adiposity in school age children.

Tracie A. BARNETT^{1,2}, PhD, Adrian E. GHENADENIK¹, PhD, Andraea VAN HULST^{2,3}, RN, PhD, Gisele CONTRERAS, MSc, Marie-Soleil Cloutier, PhD, Yan KESTENS^{3,4}, PhD, Basile CHAIX^{6,7}, PhD, Melanie HENDERSON^{2,8}, MD, PhD.

Affiliations: ¹Department of Family Medicine, McGill University, Montreal, QC, Canada; ²Centre de recherche du CHU Sainte-Justine, Montreal, Canada; ³École de santé publique de l'Université de Montréal, Montreal, Canada; ⁴Centre de Recherche du Centre Hospitalier de l'Université de Montréal (CRCHUM), Montreal, Canada; ⁵Centre Urbanisation Culture Société, ⁶Inserm, Paris, France; ⁷Université Pierre et Marie Curie-Paris6, UMR-S 707, Paris, France; ⁸Department of Pediatrics, Université de Montréal, Montreal, Canada

Address correspondence to: Tracie A. Barnett, 5858, chemin de la Côte-des-Neiges, 3rd floor, Montreal, QC H3S 1Z1

Email: tracie.barnett@mcgill.ca

Phone: (514) 398-7375

Short title: Built environment and adiposity in youth

Abbreviations: BMI, body mass index; FMI, fat mass index; GIS, Geographic Information System; QUALITY, Quebec Adipose and Lifestyle Investigation in Youth; WC, waist circumference

Key Words: adiposity, built environment, children, cluster analysis, neighborhood characteristics, obesity, childhood obesity

Funding source: The QUALITY cohort and the QUALITY Residential study were funded by the Canadian Institutes of Health Research, the Heart and Stroke Foundation of Canada and the Fonds de la recherche en santé du Québec.

Financial disclosure: none

Conflict of interest: none

What's Known on This Subject: Specific neighborhood features (e.g., infrastructures that facilitate walking) have been associated with childhood obesity. Nonetheless, findings concerning their potential to influence energy-related behaviors and outcomes in youth are not consistently supported in the literature. To date, few studies have taken into account the interrelation between multiple neighborhood dimensions in relation to childhood and adolescent adiposity.

What This Study Adds: Participants living in neighborhoods with high volumes of vehicular traffic had greater adiposity, independent of other built environment features. Neighborhood walkability may be an important target for obesity prevention, and safety from vehicular traffic may need to be prioritized.

Counts:

Abstract: 197 (AJE: max 200 words)

Main text: 3,499 words (AJE: max 3,500 words)

Contributor's Statement:

Tracie A Barnett is the principal investigator of the QUALITY Residential study. She conceptualized and designed the study, carried out the statistical analyses, co-drafted the initial manuscript, and approved the final manuscript as submitted.

Adrian E Ghenadenik contributed to the statistical analyses, revised several versions of the manuscript, and approved the final manuscript as submitted

Gisele Contreras contributed to the statistical analysis and approved the final version of the manuscript.

Andraea Van Hulst coordinated and supervised neighborhood data collection, contributed to the analytic conceptualization, co-drafted the initial manuscript, and approved the final manuscript as submitted.

Marie-Soleil Cloutier helped develop and test the neighbourhood audit tools and reviewed and approved the final manuscript.

Yan Kestens contributed to data collection for the spatial component of the study, and approved the final manuscript as submitted.

Basile Chaix contributed to the analytic conceptualization, revised the manuscript, and approved the final manuscript as submitted.

Melanie Henderson is the principal investigator of the QUALITY study, contributed to data collection, reviewed and revised the manuscript, and approved the final manuscript as submitted.

ABSTRACT

Neighborhoods are complex multidimensional systems. However, the interrelation between multiple neighborhood dimensions is seldom considered in relation to youth adiposity. We created a neighborhood typology using a range of built environment features and examined its association with adiposity in youth.

Analyses are based on data from the QUALITY cohort, an ongoing study on the natural history of obesity in Quebec youth with a history of parental obesity. Adiposity was measured at baseline and follow up approximately 8 years later. Neighborhood features were measured at baseline through in-person neighborhood assessments and geocoded administrative data and were summarized using principal components analysis. Neighborhood types were identified using cluster analysis. Associations between neighborhood types and adiposity were examined using multivariable linear regressions.

Five distinct neighborhood types characterized by levels of walkability and traffic-related safety were identified. At ages 8-10 years, children in moderate walkability/low safety neighborhoods had higher adiposity than those residing in moderate walkability/high safety neighborhoods. Attenuated associations were detected between neighborhood types and adiposity 8 years later.

Neighborhoods characterized by lower traffic safety appear to be the most obesogenic to children, regardless of other walkability-related features. Policies targeting neighborhood walkability for children may need to prioritize vehicular traffic safety.

INTRODUCTION

Obesity is a complex multi-factorial condition with wide-ranging causes encompassing genetic, biological, behavioral and socio-environmental factors (1, 2). Youth obesity has tripled over the past generation and remains a dire public health challenge (3). Obesity in childhood persists into adulthood (4), and is linked to adult disease independently of later weight status (5). Upstream, socio-environmental approaches are essential in order to impact the population-level distribution of obesity, specifically by preventing or delaying the onset of overweight in childhood (6). Although our knowledge base for developing such interventions is still limited, (7, 8) significant area-level variation in the prevalence of obesity (9-13) suggests that specific types of neighborhoods may embody an 'obesogenic potential'. Neighborhood-level features are thought to influence obesity through energy-related behaviors (e.g.: diet, physical activity, sedentary behavior) (14-16). If salient environmental features implicated in pediatric obesity can be identified, reducing the obesogenic nature of neighborhoods through even minor environmental modifications could lead to significant population-level improvements.

Neighborhood features characterizing obesogenic environments likely differ between pediatric and adult populations (17, 18) (e.g., the detrimental effect of urban sprawl in adults (19) is inverted in adolescents (20)). These differences call for tailoring research and intervention to specific populations. Policy recommendations targeting schools and neighborhoods as key environments to support healthy lifestyle behaviors in youth largely reflect this strategy (21, 22). Nonetheless, despite the theoretical appeal of different neighborhood-level features in terms of their potential to influence energy-related behaviors, their association with these outcomes in youth is not consistently supported in the literature (14, 15, 23). Among different issues underlying inconclusive findings are heterogeneous conceptual and methodological approaches (24-26). These include the definition, operationalization and measurement of neighborhood features, predictors and outcome variables, and the analytic approaches and covariates included in statistical models (14, 23, 27). Furthermore, built environment features tend to be correlated with one another, therefore modeling their association with health outcomes requires consideration for their interrelated nature (28-30).

An approach suitable to this purpose is cluster analysis. Cluster-based statistical techniques categorizing neighborhoods in terms of attribute patterns (i.e.: neighborhood typologies) have been used in relation to physical activity in adults (31-33), adolescents (34, 35), and children (36), and also in relation to adiposity in adults (19, 37) and youth (38, 39). Aside from their contribution to identify areas representing different configurations of resources, these studies highlight the importance of accounting for the presence of both health-promoting and health-detering features that contribute to shaping health area-level behaviour.

The aim of this study was to identify neighbourhood types based on objectively-assessed neighbourhood attributes, and cross-sectional and prospective associations with adiposity in children over time. The specific objectives were to: 1) create a neighborhood typology using a wide range of built environment features, including child-oriented features; 2) examine associations between neighborhood types and multiple measures of adiposity in children aged 8-10 years, and 3) examine if these types predict adiposity in these children approximately 8 years later, at ages 15-18 years.

PARTICIPANTS AND METHODS

Study design and participants

The analyses in this paper were based on data from the Quebec Adipose and Lifestyle Investigation in Youth (QUALITY) cohort, an ongoing longitudinal investigation of the natural history of obesity and cardiovascular risk factors in vulnerable children, and the complementary QUALITY Residential Study. This cohort comprises 630 children aged 8-10 years at baseline (2005-2008) and both biological parents. Families were recruited through elementary schools located within a 75-km radius of three major urban centers in the province of Quebec, Canada: Montreal, Quebec City and Sherbrooke. Eligibility criteria required participants to be Caucasian, aged 8–10 years at the time of recruitment, with both biological parents being available for the study and at least one of them being obese based on self-reported weight, height, and waist circumference (WC) (i.e., body mass index (BMI) of mother and/or father $\geq 30 \text{ kg/m}^2$, or mother's WC $\geq 88 \text{ cm}$ and/or father's WC $\geq 102 \text{ cm}$). Follow up was conducted approximately 8 years later (2012-2016) in 377 participants (retention rate=60%). A detailed description of the study design and data collection methods is available elsewhere (40). Analyses were restricted to participants for who full neighborhood environment data were available (n=485). Prospective analyses were restricted to participants with complete follow-up data (n=298). A diagram detailing the number of participants by time of survey and reasons for inclusion/exclusion from analytical samples is presented in Figure 1.

Measurements

Measurements in children

Adiposity measures were taken at both baseline and follow up. Weight was measured to the nearest 0.1kg using an electronic scale, with participants wearing lightweight indoor clothing without shoes or sweaters. Height was measured with a stadiometer, and was recorded to the nearest millimeter during maximal inspiration. Waist circumference (WC) was measured midway between the lowest rib and the superior border of the iliac crest with a non-stretchable fiberglass measurement tape. Anthropometric measurements were taken twice and averages were computed. If measurements differed by more than 0.2cm for height and WC, or 0.2kg for weight, a third measurement was taken, and the average between the two closest measurements was computed. BMI Z-scores were computed using CDC growth standards (41). Dual-energy X-ray absorptiometry was used to measure children's fat mass. Fat mass index (FMI) was computed by dividing total body fat mass by height in meters squared (kg/m^2) (42). The percentage of central body fat (computed as trunk fat mass/total fat mass x 100) was estimated using trunk fat mass obtained from the automatic region that consists of the chest, abdomen and pelvic area. Pubertal development stage was assessed by a nurse using the 5-stage Tanner scale (43, 44), and was dichotomized as pre-pubertal (Tanner 1) vs. puberty initiated (Tanner >1).

Measurements in parents

Both parents' weight and height were measured at baseline using the same instruments and protocol as in children. BMI was computed as weight (kg) divided by height squared (m^2).

Neighborhood environment features

Characteristics of the built and social environments in children's residential neighborhoods were obtained at baseline for families residing in the Montreal metropolitan area (n=512). Exact

residential addresses of each participant were geocoded. Data were collected using two sources: 1) in-person neighborhood audits, and 2) administrative data.

Audits were performed using the QUALITY audit tool, an observation grid adapted from an existing neighbourhood assessment instrument (45). This grid includes a checklist scoring 60 street-level built environment features surrounding each participant's residential address. Audits were conducted by independent pairs of trained observers. Inter-rater reliability was substantial ($\kappa > 0.60$) (46) for most of the indicators used in our analyses. Built environment features of up to 10 street segments located within the immediate residential environment were subjected to a detailed assessment. Features recorded include: 1) the presence of sidewalks (absent, present on one side only, present on both sides); 2) the presence of pedestrian aids (zebra crossing, pedestrian crossing light, pedestrian crossing sign, all-direction stop sign at intersection, widened sidewalk at intersection, paved pedestrian crossing, designated 'school corridor'); 3) the presence of traffic calming measures (speed bump, mid-street segment stop sign, speed limit 30 km/h, traffic lights, large obstacles designed to decrease the number of driving lanes); and 4) signs of social disorder (graffiti, vandalism, litter, abandoned building/construction). Indicators were created reflecting the average number of sidewalks per street segment (ranging from 0 to 2), the proportion of street segments with at least one pedestrian aid, the proportion of street segments with at least one traffic calming measure, and having at least one street segment with one or more signs of social disorder, respectively.

Administrative data were collected using the MEGAPHONE database. This is a geographic information system (GIS) that seeks to characterize social, built and natural environmental factors to understand the relationship between contextual and compositional factors and health outcomes in the Montreal region (47). The following indicators were computed for 1km street-network buffers centered on the participants' residences: 1) the number of three-way -or more- intersections; 2) a measure of land use mix (residential, commercial, industrial, recreational, or other) based on an entropy equation resulting in a score of 0 to 1, where 0 represents homogeneity (all land uses within the area are of a single type), and 1 represents heterogeneity (even distribution of all land use categories within the area) (48); 3) number of parks; 4) percentage of streets within buffers that have heavy vehicular traffic at rush hour, categorized as less than 1%, 1-5%, and $\geq 5\%$; 5) total length of streets within the buffer that have normal vehicular traffic at rush hour; 6) total length of streets within the buffer that have heavy vehicular traffic at rush hour (categorized as 0 km, >0-1 km, >1-5 km, ≥ 5 km); 7) density of private dwellings per hectare (10000m^2); 8) a normalized difference vegetation index (NDVI) based on satellite images of the amount of chlorophyll present, the index ranges from -1 to 1, with greater values indicating more vegetation; and 9) the proportion of the buffer area covered by parks. In addition, as a measure of area-level socioeconomic status, the proportion of residents aged 24-64 years who completed a university degree, was computed using 2006 Census data for each buffer.

Statistical analysis

Identification of neighborhood types

A two-step approach was used to identify neighborhood types. First, principal component analysis (PCA) with varimax rotation was used to reduce neighborhood environment measures to a parsimonious number of distinct and meaningful components. A three-factor solution was

selected based on the Kaiser criterion (Eigenvalues > 1), accounting for 62% of the total variance in the data. The first component (37% of variance) was defined by many intersections, more mixed land use, more parks, more sidewalks, more streets with low traffic and more signs of social disorder. The second component (14% of variance) was defined by meters of roadway and proportion of roadway with heavy traffic volume at rush hour, and the third component (11% of variance) was defined by the presence of traffic calming measures and pedestrian aids.

Following this step, and based on the three components from PCA, hierarchical cluster analysis using Ward's method (49) was used seeking to identify unique neighborhood types for subsequent examination in relation to child adiposity. This method starts with each multidimensional observation (neighborhood) as a single cluster and then repeatedly merges the next two closest clusters in terms of Euclidian distances between observations until a single, all-encompassing cluster remains (50). Application of this method results in a typology wherein neighborhoods that were substantively comparable on selected characteristics were grouped together despite not necessarily being geographically adjacent (51). A five-cluster solution was retained, with 68% of the variation in the data explained by the variables included in the clusters.

Neighborhood typology and adiposity

Dummy variables were created for each cluster, and multiple linear regression was used to examine associations between neighborhood types and adiposity indicators at baseline and follow-up. Models were adjusted for age, sex, pubertal development, percentage of university-educated residents, and parental BMI scores. Prospective analyses were restricted to participants who remained at the same address between baseline and follow up (n=226). Prospective models were adjusted using the same covariates as cross-sectional models, except for pubertal development, as all participants had initiated puberty at follow-up. Sensitivity analyses using different neighborhood types as referent were conducted to test the robustness of results obtained from the main models. Analyses were conducted using SAS v9.2 and Stata v15.

RESULTS

Descriptive statistics for the five neighborhood types are presented in Table 1. Patterns emerged based on two dimensions, *walkability* and *safety*. *Walkability* was characterized by number of intersections, dwelling density, total street length, sidewalks, pedestrian aids, and parks; *safety* was based on indicators of rush hour vehicular traffic and of traffic calming measures. Type 1 neighborhoods (n=132) were defined as '*moderate walkability and high safety*'; type 2 neighborhoods (n=115) as '*low walkability and high safety*'; type 3 neighborhoods (n=64) as '*moderate walkability and moderate safety*'; type 4 neighborhoods (n=108) as '*high walkability and moderate safety*'; and type 5 neighborhoods (n=66) as '*moderate walkability and low safety*'.

Characteristics of participants by neighborhood type are presented in Table 2. Mean age of children, maternal BMI, and pubertal status were similar across neighborhood types. Differences by neighborhood type were observed in terms of paternal BMI and all indicators of child adiposity, chiefly in Type 5 neighborhoods, where participants were more likely to have higher adiposity compared to those residing in other neighborhoods. Also, differences were found in

terms of child's sex, with lower proportions of boys in Type 3 neighborhoods and lower proportions of girls in Type 4 neighborhoods.

Fully-adjusted cross-sectional associations between neighborhood types and adiposity indicators are presented in Table 3. Results show that children residing in Type 5 neighborhoods (*moderate walkability and low safety*) had higher BMI Z-scores [β : 0.41 (0.11; 0.71)], FMI [β : 1.22 (0.28; 2.16)], waist circumference [β : 4.90 (1.60; 8.21)], and central fat mass percentage [β : 1.60 (0.03; 3.17)] compared to their Type 1 neighborhood counterparts (reference category). Sensitivity analyses using other neighborhood types as referents, chiefly those with relatively high safety showed similar results, highlighting the importance of this dimension in terms of its association with adiposity indicators (see Supplementary Materials for details).

Fully-adjusted prospective associations between neighborhood types and adiposity indicators are presented in Table 4, and generally extend cross-sectional findings. Although no statistically significant results between neighborhood types and adiposity indicators were detected, attenuated associations (p value <0.10) between Type 5 neighborhoods and a larger waist circumference FMI [β : 1.42 (-0.06; 2.91)], and waist circumference [β : 5.06 (-0.24; 10.36)] were observed in adolescents. Sensitivity analyses using other neighborhood types as referents showed similar results, supporting the robustness of these findings.

DISCUSSION

This study used a novel approach seeking to identify patterns of neighborhood-level environmental features (as opposed to examining individual attributes in isolation), and their association with child and youth adiposity. We used a wide range of both direct observation and GIS-derived measures of residential neighborhood characteristics to generate distinct neighborhood types. Walkability and safety emerged as the salient constructs characterizing distinct types of neighborhoods containing complex combinations of both positive and negative environmental influences. In our sample, we identified five types in the best fitting model, which we labeled as: 1) moderate walkability/high safety, 2) low walkability/high safety, 3) moderate walkability/moderate safety, 4) high walkability/moderate safety, and 5) moderate walkability/low safety.

Cross-sectional analyses, which examined associations between neighborhood types and adiposity indicators in children aged 8-10 years, showed that participants living in neighborhoods characterized by moderate walkability, and most importantly, low safety levels, had higher adiposity (BMI z-scores, FMI, waist circumference and central fat mass percentage) than their counterparts residing in other neighborhood types. These findings suggest that safety may play an important role in shaping adiposity outcomes in this age group. Studies including a recent systematic review of traffic-related environmental factors and childhood obesity have indeed reported associations between objective and subjective measures of vehicular traffic-related safety and adiposity/obesity indicators (52). Low levels of traffic-related safety may discourage active transportation and outdoor play even in areas with moderate or high walkability -such as Type 5 neighborhoods in this study, which were moderately walkable but less safe-. In these areas, destinations including schools, shops and play areas may be within walking distance, however high volumes of local traffic may discourage active transportation,

and time spent outdoors (53-57). In such neighborhoods, children may not have parental permission -or may not choose- to venture into these destinations (58). In this vein, several reviews have underscored pedestrian safety structures as one of the features most consistently associated with PA in children (14, 59, 60), and studies have reported associations between presence of pedestrian infrastructures such as sidewalks and access to paths and higher levels of physical activity (61, 62) and lower body weight (63, 64).

Prospective models detected attenuated associations between neighborhood types and adiposity in participants 8 years later (ages 15-17). Despite their lack of statistical significance, which may be partly due to lack of sufficient statistical power, these results are in line with those from cross-sectional models, and provide some support for the potential role of low-safety neighborhoods in influencing adiposity in adolescence. Very few studies have examined associations between road safety and obesity/adiposity in adolescents: most have focused their efforts on putative mediators such as physical activity (PA) and/or active transportation. In any case, the evidence concerning obesity and adiposity outcomes (52, 65), as well as PA (66, 67) in this age group remains inconclusive.

Whereas associations in adolescents are directionally similar to those detected 8 years earlier, their attenuated nature may be indicative of a less prominent role of traffic safety features as promoters or deterrents of energy expenditure in adolescents. The authors of one of the above-cited reviews (66) propose that this is because teenagers typically have less parental constraints regarding traffic safety issues, and therefore they use their immediate neighborhood features differently than children. Also, these results may point to a more limited relevance of the immediate residential environment to adolescents, as their activity spaces likely go beyond their area of residence. Therefore, our exclusive focus on surrounding residential areas may not be reflective of the full range of environmental exposures to which adolescents may be exposed to. Future studies should strive to incorporate other activity spaces to test whether non-residential areas are associated with adiposity outcomes in this age group

This study has several strengths. First, the environmental assessments using both territorial and ego-centric neighborhood definitions allowed us to capture a wide array of features to which children and adolescents are exposed to. Second, the analyses in this paper go beyond BMI as the sole measure of adiposity, to also include direct measures including waist circumference, fat mass index, and percentage of central fat mass. Of note, most studies have relied on BMI rather than on direct measures of body fat (68, 69) due to the relative ease in collecting information on weight and height. Although useful for surveillance and individual assessment purposes, BMI tends to be a more reliable measure of excess weight for height than of excess adiposity (70). Third, our findings cover an important stage of the life course, spanning from childhood to adolescence. Fourth, our methodology allowed for the examination of an array of environmental features whose effects may be difficult to disentangle using multivariable regression analyses due to multicollinearity (28). Moreover, cluster analysis methods allow to take into account the multidimensional nature of neighborhoods by which a combination of features, rather than individual ones, may influence health outcomes and behaviors (71-73). Looking at neighborhoods as combinations of multiple factors, may provide a more complete picture of the interaction of the environment with health.

Some limitations must be noted as well. First, the study sample was limited to Caucasian children at risk of obesity. Furthermore, participants in this cohort are relatively more socially advantaged than the average in the province of Quebec. These two issues potentially limit the generalizability of our findings. Second, differential loss to follow-up of families living in more disadvantaged areas, who tend to have worse adiposity outcomes, may have resulted in some selection bias. Third, and despite being a key component to this study's typology, neighborhood walkability did not emerge as a strong correlate of obesity. Although the measures used in this study incorporated several child-specific measures such as parks and 'school corridors', it is possible that the inclusion of a more focused set of indicators (e.g.: specific park features, recreational resources, and school proximity) would generate a more child-centric construct of walkability. Future research should seek to further establish more nuanced constructs of walkability relevant to children, taking into consideration their development stage and their level of independent mobility. Fourth, since the neighborhood features included in the typology were only measured at baseline, it is possible that some of these have changed over the course of the 8 years after their assessment. This may have resulted in some misclassification bias, and may partly explain the lack of significant associations between traffic safety and adiposity in adolescents. Finally, as mentioned above, this paper focused exclusively on proximal residential environments. This is likely of limited concern in younger children, as the residential neighborhood typically constitutes the most important activity space in this age group (74). Future studies should also capture activity spaces beyond the residential environment in order to capture spatial and temporal variation in exposure to contextual features (75-77).

CONCLUSION

Our study is one of the first studies to identify neighbourhood typologies and cross-sectional and prospective associations with adiposity in youth. Findings constitute an original contribution to the research literature by simultaneously examining the influence of multiple indicators of neighborhood characteristics on various measures of adiposity, at childhood and adolescence. Findings highlighting associations between areas characterized by relatively low traffic safety and a higher risk of adiposity in children suggest this may be a promising target for public health intervention. The examination of neighbourhood typologies may be a promising avenue for informing urban design through understanding how different combinations of neighbourhood environments influence health, and may provide guidance for policy and urban planners. Future research should also examine potential mechanisms of typology-health associations.

REFERENCES

1. van der Klaauw Agatha A, Farooqi IS. The Hunger Genes: Pathways to Obesity. *Cell*. 2015;161(1):119-32.
2. Spruijt-Metz D. Etiology, Treatment and Prevention of Obesity in Childhood and Adolescence: A Decade in Review. *J Res Adolesc*. 2011;21(1):129-52.
3. World Health Organization. Obesity and overweight: Key Facts 2020 [Available from: <https://www.who.int/news-room/fact-sheets/detail/obesity-and-overweight>].
4. Singh AS, Mulder C, Twisk JW, van Mechelen W, Chinapaw MJ. Tracking of childhood overweight into adulthood: a systematic review of the literature. *Obes Rev*. 2008;9(5):474-88.
5. Biro FM, Wien M. Childhood obesity and adult morbidities. *The American Journal of Clinical Nutrition*. 2010;91(5):1499S-505S.
6. Sallis JF, Cervero RB, Ascher W, Henderson KA, Kraft MK, Kerr J. An ecological approach to creating active living communities. *Annu Rev Public Health*. 2006;27:297-322.
7. Bleich SN, Segal J, Wu Y, Wilson R, Wang Y. Systematic review of community-based childhood obesity prevention studies. *Pediatrics*. 2013;132(1):e201-10.
8. Showell NN, Fawole O, Segal J, Wilson RF, Cheskin LJ, Bleich SN, et al. A systematic review of home-based childhood obesity prevention studies. *Pediatrics*. 2013;132(1):e193-200.
9. Zhang X, Onufrak S, Holt JB, Croft JB. A multilevel approach to estimating small area childhood obesity prevalence at the census block-group level. *Prev Chronic Dis*. 2013;10:E68.
10. Willms J, Tremblay M, Katzmarzyk P. Geographic and demographic variation in the prevalence of overweight Canadian children. *Obesity Research*. 2003;11(5):668-73.
11. Grow HM, Cook AJ, Arterburn DE, Saelens BE, Drewnowski A, Lozano P. Child obesity associated with social disadvantage of children's neighborhoods. *Soc Sci Med*. 2010;71(3):584-91.
12. Veugelers PJ, Fitzgerald AL. Prevalence of and risk factors for childhood overweight and obesity. *Cmaj*. 2005;173(6):607-13.
13. Bruner MW, Lawson J, Pickett W, Boyce W, Janssen I. Rural Canadian adolescents are more likely to be obese compared with urban adolescents. *Int J Pediatr Obes*. 2008;3(4):205-11.
14. Ding D, Sallis JF, Kerr J, Lee S, Rosenberg DE. Neighborhood environment and physical activity among youth a review. *Am J Prev Med*. 2011;41(4):442-55.
15. de Vet E, de Ridder DT, de Wit JB. Environmental correlates of physical activity and dietary behaviours among young people: a systematic review of reviews. *Obes Rev*. 2011;12(5):e130-42.
16. Safron M, Cislak A, Gaspar T, Luszczynska A. Micro-environmental characteristics related to body weight, diet, and physical activity of children and adolescents: a systematic umbrella review. *Int J Environ Health Res*. 2011;21(5):317-30.
17. Saelens BE, Sallis JF, Frank LD, Couch SC, Zhou C, Colburn T, et al. Obesogenic neighborhood environments, child and parent obesity: the Neighborhood Impact on Kids study. *Am J Prev Med*. 2012;42(5):e57-64.
18. Gauthier KI, Krajicek MJ. Obesogenic environment: a concept analysis and pediatric perspective. *J Spec Pediatr Nurs*. 2013;18(3):202-10.
19. Mackenbach JD, Rutter H, Compennolle S, Glonti K, Oppert JM, Charreire H, et al. Obesogenic environments: a systematic review of the association between the physical environment and adult weight status, the SPOTLIGHT project. *BMC Public Health*. 2014;14:233.

20. Seliske L, Pickett W, Janssen I. Urban sprawl and its relationship with active transportation, physical activity and obesity in Canadian youth. *Health Rep.* 2012;23(2):17-25.
21. Fenton M. Community design and policies for free-range children: creating environments that support routine physical activity. *Child Obes.* 2012;8(1):44-51.
22. Raine KD, Muhajarine N, Spence JC, Neary NE, Nykiforuk CI. Coming to consensus on policy to create supportive built environments and community design. *Canadian journal of public health = Revue canadienne de sante publique.* 2012;103(9 Suppl 3):eS5-8.
23. Dunton GF, Kaplan J, Wolch J, Jerrett M, Reynolds KD. Physical environmental correlates of childhood obesity: a systematic review. *Obes Rev.* 2009;10(4):393-402.
24. Ding D, Gebel K. Built environment, physical activity, and obesity: what have we learned from reviewing the literature? *Health Place.* 2012;18(1):100-5.
25. Diez Roux AV, Mair C. Neighborhoods and health. *Annals of the New York Academy of Sciences.* 2010;1186:125-45.
26. Carter MA, Dubois L. Neighbourhoods and child adiposity: A critical appraisal of the literature. *Health & Place.* 2010;16(3):616-28.
27. Feng J, Glass TA, Curriero FC, Stewart WF, Schwartz BS. The built environment and obesity: a systematic review of the epidemiologic evidence. *Health Place.* 2010;16(2):175-90.
28. Leal C, Bean K, Thomas F, Chaix B. Multicollinearity in associations between multiple environmental features and body weight and abdominal fat: using matching techniques to assess whether the associations are separable. *Am J Epidemiol.* 2012;175(11):1152-62.
29. Jones M, Huh J. Toward a multidimensional understanding of residential neighborhood: a latent profile analysis of Los Angeles neighborhoods and longitudinal adult excess weight. *Health Place.* 2014;27:134-41.
30. Myers CA, Denstel KD, Broyles ST. The context of context: Examining the associations between healthy and unhealthy measures of neighborhood food, physical activity, and social environments. *Prev Med.* 2016;93:21-6.
31. Adams MA, Sallis JF, Kerr J, Conway TL, Saelens BE, Frank LD, et al. Neighborhood environment profiles related to physical activity and weight status: a latent profile analysis. *Prev Med.* 2011;52(5):326-31.
32. Charreire H, Weber C, Chaix B, Salze P, Casey R, Banos A, et al. Identifying built environmental patterns using cluster analysis and GIS: relationships with walking, cycling and body mass index in French adults. *Int J Behav Nutr Phys Act.* 2012;9:59.
33. McCormack GR, Friedenreich C, Sandalack BA, Giles-Corti B, Doyle-Baker PK, Shiell A. The relationship between cluster-analysis derived walkability and local recreational and transportation walking among Canadian adults. *Health Place.* 2012;18(5):1079-87.
34. McDonald K, Hearst M, Farbakhsh K, Patnode C, Forsyth A, Sirard J, et al. Adolescent physical activity and the built environment: a latent class analysis approach. *Health Place.* 2012;18(2):191-8.
35. Norman GJ, Adams MA, Kerr J, Ryan S, Frank LD, Roesch SC. A latent profile analysis of neighborhood recreation environments in relation to adolescent physical activity, sedentary time, and obesity. *J Public Health Manag Pract.* 2010;16(5):411-9.
36. Timperio A, Crawford D, Ball K, Salmon J. Typologies of neighbourhood environments and children's physical activity, sedentary time and television viewing. *Health Place.* 2017;43:121-7.

37. Hobbs M, Griffiths C, Green MA, Jordan H, Saunders J, McKenna J. Neighbourhood typologies and associations with body mass index and obesity: A cross-sectional study. *Prev Med.* 2018;111:351-7.
38. DeWeese RS, Ohri-Vachaspati P, Adams MA, Kurka J, Han SY, Todd M, et al. Patterns of food and physical activity environments related to children's food and activity behaviors: A latent class analysis. *Health Place.* 2018;49:19-29.
39. Poulsen MN, Glass TA, Pollak J, Bandeen-Roche K, Hirsch AG, Bailey-Davis L, et al. Associations of multidimensional socioeconomic and built environment factors with body mass index trajectories among youth in geographically heterogeneous communities. *Prev Med Rep.* 2019;15:100939.
40. Lambert M, Van Hulst A, O'Loughlin J, Tremblay A, Barnett TA, Charron H, et al. Cohort profile: the Quebec adipose and lifestyle investigation in youth cohort. *Int J Epidemiol.* 2012;41(6):1533-44.
41. Kuczmarski RJ, Ogden CL, Grummer-Strawn LM, Flegal KM, Guo SS, Wei R, et al. CDC growth charts: United States. *Adv Data.* 2000(314):1-27.
42. VanItallie TB, Yang MU, Heymsfield SB, Funk RC, Boileau RA. Height-normalized indices of the body's fat-free mass and fat mass: potentially useful indicators of nutritional status. *Am J Clin Nutr.* 1990;52(6):953-9.
43. Marshall WA, Tanner JM. Variations in the pattern of pubertal changes in boys. *Arch Dis Child.* 1970;45(239):13-23.
44. Marshall WA, Tanner JM. Variations in pattern of pubertal changes in girls. *Arch Dis Child.* 1969;44(235):291-303.
45. Paquet C, Cargo M, Kestens Y, Daniel M. Reliability of an instrument for direct observation of urban neighbourhoods. *Landscape Urban Planning.* 2010;97(3):194-201.
46. Landis JR, Koch GG. The Measurement of Observer Agreement for Categorical Data. *Biometrics.* 1977;33(1):159-74.
47. CRCHUM. 2013 [Available from: <http://megaphone.crchum.qc.ca/geonetwork/srv/en/main.home>.
48. Leslie E, Coffee N, Frank L, Owend N, Bauman A, Hugo G. Walkability of local communities: Using geographic information systems to objectively assess relevant environmental attributes. *Health Place.* 2007;13:111-22.
49. Ward JH. Hierarchical Grouping to Optimize an Objective Function. *Journal of the American Statistical Association.* 1963;58(301):236-44.
50. Tan P, Steinbach M, Kumar V. *Cluster Analysis: Basic Concepts and Algorithms.* Introduction to Data Mining. Boston: Addison-Wesley; 2005.
51. Holifield R, Porter M, Walker G. Introduction Spaces of Environmental Justice: Frameworks for Critical Engagement. *Antipode.* 2009;41(4):591-612.
52. Wang Z, Zhao L, Huang Q, Hong A, Yu C, Xiao Q, et al. Traffic-related environmental factors and childhood obesity: A systematic review and meta-analysis. *Obes Rev.* 2020.
53. Pabayo R, Belsky J, Gauvin L, Curtis S. Do area characteristics predict change in moderate-to-vigorous physical activity from ages 11 to 15 years? *Soc Sci Med.* 2011;72(3):430-8.
54. Larouche R, Chaput JP, Leduc G, Boyer C, Belanger P, LeBlanc AG, et al. A cross-sectional examination of socio-demographic and school-level correlates of children's school travel mode in Ottawa, Canada. *BMC Public Health.* 2014;14:497.

55. Oluyomi AO, Lee C, Nehme E, Dowdy D, Ory MG, Hoelscher DM. Parental safety concerns and active school commute: correlates across multiple domains in the home-to-school journey. *Int J Behav Nutr Phys Act.* 2014;11(1):32.
56. Vanhelst J, Beghin L, Salleron J, Ruiz JR, Ortega FB, De Bourdeaudhuij I, et al. A favorable built environment is associated with better physical fitness in European adolescents. *Prev Med.* 2013;57(6):844-9.
57. Timperio A, Crawford D, Telford A, Salmon J. Perceptions about the local neighborhood and walking and cycling among children. *Preventive Medicine.* 2004;38:39-47.
58. Timperio A, Salmon J, Telford A, Crawford D. Perceptions of local neighbourhood environments and their relationship to childhood overweight and obesity. *Int J Obes.* 2005;29(2):170-5.
59. Larouche R. Built Environment Features that Promote Cycling in School-Aged Children. *Curr Obes Rep.* 2015;4(4):494-503.
60. Timperio A, Reid J, Veitch J. Playability: Built and Social Environment Features That Promote Physical Activity Within Children. *Curr Obes Rep.* 2015;4(4):460-76.
61. Hume C, Timperio A, Salmon J, Carver A, Giles-Corti B, Crawford D. Walking and cycling to school. Predictors of increases among children and adolescents. *American Journal of Preventive Medicine.* 2009;36(3):195-200.
62. Carver A, Timperio AF, Crawford DA. Neighborhood road environments and physical activity among youth: the CLAN study. *J Urban Health.* 2008;85(4):532-44.
63. Timperio A, Jeffery RW, Crawford D, Roberts R, Giles-Corti B, Ball K. Neighbourhood physical activity environments and adiposity in children and mothers: a three-year longitudinal study. *Int J Behav Nutr Phys Act.* 2010;7:18.
64. Ghenadenik AE, Kakinami L, Van Hulst A, Henderson M, Barnett TA. Neighbourhoods and obesity: A prospective study of characteristics of the built environment and their association with adiposity outcomes in children in Montreal, Canada. *Prev Med.* 2018;111:35-40.
65. An R, Yang Y, Hoschke A, Xue H, Wang Y. Influence of neighbourhood safety on childhood obesity: a systematic review and meta-analysis of longitudinal studies. *Obes Rev.* 2017;18(11):1289-309.
66. Nordbo ECA, Nordh H, Raanaas RK, Aamodt G. Promoting activity participation and well-being among children and adolescents: a systematic review of neighborhood built-environment determinants. *JBIEvid Synth.* 2020;18(3):370-458.
67. Van Hecke L, Ghekiere A, Veitch J, Van Dyck D, Van Cauwenberg J, Clarys P, et al. Public open space characteristics influencing adolescents' use and physical activity: A systematic literature review of qualitative and quantitative studies. *Health Place.* 2018;51:158-73.
68. Harrison F, Jones AP, van Sluijs EM, Cassidy A, Bentham G, Griffin SJ. Environmental correlates of adiposity in 9-10 year old children: considering home and school neighbourhoods and routes to school. *Soc Sci Med.* 2011;72(9):1411-9.
69. Jennings A, Welch A, Jones AP, Harrison F, Bentham G, van Sluijs EM, et al. Local food outlets, weight status, and dietary intake: associations in children aged 9-10 years. *Am J Prev Med.* 2011;40(4):405-10.
70. Prentice AM, Jebb SA. Beyond body mass index. *Obesity Reviews.* 2001;2:141-7.
71. Weden MM, Bird CE, Escarce JJ, Lurie N. Neighborhood archetypes for population health research: is there no place like home? *Health Place.* 2011;17(1):289-99.

72. Wall MM, Larson NI, Forsyth A, Van Riper DC, Graham DJ, Story MT, et al. Patterns of obesogenic neighborhood features and adolescent weight: a comparison of statistical approaches. *Am J Prev Med.* 2012;42(5):e65-75.
73. Meyer KA, Boone-Heinonen J, Duffey KJ, Rodriguez DA, Kiefe CI, Lewis CE, et al. Combined measure of neighborhood food and physical activity environments and weight-related outcomes: The CARDIA study. *Health & place.* 2015;33:9-18.
74. Dunton GF, Liao Y, Almanza E, Jerrett M, Spruijt-Metz D, Pentz MA. Locations of joint physical activity in parent-child pairs based on accelerometer and GPS monitoring. *Annals of behavioral medicine : a publication of the Society of Behavioral Medicine.* 2013;45 Suppl 1:S162-72.
75. Jones M, Pebley AR. Redefining neighborhoods using common destinations: social characteristics of activity spaces and home census tracts compared. *Demography.* 2014;51(3):727-52.
76. Palmer JR, Espenshade TJ, Bartumeus F, Chung CY, Ozgencil NE, Li K. New approaches to human mobility: using mobile phones for demographic research. *Demography.* 2013;50(3):1105-28.
77. Kwan M-P. The uncertain geographic context problem. *Annals of the Association of American Geographers.* 2012;102(4):958-68.

Table 1. Description of area features by neighborhood type in the QUALITY cohort 2005-2008

	Type 1: <i>MODERATE WALKABILITY AND HIGH SAFETY</i>	Type 2: <i>LOW WALKABILITY AND HIGH SAFETY</i>	Type 3: <i>MODERATE WALKABILITY AND MODERATE SAFETY</i>	Type 4: <i>HIGH WALKABILITY AND MODERATE SAFETY</i>	Type 5: <i>MODERATE WALKABILITY AND LOW SAFETY</i>
	(n=132)	(n=115)	(n=64)	(n=108)	(n=66)
WALKABILITY					
Number of intersections, mean (SD)	70.77 (31.08)	60.58 (25.60)	70.94 (30.05)	119.76 (30.25)	80.50 (42.62)
Land use mix, mean (SD)	0.32 (0.12)	0.30 (0.13)	0.29 (0.13)	0.55 (0.10)	0.32 (0.12)
Density of private dwellings per hectare, % (n)	2.71	2.75	3.45	4.85	2.98
High density	2.27 (3)	2.61 (3)	17.19 (11)	87.96 (95)	13.64 (9)
Average-to-high density	24.24 (32)	25.22 (29)	40.63 (26)	10.19 (11)	21.21 (14)
Average density	31.06 (41)	30.43 (35)	20.31 (13)	0.93 (1)	25.76 (17)
Average-to-low density	27.27 (36)	28.70 (33)	14.06 (9)	0.93 (1)	28.79 (19)
Low density	15.15 (20)	13.04 (15)	7.81 (5)	0	10.61 (7)
Total length of streets with normal vehicular traffic at rush hour, km, mean (SD)	204.91 (89.49)	209.03 (92.48)	317.52 (140.73)	570.24 (188.84)	225.32 (128.70)
Presence of sidewalks, % (n)	2.03	2.16	2.59	4.72	2.29
All segments with sidewalks on both sides	0.76 (1)	0	12.50 (8)	77.78 (84)	10.61 (7)
High presence of sidewalks	17.42 (23)	13.04 (15)	17.19 (11)	19.44 (21)	16.67 (11)
Moderate presence of sidewalks	18.18 (24)	33.04 (38)	23.44 (15)	0.93 (1)	13.64 (9)
Low presence of sidewalks	11.36 (15)	10.43 (12)	10.94 (7)	0.93 (1)	9.09 (6)
No segments with sidewalks	52.27 (69)	43.48 (50)	35.94 (23)	0.93 (1)	50.00 (33)
Proportion of segments with ≥ 1 pedestrians aid, mean (SD)	43.80 (28.58)	71.54 (20.57)	64.47 (24.94)	53.18 (26.72)	46.36 (28.18)
Number of parks, mean (SD)	2.11 (1.29)	0.80 (0.77)	1.41 (1.26)	3.68 (1.30)	2.80 (1.50)
Park area ratio, % (n)	1.95	1.25	1.51	2.59	2.15
< 0.01	22.73 (30)	75.65 (87)	51.56 (33)	1.85 (2)	10.61 (7)
0.1 to < 0.05	59.85 (79)	23.48 (27)	45.31 (29)	37.04 (40)	63.64 (42)
≥ 0.05	17.42 (23)	0.87 (1)	3.13 (2)	61.11 (66)	25.76 (17)

VEHICULAR TRAFFIC-RELATED SAFETY

Neighborhood streets with heavy vehicular traffic at rush hour, % (n)

< 1%	100 (132)	100 (115)	0	100 (108)	0
1 to 5%	0	0	93.75 (60)	0	48.48 (32)
≥ 5%	0	0	6.25 (4)	0	51.52 (34)
Total length of streets with heavy vehicular traffic at rush hour, % (n)	0.52	0.53	2.80	1.47	2.79
0 km	52.27 (69)	54.78 (63)	0	0.93 (1)	0
0.1 to 1 km	43.18 (57)	37.39 (43)	0	50.93 (55)	0
1.1 to 5 km	4.44 (6)	7.83 (9)	20.31 (13)	48.15 (52)	21.21 (14)
≥ 5 km	0	0	79.69 (51)	0	78.79 (52)
Proportion of streets with ≥ 1 traffic calming measure, mean (SD)	13.94 (17.30)	33.36 (24.93)	29.38 (27.53)	40.47 (25.87)	9.85 (16.11)

OTHER

NDVI, mean (SD)	0.37 (0.07)	0.33 (0.06)	0.32 (0.06)	0.24 (0.06)	0.37 (0.05)
Presence of signs of social disorder, % (n)	18.18 (24)	23.48 (27)	39.06 (25)	75.93 (82)	15.15 (10)
Neighborhood residents with university education, %, mean (SD)*	26.89 (14.61)	25.98 (13.038)	27.20 (11.35)	33.52 (13.16)	28.13 (16.24)

* Variables not used in cluster analysis to define neighborhood types
 NDVI, Normalized difference vegetation index

Table 2. Baseline characteristics of 485 children from the QUALITY cohort, by neighborhood type, 2005-2008

	Type 1: <i>MODERATE WALKABILITY AND HIGH SAFETY</i>	Type 2: <i>LOW WALKABILITY AND HIGH SAFE</i>	Type 3: <i>MODERATE WALKABILITY AND MODERATE SAFETY</i>	Type 4: <i>HIGH WALKABILITY AND MODERATE SAFETY</i>	Type5: <i>MODERATE WALKABILITY AND LOW SAFETY</i>
	(n=132)	(n=115)	(n=64)	(n=108)	(n=66)
Age, years, mean (SD)	9.58 (0.94)	9.55 (0.93)	9.68 (0.90)	9.55 (0.89)	9.83 (0.84)
Boys, % (n)	53.79 (71)	53.04 (61)	37.50 (24)	63.89 (69)	56.06 (37)
Puberty initiated, % (n)	20.61 (27)	23.48 (27)	23.44 (15)	23.15 (25)	30.30 (20)
Child's BMI Z-score, mean (SD)	0.58 (1.05)	0.66 (1.04)	0.65 (1.14)	0.72 (1.10)	1.10 (0.93)
FMI, kg/m ² , mean (SD)	5.03 (2.93)	5.30 (3.38)	5.74 (3.38)	5.59 (3.75)	6.73 (3.83)
WC, cm, mean (SD)	65.77 (10.09)	66.52 (11.85)	67.75 (12.26)	68.25 (13.24)	72.76 (14.30)
% central fat mass, mean (SD)	40.44 (4.71)	39.85 (5.48)	41.85 (6.12)	40.55 (5.76)	42.72 (5.09)
Mother's BMI, kg/m ² , mean (SD)	29.25 (6.35)	29.77 (6.61)	29.76 (7.30)	29.77 (6.03)	29.52 (7.52)
Father's BMI, kg/m ² , mean (SD)	30.59 (5.27)	30.75 (5.072)	30.54 (6.46)	29.89 (5.36)	32.84 (6.43)

BMI, body mass index; FMI, fat mass index; SD, standard deviation; WC, waist circumference

Table 3. Results of fully-adjusted linear regression models predicting adiposity at baseline among 485 children aged 8-10 years from the QUALITY cohort, using neighborhood types as predictors, 2005-2008

	Overall Adiposity				Central Adiposity			
	BMI Z-Score		FMI (kg/m ²)		Waist Circumference (cm)		% Central Fat Mass	
	β (95% CI)		β (95% CI)		β (95% CI)		β (95% CI)	
Intercept	1.53	0.45, 2.62	-6.58	-10.51, 2.65	13.49	-0.43, 27.40	26.47	19.90, 33.04
Neighborhood type								
Type 1: <i>MODERATE WALKABILITY, HIGH SAFETY</i>	Referent		Referent		Referent		Referent	
Type 2: <i>LOW WALKABILITY, HIGH SAFETY</i>	0.05	-0.20, 0.30	0.17	-0.61, 0.94	0.42	-2.32, 3.17	-0.68	-1.97, 0.62
Type 3: <i>MODERATE WALKABILITY, MODERATE SAFETY</i>	0.10	-0.19, 0.40	0.51	-0.43, 1.44	1.86	-1.43, 5.14	1.05	-0.52, 2.61
Type 4: <i>HIGH WALKABILITY, MODERATE SAFETY</i>	0.14	-0.12, 0.40	0.67	-0.14, 1.48	2.44	-0.42, 5.30*	0.41	-0.94, 1.76
Type 5: <i>MODERATE WALKABILITY, LOW SAFETY</i>	0.41	0.11, 0.71***	1.22	0.28, 2.16**	4.90	1.60, 8.21***	1.60	0.03, 3.17**
Child's age	-0.10	-0.21, 0.01	0.43	0.09, 0.77**	2.61	1.41, 3.82***	0.58	0.01, 1.14**
Sex (boys)	0.20	-0.001, 0.40	-0.63	-1.26, -0.01**	1.64	-0.56, 3.84	-1.45	-2.49, -0.40***
Puberty initiated	0.42	0.16, 0.67***	1.33	0.54, 2.12***	5.37	2.56, 8.17***	0.93	-0.39, 2.25
University-educated residents, %	-0.01	-0.08, 0.06	-0.01	-0.03, 0.01	-0.02	-0.10, 0.05	-0.01	-0.04, 0.03
Mother's BMI, kg/m ²	0.04	0.03, 0.05***	0.12	0.08, 0.16***	0.39	0.23, 0.54***	0.17	0.10, 0.25***
Father's BMI, kg/m ²	0.05	0.03, 0.06***	0.13	0.08, 0.18***	0.47	0.29, 0.64***	0.13	0.05; 0.22***

* <0.10 ** <0.05 *** <0.01

Abbreviations: BMI, body mass index; FMI, fat mass index

Table 4. Results of fully-adjusted linear regression models predicting adiposity among 298 participants aged 15-17 from the QUALITY cohort who resided at the same address both at baseline and follow-up, using neighborhood types as predictors, 2005–2016

	Overall Adiposity				Central Adiposity			
	BMI Z-Score		FMI (kg/m ²)		Waist Circumference (cm)		% Central Fat Mass	
	β (95% CI)		β (95% CI)		β (95% CI)		β (95% CI)	
Intercept	-0.64	-3.03, 1.75	-0.11	-9.05, 8.82	38.46	7.00, 69.92	26.01	13.73, 38.29
Neighborhood type								
Type 1: <i>MODERATE WALKABILITY, HIGH SAFETY</i>		Referent		Referent		Referent		Referent
Type 2: <i>LOW WALKABILITY, HIGH SAFETY</i>	-0.10	-0.43, 0.23	-0.08	-1.29, 1.13	0.09	-4.22, 4.40	-0.63	-2.29, 1.03
Type 3: <i>MODERATE WALKABILITY, MODERATE SAFETY</i>	-0.28	-0.66, 0.09	-0.71	-2.10, 0.68	-1.41	-6.36, 3.54	-1.60	-3.51, 0.31*
Type 4: <i>HIGH WALKABILITY, MODERATE SAFETY</i>	0.02	-0.33, 0.38	-0.01	-1.32, 1.31	0.47	-4.20, 5.14	-0.45	-2.26, 1.36
Type 5: <i>MODERATE WALKABILITY, LOW SAFETY</i>	0.25	-0.15, 0.65	1.42	-0.06, 2.90*	5.06	-0.24, 10.36*	0.12	-1.92, 2.16
Child's age	-0.06	-0.19, 0.07	0.11	-0.36, 0.59	0.99	-0.69, 2.67	1.04	0.39, 1.69***
Sex (boys)	0.07	-0.18, 0.31	-2.59	-3.49, -1.70***	6.54	3.35, 9.73***	2.16	0.92, 3.39***
University-educated residents, %	-0.03	-0.12, 0.06	-0.18	-0.51, 0.14	-0.22	-1.37, 0.93	-0.19	-0.64, 0.25
Mother's BMI, kg/m ²	0.05	0.03, 0.07***	0.13	0.06, 0.21***	0.38	0.11, 0.65***	0.06	-0.04, 0.16
Father's BMI, kg/m ²	0.03	0.01, 0.05**	0.10	0.01, 0.19**	0.34	0.02, 0.67**	0.06	-0.07, 0.19

* <0.10 ** <0.05 *** <0.01

Abbreviations: BMI, body mass index; FMI, fat mass index

Figure 1. Diagram detailing the number of participants by time of survey and reasons for inclusion/exclusion from analytical samples

