



**HAL**  
open science

# On Abel-Jacobi Maps of Lagrangian Families

Chenyu Bai

► **To cite this version:**

Chenyu Bai. On Abel-Jacobi Maps of Lagrangian Families. *Mathematische Zeitschrift*, 2023, 304, 10.1007/s00209-023-03292-9 . hal-04155808

**HAL Id: hal-04155808**

**<https://hal.sorbonne-universite.fr/hal-04155808>**

Submitted on 7 Jul 2023

**HAL** is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers.

L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.



Distributed under a Creative Commons Attribution - NonCommercial 4.0 International License

# On Abel-Jacobi Maps of Lagrangian Families

Chenyu Bai

## Abstract

We study in this article the cohomological properties of Lagrangian families on projective hyper-Kähler manifolds. First, we give a criterion for the vanishing of Abel-Jacobi maps of Lagrangian families. Using this criterion, we show that under a natural condition, if the variation of Hodge structures on the degree 1 cohomology of the fibers of the Lagrangian family is maximal, its Abel-Jacobi map is trivial. We also construct Lagrangian families on generalized Kummer varieties whose Abel-Jacobi map is not trivial, showing that our criterion is optimal.

## 0 Introduction

Let  $X$  be a projective hyper-Kähler manifold [1], that is, a simply connected complex projective manifold whose space of holomorphic 2-forms is generated by a nowhere degenerate 2-form  $\sigma_X$ . The dimension of  $X$  is an even number  $2n$ . A Lagrangian subvariety  $L$  of  $X$  is a dimension  $n$  irreducible possibly singular subvariety of  $X$  such that, denoting  $j: \tilde{L} \rightarrow L \hookrightarrow X$  a desingularisation of  $L$ ,  $j^*\sigma_X = 0$  in  $H^0(\tilde{L}, \Omega_{\tilde{L}}^2)$ .

In this article, we will be considering Lagrangian families of a hyper-Kähler manifold  $X$ .

**Definition 0.1.** A Lagrangian family of a hyper-Kähler manifold  $X$  is a diagram

$$\begin{array}{ccc} \mathcal{L} & \xrightarrow{q} & X \\ \downarrow p & & \\ B & & \end{array} \quad (1)$$

where  $p$  is flat and projective,  $\mathcal{L}$  and  $B$  are connected quasi-projective manifolds and  $q$  maps birationally the general fiber  $L_b := p^{-1}(b)$ ,  $b \in B$ , to a Lagrangian subvariety of  $X$ . In what follows, we will denote  $j_b$  the composition  $L_b \hookrightarrow \mathcal{L} \rightarrow X$ .

Lagrangian families were studied by Voisin in [14, 15] in different contexts, as generalizations of Lagrangian fibrations. While Lagrangian fibrations do not exist in general on projective hyper-Kähler manifolds (as this forces the Picard number to be at least 2), Lagrangian families (and even Lagrangian coverings for which  $q$  is dominant) are relatively common. See, for example, constructions in [8, 11] on Lagrangian families (coverings) on Fano varieties of lines of cubic fourfolds and on Hilbert schemes of K3 surfaces. Works of Voisin [14, 15] indicate that the existence of Lagrangian coverings implies important properties of the hyper-Kähler manifold in question. For example, it is shown in [15] that a very general projective hyper-Kähler fourfold admitting a Lagrangian covering satisfies Lefschetz standard conjecture for degree 2 cohomology. Studies of examples of Lagrangian families lead us to consider the following problem, which is motivated in the article [13].

**Problem 0.2.** Consider a Lagrangian family of a hyper-Kähler manifold  $X$  of dimension  $2n$  given by a diagram as in (1). What can be said of the map

$$\begin{array}{ccc} \psi_{\mathcal{L}}: B & \rightarrow & CH^n(X) \ ? \\ b & \mapsto & q_*(L_b) \end{array} \quad (2)$$

Let us first explain the relation between this problem and article [13]. In [13], the following conjecture is proposed and proved for some cases.

**Conjecture 0.3.** Let  $L$  and  $L'$  be two *constant cycle* Lagrangian subvarieties of a hyper-Kähler manifold  $X$ . If the cohomological classes  $[L] = [L']$  in  $H^{2n}(X, \mathbb{Q})$ , then  $L$  is rationally equivalent to  $L'$  as algebraic cycles in  $X$ .

Here, a subvariety  $Z$  of a smooth projective variety  $X$  is called a *constant cycle* subvariety, if any two points  $z, z'$  in  $Z$  are rationally equivalent in  $X$ , or equivalently, if the image of the Gysin map  $i_* : CH_0(Z) \rightarrow CH_0(X)$  is  $\mathbb{Z}$  where  $i : Z \hookrightarrow X$  is the inclusion map. The notion of constant cycle subvarieties is first proposed in [7] for curves in K3 surfaces. It is then studied in a more general setting in [13]. The condition constant cycle is a strong condition for Lagrangian subvarieties. In contrast with Lagrangian subvarieties, constant cycle Lagrangian subvarieties cannot deform into families.

**Lemma 0.4.** *Constant cycle Lagrangian subvarieties of  $X$  are rigid as constant cycle subvarieties.*

*Proof* Following the notations in [13], let

$$S_n X := \{x \in X : \text{the rational equivalence orbit of } x \text{ has dimension } \geq n\}.$$

As is shown in [13, Theorem 1.3],  $S_n X$  is a countable union of irreducible varieties of dimension  $\leq n$  and constant cycle Lagrangian subvarieties of  $X$  are exactly irreducible components of  $S_n X$  of dimension  $n$ . Hence, constant cycle Lagrangian subvarieties of  $X$  are rigid.  $\square$

One may now wonder if the condition *constant cycle* can be dropped in Conjecture 0.3. This motivates us to think about the Problem 0.2. Since the Lagrangian subvarieties in the Lagrangian family have the same cohomological class, if the map  $\Psi_{\mathcal{L}}$  defined in Problem 0.2 is nonconstant, then we cannot drop the condition *constant cycle* in Conjecture 0.3, presenting the subtlety of Conjecture 0.3.

A weaker invariant of algebraic cycles in a projective manifold is the Abel-Jacobi invariant [16, Chapter 12]. Let us denote  $\Phi_X^n : CH^n(X)_{\text{hom}} \rightarrow J^{2n-1}(X)$  the Abel-Jacobi map of  $X$ . We will recall the definitions and basic properties of the Abel-Jacobi invariant and the Abel-Jacobi maps in the beginning of Section 1. If two cycles homologous to 0 are rationally equivalent, then they have the same Abel-Jacobi invariant in the intermediate Jacobians (see Proposition 1.2). Problem 0.2 thus motivates the following question.

**Problem 0.5.** Consider a Lagrangian family of a hyper-Kähler manifold  $X$  of dimension  $2n$  given by a diagram as in (1). Let  $0 \in B$  be a point. Under which conditions is the Abel-Jacobi map

$$\begin{aligned} \Psi_{\mathcal{L}}^{AJ} : B &\rightarrow J^{2n-1}(X) \\ b &\mapsto \Phi_X^n(q_*(L_b - L_0)) \end{aligned} \quad (3)$$

trivial?

Note that  $\Psi_{\mathcal{L}}^{AJ}(b) = \Phi_X^n(\Psi_{\mathcal{L}}(b) - \Psi_{\mathcal{L}}(0))$ . Problem 0.5 can be viewed as a first step to study Problem 0.2.

In this article, we first give a criterion for the vanishing of the Abel-Jacobi map (3) for Lagrangian families of a hyper-Kähler manifold (see also Proposition 1.5).

**Proposition 0.6.** *Consider a Lagrangian family of a hyper-Kähler manifold  $X$  of dimension  $2n$  as in (1), satisfying the following condition :*

$\clubsuit$  *For general  $b \in B$ , the contraction by  $q^* \sigma_X$  gives an isomorphism  $q^* \sigma_X : T_{B,b} \xrightarrow{\cong} H^0(L_b, \Omega_{L_b})$ .*

Then the Abel-Jacobi map (3) is trivial if and only if for general  $b \in B$ , the restriction map

$$j_b^* : H^{2n-1}(X, \mathbb{Q}) \rightarrow H^{2n-1}(L_b, \mathbb{Q})$$

is zero.

The condition  $\clubsuit$  is natural. According to [11, Proposition 2.4], the deformations of a *smooth* Lagrangian subvariety are non-obstructed, and a local deformation is still Lagrangian. Therefore, if we take  $(B, b)$  to be a germ of the Hilbert scheme of deformations of a *smooth* Lagrangian subvariety  $L \subset X$ , and  $\mathcal{L} \rightarrow B$  the corresponding family, then condition  $\clubsuit$  holds since  $T_{B,b} \cong H^0(L_b, N_{L_b/X})$  by unobstructedness and  $\lrcorner\sigma_X : H^0(L_b, N_{L_b/X}) \rightarrow H^0(L_b, \Omega_{L_b})$  is an isomorphism for a smooth Lagrangian variety.

Using this criterion, we give a response to Problem 0.5.

**Theorem 0.7.** *Consider a Lagrangian family of a hyper-Kähler manifold  $X$  of dimension  $2n$  given by a diagram as in (1), satisfying condition  $\clubsuit$ . Assume that the variation of Hodge structures on the degree 1 cohomology of the fibers of  $p : \mathcal{L} \rightarrow B$  is maximal, i.e., the period map*

$$\begin{aligned} \mathcal{P} : B &\rightarrow \text{Gr}(h^{1,0}(L), H^1(L, \mathbb{C})) \\ b &\mapsto H^{1,0}(L_b) \subset H^1(L_b, \mathbb{C}) \cong H^1(L, \mathbb{C}), \end{aligned} \quad (4)$$

where  $L$  is a general fiber of  $p : \mathcal{L} \rightarrow B$ , is generically a local immersion. Then the Abel-Jacobi map (3) is trivial.

This response to Problem 0.5 is conditional. However, it can be shown that the condition “maximal variation of Hodge structures” cannot be dropped. In fact, we construct in Section 4 Lagrangian families satisfying  $\clubsuit$  for which the Abel-Jacobi map is shown to be nontrivial using Proposition 0.6. The variation of weight 1 Hodge structures of the constructed Lagrangian families is not maximal.

In Section 3, we shall explore under which conditions the variation of weight 1 Hodge structures is maximal. Let  $H^2(X, \mathbb{Q})_{ir}$  be the orthogonal complement of  $NS(X)_{\mathbb{Q}}$  in  $H^2(X, \mathbb{Q})$  with respect to the Beauville-Bogomolov-Fujiki form  $q$  of  $X$  (see [1]) and let  $b_2(X)_{ir}$  be the dimension of  $H^2(X, \mathbb{Q})_{ir}$ . We prove the following result (see also Proposition 3.2):

**Proposition 0.8.** *Consider a Lagrangian family of a hyper-Kähler manifold  $X$  of dimension  $2n$  given by a diagram as in (1), satisfying condition  $\clubsuit$ . Assume that the Mumford-Tate group of the Hodge structure  $H^2(X, \mathbb{Q})$  is maximal, i.e. it is the special orthogonal group of  $(H^2(X, \mathbb{Q})_{ir}, q)$ , and assume that  $b_2(X)_{ir} \geq 5$ . If  $h^{1,0}(L_b)$  is smaller than  $2^{\lfloor \frac{b_2(X)_{ir}-3}{2} \rfloor}$ , then the variation of weight 1 Hodge structures of  $p$  is maximal.*

**Corollary 0.9.** *Under the same assumptions as in Proposition 0.8, the Abel-Jacobi map (3) is trivial.*

Let  $p : \mathcal{L} \rightarrow B, q : \mathcal{L} \rightarrow X$  be a Lagrangian family. Up to shrinking  $B$ , we may assume that the map  $p : \mathcal{L} \rightarrow B$  is smooth. Let

$$\pi : \mathcal{A} := \text{Alb}(\mathcal{L}/B) \rightarrow B$$

be the relative Albanese variety of  $p : \mathcal{L} \rightarrow B$ . In the proof of Theorem 0.7 and Proposition 0.8, we use a similar construction to those in [9, 14] to get a holomorphic 2-form  $\sigma_{\mathcal{A}}$  on  $\mathcal{A}$ . If we assume the condition  $\clubsuit$ ,  $\pi : \mathcal{A} \rightarrow B$  is a Lagrangian fibration with respect to  $\sigma_{\mathcal{A}}$  (see Section 2). It is interesting to notice that, by this construction, under condition  $\clubsuit$ , we can translate the problem concerning Lagrangian families to a problem concerning Lagrangian fibrations. However, the total space of the Lagrangian fibration is no longer a hyper-Kähler manifold, but a completely integrable system over an open subset of the base, as studied for instance in [6, Chapter 7].

The organisation of the article is as follows. In Section 1, we prove Proposition 0.6. In Section 2, we construct a Lagrangian fibration structure on the relative Albanese variety and use it to prove Theorem 0.7. In Section 3, we discuss the condition on the maximality of the variation of Hodge structures. In Section 4, we construct Lagrangian families satisfying  $\clubsuit$  whose Abel-Jacobi map is nontrivial, showing that Theorem 0.7 is optimal.

## Acknowledgement

I would like to thank Claire Voisin for sharing her ideas with me. This paper would not be finished without her insight, instructions and encouragement. Many constructions and examples in the article are fruits of friendly and interesting discussions with her. I would also like to thank Fabrizio Anella, Peter Yi Wei, Mauro Varesco and Tim Ryan for helpful discussions. I thank the referee for his/her wonderful refereeing work and for numerous constructive comments and suggestions.

This work was done during the preparation of my PhD thesis. I would like to thank the *Institut de Mathématiques de Jussieu - Paris Rive Gauche* for marvelous research environment. The thesis is supported by the ERC Synergy Grant HyperK (Grant agreement No. 854361).

## 1 A Criterion

In this section, we establish a criterion for the vanishing of the Abel-Jacobi map (3). Let us first recall the Abel-Jacobi invariant and the Abel-Jacobi map defined by Griffiths. We follow the presentation in [16, Chapter 12]. Throughout this presentation only,  $X$  is a compact Kähler manifold.

**Definition 1.1.** The  $k$ -th intermediate Jacobian  $J^{2k-1}(X)$  is the complex torus

$$J^{2k-1}(X) = H^{2k-1}(X, \mathbb{C}) / (F^k H^{2k-1}(X) \oplus H^{2k-1}(X, \mathbb{Z})).$$

The Abel-Jacobi map

$$\begin{aligned} \Phi_X^k : CH^n(X)_{hom} &\rightarrow J^{2n-1}(X) \\ Z = \partial\Gamma &\mapsto \int_{\Gamma} \end{aligned}$$

associates to any cycle  $Z = \partial\Gamma$  homologous to zero the current of integration along  $\Gamma$  (See [16, 12.1.2]). For an algebraic cycle  $Z$  of codimension  $k$  in  $X$ , the associated an element  $\Phi_X^k(Z)$  in  $J^{2k-1}(X)$  [16, 12.1.2] is called the Abel-Jacobi invariant of the cycle  $Z$ . The triviality of Abel-Jacobi invariant is a weaker condition than the rational equivalence.

**Proposition 1.2.** *Let  $Z$  be an algebraic cycle that is rationally equivalent to 0 in  $X$ . Then the Abel-Jacobi invariant of  $Z$  is 0.*

*Proof* It is proven in [16, Lemme 21.19]. □

Let  $\mathcal{Z} \subset B \times X$  be a flat family of subvarieties of codimension  $k$  in  $X$ . More precisely,  $B$  is a connected complex manifold, and  $\mathcal{Z}$  is a subvariety of codimension  $k$  in  $B \times X$ , flat over  $B$ . Let  $p : \mathcal{Z} \rightarrow B$  and  $q : \mathcal{Z} \rightarrow X$  be the projection maps to the two components. Let  $0 \in B$  be a reference point.

**Definition 1.3.** The Abel-Jacobi map of the family  $\mathcal{Z} \subset B \times X$  with respect to the reference point  $0 \in B$  is the map

$$\begin{aligned} \Psi_{\mathcal{Z}}^{AJ} : B &\rightarrow J^{2k-1}(X) \\ b &\mapsto \Phi_X^k(q_* p^*(b) - q_* p^*(0)). \end{aligned}$$

**Theorem 1.4** (Griffiths). *We have the following properties about the Abel-Jacobi map.*

(i) *The Abel-Jacobi map  $\Psi_{\mathcal{Z}}^{AJ} : B \rightarrow J^{2k-1}(X)$  is holomorphic.*

(ii) *The image of the differential of  $\Psi_{\mathcal{Z}}^{AJ}$  at any point lies in  $H^{k-1,k}(X) \subset H^{2k-1}(X, \mathbb{C})$ .*

*Proof* (i) is proven in [16, Théorème 12.4] and (ii) is proven in [16, Corollary 12.19 and Remarque 12.20]. □

This concludes the presentation of Abel-Jacobi invariants and Abel-Jacobi maps. From now on, let  $X$  be a hyper-Kähler manifold of dimension  $2n$ . With the notation  $j_b : L_b \rightarrow X$  as in the introduction, we prove

**Proposition 1.5.** Consider a Lagrangian family of hyper-Kähler manifold  $X$  of dimension  $2n$  given by a diagram as in (1).

(a) If for general  $b \in B$ , the restriction map

$$j_b^* : H^{2n-1}(X, \mathbb{Q}) \rightarrow H^{2n-1}(L_b, \mathbb{Q})$$

is zero, then the Abel-Jacobi map (3) is trivial.

(b) If condition  $\clubsuit$  holds (see Proposition 0.6), then the converse of (a) holds.

**Remark 1.6.** The cohomology group  $H^{2n-1}(L_b, \mathbb{Q})$  has a Hodge structure of weight  $2n-1$  and level 1. By Hodge symmetry and using the fact  $j_b^* : H^{2n-1}(X, \mathbb{Q}) \rightarrow H^{2n-1}(L_b, \mathbb{Q})$  is a morphism of Hodge structures, we conclude that  $j_b^* : H^{2n-1}(X, \mathbb{Q}) \rightarrow H^{2n-1}(L_b, \mathbb{Q})$  is zero, if and only if  $j_b^* : H^{n-1,n}(X) \rightarrow H^{n-1,n}(L_b)$  is zero.

*Proof* Let  $d\Psi_{\mathcal{L},b}^{AJ} : T_{B,b} \rightarrow H^n(X, \Omega_X^{n+1})$  denote the differential of the Abel-Jacobi map  $\Psi_{\mathcal{L}}^{AJ}$  at point  $b \in B$  (Theorem 1.4). Let  $j_{b*} : H^0(L_b, \Omega_{L_b}) \rightarrow H^n(X, \Omega_X^{n+1})$  be the Gysin map, which is the Serre dual to the following composition

$$j_b^* : H^n(X, \Omega_X^{n-1}) \rightarrow H^n(L_b, \Omega_{\mathcal{L}|L_b}^{n-1}) \rightarrow H^n(L_b, \Omega_{L_b}^{n-1}). \quad (5)$$

By the above remark, the proposition follows from the following lemma and the fact that

$$\cup \sigma_X : H^n(X, \Omega_X^{n-1}) \rightarrow H^n(X, \Omega_X^{n+1})$$

is an isomorphism since  $\wedge \sigma_X : \Omega_X^{n-1} \rightarrow \Omega_X^{n+1}$  is a vector bundle isomorphism.  $\square$

**Lemma 1.7.** The following diagram is commutative:

$$\begin{array}{ccc} T_{B,b} & \xrightarrow{d\Psi_{\mathcal{L},b}^{AJ}} & H^n(X, \Omega_X^{n-1}) \\ \downarrow \cup q^* \sigma_X & & \downarrow \cup \sigma_X \\ H^0(L_b, \Omega_{L_b}) & \xrightarrow{j_{b*}} & H^n(X, \Omega_X^{n+1}). \end{array} \quad (6)$$

*Proof* We are going to show that the Serre dual of the diagram (6) is commutative.

Let  $L^\bullet \Omega_{\mathcal{L}|L_b}^i$  be the Leray filtration [16, Chapter 16] induced on the vector bundle  $\Omega_{\mathcal{L}|L_b}^i$  by the exact sequence

$$0 \rightarrow p^* \Omega_{B,b} \rightarrow \Omega_{\mathcal{L}|L_b} \rightarrow \Omega_{L_b} \rightarrow 0,$$

and defined by  $L^j \Omega_{\mathcal{L}|L_b}^i = p^* \Omega_{B,b}^j \wedge \Omega_{\mathcal{L}|L_b}^{i-j}$ . Since  $L_b$  is supposed to be Lagrangian,  $q^* \sigma_X \in H^0(L_b, L^1 \Omega_{\mathcal{L}|L_b}^2)$  and thus the cup product

$$\cup q^* \sigma_X : \Omega_{\mathcal{L}|L_b}^\bullet \rightarrow \Omega_{\mathcal{L}|L_b}^{\bullet+2}$$

sends  $L^k \Omega_{\mathcal{L}|L_b}^\bullet$  to  $L^{k+1} \Omega_{\mathcal{L}|L_b}^{\bullet+2}$ . Denoting  $\overline{q^* \sigma_X}$  the image of  $q^* \sigma_X$  in  $H^0(L_b, Gr_L^1 \Omega_{\mathcal{L}|L_b}^2) \cong H^0(L_b, \Omega_{L_b}) \otimes \Omega_{B,b}$ , this implies the existence of the following commutative diagram

$$\begin{array}{ccc} L^1 \Omega_{\mathcal{L}|L_b}^{n+1} = \Omega_{\mathcal{L}|L_b}^{n+1} & \longrightarrow & K_{L_b} \otimes p^* \Omega_{B,b} = Gr_L^1 \Omega_{\mathcal{L}|L_b}^{n+1} \\ \cup q^* \sigma_X \uparrow & & \cup \overline{q^* \sigma_X} \uparrow \\ L^0 \Omega_{\mathcal{L}|L_b}^{n-1} = \Omega_{\mathcal{L}|L_b}^{n-1} & \longrightarrow & \Omega_{L_b}^{n-1} = Gr_L^0 \Omega_{\mathcal{L}|L_b}^{n-1}, \end{array} \quad (7)$$

where  $K_{L_b}$  is the canonical bundle of  $L_b$ . Taking the  $n$ -th cohomology of (7) and combine it with  $q^* : H^n(X, \Omega_X^\bullet) \rightarrow H^n(L_b, \Omega_{\mathcal{L}|L_b}^\bullet)$ , we get the following commutative ladder

$$\begin{array}{ccccc} H^n(X, \Omega_X^{n+1}) & \xrightarrow{q^*} & H^n(L_b, \Omega_{\mathcal{L}|L_b}^{n+1}) & \longrightarrow & H^n(L_b, K_{L_b} \otimes p^* \Omega_{B,b}) \cong \Omega_{B,b} \\ \cup \sigma_X \uparrow & & \cup q^* \sigma_X \uparrow & & \cup \overline{q^* \sigma_X} \uparrow \\ H^n(X, \Omega_X^{n-1}) & \xrightarrow{q^*} & H^n(L_b, \Omega_{\mathcal{L}|L_b}^{n-1}) & \longrightarrow & H^n(L_b, \Omega_{L_b}^{n-1}). \end{array} \quad (8)$$

**Lemma 1.8.** *The composite in the first row of the diagram (8) coincides with the dual of  $d\Psi_{\mathcal{L},b}$ .*

*Proof* Let  $\bar{p} : \bar{\mathcal{L}} \rightarrow \bar{B}$ ,  $\bar{q} : \bar{\mathcal{L}} \rightarrow X$  be a relative completion of  $p : \mathcal{L} \rightarrow B$  with respect to the morphism  $q : \mathcal{L} \rightarrow X$ . More precisely,  $\bar{B}$  is a smooth projective variety,  $\bar{p}$  is a flat morphism extending  $p$ , and  $\bar{q}$  is a morphism extending  $q$ . The extended family has a Abel–Jacobi map  $\Psi_{\bar{\mathcal{L}}}^{AJ} : \bar{B} \rightarrow J^{2n-1}(X)$  that induces a morphism of complex tori  $\psi : \text{Alb}(\bar{B}) \rightarrow J^{2n-1}(X)$  whose differential is given by the morphism of Hodge structures ([16, Théorème 12.17])

$$\bar{q}_* \bar{p}^* : H^{2d-1}(\bar{B}, \mathbb{Z}) \rightarrow H^{2n-1}(X, \mathbb{Z}),$$

where  $d = \dim B$ . It is well-known that the differential of the Albanese map  $\text{alb} : \bar{B} \rightarrow \text{Alb}(\bar{B})$  at a point  $b \in B$  is given by the dual of the evaluation map  $ev_b : H^0(\bar{B}, \Omega_{\bar{B}}) \rightarrow \Omega_{\bar{B},b}$ . Hence, the dual of  $d\Psi_{\bar{\mathcal{L}},b}^{AJ}$  is given by the correspondance  $\bar{p}_* \bar{q}^* : H^n(X, \Omega_X^{n+1}) \rightarrow H^0(\bar{B}, \Omega_{\bar{B}})$  composed with the evaluation map  $ev_b : H^0(\bar{B}, \Omega_{\bar{B}}) \rightarrow \Omega_{\bar{B},b} \cong \Omega_{B,b}$ . The domain  $\bar{B}$  can be restricted to  $B$  before the evaluation map  $ev_b : H^0(\bar{B}, \Omega_{\bar{B}}) \rightarrow \Omega_{\bar{B},b}$ . Therefore, the dual of  $d\Psi_{\mathcal{L},b}^{AJ}$  is given by the correspondance  $p_* q^* : H^n(X, \Omega_X^{n+1}) \rightarrow H^0(B, \Omega_B)$  composed with the evaluation map  $ev_b : H^0(B, \Omega_B) \rightarrow \Omega_{B,b}$ . The Gysin map  $p_* : H^n(\mathcal{L}, \Omega_{\mathcal{L}}^{n+1}) \rightarrow H^0(B, \Omega_B)$  is given by the Leray filtration. Taken together, the dual of  $d\Psi_{\mathcal{L},b}^{AJ}$  is given by

$$H^n(X, \Omega_X^{n+1}) \xrightarrow{q^*} H^n(\mathcal{L}, \Omega_{\mathcal{L}}^{n+1}) \rightarrow H^n(\mathcal{L}, K_{\mathcal{L}/B} \otimes p^* \Omega_B) \rightarrow H^0(B, R^n p_* \Omega_{\mathcal{L}/B} \otimes \Omega_B) \xrightarrow{ev_b} \Omega_{B,b}.$$

Since the restriction to the fiber  $L_b$  and taking the Leray filtration are commutative processes, the above composite of maps is equal to the one that takes the restriction to the fiber  $L_b$  first and then takes the Leray filtration. The latter one is exactly the first row of the diagram (8).  $\square$

As in (5), the composite in the second row is  $j_b^*$ . Taking into account of Lemma 1.8, the diagram 8 is indeed the Serre dual of the diagram (6). This concludes the proof of Lemma 6.  $\square$

## 2 Lagrangian Fibrations

In this section, we associate to any Lagrangian family satisfying condition  $\clubsuit$  a Lagrangian fibration with the help of a construction appeared in [9, 14], and use this Lagrangian fibration to prove Theorem 0.7.

### General Constructions

Let (1) be a Lagrangian family of a hyper-Kähler manifold  $X$  of dimension  $2n$ . We fix a relative polarization of  $\mathcal{L} \rightarrow B$  given by a hyperplane section of  $X$ . Let

$$\pi : \mathcal{A} := \text{Alb}(\mathcal{L}/B) \rightarrow B$$

be the relative Albanese variety of  $p : \mathcal{L} \rightarrow B$ .

**Lemma 2.1.** *Let  $l$  be the relative dimension of  $p : \mathcal{L} \rightarrow B$ . Then there exist an open dense subset  $B_0 \subset B$  and a finite covering  $B'_0 \rightarrow B_0$  such that, denoting  $p'_0 : \mathcal{L}'_0 \rightarrow B'_0$  the base change of  $p$  under  $B'_0 \rightarrow B_0 \hookrightarrow B$  and  $\pi'_0 : \mathcal{A}'_0 \rightarrow B'_0$  the relative Albanese variety of  $p'_0$ , there is a cycle  $Z_0 \in CH^1(\mathcal{A}'_0 \times_{B'_0} \mathcal{L}'_0)$  such that*

$$[Z_0]^* : p'_{0*} \Omega_{\mathcal{L}'_0/B'_0} \rightarrow \pi'_{0*} \Omega_{\mathcal{A}'_0/B'_0}$$

*is an isomorphism.*

*Proof* Let  $B_0 \subset B$  be the subset of regular points of  $p : \mathcal{L} \rightarrow B$ . For  $b \in B_0$ , let  $C_b \subset L_b$  be a complete intersection curve and  $J_{C_b}$  the Jacobian variety of  $C_b$ . By Lefschetz theorem on hyperplane sections,  $j_* : J_{C_b} \rightarrow$

$A_b := \text{Alb}(L_b)$  is surjective. By the semi-simplicity of polarized Hodge structures, there exists a  $\mathbb{Q}$ -section  $s : A_b \rightarrow J_{C_b}$  of  $j_*$ , i.e., there exists  $N > 0$  such that  $j_* \circ s = N \cdot \text{id}_{A_b}$ . On  $J_{C_b} \times C_b$ , we have the Poincaré divisor  $d_b \in CH^1(J_{C_b} \times C_b)$  such that  $[d_b]^* : H^1(C_b, \mathbb{Q}) \rightarrow H^1(J_{C_b}, \mathbb{Q})$  is an isomorphism of Hodge structures, exhibiting the inverse of the pull-back of the Jacobi map  $\text{jac}_b : C_b \rightarrow J_{C_b}$ . Let us consider

$$\begin{array}{ccc} A_b \times C_b & \xrightarrow{(s, \text{id}_{C_b})} & J_{C_b} \times C_b \\ & \downarrow (id_{A_b}, j) & \\ & A_b \times L_b & \end{array}$$

and define  $Z_b := (id_{A_b}, j)_*(s, \text{id}_{C_b})^*(d_b) \in CH^1(A_b \times L_b)$ . Then  $[Z_b]^* : H^1(L_b, \mathbb{Q}) \rightarrow H^1(A_b, \mathbb{Q})$  is given by the composition

$$H^1(L_b, \mathbb{Q}) \xrightarrow{j^*} H^1(C_b, \mathbb{Q}) \xrightarrow{d_b^*} H^1(J_{C_b}, \mathbb{Q}) \xrightarrow{s^*} H^1(A_b, \mathbb{Q}), \quad (9)$$

which is an isomorphism by the definition of  $s$ . In fact, let  $(j_*)^* : H^1(A_b, \mathbb{Q}) \rightarrow H^1(J_{C_b}, \mathbb{Q})$  be the pull-back map of  $j_* : J_{C_b} \rightarrow A_b = \text{Alb}(L_b)$ . Precomposing the left-hand-side of  $(j_*)^* : H^1(A_b, \mathbb{Q}) \rightarrow H^1(J_{C_b}, \mathbb{Q})$  with the natural identification  $H^1(\text{Alb}(L_b), \mathbb{Q}) \cong H^1(L_b, \mathbb{Q})$ , and post-composing the right hand side with the pull-back of the Jacobi map  $\text{jac}_b^* : H^1(J_{C_b}, \mathbb{Q}) \rightarrow H^1(C_b, \mathbb{Q})$ , we would get the pull-back map  $j^* : H^1(L_b, \mathbb{Q}) \rightarrow H^1(C_b, \mathbb{Q})$ . Since  $\text{jac}_b^* : H^1(J_{C_b}, \mathbb{Q}) \rightarrow H^1(C_b, \mathbb{Q})$  is the inverse of  $d_b^* : H^1(C_b, \mathbb{Q}) \rightarrow H^1(J_{C_b}, \mathbb{Q})$ , the composite  $d_b^* \circ j^* : H^1(L_b, \mathbb{Q}) \rightarrow H^1(J_{C_b}, \mathbb{Q})$  of the first two maps of the composition (9) is exactly the pull-back map of  $j_* : J_{C_b} \rightarrow A_b = \text{Alb}(L_b)$  at the level of cohomology, after the natural identification  $H^1(\text{Alb}(L_b), \mathbb{Q}) \cong H^1(L_b, \mathbb{Q})$ . Since  $j_* \circ s = N \cdot \text{id}_{A_b}$ , we get the desired isomorphism.

The cycles  $Z_b$  are defined fiberwise, but standard arguments [17, Chapter 3] show that they can be constructed in family over a smooth generically finite cover  $B'_0 \rightarrow B_0$ . Let us spell out the standard arguments. By the theory of Hilbert schemes, there are countably many connected projective  $B_0$ -schemes  $H_1, \dots, H_i, \dots$  together with the universal families of cycles  $\mathcal{Z}_1, \dots, \mathcal{Z}_i, \dots$  such that for each  $i \in \mathbb{N}$ , every fiber of  $\mathcal{Z}_i \rightarrow H_i$  is a cycle  $Z_b \in \mathcal{Z}^1(A_b \times L_b)$  such that  $[Z_b]^* : H^1(L_b, \mathbb{Q}) \rightarrow H^1(A_b, \mathbb{Q})$  is an isomorphism. By the construction of the previous paragraph, the structure map  $\bigcup_{i \in \mathbb{N}} H_i \rightarrow B_0$  is surjective, thus there is  $i \in \mathbb{N}$ , such that  $H_i \rightarrow B_0$  is surjective. Since  $H_i \rightarrow B_0$  is projective, we may take a multisection of the map  $B'_0 \rightarrow B_0$ , and the universal family  $\mathcal{Z}_i$  restricted to  $B_0$  gives the desired 1-cycle.  $\square$

For the sake of simplicity, we shall note  $B_0, \mathcal{L}_0$  and  $\mathcal{A}_0$  instead of  $B'_0, \mathcal{L}'_0$  and  $\mathcal{A}'_0$ . We define a holomorphic 2-form  $\sigma_{\mathcal{A}_0}$  on  $\mathcal{A}_0$  by setting

$$\sigma_{\mathcal{A}_0} := [Z_0]^* q_0^* \sigma_X, \quad (10)$$

where  $q_0 : \mathcal{L}_0 \rightarrow X$  is the natural map.

**Proposition 2.2.** (a) The 2-form  $\sigma_{\mathcal{A}_0}$  is closed.

(b)  $\sigma_{\mathcal{A}_0}$  vanishes on fibers of  $\pi_0 : \mathcal{A}_0 \rightarrow B_0$ .

(c) The composite morphism  $\kappa : T_{B_0} \xrightarrow{\lrcorner q_0^* \sigma_X} p_{0*} \Omega_{\mathcal{L}_0/B_0} \xrightarrow{[Z_0]^*} \pi_{0*} \Omega_{\mathcal{A}_0/B_0}$  is given by the contraction  $\lrcorner \sigma_{\mathcal{A}_0}$ .

*Proof* (a) Let  $Z_q := (id, q_0)_* Z_0 \in CH(\mathcal{A}_0 \times X)$ . Then by the projection formula,  $\sigma_{\mathcal{A}_0} = [Z_q]^* \sigma_X$ . Let  $\mathcal{A}'$  be a projective completion of  $\mathcal{A}_0$ . Then  $Z_q$  extends to a cycle  $\bar{Z}_q$  of  $\mathcal{A}' \times X$ .  $\sigma_{\mathcal{A}_0}$  extends to a 2-form  $\sigma_{\mathcal{A}'} := [\bar{Z}_q]^* \sigma_X$  which is automatically closed since  $\mathcal{A}'$  is projective. Thus,  $\sigma_{\mathcal{A}_0} = \sigma_{\mathcal{A}'/\mathcal{A}_0}$  is also closed.

(b) Since  $Z_0$  is a cycle in  $\mathcal{A}_0 \times_{B_0} \mathcal{L}_0 \subset \mathcal{A}_0 \times \mathcal{L}_0$ , the morphism  $[Z_0]^* : H^*(\mathcal{L}_0) \rightarrow H^*(\mathcal{A}_0)$  preserves the Leray filtrations on both sides. Therefore,  $\sigma_{\mathcal{A}_0} \in H^0(\mathcal{A}_0, \pi_0^* \Omega_{B_0} \wedge \Omega_{\mathcal{A}_0}) \subset H^0(\mathcal{A}_0, \Omega_{\mathcal{A}_0}^2)$  since  $q_0^* \sigma_X \in H^0(\mathcal{L}_0, p_0^* \Omega_{B_0} \wedge \Omega_{\mathcal{L}_0})$  by the definition of Lagrangian families. Therefore,  $\sigma_{\mathcal{A}_0}$  vanishes on the fibers of  $\pi_0 : \mathcal{A}_0 \rightarrow B_0$ .

(c) By Lemma 2.1,  $[Z_0]^*$  induces an isomorphism  $H^0(B_0, \Omega_{B_0} \otimes p_{0*} \Omega_{\mathcal{L}_0/B_0}) \rightarrow H^0(B_0, \Omega_{B_0} \otimes \pi_{0*} \Omega_{\mathcal{A}_0/B_0})$  which sends  $\lrcorner q_0^* \sigma_X$  to  $\lrcorner \sigma_{\mathcal{A}_0}$ .  $\square$

By (b) and (c) of the above proposition, we get the following diagram that is commutative up to a sign:

$$\begin{array}{ccccccc}
0 & \longrightarrow & T_{\mathcal{A}_0/B_0} & \longrightarrow & T_{\mathcal{A}_0} & \longrightarrow & \pi_0^* T_{B_0} \longrightarrow 0 \\
& & \downarrow (\pi_0^* \kappa)^* & & \downarrow \lrcorner \sigma_{\mathcal{A}_0} & & \downarrow \pi_0^* \kappa \\
0 & \longrightarrow & \pi_0^* \Omega_{B_0} & \longrightarrow & \Omega_{\mathcal{A}_0} & \longrightarrow & \Omega_{\mathcal{A}_0/B_0} \longrightarrow 0.
\end{array} \tag{11}$$

Here, the commutativity of the second square is dual to Proposition 2.2 (c). Since the dual of  $\lrcorner \sigma_{\mathcal{A}_0} : T_{\mathcal{A}_0} \rightarrow \Omega_{\mathcal{A}_0}$  is given by  $-\lrcorner \sigma_{\mathcal{A}_0} : T_{\mathcal{A}_0} \rightarrow \Omega_{\mathcal{A}_0}$ , the first square is anti-commutative.

**Lemma 2.3.** *If condition  $\clubsuit$  (see Proposition 0.6) holds for all  $b \in B_0$ , then  $\sigma_{\mathcal{A}_0}$  is nowhere degenerate on  $\mathcal{A}_0$ .*

*Proof* If condition  $\clubsuit$  holds, then  $\pi_0^* \kappa : \pi_0^* T_{B_0} \rightarrow \Omega_{\mathcal{A}_0/B_0}$  is an isomorphism. By the commutativity of (11) and the five lemma,  $\lrcorner \sigma_{\mathcal{A}_0} : T_{\mathcal{A}_0} \rightarrow \Omega_{\mathcal{A}_0}$  is an isomorphism, which means that  $\sigma_{\mathcal{A}_0}$  is nowhere degenerate.  $\square$

## Symmetry

Let (1) be a Lagrangian family of a hyper-Kähler manifold  $X$  of dimension  $2n$ . We fix a relative polarization of  $\mathcal{L} \rightarrow B$  given by a hyperplane section of  $X$ . Let  $b \in B$  be a general point. The infinitesimal variation of Hodge structures on degree 1 cohomology of the fibers of  $p : \mathcal{L} \rightarrow B$  at  $b$  is given by (see [16, Lemme 10.19])

$$\bar{\nabla} : T_{B,b} \rightarrow \text{Hom}(H^0(L_b, \Omega_{L_b}), H^1(L_b, \mathcal{O}_{L_b})).$$

Precomposed with the map  $\lrcorner q^* \sigma_X : T_{B,b} \rightarrow H^0(L_b, \Omega_{L_b})$ , the map  $\bar{\nabla}$  induces a bilinear map

$$\begin{array}{ccc}
S : T_{B,b} \times T_{B,b} & \rightarrow & H^1(L_b, \mathcal{O}_{L_b}) \\
(u, v) & \mapsto & \bar{\nabla}_u(v \lrcorner q^* \sigma_X).
\end{array} \tag{12}$$

**Proposition 2.4.** *The bilinear map  $S$  is symmetric in the sense that  $S(u, v) = S(v, u)$  for any  $u, v \in T_{B,b}$ .*

*Proof* By Griffiths' transversality [16, Chapter 17],  $S(u, v) = \rho(u) \lrcorner (v \lrcorner q^* \sigma_X)$ , where  $\rho : T_{B,b} \rightarrow H^1(L_b, T_{L_b})$  is the Kodaira-Spencer map. Therefore, we need to show that the following diagram is commutative

$$\begin{array}{ccc}
T_{B,b} \otimes T_{B,b} & \xrightarrow{\rho \otimes id} & H^1(L_b, T_{L_b}) \otimes T_{B,b} \\
\downarrow \lrcorner q^* \sigma_X \otimes id & & \downarrow \beta \\
H^0(L_b, \Omega_{L_b}) \otimes T_{B,b} & \xrightarrow{\alpha} & H^1(L_b, \mathcal{O}_{L_b}),
\end{array} \tag{13}$$

where  $\alpha(\omega, v) = \rho(v) \lrcorner \omega$  and  $\beta(u, \chi) = u \lrcorner (\chi \lrcorner q^* \sigma_X)$ .

To see the commutativity of (13), restrict the commutative ladder (11) to  $L_b$  and apply the cohomology on  $L_b$ , then the commutativity of (11) implies the commutativity of (13). Indeed, (13) is the connecting homomorphism of the cohomology of (11) tensored by  $T_{B,b}$ .  $\square$

**Remark 2.5.** When condition  $\clubsuit$  is satisfied, the symmetry of  $S$  comes from the completely integrable system structure on  $(\mathcal{A}_0, \sigma_{\mathcal{A}_0})$ . What we proved is in fact the symmetry of

$$\begin{array}{ccc}
S' : T_{B,b} \times T_{B,b} & \rightarrow & H^1(A_b, \mathcal{O}_{A_b}) \\
(u, v) & \mapsto & \bar{\nabla}_u(v \lrcorner \sigma_{\mathcal{A}_0}).
\end{array} \tag{14}$$

Fixing a relative polarisation on  $\mathcal{A}_0 \rightarrow B_0$ , we have natural isomorphisms (always under  $\clubsuit$ ):  $H^1(A_b, \mathcal{O}_{A_b}) \cong H^0(A_b, \Omega_{A_b})^* \cong T_{B,b}^*$  and we can thus view  $S'$  as an element in  $T_{B,b}^* \otimes T_{B,b}^* \otimes T_{B,b}^*$ . If this relative polarisation is principal, Donagi and Markman proved in [6, Lemma 7.2] that  $S'$  lies in  $\text{Sym}^3 T_{B,b}^*$ . This result is called ‘‘weak cubic condition’’ in [6, Lemma 7.2]. See also [12, Theorem 4.4]

Now we are ready to prove the main theorem.

*Proof of Theorem 0.7.* Under the assumptions of Theorem 0.7, assume by contradiction that the Abel–Jacobi map (3) is not constant. In what follows, we fix a relative polarization on  $p_0 : \mathcal{L}_0 \rightarrow B_0$  induced from a hyperplane section of  $X$ , so that  $R^{2n-1}p_{0*}\mathbb{Q} \cong R^1p_{0*}\mathbb{Q}$ . By Proposition 0.6, the morphism

$$j^* : H^{2n-1}(X, \mathbb{Q}) \rightarrow R^{2n-1}p_{0*}\mathbb{Q} \cong R^1p_{0*}\mathbb{Q}$$

of variations of Hodge structures on an open subset  $B_0 \subset B$  containing  $b$  is not zero. Hence, there is a non-zero locally constant sub-variation of Hodge structures  $I := \text{Im}j^* \subset R^1p_{0*}\mathbb{Q}$ . Since  $I$  is locally constant, for any  $\omega \in I_b^{1,0}$  and  $u \in T_{B,b}$ ,  $\nabla_u(\omega) = 0$ . Recall that  $\clubsuit$  means that  $\lrcorner q^* \sigma_X : T_{B,b} \rightarrow H^0(L_b, \Omega_{L_b})$  is bijective. Let  $F := (\lrcorner q^* \sigma_X)^{-1}(I^{1,0}) \subset T_{B,b}$ . Then by the symmetry of  $S$  given by Proposition 2.4,  $F$  lies in the kernel of  $\bar{\nabla}$ , which contradicts our assumption that the variation of Hodge structures is maximal.  $\square$

### 3 Maximal Variations

In this section, we study under what conditions could the variation of Hodge structures of a Lagrangian family be maximal. Consider a Lagrangian family of a hyper-Kähler manifold  $X$  of dimension  $2n$  satisfying the condition  $\clubsuit$  given by the diagram as in (1). Let  $U \subset B$  be a simply connected open subset of  $B_0 \subset B$  and let

$$\begin{aligned} \mathcal{P} : U &\rightarrow Gr(h^{1,0}(L), H^2(L, \mathbb{C})) \\ b &\mapsto H^{1,0}(L_b) \subset H^1(L_b, \mathbb{C}) \cong H^1(L, \mathbb{C}), \end{aligned} \quad (15)$$

be the local period map of the Lagrangian family.

In what follows, we are going to use a universal property of the Kuga–Satake construction proved in [10, Proposition 6].

**Theorem 3.1** ([10]). *Let  $(H^2, q)$  be a polarized Hodge structure of hyper-Kähler type of dimension  $\geq 5$ . Assume that the Mumford–Tate group of the Hodge structure on  $H^2$  is maximal, namely the special orthogonal group of  $(H^2, q)$ . Let  $H$  be a simple effective weight-1 Hodge structure, such that there exists an injective morphism of Hodge structures of bidegree  $(-1, -1)$*

$$H^2 \hookrightarrow \text{Hom}(H, A)$$

for some weight-1 Hodge structure  $A$ . Then  $H$  is a direct summand of the Kuga–Satake Hodge structure  $H_{KS}^1(H^2, q)$ . In particular,  $\dim H \geq 2 \lfloor \frac{\dim H^2 - 1}{2} \rfloor$ .

With the same notations as in the introduction, we prove

**Proposition 3.2.** *Assume that the Mumford–Tate group of the Hodge structure  $H^2(X, \mathbb{Q})$  is maximal, i.e. it is the special orthogonal group of  $(H^2(X, \mathbb{Q})_{\text{tr}}, q)$  and assume  $b_2(X)_{\text{tr}} \geq 5$ . If the dimension of  $H^{0,1}(L_b)$  is smaller than  $2 \lfloor \frac{b_2(X)_{\text{tr}} - 3}{2} \rfloor$  for a general fiber  $L_b$  of  $p : \mathcal{L} \rightarrow B$ , then the variation of weight 1 Hodge structure of  $p$  is maximal.*

*Proof* We use the same argument as in [10] where similar results were proved for Lagrangian fibrations. Assuming that the period map (15) is not generically an immersion, we are going to prove that  $\dim H^{0,1}(L_b) \geq 2 \lfloor \frac{b_2(X)_{\text{tr}} - 3}{2} \rfloor$ . By assumption, the nonempty general fibers of  $\mathcal{P}$  are of dimension  $\geq 1$ . Let  $b \in U$  be a general point and let  $B_b$  the fiber of  $\mathcal{P}$  passing through  $b$ . Let  $U_b = B_b \cap U$ . Then the fibers of  $\pi|_{U_b} : \mathcal{A}_{U_b} \rightarrow U_b$  are isomorphic with each other. Thus, up to a base change by a finite covering of  $U_b$ , we may assume  $\pi|_{U_b} : \mathcal{A}_{U_b} \rightarrow U_b$  is trivial, i.e.,  $\mathcal{A}_{U_b} = U_b \times A_b$ . Let  $\pi_{F_b} : \mathcal{A}_{F_b} \rightarrow F_b$  be a smooth completion of  $\pi|_{U_b}$ , then  $\mathcal{A}_{F_b}$  is birational to  $F_b \times A_b$ , which

gives a morphism  $H^2(\mathcal{A}_{F_b}) \rightarrow H^2(F_b \times A_b)$ . Recall by Lemma 2.1, we get a morphism  $[Z]^* : H^2(X) \rightarrow H^2(\mathcal{A})$  that sends  $\sigma_X$  to a holomorphic 2-form which is non-degenerate on  $\mathcal{A}_U$ . Finally, the rational map  $\mathcal{A}_{F_b} \dashrightarrow \mathcal{A}$  induces  $H^2(\mathcal{A}) \rightarrow H^2(\mathcal{A}_{F_b})$ . Compositing all these maps, we get a morphism

$$\alpha : H^2(X)_{ir} \hookrightarrow H^2(X) \rightarrow H^2(\mathcal{A}) \rightarrow H^2(\mathcal{A}_{F_b}) \rightarrow H^2(F_b \times A_b) \rightarrow H^1(F_b) \otimes H^1(A_b), \quad (16)$$

where the last map is given by the projection in the Künneth decomposition.

**Lemma 3.3.**  $\alpha : H^2(X)_{ir} \rightarrow H^1(F_b) \otimes H^1(A_b)$  is injective.

*Proof* Since  $h^{2,0}(X) = 1$  and  $H^{2,0}(X)$  is orthogonal to  $NS(X)$  with respect to the Beauville-Bogomolov-Fujiki form,  $H^2(X)_{ir}$  is a simple Hodge structure. Therefore, to show the injectivity of  $\alpha$  it suffices to show that  $\alpha$  is not zero. We claim that  $\alpha(\sigma_X) \neq 0$ . Indeed, Since  $A_b$  is Lagrangian with respect to  $\sigma_{\mathcal{A}}$  (Proposition 2.2 (b)), in the Künneth's decomposition of  $H^2(A_b \times F_b)$ , the image of  $\sigma_X$  in  $H^0(F_b) \otimes H^2(A_b)$  is zero. If furthermore  $\alpha(\sigma_X) = 0$  in  $H^1(F_b) \otimes H^1(A_b)$ , then the image of  $\sigma_X$  on  $F_b \times A_b$  comes from a 2-form on  $F_b$ , which has rank  $\leq \dim F_b$ . Therefore, the rank of  $\sigma_{\mathcal{A}}$  has rank  $\leq \dim F_b$  on  $\mathcal{A}_{U_b}$ . On the other hand, the codimension of  $\mathcal{A}_{U_b}$  in  $\mathcal{A}_U$  is  $\dim B - \dim F_b$ , and thus the non-degeneration of  $\sigma_{\mathcal{A}_U}$  implies that  $\sigma_{\mathcal{A}}$  has rank  $\geq 2 \dim F_b$  on  $\mathcal{A}_{U_b}$ . This is a contradiction since we are assuming  $\dim F_b \geq 1$ .  $\square$

We are now in the position to use the universal property of the Kuga-Satake construction (see Theorem 3.1 above). Since  $\alpha : H^2(X)_{ir} \rightarrow H^1(F_b) \otimes H^1(A_b)$  is nonzero, there is at least one simple direct factor  $A$  of  $A_b$  such that  $H^2(X)_{ir} \rightarrow H^1(F_b) \otimes H^1(A)$  is nonzero thus injective. Taking  $H^2$  as  $H^2(X)_{ir}$ , we conclude by Theorem 3.1 that

$$\dim H^{0,1}(L_b) = \dim A_b \geq \dim A \geq \frac{1}{2} \times 2^{\lfloor \frac{\dim H^2(X)_{ir}-1}{2} \rfloor} = 2^{\lfloor \frac{b_2(X)_{ir}-3}{2} \rfloor},$$

as desired.  $\square$

## 4 Example of a Lagrangian Family with Nontrivial Abel-Jacobi Map

Recall the construction of generalized Kummer varieties introduced in [1]. Let  $A$  be an abelian surface and  $A^{[n+1]}$  the Hilbert scheme of length  $n+1$  subschemes of  $A$ . Let  $\text{alb} : A^{[n+1]} \rightarrow A$  be the composition of the Hilbert-Chow morphism and the summation map

$$A^{[n+1]} \rightarrow A^{(n+1)} \rightarrow A.$$

Note that  $\text{alb}$  is an isotrivial fibration. The generalized Kummer variety  $K_n(A)$  is defined to be the fiber of  $\text{alb}$  over  $0 \in A$ . As is shown in [1],  $K_n(A)$  is a hyper-Kähler manifold of dimension  $2n$ .

In this section, we are going to construct Lagrangian families of  $X := K_n(A)$  for  $n \geq 2$ , satisfying condition  $\clubsuit$  and whose Abel-Jacobi map is *not* trivial.

For any  $x \in A$ , one defines a subvariety  $Z_x$  of  $K_n(A)$  consisting of Artinian subschemes of  $A$  of length  $n+1$  supported on  $x$  and  $-nx$ , with multiplicities  $n$  and  $1$ , respectively. By [3, Proposition VI.1.1],  $Z_x$  is a *rational* variety of dimension  $n-1$  if  $x$  is *not* an  $(n+1)$ -torsion point. Let  $Z = \bigcup_{x \in A} Z_x$  and let  $\pi : Z \rightarrow A$  send elements in  $Z_x$  to  $x$ . For any curve  $C \subset A$ , define  $Z_C = \bigcup_{x \in C} Z_x$ .

Now let  $B$  be a connected open subset of the Hilbert scheme of deformations of a smooth curve  $C \subset A$  and  $\mathcal{C} \rightarrow B$  the corresponding family.

**Lemma 4.1.**  $\{Z_C\}_{C \in B}$  is a Lagrangian family of  $K_n(A)$  satisfying condition  $\clubsuit$ .

*Proof* Since for general  $C$ ,  $Z_C$  is a fibration over a curve  $C$  whose general fibers are rational, any holomorphic 2-form on  $Z_C$  is 0. Furthermore,  $\dim Z_C = n = \dim K_n(A)/2$ . These imply that  $\{Z_C\}_{C \in B}$  is a Lagrangian family.

We now show that this family satisfies condition  $\clubsuit$ . Denoting  $\mathcal{L}$  the total space of the family  $\{Z_C\}_{C \in B}$  and  $L$  a general fiber, and using as before the following notation

$$\begin{array}{ccc} \mathcal{L} & \xrightarrow{q} & X \\ \downarrow p & & \\ B & & \end{array},$$

we need to show that  $\lrcorner q^* \sigma_{K_n(A)} : H^0(L, N_{L/Z}) = H^0(L, N_{L/\mathcal{L}}) \rightarrow H^0(L, \Omega_L)$  is an isomorphism. Since the general fibers of  $\pi$  are rational,  $q^* \sigma_{K_n(A)} = \pi^* \sigma_A$ , where  $\sigma_A$  is the unique (up to coefficients) holomorphic 2-form on  $A$ . Therefore, we can conclude by the commutativity of the following diagram

$$\begin{array}{ccc} H^0(C, N_{C/A}) & \xrightarrow{-\sigma_A} & H^0(C, \Omega_C) \\ \downarrow \pi^* & & \downarrow \pi^* \\ H^0(L, N_{L/Z}) & \xrightarrow{-\pi^* \sigma_A} & H^0(L, \Omega_L) \end{array}$$

noting that the two vertical arrows are isomorphisms since the fibers of  $\pi$  are rational, and that  $\lrcorner \sigma_A : H^0(C, N_{C/A}) \rightarrow H^0(C, \Omega_C)$  is an isomorphism since  $\sigma_A$  is nondegenerate.  $\square$

**Proposition 4.2.** *The Abel-Jacobi map of the Lagrangian family  $\{Z_C\}_{C \in B}$  is not trivial.*

*Proof* Let  $i : C \hookrightarrow A$  be a general curve in the family  $\mathcal{C} \rightarrow B$ . By Proposition 1.5(a), it suffices to show that the restriction map  $H^{2n-1}(X, \mathbb{Q}) \rightarrow H^{2n-1}(Z_C, \mathbb{Q})$  is nonzero.

Define an injective morphism

$$\begin{aligned} \beta : A \times A &\hookrightarrow A^{(n+1)} \\ (x, y) &\mapsto n\{x\} + \{y\}, \end{aligned}$$

where we use the notation  $\{x\} \in \mathcal{Z}_0(A)$  the 0-cycle of the point  $x \in A$ . Consider the following pull-back diagram defining a subvariety  $Z' \subset A^{[n+1]}$

$$\begin{array}{ccc} Z' & \xleftarrow{\alpha} & A^{[n+1]} \\ \downarrow \pi' & & \downarrow c \\ A \times A & \xleftarrow{\beta} & A^{(n+1)} \end{array},$$

where  $c : A^{[n+1]} \rightarrow A^{(n+1)}$  is the Hilbert-Chow morphism. Then  $Z = Z' \cap K_n(A) \subset A^{[n+1]}$ . We have the following commutative diagram where all three squares are pull-back diagrams

$$\begin{array}{ccccc} Z & \xrightarrow{\quad} & X = K_n(A) & & \\ \downarrow \pi & \searrow & \downarrow & \searrow & \\ A & & A^{[n+1]} & \xrightarrow{\text{alb}} & A \\ \downarrow f & & \downarrow \pi' & & \downarrow c \\ A \times A & \xleftarrow{\beta} & A^{(n+1)} & \xrightarrow{\Sigma} & A \end{array},$$

Here  $f : A \rightarrow A \times A$  defined by  $x \mapsto (x, -nx)$  is the fiber over  $0 \in A$  of the trivial fibration  $\Sigma \circ \beta : A \times A \rightarrow A$ .

By [5, Corollary 5.1.5],  $[Z']^* : H^{2n-1}(A^{[n+1]}, \mathbb{Q}) \rightarrow H^1(A \times A, \mathbb{Q})$  is surjective. Furthermore, the restriction map  $f^* : H^1(A \times A, \mathbb{Q}) \rightarrow H^1(A, \mathbb{Q})$  is surjective since  $f$  is the fiber of a trivial fibration. These imply that  $[Z]^* : H^{2n-1}(X, \mathbb{Q}) \rightarrow H^1(A, \mathbb{Q})$  is surjective. Finally, since the restriction map  $i^* : H^1(A, \mathbb{Q}) \rightarrow H^1(C, \mathbb{Q})$  is injective by Lefschetz hyperplane theorem, the composition map  $i^* \circ [Z]^* : H^{2n-1}(X, \mathbb{Q}) \rightarrow H^1(C, \mathbb{Q})$  is nonzero. This implies that the restriction map  $H^{2n-1}(X, \mathbb{Q}) \rightarrow H^{2n-1}(Z_C, \mathbb{Q})$  is nonzero, as desired.  $\square$

## References

- [1] A. Beauville, *Variétés kählériennes dont la première classe de Chern est nulle*, J. Differential Geometry 18(1983), 755-782.
- [2] A. Beauville, R. Donagi, *The variety of lines of a cubic fourfold*, C. R. Acad. Sci. Paris 301 Séries I, (1985), 703-706.
- [3] J. Briançon. *Description de  $\text{Hilb}^n\mathbb{C}\{x,y\}$* , Invent. Math., 41:45–89, 1977.
- [4] H. Clemens, P. Griffiths, *The intermediate Jacobian of the cubic threefold*, Ann. of Math. (2) 95 (1972), 281–356.
- [5] M. A. de Cataldo, L. Migliorini, *The Chow groups and the motive of the Hilbert scheme of points on a surface*, J. algebra 251 (2002), 824-848.
- [6] R. Donagi, E. Markman, *Spectral covers, algebraically completely integrable, Hamiltonian systems, and moduli of bundles*, Integrable Systems and Quantum Groups (Montecatini Terme 1993), Lecture Notes in Math., 1620, pp. 1-119. Springer, Berlin-Heidelberg, 1998.
- [7] D. Huybrechts, *Curves and cycles on K3 surfaces (with an appendix by C. Voisin)*, Algebraic Geometry 1 (2014), 69-106.
- [8] A. Iliev, L. Manivel, *Cubic hypersurfaces and integrable systems*, Amer. J. Math. 130 (2008), no. 6, 1445–1475.
- [9] R. Laza, G. Saccà, C. Voisin, *A hyper-Kähler compactification of the Intermediate Jacobian fibration associated with a cubic fourfold*, Acta Mathematica, 218:1 (2017), 55-135.
- [10] B. van Geemen, C. Voisin, *On a conjecture of Matsushita*, Int. Math. Res. Notices (2016) Vol. 2016, 3111-3123.
- [11] C. Voisin, *Sur la stabilité des sous-variétés lagrangiennes des variétés symplectiques holomorphes*, Complex projective geometry (Trieste, 1989/Bergen, 1989), 294-303, London Math. Soc. Lecture Note Ser., 179, Cambridge Univ. Press, Cambridge, 1992.
- [12] C. Voisin, *Torsion points of sections of Lagrangian torus fibrations and the Chow ring of hyper-Kähler manifolds*, The Abel Symposium Abelsymposium 2017: Geometry of Moduli, pp 295-326.
- [13] C. Voisin, *Remarks and questions on coisotropic subvarieties and 0-cycles of hyper-Kähler varieties*, in “K3 Surfaces and Their Moduli”, Proceedings of the Schiermonnikoog conference 2014, C. Faber, G. Farkas, G. van der Geer, Editors, Progress in Math 315, Birkhäuser (2016), 365-399.
- [14] C. Voisin, *Triangle varieties and surface decomposition of hyper-Kähler manifolds*, Recent Developments in Algebraic Geometry: To Miles Reid for his 70th Birthday, edited by Hamid Abban; Gavin Brown; Alexander Kasprzyk; Shigefumi Mori, London Mathematical Society Lecture Note Series Cambridge University Press.
- [15] C. Voisin, *On the Lefschetz standard conjecture for Lagrangian covered hyper-Kähler varieties*, to appear Advances in Mathematics.
- [16] C. Voisin. *Théorie de Hodge et géométrie algébrique complexe*, Cours spécialisés, SMF 2003.
- [17] C. Voisin, *Chow rings, decomposition of the diagonal and the topology of families*, Annals of Math. Studies 187, Princeton University Press 2014.

Sorbonne Université and Université de Paris, CNRS, IMJ-PRG, F-75005 Paris, France.

*Email adress:* `chenyu.bai@imj-prg.fr`.