

		Total patients (n=486)	Patients with UIA (n=243)	Patients with RIA (n=243)	p-value
<i>Clinical and demographic data</i>					
	Males	177/486 (36.4%)	90/243 (37.0%)	87/243 (35.8%)	0.78
	Hypertension	210/482 (43.6%) [§]	119/240 (49.6%) [§]	91/242 (37.6%) [§]	0.008*
	Age : mean (+/- SD)	57 (± 12.6)	57 (± 11.9)	57 (± 13.3)	0.84
	History of subarachnoid hemorrhage	16/486 (3.3%)	7/243 (2.9%)	9/243 (3.7%)	0.61
	Smoking	162/458 (35.4%) [§]	84/239 (35.1%) [§]	78/219 (35.6%) [§]	0.92
	Multiple aneurysms	141/480 (29.4%) [§]	79/243 (32.5%) [§]	62/247 (25.1%) [§]	0.13
		Total aneurysms (n=604)	UIA (n=355)	RIA (n=249)	
<i>Aneurysm characteristics</i>					
	Size (mm): mean (+/- SD)	6.0 (± 3.4)	5.3 (± 2.9)	7.0 (± 3.8)	<0.0001*
	Irregular shape	314/604 (52.0%)	131/355 (36.9%)	183/249 (73.5%)	<0.0001*

Supplemental Table 1 – General characteristics of aneurysm population.

UIA: unruptured intracranial aneurysm; RIA: ruptured intracranial aneurysm; SD: standard deviation.

* p-value < 0.05.

[§] missing values for some patients.