



HAL
open science

Characterization of extreme Gibbs measures for a Categorical Approach to Statistical Mechanics

Grégoire Sergeant-Perthuis

► **To cite this version:**

Grégoire Sergeant-Perthuis. Characterization of extreme Gibbs measures for a Categorical Approach to Statistical Mechanics. 2024. hal-04456412v2

HAL Id: hal-04456412

<https://hal.sorbonne-universite.fr/hal-04456412v2>

Preprint submitted on 22 Feb 2024

HAL is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers.

L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.



Distributed under a Creative Commons Attribution - NonCommercial - NoDerivatives 4.0
International License

Characterization of extreme Gibbs measures for a Categorical Approach to Statistical Mechanics

Grégoire Sergeant-Perthuis^{*1}

¹Laboratoire de Biologie Computationnelle et Quantitative, Sorbonne
Université, Paris , France, , ORCID: 0000-0002-2079-3410

February 22, 2024

Abstract

In recent work we proposed to follow a categorical approach to Statistical Mechanics. In this paper we continue in this direction and give a categorical formulation of the decomposition of Gibbs measures by extending the characterization of extreme Gibbs measures to the categorical setting.

Keywords: Statistical mechanics, category theory, Extreme Gibbs measures, Zero-one laws

MSC: 18D99,60F20,60A99

1 Introduction

Statistical Mechanics is the field interested in the statistical study of interacting particles ($X_i \in E_i, i \in \mathbb{N}$). A landmark rigorous formulation of Statistical Mechanics can be found in Georgii's *Gibbs Measures and Phase Transition* [1]. Such a framework is needed to define phases or, more generally, *Gibbs measures of statistical systems*. Phases only exist for infinitely many interacting particles and do not exist for finite-size systems. The main constructions of such a framework rely on the necessity to have a *universe*, denoted Ω , that encompasses all possible configurations of the system, i.e., all the possible joint configurations of the particles $\Omega := \prod_{i \in \mathbb{N}} E_i$. Not referring to a global space is problematic as exhibited by Girya's seminal work on *A Categorical Approach to Probability Theory* [2] and can be summarized by the fact that the Girya monad, \mathbb{P} , does not commute to limits. We are interested in describing statistical systems for which it is not possible to have

^{*}gregoireserper@gmail.com

complete knowledge of the states of the particles at the same time; we want to forget about the ‘ Ω ’ and we want to study statistical systems ‘locally’.

In [3, 4], we proposed to reformulate Statistical Mechanics and Gibbs measures introducing an appropriate category. Among others, in [5], we gave a characterization of independent variables in terms of projective objects in this category and an easy-to-verify condition that characterizes such objects (for the injective case see [6–9]). In [10], we proposed an Entropy functional for the categorical version of statistical systems. In this article, we show how the characterization of extreme Gibbs measures (Theorem 7.7 [1]), one of the steps for proving a zero-one law for the extreme Gibbs measures, transfers in the categorical setting. In the classical theory of rigorous statistical mechanics, the tail σ -algebra generates the observables for which a ‘generalized’ law of large numbers (zero-one law) holds. In this article, we give a candidate for such a σ -algebra in the categorical setting and show the associated extreme Gibbs measures decomposition. We will discuss in a follow-up paper the generalization of the zero-one law of specifications to \mathcal{A} -specifications.

2 \mathcal{A} -specifications and their Gibbs measures

In classical theory, a statistical system is defined by a specification (Definition 1.23 [1]). We proposed in [3, 4] to reformulate such constructions as a couple of a presheaf and a functor over a poset, denoted as \mathcal{A} ; we call such a couple an \mathcal{A} -specification. Let us recall their definition. Let us denote **Mes** the category that has as objects measurable spaces and as morphisms measurable maps, and **Kern** the category that has as objects measurable spaces and as morphisms Markov kernels.

Definition 2.1 (\mathcal{A} -specifications (Definition 7 [3])). Let \mathcal{A} be a poset. An \mathcal{A} -specification is a couple (G, F) of a presheaf and a functor $G : \mathcal{A}^{op} \rightarrow \mathbf{Mes}$ and $F : \mathcal{A} \rightarrow \mathbf{Kern}$ such that for any $a, b \in \mathcal{A}$ with $b \leq a$,

$$G_b^a \circ F_a^b = \text{id} \quad (1)$$

We denote $\alpha \circ \beta$ simply as $\alpha\beta$ in what follows.

Let us now recall the categorical generalization of the classical Gibbs measures (Definition 1.23 [1]) that we propose.

Definition 2.2 (Gibbs measure for \mathcal{A} -specifications Definition 8 [3]). Let $\gamma := (G, F)$ be an \mathcal{A} -specification,

$$\mathcal{G}(\gamma) := \{(p_a \in \mathbb{P}(G(a)), a \in \mathcal{A}) \mid \forall b \leq a, F_a^b p_b = p_a\} \quad (2)$$

with $\mathbb{P}(G(a))$ the (measurable) space of probability measures over $G(a)$.

3 Categorical Version of the Tail σ -Algebra

Measurable maps are particular Markov kernels. Let $F : E_b \rightarrow E_a$ be a Markov kernel; let us denote it also as $F(\omega_a|\omega_b)$, where $\omega_b \in E_b$ and $\omega_a \in E_a$. One associates to a Markov kernel $F : E_b \rightarrow E_a$ a linear map $\pi : L^\infty(E_a) \rightarrow L^\infty(E_b)$ defined as follows: for any $f \in L^\infty(E_a)$,

$$\forall \omega_b \in E_b, \quad \pi(f)(\omega_b) = \int f(\omega_a)F(d\omega_a|\omega_b) \quad (3)$$

This association is ‘functorial’, we may denote the underlying functor $L^\infty : \mathbf{Kern}^{op} \rightarrow \mathbf{Vect}$ which is presheaf from the category of Markov kernels to the category of vector spaces. It is the presheaf that associates spaces to their space of observables. Let us denote $L^\infty \circ G : \mathcal{A} \rightarrow \mathbf{Vect}$ as i and $L^\infty \circ F : \mathcal{A}^{op} \rightarrow \mathbf{Vect}$ as π . In these notation one has that for any $a, b \in \mathcal{A}$ such that $b \leq a$ then $\pi_b^a \circ i_a^b = \text{id}$.

Let us recall the definition of the tail σ -algebra in the classical formulation. Let $\Omega = \prod_{i \in \mathbb{N}} E_i$, where E_i are measurable spaces; let us denote $\mathcal{F}_{\geq k}$ as the σ -algebra generated by the cylinders $\prod_{n \geq k} E_n$. The tail σ -algebra is defined as $\bigcap_{k \in \mathbb{N}} \mathcal{F}_{\geq k}$. A more general definition holds when I is any set and $\mathcal{F}_{\supseteq a}$ is indexed by a subset $a \subseteq I$ that is co-finite, i.e., which has a finite complement. For a functor from \mathcal{A} to \mathbf{Mes} , let us denote $\sigma(G(a))$ as the sigma algebra of the measurable space $G(a)$, where $a \in \mathcal{A}$, and $\sigma(G)$ as the underlying functor defined as $\sigma(G)_a^b A_b := G_b^{a-1} A_b$, with $b \leq a$. We propose that one candidate that plays the role of the tail σ -algebra for a given specification $\gamma = (G, F)$ is $\lim \sigma(G)$ defined as,

$$\lim \sigma(G) := \{(A_a \in \sigma(G(a)), a \in \mathcal{A}) | \forall a, b \in \mathcal{A}, \quad A_a = G_b^{a-1} A_b\} \quad (4)$$

Let us denote $1_A : E \rightarrow \{0, 1\}$ the indicator function over the set A that sends $\omega \in A$ to 1 and $\omega \notin A$ to 0. Let us remark that $1_{A_b} \circ G_b^a = 1[G_b^a(\omega_a) \in A_b] = 1_{G_b^{a-1} A_b}$. Remark that $A \in \lim \sigma(G)$ is equivalent to $1_A \in \lim i$; in other words, $\lim \sigma(G)$ is the restriction of $\lim i$ to indicator functions of the form 1_{A_a} , $a \in \mathcal{A}$.

Finally, we also need to recall that for any $f \in L^\infty(E)$ and $\mu \in \mathbb{P}(E)$, one can define a measure $f \cdot \mu$ as $f \cdot \mu(d\omega) = f(\omega)d\omega$.

The key proposition of this document is Proposition 3.1; the proof of that proposition is given for $G(a)$, $a \in \mathcal{A}$ finite measurable sets. Therefore, we assume in what follows that the measurable sets are finite. However, there is no finiteness constraint on \mathcal{A} . We will say that $F > 0$ when for any $a, b \in \mathcal{A}$, such that $b \leq a$, $F(\omega_a|\omega_b) > 0$ for any ω_b such that $G_b^a(\omega_a) = \omega_b$; $G \circ F = \text{id}$ requires that $F(\omega_a|\omega_b) = 0$ when $G_b^a(\omega_a) \neq \omega_b$.

The following lemma is an extension of the classical result that states that conditioning over a σ -subalgebra $\mathcal{F}_1 \subseteq \mathcal{F}$ defines a morphism of modules when finer (\mathcal{F} measurable) observables are seen as modules over the coarser (\mathcal{F}_1 measurable observables).

Lemma 3.1. *Let E_1, E_2 be two measurable spaces, let $g : E_2 \rightarrow E_1$ be a measurable map and $f : E_1 \rightarrow E_2$ be a Markov kernel so that, $f \circ g = \text{id}$. Let us denote respectively i and π the induced linear maps on $L^\infty(E_1), L^\infty(E_2)$. Let $h \in L^\infty(E_2)$ and $k \in L^\infty(E_1)$, then,*

$$\pi(h).k = \pi(h.i(k)) \quad (5)$$

Proof. Let us first prove the result in the particular case when $h = 1_B$ with $B \in \sigma(E_2)$ and $k = 1_A$ with $A \in \sigma(E_1)$. Let us denote \bar{A} the complement of A , then $1_A + 1_{\bar{A}} = 1$ and $1_A.1_{\bar{A}} = 0$. Furthermore $i(1_A) = g^{-1}A$

$$\pi(1_B) = \pi(1_B.1_{g^{-1}A}) + \pi(1_B.1_{\overline{g^{-1}A}}) \quad (6)$$

and

$$\pi(1_B.1_{g^{-1}A}) \leq \pi(1_{g^{-1}A}) = \pi \circ i(1_A) = 1_A \quad (7)$$

$$\pi(1_B.1_{\overline{g^{-1}A}}) \leq \pi(1_{\overline{g^{-1}A}}) = 1_{\bar{A}} \quad (8)$$

Therefore,

$$\pi(1_B)1_A = \pi(1_B.1_{g^{-1}A})1_A + \pi(1_B.1_{\overline{g^{-1}A}})1_A \quad (9)$$

But, $\pi(1_B.1_{\overline{g^{-1}A}})1_A \leq 1_{\bar{A}}.1_A = 0$ so, $\pi(1_B)1_A = \pi(1_B.1_{g^{-1}A})1_A$. Furthermore, $\pi(1_B.1_{g^{-1}A}) = \pi(1_B.1_{g^{-1}A})1_A + \pi(1_B.1_{g^{-1}A})1_{\bar{A}}$ therefore,

$$\pi(1_B.1_{g^{-1}A})1_{\bar{A}} \leq \pi(i(1_A)).1_{\bar{A}} = 0 \quad (10)$$

We just showed that,

$$\pi(1_B.1_{g^{-1}A}) = \pi(1_B.1_{g^{-1}A})1_{\bar{A}} \quad (11)$$

So $\pi(1_B).1_A = \pi(1_B.i(1_A))$. The result then extends by linearity directly to $h = \sum_{k \leq n} 1_{B_k}$ and $k = \sum_{k \leq n_1} 1_{A_k}$, which ends the proof. \square

Let us remark that if $A \in \lim \sigma(G)$ then $\bar{A} := (\bar{A}_a, a \in \mathcal{A})$ is also in $\lim i$.

Proposition 3.1. *Let $\gamma = (G, F)$ be a specification, let $G(a)$ be finite sets for any $a \in \mathcal{A}$, let $F > 0$. Let $\mu \in \mathcal{G}(\gamma)$, for any $f \in \prod_{a \in \mathcal{A}} L^\infty(G(a))$, such that $\forall a \in \mathcal{A}, \mu_a(f_a) = 1$,*

$$f.\mu \in \mathcal{G}(\gamma) \iff \exists \tilde{f} \in \lim i, \text{ s.t. } f.\mu = \tilde{f}.\mu \quad (12)$$

Proof. Let us assume that $f.\mu \in \mathcal{G}(\gamma)$, then for any $a, b \in \mathcal{A}$ such that $b \leq a$, and for any $g_a \in L^\infty(G(a))$, by hypothesis, $(f.\mu)_b \pi_b^a(g_a) = (f.\mu)_a(g_a)$; it can be rewritten as,

$$\mu_b(\pi_b^a(g_a).f_b) = \mu_a(f_a.g_a) \quad (13)$$

By Lemma 3.1, $\pi_b^a(g_a).f_b = \pi_b^a(g_a.i_a^b f_b)$; therefore,

$$\mu_b \pi_b^a(g_a.i_a^b f_b) = \mu_a(f_a.g_a) \quad (14)$$

Therefore $f_a = i_a^b f_b$ μ_a -almost surely.

We will now show that there is $\tilde{f} \in \lim i$ such that $\tilde{f}.\mu = f.\mu$. It is in this part of the proof that we assume that $G(a)$ are finite sets and that $F > 0$. Let's call $S_a := \text{supp} \mu_a$, the support of μ_a , i.e., the set $\text{supp} \mu_a := \{\omega_a \in G(a) \mid \mu_a(\omega_a) > 0\}$. Let us denote $N_a := \overline{S_a}$ its complement and $M_a = 1_{\overline{N_a}}$. We will now show that $(N_a, a \in \mathcal{A}) \in \lim \sigma(G)$.

For any $b, a \in \mathcal{A}$ such that $b \leq a$, $\mu_a i_a^b = \mu_b$; therefore, as $\mu_b(M_b) = 0$, one has that $\mu_a i_a^b(M_b) = 0$. Recall that $i_a^b(M_b)$ is the indicator function of the set $G_b^{a-1} N_b$; the previous remark implies that $i_a^b(M_b) \leq M_a$. Furthermore, $\mu_b \pi_b^a(M_a) = \mu_a(M_a) = 0$; therefore, $\pi_b^a(M_a) \leq M_b$.

Hence, as $\pi_b^a(i_a^b M_b) = M_b$ and $i_a^b M_b \leq M_a$, then by applying π_b^a on both sides, $M_b \leq \pi_b^a(M_a)$. And so $\pi_b^a(M_a) = M_b$.

Recall that we showed that $\pi_b^a(M_a) = M_b$ and $i_a^b(M_b) \leq M_a$. In particular, $M_a - i_a^b(M_b) \geq 0$; furthermore, $\pi_b^a(M_a - i_a^b(M_b)) = 0$ so $M_a = i_a^b(M_b)$. To be more explicit: $\forall \omega_b \in G(b)$,

$$\pi_b^a(M_a - i_a^b(M_b))(\omega_b) = \sum_{\omega_a \in G(a)} F(\omega_a \mid \omega_b) [M_a - i_a^b(M_b)](\omega_a) \quad (15)$$

As $F(\omega_a \mid \omega_b) > 0$ by hypothesis, then $M_a = i_a^b(M_b)$. This implies that $M \in \lim i$ and $N \in \lim \sigma(G)$. This also implies that $S \in \lim \sigma(G)$.

Let $\tilde{f} = f 1_S$. Then $f 1_S \in \lim F$ and for any $a \in \mathcal{A}$, $f_a = \tilde{f}_a$ μ_a -a.s., which ends the proof. \square

One remarks that $\lim i$ is a subset of $\lim \pi$: for any $f \in \lim i$, by definition for any $a, b \in \mathcal{A}$ such that $b \leq a$, $i_a^b f_b = f_a$ and so $\pi_b^a i_a^b f_b = \pi_b^a f_a$ so $f_b = \pi_b^a f_a$.

Let us also remark that for any $b \leq a \in \mathcal{A}$, $\mu_a(A_a) = \mu_b(A_b)$ for $\mu \in \mathcal{G}(\gamma)$, $A \in \lim G$.

Theorem 3.1 (Extreme measure characterisation (Generalisation of Theorem 7.7 [1])). *Let $\gamma = (G, F)$ be a specification, let $G(a)$ be finite sets for any $a \in \mathcal{A}$, let $F > 0$. $G(\gamma)$ is a convex set. Each $\mu \in \mathcal{G}(\gamma)$ is uniquely determined by its restriction to $\lim \sigma(G)$. Furthermore μ is extreme in $\mathcal{G}(\gamma)$ if and only if for any $A \in \lim \sigma(G)$, $\forall a \in \mathcal{A}$, $\mu_a(A_a) = 0$ or 1.*

Proof. Let us denote π^* the functor from \mathcal{A} to \mathbf{Vect} for which for any $b \leq a$, $\pi_a^{*b} : (L^\infty F(b))^* \rightarrow (L^\infty F(a))^*$ is the dual of π_b^a that send linear forms to linear forms. Then $\mathcal{G}(\gamma)$ is a subspace of the vector space $\lim F^*$ and furthermore for any $a \in \mathcal{A}$ and $p \in [0, 1]$, $p\mu_a + (1-p)\nu_a \in \mathbb{P}(G(a))$ whenever $\mu_a, \nu_a \in \mathbb{P}(G(a))$. Therefore $\mathcal{G}(\gamma)$ is a convex set.

Proposition 3.1, allows us to apply a similar proof, when done with caution, to the one found of Theorem 7.7 in [1]. Let us recall the proof. Let $\mu, \nu \in \mathbb{G}(\gamma)$ such that $\mu|_{\lim i} = \nu|_{\lim i}$. Let $\bar{\mu} = \frac{\mu + \nu}{2}$, then $\bar{\mu} \in \mathcal{G}(\gamma)$. But μ and ν are absolutely continuous with respect to $\bar{\mu}$ therefore for any $a \in \mathcal{A}$ there is $f_a, g_a \in L^\infty(G(a))$ such that $\mu_a = f_a \bar{\mu}_a$ and $\nu_a = g_a \bar{\mu}_a$. By Proposition 3.1, $f, g \in \lim i$. By hypothesis, for any $h \in \lim i$ $\bar{\mu}_a(h_a) = \mu_a(h_a) = \nu_a(h_a)$. Importantly i_a^b is a ring morphism of $L^\infty(G(b))$, i.e. $i_a^b(k_b \cdot h_b) = i_a^b(k_b) \cdot i_a^b(h_b)$. Therefore for any $h, k \in \lim i$, $k \cdot h \in \lim i$; as $f - g \in \lim i$, then it is also the case that $(f - g)^2 \in \lim i$; but for any $a \in \mathcal{A}$,

$$\bar{\mu}_a[(f_a - g_a)^2] = 0 \quad (16)$$

so $f_a = g_a$ $\bar{\mu}_a$ - a.s. Therefore $f\bar{\mu} = g\bar{\mu}$ and $\mu = \nu$.

Showing that $\mu \in \mathcal{G}(\gamma)$ is extreme is equivalent to μ being trivial on $\lim i$ is a direct generalization of Corollary 7.4 [1] thanks to Proposition 3.1. Let $\mu \in \mathcal{G}(\gamma)$ be not trivial on $\lim \sigma(G)$ then there is $A = (A_a, a \in \mathcal{A}) \in \lim \sigma(G)$ such that,

$$\forall a \in \mathcal{A}, \quad 0 < \mu_a(A_a) < 1 \quad (17)$$

Therefore for any $a \in \mathcal{A}$,

$$\mu_a = \mu_a(A_a) \frac{1_{A_a} \cdot \mu_a}{\mu_a(A_a)} + \mu_a(\bar{A}_a) \frac{1_{\bar{A}_a} \cdot \mu_a}{\mu_a(\bar{A}_a)} \quad (18)$$

Furthermore, for any $b \leq a$,

$$i_a^b \frac{1_{A_b}}{\mu_a(A_b)} = \frac{1_{A_a}}{\mu_a(A_b)} = \frac{1_{A_a}}{\mu_a(A_a)} \quad (19)$$

Therefore $\frac{1_{A_a}}{\mu_a(A_a)}, a \in \mathcal{A}$ is in $\lim i$ and so by Lemma 3.1, $\left(\frac{1_{A_a} \cdot \mu_a}{\mu_a(A_a)}, a \in \mathcal{A}\right) \in \mathcal{G}(\gamma)$. Similarly $\left(\frac{1_{\bar{A}_a} \cdot \mu_a}{\mu_a(\bar{A}_a)}, a \in \mathcal{A}\right) \in \mathcal{G}(\gamma)$. In particular there is $0 < p < 1$ so that $\mu = p\nu + (1-p)\nu_1$ with $\nu, \nu_1 \in \mathcal{G}(\gamma)$. Therefore μ is not an extreme measure.

Assume now that $\mu \in \mathcal{G}(\gamma)$ is such that for any $A \in \lim \sigma(A)$, and any $a \in \mathcal{A}$, $\mu_a(A_a) = 0$ or 1 . Suppose that there is $0 < p < 1$ such that $\mu = p\nu + (1-p)\nu_1$ with $\nu, \nu_1 \in \mathcal{G}(\gamma)$. Then for any $a \in \mathcal{A}$, ν_a, ν_{1a} is absolutely continuous with respect to μ_a . Therefore, there are $(f_a \geq 0, a \in \mathcal{A}), (g_a \geq 0, a \in \mathcal{A})$ both in $\prod_{a \in \mathcal{A}} L^\infty(G(a))$ such that $\nu = f\mu$ and $\nu_1 = g\mu$. As $\nu, \nu_1 \in \mathcal{G}(\gamma)$, then by Lemma 3.1, $f, g \in \lim i$. Therefore, for all $a \in \mathcal{A}$,

$\mu_a(f_a) = 0$ or for all $a \in \mathcal{A}$, $\mu_a(g_a) = 0$. So, $\mu = \nu$ or $\mu = \nu_1$ and μ is extreme in $\mathcal{G}(\gamma)$. □

Let us remark that if \mathcal{A} has only one connected component, then for $A \in \lim \sigma(G)$, satisfying $\forall a \in \mathcal{A}, \mu_a(A_a) = 0$ or 1 is equivalent to $\exists a \in \mathcal{A}, \mu_a(A_a) = 0$ or 1 . Indeed, if a, b are in the same connected component, i.e., $a \leq b$ or $b \leq a$, then $\mu_a(A_a) = \mu_b(A_b)$.

References

- [1] H.-O. Georgii, *Gibbs Measures and Phase Transitions*. de Gruyter, 1988.
- [2] M. Giry, “A categorical approach to probability theory,” in *Categorical aspects of topology and analysis*, vol. 915 of *Lecture notes in Mathematics*, pp. 68–85, Springer, 1982.
- [3] G. Sergeant-Perthuis, “A categorical approach to statistical mechanics,” in *Geometric Science of Information* (F. Nielsen and F. Barbaresco, eds.), (Cham), pp. 258–267, Springer Nature Switzerland, 2023.
- [4] G. Sergeant-Perthuis, *Intersection property, interaction decomposition, regionalized optimization and applications*. PhD thesis, Université de Paris, 2021.
- [5] G. Sergeant-Perthuis, “Interaction decomposition for presheaves,” 2021.
- [6] G. Sergeant-Perthuis, “Bayesian/graphoid intersection property for factorisation spaces,” 2021.
- [7] G. Sergeant-Perthuis, “Intersection property and interaction decomposition,” 2021.
- [8] G. Sergeant-Perthuis, “Interaction decomposition for hilbert spaces,” 2021.
- [9] D. Bennequin, O. Peltre, G. Sergeant-Perthuis, and J. P. Vigneaux, “Extra-fine sheaves and interaction decompositions,” 2020.
- [10] G. Sergeant-Perthuis, “Regionalized optimization,” 2022.