



HAL
open science

Timing neurogenesis: a clock or an algorithm?

Julien Pigeon, Bassem A Hassan

► **To cite this version:**

Julien Pigeon, Bassem A Hassan. Timing neurogenesis: a clock or an algorithm?. Current Opinion in Genetics and Development, 2024, 85, pp.102156. 10.1016/j.gde.2024.102156 . hal-04461382

HAL Id: hal-04461382

<https://hal.sorbonne-universite.fr/hal-04461382>

Submitted on 16 Feb 2024

HAL is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers.

L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.

1 **Timing neurogenesis: a clock or an algorithm?**

2 Julien Pigeon¹ and Bassem A. Hassan^{1*}

3 ¹ Institut du Cerveau-Paris Brain Institute (ICM), Sorbonne Université, Inserm, CNRS,
4 Hôpital Pitié-Salpêtrière, Paris, France.

5 *Corresponding author. Email: bassem.hassan@icm-institute.org

7 **Abstract**

8 Emerging evidence supports the existence of dedicated molecular mechanisms under
9 evolutionary selection to control time during neurogenesis. Here, we briefly review these
10 mechanisms and discuss a potentially useful conceptual framework inspired by Computer
11 Science to think about how these biological mechanisms operate during brain development
12 and evolution.

13 **Main text**

14 In a fundamental sense, time represents the continuous progression of existence and events. It
15 unfolds moment by moment and permits a structured chronology where events occur in a
16 specific sequence, not randomly or all at once. This framework allows events to be ordered
17 from the past through the present into the future, which in turn helps us understand causal
18 chains where causes precede effects.

19 In developmental biology, the concept of time is closely intertwined with the pace of
20 embryonic development. This connection can be viewed in at least two ways: the specific
21 sequence of events that occur in a particular chronological order as well as the total amount of
22 time this sequence occupies. Both of these dimensions are found in one of the most studied
23 and evolutionarily conserved biological mechanisms, neurogenesis, the generation of neurons
24 from neural progenitor cells (NPCs). Not only are different neuronal subtypes generated at
25 different time points in a specific order, but the total amount of time dedicated to
26 neurogenesis is itself a species-feature of brain development. Intriguingly, while the
27 sequential stages of brain development share ancestral blueprints, the temporal dynamics can

28 diverge considerably among species. The human brain, for example, undergoes a particularly
29 extended period of neurogenesis compared to rodents and even other primates, allowing for a
30 disproportional increase in the number of neurons, especially in the cerebral neocortex [1],
31 [2]. Thus, there is a correlation between differences in timing and brain size, presumably
32 intimately linked to differences in cognitive capabilities.

33 This review will discuss the temporal dynamics of neurogenesis and the factors that drive
34 neural progenitor cells (NPCs) to their transition into post-mitotic neurons. We will explore a
35 conceptual framework for understanding how and why changes in timing arose during
36 evolution and what this framework predicts as possible future experimental avenues.

37 • **Time in cell state transitions and its control through intrinsic and extrinsic cues**

38 Classical analyses using immunohistochemistry and, more recently, single-cell
39 transcriptomics have revealed that distinct cellular states are associated with specific gene
40 expression patterns. Therefore, the acquisition of a specific cellular identity requires a
41 remodeling of these gene regulatory networks (GRNs) that underlie cell state-specific
42 expression patterns. Studies across species and neural tissue types, from fruit flies to human *in*
43 *vitro* models, have shown that the commitment of a NPC to neurogenesis is influenced by
44 internal and external cellular cues that act on its GRN to determine whether it will continue to
45 self-renew as a progenitor or differentiate into a neuron. One of the causal molecular
46 mechanisms of this dynamic equilibrium between the progenitor and neuron states is
47 mediated by feedback regulation between the Notch signaling pathway and a class of basic
48 Helix-Loop-Helix (bHLH) transcription factors called proneural proteins [3]. When the
49 proneural protein is maintained and stabilized in a NPC, it will overcome the anti-
50 differentiation signal of the Notch pathway and promote neurogenesis ([3]–[5]). Additionally,
51 in the local environment of these NPCs, many secreted morphogens by specific brain regions
52 and the NPCs themselves with their neuronal progeny will weigh on this balance. For
53 example, WNT, FGF, and Shh, play a significant role in maintaining progenitors' self-
54 renewal through different processes to repress proneural genes (reviewed in [6]). In other
55 words, controlling the temporal dynamics of the expression and activity of transcription
56 factors by intrinsic and extrinsic cues can significantly impact the timing of the switch from
57 progenitor self-renewal to differentiation and, thus, the time available for neurogenesis. In the
58 case of the particularly slow pace of human cortical neurogenesis, it has been shown that a
59 human-specific variant of the Notch protein contributes to the developmental tempo by

60 intrinsically promoting increased levels of Notch activity in progenitors, presumably
61 inhibiting proneural protein expression [7**]. On the other hand, the conserved Amyloid
62 Precursor Protein involved in Alzheimer's disease is required in human cortical NPCs to
63 mediate extrinsic WNT signaling to delay the NPC to neuron transition, and this correlates
64 with reduced expression of the proneural protein Neurogenin2 [8**]. Finally, evidence shows
65 that the rate of degradation of transcription factors involved in neurogenesis is slower in
66 humans than in mouse NPCs [9**]. There is also evidence that more stable or longer-lived
67 proneural proteins correlate with [10] and can cause [11*] an increase in neuronal production.
68 Emerging evidence also causally implicates mitochondrial metabolism in the NPC-neuron
69 fate decision [12]. Whether this is mechanistically linked to changes in transcription factor
70 activity or stability remains unclear. Interestingly, however, stem cell programming into
71 neurons by Neurogenin2 overexpression results in large-scale remodeling of cellular
72 organelles and proteomes, suggesting that a link may also exist during normal neurogenesis
73 [13*].

74 Overall, these examples support the view that the temporal dynamics of neurogenesis is
75 controlled mainly by cell-intrinsic mechanisms that appear to converge on regulating the
76 expression and activity levels of transcription factors and that these mechanisms operate on
77 different time scales in different species. However, this only pushes the question from why
78 the phenotypes (i.e. neurogenesis) emerge on different time scales to why the intrinsic cellular
79 mechanisms operate on different time scales. This question is particularly intriguing because
80 the operant mechanisms discovered so far are highly conserved across eukaryotes. Although
81 mitochondrial homeostasis and metabolism or protein turnover seem to be slower in humans
82 than in mice, for example, there is no evidence that the highly conserved enzymes that carry
83 out these reactions have intrinsically slower substrate processing rates. Finally, given that all
84 these mechanisms appear to converge on the activity of transcription factors that are also
85 themselves highly conserved, the conundrum seems even more perplexing. One way to think
86 about the problem from a broader perspective is to shift from a mechanistic view to a
87 theoretical view in order to provide a general framework into which the currently understood
88 mechanisms would fit. This would guide future hypotheses that could be tested with
89 mechanistic approaches. What might such a theoretical framework look like?

90 **Time from Computer Science to Biology**

91 Perhaps our human conception of time as something to be measured with clocks misleads our
92 understanding of developmental time. In Computer Science, time is a measure of
93 computational complexity. It describes the number of elementary operations an algorithm
94 uses to process a set of input data. Interestingly, this measurement, known as time complexity,
95 characterizes how the execution time of an algorithm increases as a function of the size of the
96 input. The larger the input data set, the longer an algorithm will take to complete its
97 processing.

98 To translate this concept to biology, one needs to begin by defining what constitutes an
99 algorithm and what constitutes the source of data. We propose that an operationally useful
100 way of thinking about this is to consider the sum of all cellular biological reactions as the
101 algorithm. After all, during development, various proteins literally “process” input in the
102 shape of other biomolecules or metabolites to produce a cell state transition as a phenotypic
103 output. More importantly, the genome is the only source of biological data for which we have
104 evidence supporting a causal role in generating evolutionary change. As we have argued
105 above, and as decades of molecular genetic mechanistic studies have shown, the biochemistry
106 of fundamental cellular processes, that is to say, the algorithm in this metaphor, is highly
107 conserved. What is changing is the data in the shape of evolutionary innovations in the
108 content of genomic information. A particularly striking example in the context of
109 neurogenesis and neuronal differentiation comes from the transplantation of human cortical
110 pyramidal neurons into the neonatal mouse brain, which develops much faster than the human
111 brain. What these studies show is that the maturation of the human neurons in the mouse brain
112 takes up to a year [14], [15] suggesting that the mechanisms underlying the developmental
113 timing are cell intrinsic and encoded by the human genome. A last important angle to consider
114 is the processing modality of the cellular algorithm. Most, if not all, cellular interaction
115 networks are partially promiscuous. This is referred to as “many-to-many” interaction
116 networks, and they are very prevalent in ligand-receptor and transcription factor dimerization
117 networks, for example, including for key neurogenesis pathways such as WNT, Notch and the
118 proneural proteins. It has recently been argued that such networks are both more robust and
119 have greater computational power [16*] than highly specific, one-to-one interaction networks,
120 likely explaining why they have been favored by evolution. Furthermore, just like for an
121 algorithm, several biological processes run in parallel but converge on a terminal step that
122 induces the transition to the next state, such as proneural protein levels and activity in NPCs
123 (see next paragraph). This terminal step acts as a coincidence detecting gatekeeper ensuring

124 that different aspects of cellular metabolism are all compatible with the impending state
125 transition.

126 In summary, a potentially helpful way to think about how the temporal dynamics of biological
127 processes operate is to assume that a highly conserved algorithm of flexible biochemical
128 interactions processes a variable amount of data encoded by the genome. In this conception,
129 the differences in developmental timing would arise principally from differences in the
130 amount of information encoded by the genome.

131

132 • **Sources of differences in the amount of genomic information**

133 One important source of increase in genomic information during evolution is gene
134 duplication, a major evolutionary driving force [17]. Interestingly, in *Drosophila*, where
135 proneural genes were discovered, there are 58 bHLH genes whereas there are 39 in *C. elegans*
136 and 125 in humans [18], [19]. Moreover, these transcription factors induce the expression of
137 Notch ligands such as Delta and Serrate in flies. Here again, we observe an evolutionary
138 change in the number of proteins. In flies, there are two ligands of Notch, Delta and Serrate,
139 while in vertebrates there are 5 known ligands: Delta like DLL1, DLL3, DLL4 and the Jagged
140 family composed of Jagged1 and Jagged2. In addition, the number of Notch receptors has
141 been multiplied. In flies, there is a single Notch receptor, while in humans, there are 4
142 (Notch1-4) [20]. These ligands and receptors are expressed in complex overlapping spatio-
143 temporal patterns. Altogether, these duplications of many genes multiply the possible
144 interactions between ligands and receptors. The increased complexity in these ligand-receptor
145 interactions carrying both activation and inhibition signals can create delays in signal
146 interpretation into a specific pattern of effector activation [21]. Furthermore, different ligands
147 can transmit opposing patterning information [22] and thus their co-expression would also
148 require more time for that information to be resolved into a fate decision. Finally, many
149 molecular interactions, especially ligand-receptor interactions and gene activation by
150 proneural proteins, are quantitative in that they depend on the absolute amount of active
151 protein. Because levels of any biomolecule at any time are subject to variation due to both
152 stochastic and random noise, the greater the number of proteins in an interaction network, the
153 more noise in the network and, thus, the longer it may take for efficient signal transmission
154 [23]. Having a single Notch receptor and a single ligand makes the signal transmission direct.

155 As discussed above, gene duplication in humans appears to have been taken to a new level
156 with entire families of rapidly evolving genes dedicated to the attenuation of neurogenic
157 signals, as in the example of Noch2NL [24**].

158 Other potential sources of increase in information are gene regulatory elements, notably
159 highly conserved enhancer/repressor sequences that have been subject to variation only in the
160 human lineage called Human Accelerated Regions (HARs). Interestingly, more than half of
161 these regions tested in vitro displayed enhancer activity, specifically in NPCs [25]–[27] such
162 as HAR5 that enhances WNT signaling through FZD8 to promote NPCs self-renewal [28].
163 Another source of quantitative regulation of protein levels is post-transcriptional control by
164 noncoding RNAs, considered as another driver of brain evolution. They represent 10 to 15%
165 of the human genome [29]. There is almost twice the number of miRNAs in humans than in
166 mice and six times that in *Drosophila* [30]. For example, the miR-2115, a great ape-specific
167 miRNA, promotes NPCs proliferation by targeting the ancient gene *ORC4* mutations which
168 cause the Meier-Gorlin microcephalic syndrome [31*], [32]. Furthermore, a primate lncRNA
169 mediates Notch signaling during neuronal development by sequestering miRNA, promoting
170 NPCs self-renewal [33]. The increase in the number of quantitative post-transcriptional
171 inhibitors of neurogenic proteins would be expected to delay the accumulation of these
172 proteins to functional levels and thus delay cell state transitions that depend on these proteins.

173 In summary, the combination of gene duplication and increase in the number of regulators
174 increases both the total amount of information the cellular algorithm must process as well as
175 the amount of noise in this information. The combined effect is to lengthen processing time
176 and thus delay the time interval of cell state transitions during neurogenesis.

177 • **Time, robustness, and size**

178 Why would natural selection have favored these temporal delays over evolutionary time, in
179 the primate and specifically human lineage for example? Needless to say, evolution does not
180 have a plan to produce humans. It must, therefore, be that such a delay in neurogenesis
181 confers a selective advantage for the fitness of the organism. One argument for which there is
182 both experimental and theoretical evidence is that the increase in the number of genes, the
183 promiscuity of protein interactions, and the stochastic noise in molecular processes all confer
184 greater robustness on cell state transitions and developmental patterning [16*], [23] in this
185 scenario, the principal selection pressure is on robustness and the increase in neurogenic time

186 is an unintended but inevitable consequence; an exaptation [34] in the “Gouldian” sense,
187 rather than an adaptation. However, it can also be argued that increasing the amount of time a
188 neural progenitor remains in a progenitor state increases the number of neurons it can
189 generate, and this, in turn, results in increasing the brain’s adaptative cognitive capabilities. In
190 this scenario, the random accident of gene duplication, for example, gave rise to time delay,
191 which gave rise to an increase in brain size with increased cognitive capabilities, and it is this
192 that confers the selective advantage [35]. The truth most probably is that both views are
193 correct. Exaptation and adaptation are synergetic, and both processes can be seen at work at
194 organismal, cellular, and subcellular levels in evolutionary history [36].

195

196 **Conclusion**

197 In this review, we hypothesized that the amount, rather than the type, of information encoded
198 by the genome controls developmental time, with a focus on neurogenesis and its mechanisms
199 as an example. Using a concept borrowed from computer science, we suggest that
200 fundamental cellular metabolism, the *type* of information in this metaphor, constitutes a
201 conserved algorithm that processes information at a conserved pace. In contrast, the total
202 amount of molecules and their interactions constitute the data that this algorithm must
203 process, and the *amount* of this data dictates the time necessary for a developmental process
204 like neurogenesis to be completed. The emerging evidence that conserved processes and
205 genes can have different effects on neurogenesis in different species [8**], [9**] by altering
206 time is testament to the key importance of temporal regulation in brain evolution. If the
207 genomic changes that drive brain evolution have the main effect of altering developmental
208 time, it can be reasonably argued that temporal patterning is the main feature of the genetic
209 code, and it is related to the total amount of “data” encoded by the genome. To test this idea,
210 future experiments and modeling efforts should be directed towards greater accuracy in the
211 measurement of time in molecular processes at high resolution during cell state transitions and
212 deciphering the causal effects of quantitative changes in protein levels on the timing, size and
213 complexity of neurogenic processes.

214 **Acknowledgments:**

215 We thank members of the Hassan team for stimulating discussions. Work relevant to this
216 review has been funded by the Paris Brain Institute-ICM core funding, the Agence Nationale
217 de la Recherche (ANR, 21-CE13-0041-01), Fondation Neuro Glia (2003009NA), Fondation
218 France Alzheimer (2112002NA) and the Roger De Spoelberch Prize.

219 **Author contributions:**

220 J.P. and B.A.H conceived of and wrote the manuscript. J.P. came up with the idea of
221 comparing biological processes to an algorithm.

222 **Competing interest:**

223 The authors declare no competing interests.

224

225 • **Bibliography:**

226 [1] P. Rakic, ‘Evolution of the neocortex: Perspective from developmental biology’, *Nat Rev*
227 *Neurosci*, vol. 10, no. 10, pp. 724–735, Oct. 2009, doi: 10.1038/nrn2719.

228 [2] P. Vanderhaeghen and F. Polleux, ‘Developmental mechanisms underlying the evolution of
229 human cortical circuits’, *Nat Rev Neurosci*, vol. 24, no. 4, Art. no. 4, Apr. 2023, doi:
230 10.1038/s41583-023-00675-z.

231 [3] N. Bertrand, D. S. Castro, and F. Guillemot, ‘Proneural genes and the specification of neural cell
232 types’, *Nature Reviews Neuroscience*, vol. 3, no. 7, Art. no. 7, Jul. 2002, doi: 10.1038/nrn874.

233 [4] R. Kageyama, T. Ohtsuka, H. Shimojo, and I. Imayoshi, ‘Dynamic regulation of Notch signaling
234 in neural progenitor cells’, *Current Opinion in Cell Biology*, vol. 21, no. 6, pp. 733–740, Dec.
235 2009, doi: 10.1016/j.ceb.2009.08.009.

236 [5] F. Guillemot and B. A. Hassan, ‘Beyond proneural: emerging functions and regulations of
237 proneural proteins’, *Current Opinion in Neurobiology*, vol. 42, pp. 93–101, Feb. 2017, doi:
238 10.1016/j.conb.2016.11.011.

239 [6] L. Tiberi, P. Vanderhaeghen, and J. van den Aamele, ‘Cortical neurogenesis and morphogens:
240 diversity of cues, sources and functions’, *Current Opinion in Cell Biology*, vol. 24, no. 2, pp. 269–
241 276, Apr. 2012, doi: 10.1016/j.ceb.2012.01.010.

242 [7**] I. K. Suzuki *et al.*, ‘Human-Specific NOTCH2NL Genes Expand Cortical Neurogenesis
243 through Delta/Notch Regulation’, *Cell*, vol. 173, no. 6, pp. 1370-1384.e16, May 2018, doi:
244 10.1016/j.cell.2018.03.067.

245 - This study identifies recently evolved genes active during human corticogenesis and
246 demonstrates a role for human-specific NOTCH paralogs regulate human cortical neurogenesis.

- 247 [8**] K. Shabani *et al.*, ‘The temporal balance between self-renewal and differentiation of human
248 neural stem cells requires the amyloid precursor protein’, *Sci Adv*, vol. 9, no. 24, p. eadd5002, Jun.
249 2023, doi: 10.1126/sciadv.add5002.
- 250 - This study provides evidence that the temporal balance between self-renewal and differentiation
251 of human cortical NPCs is regulated by the Amyloid Precursor Protein, a causal factor of
252 Alzheimer’s disease.
- 253 [9**] T. Rayon *et al.*, ‘Species-specific pace of development is associated with differences in protein
254 stability’, *Science*, vol. 369, no. 6510, Sep. 2020, doi: 10.1126/science.aba7667.
- 255 - This study identifies a two fold increase in protein stability in human compared to mouse stem
256 cells suggesting that this partially accounts for the slower pace of human development.
- 257 [10]T. Zhang *et al.*, ‘Generation of excitatory and inhibitory neurons from common progenitors via
258 Notch signaling in the cerebellum’, *Cell Reports*, vol. 35, no. 10, p. 109208, Jun. 2021, doi:
259 10.1016/j.celrep.2021.109208.
- 260 [11*] S. Weinberger *et al.*, ‘Evolutionary changes in transcription factor coding sequence
261 quantitatively alter sensory organ development and function’, *eLife*, vol. 6, p. e26402, Apr. 2017,
262 doi: 10.7554/eLife.26402.
- 263 - This study demonstrates that coding sequence evolution contributes to differences in proneural
264 proteins stability which in turn underlie quantitative variation in neuronal development.
- 265 [12]R. Iwata *et al.*, ‘Mitochondria metabolism sets the species-specific tempo of neuronal
266 development’, *Science*, vol. 379, no. 6632, p. eabn4705, Jan. 2023, doi: 10.1126/science.abn4705.
- 267 [13*] A. Ordureau *et al.*, ‘Temporal proteomics during neurogenesis reveals large-scale proteome
268 and organelle remodeling via selective autophagy’, *Molecular Cell*, vol. 81, no. 24, pp. 5082-
269 5098.e11, Dec. 2021, doi: 10.1016/j.molcel.2021.10.001.
- 270 - This study reveals broad organelle remodeling and changes in autophagy during stem cell
271 reprogramming to neurons by the proneural protein Neurogenin2 suggesting a role for these
272 processes in cell fate changes.
- 273 [14]I. Espuny-Camacho *et al.*, ‘Pyramidal Neurons Derived from Human Pluripotent Stem Cells
274 Integrate Efficiently into Mouse Brain Circuits In Vivo’, *Neuron*, vol. 77, no. 3, pp. 440–456, Feb.
275 2013, doi: 10.1016/j.neuron.2012.12.011.
- 276 [15]D. Linaro *et al.*, ‘Xenotransplanted Human Cortical Neurons Reveal Species-Specific
277 Development and Functional Integration into Mouse Visual Circuits’, *Neuron*, vol. 104, no. 5, pp.
278 972-986.e6, Dec. 2019, doi: 10.1016/j.neuron.2019.10.002.
- 279 [16*] H. E. Klumpe, J. Garcia-Ojalvo, M. B. Elowitz, and Y. E. Antebi, ‘The computational
280 capabilities of many-to-many protein interaction networks’, *Cell Systems*, vol. 14, no. 6, pp. 430–
281 446, Jun. 2023, doi: 10.1016/j.cels.2023.05.001.
- 282 - This paper, discusses experimental and theoretical evidence for the role of promiscuous
283 architectures in protein-protein interaction networks in the control of cellular behavior.
- 284 [17]S. Ohno, ‘Gene duplication and the uniqueness of vertebrate genomes circa 1970-1999’, *Semin
285 Cell Dev Biol*, vol. 10, no. 5, pp. 517–522, Oct. 1999, doi: 10.1006/scdb.1999.0332.

- 286 [18]E. Simionato *et al.*, ‘Origin and diversification of the basic helix-loop-helix gene family in
287 metazoans: insights from comparative genomics’, *BMC Evol Biol*, vol. 7, p. 33, Mar. 2007, doi:
288 10.1186/1471-2148-7-33.
- 289 [19]N. E. Baker and N. L. Brown, ‘All in the family: proneural bHLH genes and neuronal diversity’,
290 *Development*, vol. 145, no. 9, May 2018, doi: 10.1242/dev.159426.
- 291 [20]T. Pierfelice, L. Alberi, and N. Gaiano, ‘Notch in the Vertebrate Nervous System: An Old Dog
292 with New Tricks’, *Neuron*, vol. 69, no. 5, pp. 840–855, Mar. 2011, doi:
293 10.1016/j.neuron.2011.02.031.
- 294 [21]R. Kuintzle, L. A. Santat, and M. B. Elowitz, ‘Diversity in Notch ligand-receptor signaling
295 interactions’. bioRxiv, p. 2023.08.24.554677, Aug. 25, 2023. doi: 10.1101/2023.08.24.554677.
- 296 [22]N. Nandagopal, L. A. Santat, L. LeBon, D. Sprinzak, M. E. Bronner, and M. B. Elowitz,
297 ‘Dynamic Ligand Discrimination in the Notch Signaling Pathway’, *Cell*, vol. 172, no. 4, pp. 869-
298 880.e19, Feb. 2018, doi: 10.1016/j.cell.2018.01.002.
- 299 [23]J. E. Ladbury and S. T. Arold, ‘Noise in cellular signaling pathways: causes and effects’, *Trends*
300 *Biochem Sci*, vol. 37, no. 5, pp. 173–178, May 2012, doi: 10.1016/j.tibs.2012.01.001.
- 301 [24**] I. T. Fiddes *et al.*, ‘Human-Specific NOTCH2NL Genes Affect Notch Signaling and Cortical
302 Neurogenesis’, *Cell*, vol. 173, no. 6, pp. 1356-1369.e22, May 2018, doi:
303 10.1016/j.cell.2018.03.051.
- 304 - This paper describes the contribution of human-specific NOTCH2NL to the rapid evolution of
305 the larger human neocortex and suggests a role for the loss of genomic stability at the 1q21.1
306 locus in neurodevelopmental disorders.
- 307 [25]S. Whalen *et al.*, ‘Machine learning dissection of human accelerated regions in primate
308 neurodevelopment’, *Neuron*, vol. 111, no. 6, pp. 857-873.e8, Mar. 2023, doi:
309 10.1016/j.neuron.2022.12.026.
- 310 [26]S. Uebbing *et al.*, ‘Massively parallel discovery of human-specific substitutions that alter
311 enhancer activity’, *Proceedings of the National Academy of Sciences*, vol. 118, no. 2, p.
312 e2007049118, Jan. 2021, doi: 10.1073/pnas.2007049118.
- 313 [27]K. M. Girskis *et al.*, ‘Rewiring of human neurodevelopmental gene regulatory programs by
314 human accelerated regions’, *Neuron*, vol. 109, no. 20, pp. 3239-3251.e7, Oct. 2021, doi:
315 10.1016/j.neuron.2021.08.005.
- 316 [28]J. L. Boyd *et al.*, ‘Human-Chimpanzee Differences in a FZD8 Enhancer Alter Cell-Cycle
317 Dynamics in the Developing Neocortex’, *Current Biology*, vol. 25, no. 6, pp. 772–779, Mar. 2015,
318 doi: 10.1016/j.cub.2015.01.041.
- 319 [29]D. H. Geschwind and P. Rakic, ‘Cortical Evolution: Judge the Brain by Its Cover’, *Neuron*, vol.
320 80, no. 3, pp. 633–647, Oct. 2013, doi: 10.1016/j.neuron.2013.10.045.
- 321 [30]E. Berezikov, ‘Evolution of microRNA diversity and regulation in animals’, *Nat Rev Genet*, vol.
322 12, no. 12, pp. 846–860, Nov. 2011, doi: 10.1038/nrg3079.

- 323 [31*] T. J. Nowakowski *et al.*, 'Regulation of cell-type-specific transcriptomes by microRNA
324 networks during human brain development', *Nat Neurosci*, vol. 21, no. 12, pp. 1784–1792, Dec.
325 2018, doi: 10.1038/s41593-018-0265-3.
- 326 - This study identifies a role for a great-ape-specific miRNA in the control of NPC proliferation
327 rates during human brain development.
- 328 [32]D. L. Guernsey *et al.*, 'Mutations in origin recognition complex gene ORC4 cause Meier-Gorlin
329 syndrome', *Nat Genet*, vol. 43, no. 4, pp. 360–364, Feb. 2011, doi: 10.1038/ng.777.
- 330 [33]N. Rani *et al.*, 'A Primate lncRNA Mediates Notch Signaling during Neuronal Development by
331 Sequestering miRNA', *Neuron*, vol. 90, no. 6, pp. 1174–1188, Jun. 2016, doi:
332 10.1016/j.neuron.2016.05.005.
- 333 [34]S. J. Gould and E. S. Vrba, 'Exaptation-A Missing Term in the Science of Form', *Paleobiology*,
334 vol. 8, no. 1, pp. 4–15, 1982.
- 335 [35]A. M. Boddy *et al.*, 'Evidence of a Conserved Molecular Response to Selection for Increased
336 Brain Size in Primates', *Genome Biol Evol*, vol. 9, no. 3, pp. 700–713, Mar. 2017, doi:
337 10.1093/gbe/evx028.
- 338 [36]M. Frenkel-Pinter, A. S. Petrov, K. Matange, M. Travisano, J. B. Glass, and L. D. Williams,
339 'Adaptation and Exaptation: From Small Molecules to Feathers', *J Mol Evol*, vol. 90, no. 2, pp.
340 166–175, Apr. 2022, doi: 10.1007/s00239-022-10049-1.

341