



HAL
open science

Encephalopathies with KCNC1 variants: genotype-phenotype-functional correlations

Jillian M Cameron, Snezana Maljevic, Umesh Nair, Ye Htet Aung, Benjamin Cogné, Stéphane Bézieau, Edward Blair, Bertrand Isidor, Christiane Zweier, André Reis, et al.

► To cite this version:

Jillian M Cameron, Snezana Maljevic, Umesh Nair, Ye Htet Aung, Benjamin Cogné, et al.. Encephalopathies with KCNC1 variants: genotype-phenotype-functional correlations. *Annals of Clinical and Translational Neurology*, 2019, 6, pp.1263 - 1272. 10.1002/acn3.50822 . hal-04513286

HAL Id: hal-04513286

<https://hal.sorbonne-universite.fr/hal-04513286>

Submitted on 20 Mar 2024

HAL is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers.



L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.



Distributed under a Creative Commons Attribution - NonCommercial - NoDerivatives 4.0 International License

RESEARCH ARTICLE

Encephalopathies with *KCNC1* variants: genotype-phenotype-functional correlations

Jillian M. Cameron¹ , Snezana Maljevic², Umesh Nair², Ye Htet Aung², Benjamin Cogné^{3,4}, Stéphane Bézieau^{3,4}, Edward Blair⁵, Bertrand Isidor^{3,4}, Christiane Zweier⁶, André Reis⁶, Mary Kay Koenig⁷, Timothy Maarup⁸, Dean Sarco⁸, Alexandra Afenjar⁹, A. H. M. Mahbubul Huq¹⁰, Mary Kukulich¹¹, Thierry Billette de Villemeur¹², Caroline Nava¹³, Bénédicte Héron¹⁴, Steven Petrou² & Samuel F. Berkovic¹ 

¹Epilepsy Research Centre, Department of Medicine, University of Melbourne, Austin Health, Heidelberg, Melbourne, Australia

²Florey Institute of Neuroscience and Mental Health, University of Melbourne, Melbourne, Australia

³L'institut du thorax, INSERM, CNRS, UNIV Nantes, Nantes, France

⁴Service de génétique médicale, Centre Hospitalier, Université de Nantes, Nantes, France

⁵Oxford Centre for Genomic Medicine, ACE Building, Nuffield Orthopaedic Centre, Windmill Road, Oxford, United Kingdom

⁶Institute of Human Genetics, Friedrich-Alexander-Universität Erlangen-Nürnberg (FAU), Erlangen, Germany

⁷Department of Paediatrics, Division of Child & Adolescent Neurology, The University of Texas McGovern Medical School, Houston, Texas

⁸Southern California Permanente Medical Group, Pasadena, California

⁹Centre de référence des malformations et maladies congénitales du cervelet, Département de génétique médicale, Sorbonne Université, GRC ConCer-LD, AP-HP, Hôpital Armand Trousseau, F-75012Paris, France

¹⁰Wayne State University, Detroit, Michigan

¹¹Genetics Department, Cook Children's Health Care System, Fort Worth, Texas

¹²Sorbonne Université, GRC ConCer-LD, Neuropédiatrie, Trousseau Hospital, AP-HP, Inserm U1141, Paris, France

¹³Département de Génétique, Sorbonne Universités, Institut du Cerveau et de la Moelle épinière, ICM, Inserm U1127, CNRS UMR 7225, AP-HP, Hôpital de la Pitié Salpêtrière, F-75013Paris, France

¹⁴Sorbonne Université, GRC N°19, Service de Neuropédiatrie, Hôpital Trousseau La Roche Guyon (APHP), La Roche Guyon, France

Correspondence

Samuel F Berkovic, Epilepsy Research Centre, 245 Burgundy St., Heidelberg, Melbourne 3084, Australia. Tel: (+613) 9035 7093; Fax: (+613) 9496 2291; E-mail: s.berkovic@unimelb.edu.au

Funding information

We thank Katherine Helbig for facilitating referral of a case and Karen Oliver for assistance. Supported by an NHMRC Program Grant (ID: 1091593) to Prof Berkovic and Petrou.

Received: 1 June 2019; Accepted: 4 June 2019

Annals of Clinical and Translational Neurology 2019; 6(7): 1263–1272

doi: 10.1002/acn3.50822

Abstract

Objective: To analyze clinical phenotypes associated with *KCNC1* variants other than the Progressive Myoclonus Epilepsy-causing variant p.Arg320His, determine the electrophysiological functional impact of identified variants and explore genotype-phenotype-physiological correlations. **Methods:** Ten cases with putative pathogenic variants in *KCNC1* were studied. Variants had been identified via whole-exome sequencing or gene panel testing. Clinical phenotypic data were analyzed. To determine functional impact of variants detected in the K_v3.1 channel encoded by *KCNC1*, *Xenopus laevis* oocyte expression system and automated two-electrode voltage clamping were used. **Results:** Six unrelated patients had a Developmental and Epileptic Encephalopathy and a recurrent *de novo* variant p.Ala421Val (c.1262C > T). Functional analysis of p.Ala421Val revealed loss of function through a significant reduction in whole-cell current, but no dominant-negative effect. Three patients had a contrasting phenotype of Developmental Encephalopathy without seizures and different *KCNC1* variants, all of which caused loss of function with reduced whole-cell currents. Evaluation of the variant p.Ala513Val (c.1538C > T) in the tenth case, suggested it was a variant of uncertain significance. **Interpretation:** These are the first reported cases of Developmental and Epileptic Encephalopathy due to *KCNC1* mutation. The spectrum of phenotypes associated with *KCNC1* is now broadened to include not only a Progressive Myoclonus Epilepsy, but an infantile onset Developmental and Epileptic Encephalopathy, as well as Developmental Encephalopathy without seizures. Loss of function is a key feature, but definitive electrophysiological separation of these phenotypes has not yet emerged.

Introduction

Humans have over 70 genes encoding for different potassium channel subunits. Pathogenic variants in many of these genes have been associated with a variety of neurological, cardiac, and other diseases. Voltage-gated potassium channels are important in setting membrane resting potential, regulating firing patterns and modifying action potential duration and neurotransmitter release.^{1,2} Mutations in genes encoding voltage-gated potassium channels have been implicated in several conditions including spinocerebellar ataxia, paroxysmal movement disorders, and various epilepsies.^{3–5}

KCNC1, which encodes the K_v3.1 subunit, was not known to be associated with human disease until the recent description of Myoclonus Epilepsy and Ataxia due to *KCNC1* mutation (MEAK).^{6,7} MEAK is a type of Progressive Myoclonus Epilepsy (PME) due to a specific recurrent heterozygous missense mutation, p.Arg320His. K_v3.1 expression is largely restricted to the central nervous system, with predominant expression in GABAergic interneurons.⁸ When analyzed *in vitro*, the PME causing mutation was shown to have a dominant-negative loss of function effect.⁶ Subsequently, a truncating variant in *KCNC1*, p.Arg339X, was described in three related individuals with intellectual disability (ID) and dysmorphic features without epilepsy.⁹ This raises the possibility of a wider spectrum of phenotypes associated with pathogenic *KCNC1* variants. We thus sought cases with other variants in *KCNC1* and analyzed the clinical, electrographic, and radiological features.

Here, we describe an emerging genotype-phenotype correlation with pathogenic variants in *KCNC1*. Analysis of ten patients with putative pathogenic variants in *KCNC1* facilitated identification of nine patients with novel pathogenic *KCNC1* variants. Six of these patients carry a recurrent *KCNC1* variant and present with a developmental and epileptic encephalopathy (DEE).¹⁰ Three patients had a contrasting phenotype of developmental encephalopathy *without* seizures and different *KCNC1* variants. The cellular electrophysiological impact of these variants is also presented.

Methods

Ascertainment

We studied 10 cases with putative pathogenic variants in *KCNC1*. Eight cases were referred to the University of Melbourne by direct personal correspondence given our previously published work. Two cases were identified employing the web-based matching platform DECIPHER.¹¹

Variant identification

Variants were identified via clinical whole-exome sequencing (WES) or gene panel testing via the subjects' home institution. The variant in one patient (patient 1) was identified through research WES testing as part of the Deciphering Developmental Disorders study.¹² Consent for genetic testing was obtained through treating institutions using local protocols.

Functional analysis

Mutagenesis and RNA preparation

Mutations were introduced into human *KCNC1* cDNA encoding transcript 1 (NM_001112741; Kv3.1b) or transcript 2 (NM_004976; Kv3.1a) cloned into a pCMV-Entry vector that were obtained from OriGene Technologies using QuikChange Lightning Site-Directed Mutagenesis Kit (Agilent Technologies) according to the manufacturer's instructions. Insertion of mutation was confirmed and additional mutations excluded by Sanger sequencing. cRNA was prepared using the T7 mMessage mMachine kit from Ambion (Austin, TX).

Oocyte preparation and injection

Xenopus oocytes (Dumont stage V or VI) were surgically removed from *Xenopus laevis* and prepared as described previously.¹³ Oocytes were kept in ND96 solution and stored at 17°C. Equal volumes (50 nL) of cRNAs with the concentrations adjusted to 0.2–0.5 µg/µL were injected using Robooinject® (Multi Channel Systems) in the same batch of oocytes plated in 96 well-plates. Recordings were performed in parallel two days after injection. Amplitudes of interest for all currents recorded on the same day were normalized to the mean value obtained for the K_v3.1 wild-type on that day, so that the normalized data from different experiments could be pooled.

Automated oocyte two-microelectrode voltage clamp

Potassium currents in oocytes were recorded at room temperature (20–22°C) on Roboocyte2® (Multi Channel Systems) using prepulled and prepositioned intracellular glass microelectrodes with a resistance of 0.3–1 MΩ when filled with 0.5 mol/L KCl/ 1.5 mol/L KAc. Oocytes were held at –90 mV and perfused with bath solution containing (in mmol/L) 96 NaCl, 2 KCl, 1.8 CaCl₂, 1 MgCl₂ and 5 HEPES, pH 7.5. Currents were sampled at 5 kHz. For the analysis of channel activation, we used 0.5-s long depolarizing steps applied at 10 mV increments,

from -60 mV to $+60$ mV, followed by a step to -90 mV for 0.5 sec to analyze tail currents.⁶

Data analysis

Electrophysiological recordings from the Robocyte2® were extracted using Robocyte2+ (Multi Channel Systems) and analyzed using AxoGraph (AxoGraph Scientific, Sydney, Australia). Plotting of graphs and statistical analysis was performed using GraphPad Prism 7 (GraphPad Software, La Jolla, CA). Maximum current amplitudes were compared at the end of a 0.5-s test pulse to $+60$ mV. The voltage-dependence of channel activation was derived from tail current amplitudes recorded at -90 mV. A Boltzmann function was fit to the current-voltage relationships, $I(V) = I_{\max}/(1 + \exp[(V - V_{0.5})/k]) + C$, where I_{\max} is the maximum tail current amplitude at test potential V , $V_{0.5}$ the half-maximal activation potential, k a slope factor reflecting characteristics of voltage-dependent channel gating and C a constant. All data are shown as mean values \pm SEM. Statistically significant differences were determined using one way ANOVA with Dunnett's or Tukey's multicomparisons correction or with non-parametric Mann-Whitney test.

Results

KCNC1 variants associated with developmental and epileptic encephalopathy (DEE)

We identified six unrelated patients with DEE and a recurrent missense variant p.Ala421Val (c.1262C > T). All were confirmed *de novo* variants. The variant p.Ala421Val is absent from the gnomAD database and predicted as deleterious using *in silico* methods. The six patients were from the United Kingdom, Germany, France, and North America. Patient 1 had a family history of epilepsy on both maternal and paternal sides; the remaining five patients had no significant family history. All presented between birth and 10 months with predominantly generalized seizure types (Table 1); myoclonic seizures were reported in all but one case (patient 2).

Developmental delay without regression was seen in all patients, however, they were able to walk and achieved intelligible speech. Associated neurological features included mild non-progressive ataxia ($n = 4$), as well as hypotonia ($n = 4$). Three patients (3, 4, and 6) exhibited similar dysmorphic features including a large mouth, smooth philtrum, up-slanting palpebral fissure and dental enlargement (Fig. 1). Three patients were noted to have a combination of respiratory co-morbidities (central sleep apnoea, sleep disturbance) and musculoskeletal

abnormalities (pectus excavatum, hypermobile joints). Two of these patients also had gastrointestinal co-morbidities.

All patients had abnormal EEGs. Five patients demonstrated slowing of background rhythms. Four patients demonstrated interictal generalized spike-and-wave discharges, and EEG in the other two patients captured electroclinical seizures with bi-frontal onset. Myoclonus without electrographic correlate was noted in two patients, and normalization of electrographic abnormalities over a period of years was noted in two patients. MRI Brain imaging showed no specific abnormalities.

All patients had moderate-severe intellectual disability. Long-term seizure frequency and response to treatment was variable. Two patients achieved improved seizure control in late childhood (at 9 and 15 years of age, respectively) with combination antiepileptic therapy, and one patient had a good clinical response to vagal nerve stimulator insertion. The remaining three patients had ongoing refractory seizures despite multiple antiepileptic medications.

A single patient with DEE and *KCNC1* missense variant p.Ala513Val (c.1538C > T) was also identified. The electro-clinical phenotype was of Epilepsy of Infancy with Focal Migrating Seizures (EIFMS), and the patient died at 6 months of age from cardiorespiratory arrest. Recognized genetic causes of EIFMS were not detected on a 77 gene panel testing. The *KCNC1* variant was confirmed *de novo* but it appears once in gnomAD, and thus was considered a variant of uncertain significance.

KCNC1 variants associated with developmental encephalopathy (DE) without seizures

Three patients, including a mother-son pair, presented with a developmental encephalopathy without seizures (Table 2). The mother-son pair with a truncating variant p.Gln492X (c.1474C > T) had mild-moderate intellectual disability, no dysmorphic features and the son developed a nephroblastoma at four years of age. No other causative variants were identified on WES. The third case had missense variant p.Arg317His (c.950G > A), and presented with mild intellectual disability and autism. The patient had never been noted to have a seizure. A 23-hour EEG video to investigate episodes of staring revealed rare irregular generalized spike wave discharges, with no clinical seizures. Brain MRI revealed cerebellar and posterior pontine atrophy. Trio WES also revealed a maternally inherited *MAOA* variant.

Functional analysis

To determine functional impact of variants detected in the $K_v3.1$ channel (Fig. 2A), encoded by *KCNC1*, we used

Table 1. Clinical features of patients with Developmental and Epileptic Encephalopathy and recurrent *KCNC1* missense variant p.Ala421Val.

Patient	1	2	3	4	5	6
Inheritance	De novo	De novo	De novo	De novo	De novo	De novo
Gender	F	M	F	M	F	M
Age at onset	Birth	3 months	10 months	9 months	9.5 months	5 months
Seizure types	Myoclonic Absence GTCS	Focal to bilateral tonic clonic Focal	GTCS Focal Myoclonic	Myoclonic Atypical absence Atonic GTCS	Myoclonic	Myoclonic GTCS Absence with myoclonus Focal to bilateral tonic clonic
Response to treatment	Refractory	Response to LMT and VPA	Refractory then response to VPA Clobazam Stiripentol	Refractory to medication, response to VNS	Response to LEV and LMT	Response to VPA
Age at walking	2.5 years	3 years	2.5 years	2 years	2.5 years	3 years
Intelligible speech	Yes	Yes	Yes	Yes	Yes	Yes
Cognitive Function	Severe ID	Moderate-severe ID	Moderate-severe ID	Moderate-severe ID	Global developmental delay ²	Moderate ID
Neurological Features	Ataxia ¹	Ataxia ¹ , severe hypotonia	Nil	Ataxia ¹ , mild decreased power, hypotonia	Ataxia ¹ , mild athetosis, hypotonia	Unsteady gait, hypotonia
Co-morbidities	Behavioural disturbance	Esophagitis Sleeping disturbance	Nil	GERD, Chronic diarrhoea, Failure to thrive,	Nil	Sleep disturbance Hypermobile joints Stereotypies
Clinical course	Refractory seizures age 16	Monthly seizures at 3 years	Seizure control improved age 15	<1 seizure every 3 months post VNS	Seizures every 2–3 months	Seizure control improved at age 9

Abbreviations: GERD, gastro-esophageal reflux disease; ID, intellectual disability; GTCS, generalized tonic clonic seizures; LEV, levetiracetam; LMT, lamotrigine; VNS, vagal nerve stimulator; VPA, sodium valproate.

¹Ataxia, where noted, was mild and non-progressive.

²Child too young for grading of intellectual disability.



Figure 1. Facial features of patient 3 showing a large mouth, smooth philtrum, up-slanting palpebral fissure, and dental enlargement.

Table 2. Clinical features of patients with *KCNC1* variants and Developmental Encephalopathy without seizures.

Patient	1	2	3
<i>KCNC1</i> variant	p.Arg317His	p.Gln492X	p.Gln492X
Gender	M	F	M
Cognitive Function	Mild ID	Mild-Moderate ID	Mild-moderate ID
Seizures	No	No	No
Dysmorphic Features	Nil	Nil	Nil
Neurological Features	Autism	Nil	Nil
MRI	Cerebellar and posterior pontine atrophy	Not performed	Not performed
EEG	Rare generalized spike-wave	Not performed	Not performed

Xenopus laevis oocyte expression system and automated two-electrode voltage clamping. Current traces obtained from a set of depolarizing pulses from -60 to $+60$ mV (Fig. 2B) revealed a clear loss of function for all but the Ala513Val variant, being the variant observed once in gnomAD.

Current amplitudes analyzed at the end of the current trace were obtained at maximum voltage ($+60$ mV) and normalized for all variants to the corresponding wild type ($K_v3.1a$ for p.Ala421Val, p.Arg317His, p.Gln492X and p.Arg339X and $K_v3.1b$ for p.Ala513Val) recorded on the

same day, therefore providing the relative current amplitude (Fig. 2C and D). Voltage dependence of activation determined from tail current amplitudes recorded at -90 mV for the variants showing current amplitudes above the background level (p.Gln492X and p.Ala513Val) showed no significant difference compared to the corresponding WT channels (Fig. 2E).

To determine potential dominant-negative effect, we performed coexpression experiments, where both WT and mutant cRNA of the same concentration were injected at a 1:1 ratio. Two variants that showed very small or no




Figure 2. Functional expression of *Kv3.1* variants in *Xenopus laevis* oocytes (A) Schematic of the *Kv3.1* channel with putative positions of variants reported in this study (Orange - Developmental and Epileptic Encephalopathy [DEE] variant; Green - Developmental Encephalopathy variants [DE]; Light blue - variant of uncertain significance). The dashed line shows the distal part of the C-terminus that is present only in the longer transcript variant (*Kv3.1b*) indicating that the A513V variant is only found in this transcript. (B) Representative traces of whole-cell currents recorded from *Xenopus laevis* oocytes injected with the same amount of cRNA encoding *Kv3.1* wild-type (*Kv3.1a* and *Kv3.1b*) and different variants during 0.5 sec voltage steps ranging from -60 mV to $+60$ mV. Insets show blown up tail currents, which were analyzed to generate conductance-voltage relationships for WT and mutant channels shown in E. (C, D) Current amplitudes analyzed at the end of the voltage step to $+60$ mV and normalized to the mean current amplitude of the corresponding WT recorded on the same day; *Kv3.1a* ($n = 136$), A421V ($n = 62$), R317H ($n = 32$), Q492X ($n = 30$), R339X ($n = 48$), and water ($n = 35$); *Kv3.1b* ($n = 15$), A513V ($n = 17$). **** $P < 0.0001$, using one-way ANOVA with Dunnett's multiple comparisons test (C). Mann-Whitney non-parametric test (D) revealed $P = 0.5$. (E) Conductance-voltage relationships for the WT transcripts and variants showing current amplitudes above the background level. $V_{0.5}$ and k values were as follows: for *Kv3.1a* 23 ± 2 mV, 12.8 ± 0.6 ($n = 37$), for *Kv3.1b* 33.8 ± 1.1 mV, 12.80 ± 1.04 ($n = 18$), for Q492X 15 ± 3 mV, 12.7 ± 0.8 ($n = 19$), and for A513V 29 ± 2 mV, 11.6 ± 0.6 ($n = 21$); $V_{0.5}$ *Kv3.1a* vs $V_{0.5}$ *Kv3.1b*, ANOVA with Tukey's multiple comparisons test.

currents, (Arg317His and Arg339X) caused a significant reduction compared to the WT current amplitude indicating a dominant-negative effect (Fig. 3A and 3). For the coexpression of the Gln492X with the WT, the normalized current amplitudes were as follows: WT + H₂O -1.00 ± 0.03 ($n = 111$) and WT + Gln492X -1.02 ± 0.12 ($n = 21$). In both cases, the injected amount of WT cRNA was the same, so that the resulting current amplitude of the coexpression should be the sum of the expressed WT + H₂O (1.00 ± 0.03) and half of the amplitude recorded for the Gln492X alone (0.7 ± 0.1 ; Fig. 2C), that is, with a value of about 1.3. Since this predicted value is larger than the recorded one this might suggest the dominant-negative effect of the Gln492X.

The conductance-voltage relationships for Ala421Val + WT, Arg317His + WT, Gln492X + WT, and Arg339X + WT did not differ from the wild type alone

(Fig. 3C). With no changes seen in the voltage dependence of activation, the observed reduction in current amplitudes may be explained by other mechanisms such as reduced number of mutant channels reaching the membrane or reduced open probability of these channels.

Discussion

We describe the phenotypic features of 10 patients with putative pathogenic variants in *KCNK1*, and also present the electrophysiological impact of these variants. These are summarized in Table 3 and compared to the previously described recurrent p.Arg320His variant that causes progressive myoclonus epilepsy.^{6,7}

Six patients carry the recurrent *KCNK1* missense variant p.Ala421Val and present with a DEE. These are the first reported cases of DEE due to *KCNK1* mutation, and




Figure 3. Dominant-negative effect of Kv3.1 variants. (A) Representative current traces recorded from *Xenopus laevis* oocytes injected with the same amount of cRNA encoding Kv3.1a wild-type with addition of either H₂O or the same amount cRNA encoding Kv3.1 variants in a 1:1 ratio. (B) Current amplitudes analyzed at the end of the voltage step to +60 mV and normalized to the mean current amplitude of WT + H₂O recorded on the same day revealed a significant reduction for R317H and R339X but not for the A421V coexpression, ****P* < 0.001, ***P* < 0.01 using one-way ANOVA with Dunnett's multiple comparisons test; WT + H₂O (*n* = 111), WT + A421V (*n* = 62); WT + R317H (*n* = 11); WT + Q492X (*n* = 21); WT + R339X (*n* = 17). (C) Conductance-voltage relationships of the WT and its coexpressions with A421V, R317H and R339X. *V*_{0.5} and slope factor (*k*) values were as follows: for WT + H₂O 18.2 ± 1.4 mV, 12.4 ± 0.4 (*n* = 27), for WT + A421V 16.6 ± 1.8, 15.2 ± 0.4 (*n* = 36), for WT + R317H 21.1 ± 4.5 mV, 17.8 ± 1.9 (*n* = 8), for WT + Q492X 25 ± 3 mV, 13.9 ± 0.5 (*n* = 12), and for WT + R339X 15.8 ± 3.1 mV, 12.2 ± 1.1 (*n* = 15).

thus broaden the spectrum of phenotypes associated with *KCNC1* missense variants to include infantile onset DEE. The K_v3 subfamily exhibits very fast activation and deactivation kinetics, enabling the generation of very high frequency action potentials.¹ K_v3.1 expression is largely restricted to the central nervous system, predominantly in inhibitory GABAergic interneurons.⁸ An impact of *KCNC1* variants on early development is supported by the fact that embryonic expression of K_v3.1 channels in animal models is well-established,^{14–16} with gradual increases in transcript levels noted after birth in murine models, especially postnatal days 8–14.¹⁷ Embryonic expression when the fast spiking phenotype is not present suggests that these channels may also have other critical roles in early development, including in cell proliferation, migration, and regulation of neuronal and glial properties.^{17,18}

In this cohort of patients with DEE and *KCNC1* variant p.Ala421Val, onset was within the first 10 months of life, with predominantly generalized seizure types seen. Myoclonus was a prominent feature in many patients but the

pattern was not that of a PME. Seizures were generally refractory to treatment, persistent developmental delay without regression and moderate-severe intellectual disability were seen. Mild non-progressive ataxia was a common associated neurological feature. No consistent associated co-morbidities were noted in this cohort, however musculoskeletal, respiratory, and gastrointestinal features were seen in several patients. Functional analysis of *KCNC1* variant p.Ala421Val revealed loss of function through a significant reduction in whole-cell current, but no dominant-negative effect. This loss of function variant may cause disinhibition due to impaired firing of the fast-spiking GABAergic interneurons, conceptually providing a plausible mechanism for the development of seizures.

One patient had a different missense *KCNC1* variant p.Ala513Val, and had the more severe clinical phenotype of EIMFS. This variant appears once in gnomAD, and a different variant at the same amino acid location (p.Ala513Thr) appears three times. This, coupled with the fact that this variant did not show definitive *in vitro*

Table 3. Phenotypes and electrophysiological changes associated with *KCNC1* variants.

Phenotype	Progressive myoclonus epilepsy	Developmental and epileptic encephalopathy	Developmental encephalopathy without seizures		
<i>KCNC1</i> variant	p.Arg320His	p.Ala421Val	p.Arg339X	p.Gln492X	p.Arg317His
Seizure types	Myoclonic tonic-clonic (infrequent)	Myoclonic other generalized focal	Nil	Nil	Nil
Cognitive function	Normal with mild late decline in some	Moderate-severe ID	Moderate ID	Mild-moderate ID	Mild ID
Electrophysiological characteristics					
Whole-cell current	Marked reduction	Marked reduction	Marked reduction	Moderate reduction	Marked reduction
Current-voltage relationship	Gain of function	No change	No change	No change	No change
Dominant negative effect	Yes	No	Yes	Unknown ¹	Yes

¹Not assessed as current with mutant was not markedly reduced (see Methods).

changes, suggests that it is unlikely to be pathogenic even though it occurred *de novo* and other known causes of EIMFS (e.g., *SCN2A* and *KCNT1* variants) were not detected. Although *de novo* mutation is the usual mechanism in DEEs, this highlights caution is necessary in attributing pathogenesis to a novel *de novo* variant in an otherwise persuasive clinico-molecular context.

The other three patients with novel variants in *KCNC1* presented with a different phenotype of developmental encephalopathy (DE) *without* seizures. These cases add to the published literature of three related patients from one family with a different *KCNC1* truncation variant p.Arg339X, with DE but without epilepsy.⁹ Of note, the previously reported cases also had dysmorphic features (including prognathism, dysplastic ears, protruding tongue), which our patients did not.

Interestingly, quantification of mRNA in fibroblasts from a patient with Arg339X variant showed > 50% reduction of *KCNC1* transcript compared to the control data, possibly due to nonsense mediated decay (NMD).⁹ This effect may explain why the observed dominant-negative effect for this mutant was not more pronounced in our experiments. The pathogenicity of this variant may thus be a consequence of both reduced expression of *KCNC1* and a damaging effect of the mutant protein that escaped NMD and interacted with the WT subunits. We were not able to obtain fibroblasts from Gln492X variant carriers. However, given that this variant produces a longer channel protein yielding potassium currents reaching about 70% of the WT, we do not expect the NMD effect would be the critical mechanism. Our functional analysis of the three different *KCNC1* variants associated with this phenotype of DE without seizures reveals that again all variants cause a loss of function, all with reduced whole-cell currents. Moreover, two of the three variants caused a dominant-negative effect, which was not seen for the DEE-causing variant

p.Ala421Val. However, p.Gln492X showed a milder effect so that it is difficult to make conclusions about the electrophysiological features which distinguish these variants from the DEE-causing variant.

Of note, the p.Arg317His variant resulted in functional changes similar to the PME-causing p.Arg320His variant but the phenotype seen in this patient is a DE without epilepsy. The proximity of this variant to the PME causing one is noted, (Arg317 is the third arginine in the S4 segment of K_v3.1), and somewhat challenging to reconcile. However, we did observe that the loss of function and the dominant negative effect were not as prominent as seen for p.Arg320His. Moreover, no shift in the voltage-dependence of activation was found for the Arg317His expressed with the WT. This is consistent with the findings reported for two S4 segment disease variants at the same position in the related K_v3.3. Both of these variants caused a dominant negative effect but only the alteration of the fourth arginine lead to a hyperpolarizing shift of the activation curve.¹⁹ The third and fourth arginine have been shown critical for the voltage sensor domain conformation. Analogous arginines in the crystal structure of the K_v1.2 channel are shown to face S1 and S2 helices, and establish salt bridge interactions with the acidic amino acids.²⁰ Interestingly, removal of the positive charge within the S4 segment also affects protein turnover and surface expression of K_v3.3.²¹ We may thus propose that a similar mechanism may underlie the pathogenicity of Arg317His. This patient was also noted to have a maternally inherited *MAOA* variant. Mutation in *MAOA* has been implicated in DE, autism, and behavioral disturbances,²² with no reports of EEG abnormalities. The patient's mother was unaffected.

A lack of clear distinguishing electrophysiological features between the DEE-causing variants and those associated with DE without epilepsy may, at least in part, be a

reflection of the fact that we have not examined the variants' impact in the context of co-expression of other K_v3 subunits. K_v3 subunits assemble either as homomers or heteromers to form voltage-gated tetrameric potassium ion channels. In vitro, $K_v3.1$ can assemble with $K_v3.3$ and $K_v3.4$, and $K_v3.1$ and $K_v3.2$ co-immunoprecipitate in vivo.^{23,24} $K_v3.1$ and $K_v3.3$ have areas of overlapping expression, especially in the cerebellum, and a degree of functional redundancy is thought to explain why *KCNC1* and *KCNC3* knockout mice have relatively mild phenotypes, but double knockout mice exhibit myoclonus, tremor, and ataxia.^{25–27} Clearly there is significant interaction between subunits, and it is feasible that this may be relevant when considering an emerging spectrum of disease seen with different mutations in *KCNC1*.

The description of these cases expands the phenotypic spectrum seen with mutation in *KCNC1* to now not only include a type of Progressive Myoclonus Epilepsy, but to also include an infantile onset DEE, as well as a DE without seizures. A genotype-phenotype correlation is emerging with respect to pathogenic variants in *KCNC1*, whereby the recurrent missense mutation p.Arg320His causes MEAK, the recurrent missense variant p.Ala421Val gives rise to the infantile onset DEE, and truncation variants result in a developmental encephalopathy without seizures. At this stage, definitive electrophysiological functional correlations for these phenotypes have not emerged, which is a target for future investigation.

Acknowledgments

We thank Katherine Helbig for facilitating referral of a case and Karen Oliver for assistance. Supported by an NHMRC Program Grant (ID: 1091593) to Prof Berkovic and Petrou.

Author Contributions

J.C, S.M, S.P, and S.B were involved in conception and design of the study, acquisition and analysis of data and manuscript drafting and completion. All remaining authors were involved in acquisition and analysis of data and manuscript review.

Conflict of interest

The authors declare no conflicts of interest.

References

- Rudy B, McBain CJ. K_v3 channels: voltage-gated K^+ channels designed for high-frequency repetitive firing. *Trends Neurosci* 2001;24:517–526.

- Kaczmarek LK, Aldrich RW, Chandy KG, et al. International union of basic and clinical pharmacology. C. Nomenclature and properties of calcium-activated and sodium-activated potassium channels. *Pharmacol Rev* 2017;69:1–11.
- Zhang Y, Kaczmarek LK. $K_v3.3$ potassium channels and spinocerebellar ataxia. *J Physiol* 2016;594:4677–4684.
- Kohling R, Wolfart J. Potassium channels in epilepsy. *Cold Spring Harb Perspect Med* 2016;6:a022871.
- Rajakulendran S, Schorge S, Kullmann DM, et al. Episodic ataxia type 1: a neuronal potassium channelopathy. *Neurotherapeutics* 2007;4:258–66.
- Muona M, Berkovic SF, Dibbens LM, et al. A recurrent de novo mutation in *KCNC1* causes progressive myoclonus epilepsy. *Nat Genet* 2015;47:39–46.
- Oliver KL, Franceschetti S, Milligan CJ, et al. Myoclonus epilepsy and ataxia due to *KCNC1* mutation: analysis of 20 cases and K^+ channel properties. *Ann Neurol* 2017;81:677–689.
- Gan L, When Kaczmarek LK. When, where, and how much? Expression of the $K_v3.1$ potassium channel in high-frequency firing neurons. *J Neurobiol* 1998;37:69–79.
- Poirier K, Viot G, Lombardi L, et al. Loss of function of *KCNC1* is associated with intellectual disability without seizures. *Eur J Hum Genet* 2017;25:560–564.
- Scheffer IE, Berkovic S, Capovilla G, et al. ILAE classification of the epilepsies: position paper of the ILAE commission for classification and terminology. *Epilepsia* 2017;58:512–521.
- Firth HV, Richards SM, Bevan AP, et al. DECIPHER: database of chromosomal imbalance and phenotype in humans using ensembl resources. *Am J Hum Genet* 2009;84:524–533.
- Firth HV, Wright CF, Deciphering Developmental Disorders Study. The deciphering developmental disorders (DDD) study. *Dev Med Child Neurol* 2011;53:702–703.
- Petrou S, Ugur M, Drummond RM, et al. P2X7 purinoceptor expression in *Xenopus* oocytes is not sufficient to produce a pore-forming P2Z-like phenotype. *FEBS Lett* 1997;411:339–345.
- Perney TM, Marshall J, Martin KA, et al. Expression of the mRNAs for the $K_v3.1$ potassium channel gene in the adult and developing rat brain. *J Neurophysiol* 1992;68:756–766.
- Feng JJ, Morest DK. Development of synapses and expression of a voltage-gated potassium channel in chick embryonic auditory nuclei. *Hear Res* 2006;216–217:116–126.
- Kuenzel T, Wirth MJ, Luksch H, et al. Increase of $K_v3.1b$ expression in avian auditory brainstem neurons correlates with synaptogenesis in vivo and in vitro. *Brain Res* 2009;1302:64–75.
- Boda E, Hoxha E, Pini A, et al. Brain expression of K_v3 subunits during development, adulthood and aging and in

- a murine model of Alzheimer's disease. *J Mol Neurosci* 2012;46:606–615.
18. Yasuda T, Cuny H, Adams DJ. Kv3.1 channels stimulate adult neural precursor cell proliferation and neuronal differentiation. *J Physiol* 2013;591:2579–2591.
 19. Minassian NA, Lin MC, Papazian DM. Altered Kv3.3 channel gating in early-onset spinocerebellar ataxia type 13. *J Physiol* 2012;590:1599–1614.
 20. Long SB, Campbell EB, Mackinnon R. Voltage sensor of Kv1. 2: structural basis of electromechanical coupling. *Science* 2005;309:903–908.
 21. Zhao J, Zhu J, Thornhill WB. Spinocerebellar ataxia-13 Kv3.3 potassium channels: arginine-to-histidine mutations affect both functional and protein expression on the cell surface. *Biochem J* 2013;454:259–265.
 22. Piton A, Poquet H, Redin C, et al. 20 ans apres: a second mutation in MAOA identified by targeted high-throughput sequencing in a family with altered behavior and cognition. *Eur J Hum Genet* 2014;22:776–783.
 23. Baranauskas G, Tkatch T, Nagata K, et al. Kv3.4 subunits enhance the repolarizing efficiency of Kv3.1 channels in fast-spiking neurons. *Nat Neurosci* 2003;6:258–266.
 24. Weiser M, Vega-Saenz de Miera E, Kentros C, et al. Differential expression of Shaw-related K⁺ channels in the rat central nervous system. *J Neurosci* 1994;14:949–972.
 25. Ho CS, Grange RW, Joho RH. Pleiotropic effects of a disrupted K⁺ channel gene: reduced body weight, impaired motor skill and muscle contraction, but no seizures. *Proc Natl Acad Sci USA* 1997;94:1533–1538.
 26. Espinosa F, McMahon A, Chan E, et al. Alcohol hypersensitivity, increased locomotion, and spontaneous myoclonus in mice lacking the potassium channels Kv3.1 and Kv3.3. *J Neurosci* 2001;21:6657–6665.
 27. McMahon A, Fowler SC, Perney TM, et al. Allele-dependent changes of olivocerebellar circuit properties in the absence of the voltage-gated potassium channels Kv3.1 and Kv3.3. *Eur J Neurosci* 2004;19:3317–3327.