



HAL
open science

Narcolepsy risk loci outline role of T cell autoimmunity and infectious triggers in narcolepsy

Hanna M Ollila, Eilon Sharon, Ling Lin, Nasa Sinnott-Armstrong, Aditya Ambati, Selina M Yogeshwar, Ryan P Hillary, Otto Jolanki, Juliette Faraco, Mali Einen, et al.

► To cite this version:

Hanna M Ollila, Eilon Sharon, Ling Lin, Nasa Sinnott-Armstrong, Aditya Ambati, et al.. Narcolepsy risk loci outline role of T cell autoimmunity and infectious triggers in narcolepsy. *Nature Communications*, 2023, 14, pp.2709. 10.1038/s41467-023-36120-z . hal-04523091

HAL Id: hal-04523091

<https://hal.sorbonne-universite.fr/hal-04523091>

Submitted on 27 Mar 2024

HAL is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers.

L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.

Narcolepsy risk loci outline role of T cell autoimmunity and infectious triggers in narcolepsy

Received: 6 July 2021

Accepted: 17 January 2023

Published online: 15 May 2023

 Check for updates

A list of authors and their affiliations appears at the end of the paper

Narcolepsy type 1 (NT1) is caused by a loss of hypocretin/orexin transmission. Risk factors include pandemic 2009 H1N1 influenza A infection and immunization with Pandemrix®. Here, we dissect disease mechanisms and interactions with environmental triggers in a multi-ethnic sample of 6,073 cases and 84,856 controls. We fine-mapped GWAS signals within HLA (DQ0602, DQB1*03:01 and DPB1*04:02) and discovered seven novel associations (CD207, NAB1, IKZF4-ERBB3, CTSC, DENND1B, SIRPG, PRF1). Significant signals at TRA and DQB1*06:02 loci were found in 245 vaccination-related cases, who also shared polygenic risk. T cell receptor associations in NT1 modulated TRAJ*24, TRAJ*28 and TRBV*4-2 chain-usage. Partitioned heritability and immune cell enrichment analyses found genetic signals to be driven by dendritic and helper T cells. Lastly comorbidity analysis using data from FinnGen, suggests shared effects between NT1 and other autoimmune diseases. NT1 genetic variants shape autoimmunity and response to environmental triggers, including influenza A infection and immunization with Pandemrix®.

The sleep disorder type 1 narcolepsy (NT1) affects ~0.03% individuals across ethnic groups and populations^{1–3} and onset manifestation most commonly occurs in childhood or adolescence. As one of the symptoms, cataplexy, is almost entirely specific to NT1, diagnosis is clinical, although sleep recordings are performed for confirmation. Symptoms are caused by the autoimmune destruction of hypocretin/orexin (HCRT) neurons in the hypothalamus⁴. NT1 is 97% associated with HLA-DQA1*01:02-DQB1*06:02, alleles encoding the DQ heterodimer DQ0602^{5,6}. Other predisposing loci include T-cell receptor (TCR) loci *TRA* and *TRB*, a type 1 interferon response receptor gene, *IFNARI*, as well as other autoimmune-associated genes (*CTSH*, *P2RY11*, *ZNF365*, and *TNFSF4*)^{7–10}. Recent studies identified HCRT, notably C-amidated fragments of secreted HCRT peptides (HCRT_{NH2}), as CD4⁺ T-cell autoantigens^{11–15}.

Triggers of NT1 autoimmunity point to Influenza-A^{9,16,17} and, secondarily, *Streptococcus pyogenes* infections^{18,19}. Onset in children is abrupt and seasonal peaking between spring and summer¹⁶, presumably following a winter infection. Further, multiple countries have reported increased incidence of NT1 4–6 months following the

2009 H1N1 (pH1N1) “swine flu” pandemic^{16,17,20}. Finally, immunization with Pandemrix®, a pH1N1 vaccine created to prevent the 2009 pandemic, is an established trigger for NT1^{17,20,21}. Increased incidence following Pandemrix® was first seen in Northern Europe, with incidence in children increasing from 0.79/100,000 to 6.3/100,000²¹. Specificity is striking, as increased NT1 was later detected in all European countries where Pandemrix® was used, whereas countries using other vaccine brands did not display vaccination-associated increases in incidence^{17,20}. The reason for the vaccine brand specificity may involve differences in flu antigen preparations and/or timing of vaccination when infections peaked in some countries^{17,20}. Frequency of other autoimmune diseases did not increase following Pandemrix® vaccination²⁰.

In this study, we characterize novel genetic factors for NT1 across multiple ethnic groups, performing computational and functional fine mapping. Our findings establish a compelling pathophysiological mechanism for the disease that implicate antigen presentation by DQ0602 to specific CD4⁺ T cells and subsequent CD8⁺ T-cell activation, with applications in the autoimmune disease and vaccination fields.

✉ e-mail: mignot@stanford.edu

Results

GWAS discovers novel risk loci for NT1

5,848 cases and 61,153 controls derived from ten cohorts were used as the initial discovery GWAS sample (Table S1). We found associations in HLA ($P < 10^{-216}$), confirmed previously identified loci (*TRA*, *TRB*, *CTSH*, *IFNAR1*, *ZNF365*, *TNFSF4*) and found 7 novel loci near *CD207*, *NAB1*, *IKZF4-ERBB3*, *CTSC*, *DENND1B*, *SIRPG* and *PRF1* (Figs. 1A and 2A, B, Table 1; Supplementary Figs. 1 and 2). We observed that most associations were shared across all ethnic groups. Significance between-cohort heterogeneity was observed with *TRA*, *SIRPG* and *DENND1B* (Table S2). Finally, as both influenza infections and, in rare cases, immunization with Pandemrix®, associates with NT1²⁰, 245 vaccination induced NT1 cases identified in four countries were also studied. In this sub-sample, we found GWAS significant signals with HLA-DQB1*06:02 and *TRA* rs1154155, as well as shared polygenic risk (Table 1 and Supplementary Fig. 3). The lack of association of other loci is likely due to the small number of individuals with vaccination-related narcolepsy.

Interestingly, GWAS results are unusually rich in missense variants. In addition to HLA polymorphisms, these include a TRAJ24 (F8V) substitution, polymorphisms in langerin (CD207 N288D and K313I), as well as variants in CTSC (I453L) and PRF1 (A91V); the last two are known hypomorphs involved in autosomal recessive conditions with

abnormal sensitivity to viral infections. Finally, we found that the effects of some of these variants colocalized between NT1 and type 1 diabetes (CTSH G11R and SIRPG S286L, posterior probability = 1.0). Functional effects of these missense variants are detailed in Supplementary Data 1 and Supplementary Fig. 2.

Other polymorphisms found in the GWAS and associated with other autoimmune diseases include *ZNF365* (atopic eczema, ankylosing spondylitis, Crohn’s disease, psoriasis, primary sclerosing cholangitis, ulcerative colitis, posterior probability = 1.0), *TNFSF4* (eczema, asthma and allergic diseases, posterior probability = 1.0), *NAB1* (primary biliary cholangitis $r^2 = 0.48$, rheumatoid arthritis, $r^2 = 0.15$) and *IKZF4-ERBB* (vitiligo and alopecia areata $r^2 = 0.36$) (see Table Supplementary Data 1 and Supplementary Fig. 2 for functional descriptions and associations).

NT1 shares variants with other autoimmune diseases

Heritability in NT1 is similar to other pediatric autoimmune diseases²²; GCTA estimated observed scale heritability to be $h^2_{SNP}[ci] = 0.403$ [0.015]. Using a prevalence of 0.03%^{1,3}, we estimate population heritability at $h^2_{SNP}[ci] = 0.231$ [0.0088], consistent with twin studies²³. We found that shared heritability was largest with sleepiness and daytime napping and with autoimmune




Fig. 1 | Multi-ethnic genetic analysis of NT1. A Multi-ethnic analysis conducted in 6073 cases and 84,856 controls reveals significant associations in 13 loci in addition to HLA. The x-axis shows genomic location by chromosome and the y-axis shows $-\log_{10} P$ -values. The red horizontal line indicates the genome-wide significant P -value threshold of 5×10^{-8} . P -values larger than 10^{-75} were set to 10^{-75} (HLA locus has many SNPs with P -value $< 10^{-216}$). Variants are shared at individual level with known autoimmune traits, with notable exception at variants within the *TRA* and *TRB* loci and variants within *CD207* and *IFNAR1* (see Supplementary Data 1). Raw P -values are reported using two-sided fixed-effects meta-analysis. Multiple testing correction has been done at genome-wide level so that variants with nominal P -value under 5×10^{-8} were considered statistically significant. **B** Associated variants are located on chromosome positions that have active eQTLs in blood samples, as evidenced by an analysis using GARFIELD. Enrichment analysis has been binned by P -value threshold

and raw two-sided P -values are reported. **C** When using stratified LD score regression, association within individual blood cell types implicate NK cells, CD4+ T and CD8+ T cells. Statistically significant enrichment is marked with a line corresponding to enrichment P -value = 0.05 (dashed line) and FDR corrected P -value = 0.05 (dotted line). Raw two-sided P -values are reported and we have show significance also by false-discovery rate of 0.05 (dotted line). **D** Global enrichment is seen with autoimmune traits in general using variants that were genome-wide significant. Raw two-sided P -values from hypergeometric test are reported. Fig. 2A, B, E: Raw P -values are reported using two-sided fixed-effects meta-analysis. **E** Overall enrichment was seen with MS and SLE using LDSC. **F** PheWAS with narcolepsy risk variants showed association across different autoimmune traits. positive beta is depicted with square and negative with triangle. For details, see methods.




Fig. 2 | TRA rs1154155 is associated with multiple functional SNPs affecting TRAJ24 and TRAJ28; TRB lead variant affects TRBV 4-2 usage. **A** NT1 association with rs1154155 within the T-cell receptor alpha chain (*TRA*) locus. **B** NT1 association with rs7458379 within the T-cell receptor beta chain (*TRB*) locus. **C** The NT1 association within the *TRA* locus spans a region containing ~30 *TRAJ* genes that contains 4 SNPs in almost perfect LD (rs1154155, rs1483979, rs3764159, rs3764160) over a 18 kb region. **D** Exemplary TCR receptor structure of a TCR containing TCR J24. rs1483979 encodes a leucine (J24*02) to phenylalanine (J24*01) substitution (F5L) within the J24 segment. The substitution is projected within the Complementary Determining Region (CDR) 3 of the corresponding family of receptors, in an area anticipated to interact directly with the peptide presented by HLA. **E** Usage of

TRAJ28*01 in 895 individuals shows similar association with NT1 lead variant rs1154155, with posterior probability of 0.958 between NT1 and increased TRAJ28 usage. **F** T-cell receptor sequencing in CD4+ T memory cells in 60 NT1 patients and 42 matched controls confirmed the effect of rs1483979 on usage of TRAJ28*01 with similar effect seen in the NT1 cases. It also shows a decreasing effect of rs1483979C on TRAJ24 expression, with a ratio of 24 F/L (associated allele over non-associated allele in -0.4 in heterozygous subjects, see Supplementary Fig. 5). The center of the boxplot corresponds to the median corresponding to 50th percentile, the box indicates the upper and lower bounds of the interquartile range corresponding to 25th and 75th percentile, and the min and max values correspond to the plus/minus two interquartile ranges.

traits including psoriasis, multiple sclerosis and systemic lupus erythematosus in particular ($P < 0.05$, Table S3, Fig. 1E). We computed enrichment of NT1-associated genes with publicly available GWAS data at gene and variant level, identifying overlap with immune and infectious traits such as asthma, type 1 diabetes, primary biliary cholangitis, plantar warts and hepatitis B ($P < 0.001$, Fig. 1D–F and Table S4). Almost all variants identified (*ZNF365*, *TNFSF4*, *NAB1*, *IKZF4-ERBB3*, *CTSC*, *DENND1B*, *SIRPG*, and *PRF1*) are the same or strongly linked with markers associated with other autoimmune diseases (Supplementary Data 1).

In case/control studies, NT1 has been associated with various autoimmune diseases in some^{24,25}, but not all²⁶ studies. For example, Chen et al. (2021) found association of narcolepsy with asthma in the general Taiwanese population²⁷. To further examine if autoimmune traits are associated with NT1 in epidemiological samples, we explored association between narcolepsy and autoimmune diseases and asthma in 342,499 participants of the FinnGen cohort, retrieving diagnosis of autoimmune diseases, asthma and narcolepsy. In this sample, narcolepsy (157 participants) was associated with psoriasis (OR = 2.29 [1.07–4.90], $P = 0.033$), hypothyroidism (OR = 4.61[2.39–8.93], $P = 5.19 \times 10^{-6}$), rheumatoid diseases (OR = 2.20[1.00–4.82], $P = 0.049$), asthma (OR = 4.58[3.04–6.89], $P = 3.45 \times 10^{-13}$) and “any” autoimmune disease (OR = 2.07[1.50–2.84], $P = 8.31 \times 10^{-6}$). Of note, since DQ0602 is extremely (OR = 0.03) protective against type 1 diabetes²⁸ and strongly protective (OR = 0.64) against primary biliary cholangitis²⁹; no narcolepsy cases had these dual pathologies. Taken together, these findings suggest shared effects between narcolepsy and other autoimmune diseases at both the epidemiological level and at multiple genetic loci, modulated by HLA genotypes.

Variants involved in antigenic stimulation and infections

The specific polymorphisms in langerin (CD207) we found associated with NT1 have previously been linked to interferon stimulus and influenza uptake by dendritic cells (DC) (Supplementary Data 1). Langerin is a type II transmembrane C-type lectin receptor expressed in Langerhans cells, a specialized type of dendritic cells located exclusively in the respiratory tract and the epidermis, and it recognizes mannose-rich sugars expressed by bacterial, fungal or viral pathogens, including HIV-1 and Influenza-A (Supplementary Data 1).

Our leading variant, rs3815556G, a rare allele, is in complete linkage disequilibrium ($r^2 = 1$) with two coding variants, rs13383830C (N288D) and rs57302492A (K313I) that modulate recognition of bacterial versus viral antigens. The rare Asp-288/Ile-313 haplotype has langerin molecules with enhanced affinity for GlcNAc, present in influenza and other viruses, whereas the other haplotype has higher affinity for high-mannose structures and fucosylated glycans, as well as 6SO4-Gal binding activity. This may potentially allow for protection against a wider range of microorganisms, notably bacterial ones as well (Supplementary Data 1). The NT1-associated variants may thus affect disease predisposition by increasing influenza viral (as opposed to bacterial) uptake and antigen presentation to CD4+ T cells.

In addition to langerin, we identified a regulatory variant near *IL10RB-IFNARI*, rs2096464T. The SNP is a strong eQTL for *IFNARI* expression in various tissues in GTEx. *IFNARI* controls dendritic cell responses to viral infections, notably Influenza-A. We therefore examined *IFNARI* expression in DC following H1N1 infection (PR8 delta NS1), finding that the exact NT1 predisposing SNP (rs2096464) is the major eQTL for this effect ($P = 1.92 \times 10^{-25}$, $\beta = 0.140$), as well as for interferon stimulation ($P = 10^{-33}$, $\beta = 0.215$), as it is in perfect

Table 1 | Association of genetic variants with narcolepsy. Raw P-values are reported using two-sided fixed-effects meta-analysis

Closest gene	chr	rsid	Position	Other Allele	Effect Allele	Allele Frequency	Meta-analysis of transethnic cohorts,				Meta-analysis of vaccination and non-vaccination cohorts							
							No vaccination cases				Meta-analysis of vaccination cases				Meta-analysis of vaccination and non-vaccination cohorts			
							P-value	beta	SE	P-value Vaccination	beta	se	P-value	beta	se	n cases	n controls	
TNFSF4	1	rs10158467	173131493	A	G	0.29	6.64×10^{-8}	0.159	0.029	0.108	0.109	1.77×10^{-08}	0.16	0.028	5601	86,675		
DEKND1B	1	rs12131588	197666111	G	A	0.19	3.52×10^{-9}	0.17	0.029	0.191	0.126	1.46×10^{-09}	0.17	0.028	5601	86,675		
CD207	2	rs3815556	71059153	A	G	0.062	3.33×10^{-10}	0.246	0.039	0.05	0.22	6.62×10^{-11}	0.251	0.039	6073	87,377		
NAB1	2	rs7291718	191486081	C	T	0.192	1.03×10^{-09}	-0.205	0.034	0.039	0.139	1.38×10^{-10}	-0.209	0.033	5601	86,675		
TRB	7	rs7456379	142038166	C	T	0.394	8.03×10^{-12}	0.155	0.023	0.019	0.13	9.40×10^{-13}	0.16	0.022	5998	84,581		
ZNF365	10	rs10995245	64391375	G	A	0.348	4.15×10^{-12}	0.148	0.021	0.302	0.103	2.62×10^{-12}	0.147	0.021	6073	87,377		
CTSC	11	rs7112455	88051293	T	A	0.052	5.58×10^{-09}	0.244	0.042	0.06	0.199	1.12×10^{-09}	0.249	0.041	6073	87,377		
IKZF4-ERBB3	12	rs11171731	56443342	C	T	0.33	5.72×10^{-09}	0.147	0.025	0.019	0.101	4.97×10^{-10}	0.152	0.024	5601	86,675		
TRA	14	rs1154155	23002684	T	G	0.17	2.28×10^{-26}	0.463	0.025	1.65×10^{-13}	0.115	7.31×10^{-06}	0.48	0.024	6073	87,377		
CTSH	15	rs34593439	79234957	G	A	0.11	2.56×10^{-09}	0.215	0.036	0.003	0.147	8.24×10^{-11}	0.227	0.035	6073	87,377		
SIRPB	20	rs6034239	1616137	G	A	0.575	1.21×10^{-08}	0.123	0.022	0.056	0.113	2.61×10^{-09}	0.126	0.021	5992	86,957		
IFNAR1	21	rs2096464	34686049	G	T	0.301	4.55×10^{-14}	-0.179	0.024	0.653	0.115	7.40×10^{-14}	-0.174	0.023	6073	87,377		
Novel loci in Z-score-based analysis																		
PRF1	10	rs78325861	72378489	C	G	0.046	2.27×10^{-06}	-0.338	0.072	0.717	0.432	3.4×10^{-09}	-0.333	0.071	5605	85,481		

Allele frequencies are given for effect allele, position reflects hg19.

LD with the leading variant for the signal (rs6517159, $r^2 = 0.93$, Supplementary Fig. 4). Taken together, these findings suggest that NT1-associated variants may affect disease predisposition by increasing influenza viral (as opposed to bacterial) uptake and antigen presentation to CD4⁺ T cells, although additional mechanisms could be involved.

Fine mapping of multi-loci association in the HLA region

To further fine map the HLA association, we imputed classical HLA class I (HLA-A, HLA-B, HLA-C) and class II (HLA-DRB1, HLA-DQA1, HLA-DQB1, HLA-DPA1 and HLA-DPB1) genes using HIBAG³⁰ and HLA IMP:02³¹ and examined allele associations with NT1. As expected^{5,6}, the strongest association was with *DQA1*01:02-DQB1*06:02* (DQ0602). To delineate additional signals, we performed conditional analysis using stepwise forward regression. We discovered protective associations with *DQA1*01:01* and *DQA1*01:03* (OR = 0.30, $P < 10^{-15}$ and OR = 0.30, $P < 10^{-20}$, respectively) and confirmed predisposing effects of *DQB1*03:01* and *DQA1*01:02* across ethnic groups, as shown before^{5,6,32,33} (OR = 1.23, $P < 0.001$ and OR = 1.47 $P < 1 \times 10^{-6}$, respectively) (Table S5). Finally, controlling for both *DQB1* and *DQA1* effects, a protective association was seen with the *DPB1*04:02* allele (P -value $< 10^{-20}$), whereas smaller predisposing effects were found with *DPB1*05:01* and at HLA class I with *A*11:01*, *B*51:01*, *B*35:01* and *B*35:03*, and protective association with *A*03:01* ($P < 0.01$, Table S5). Taken together, these findings confirm and extend results from previous studies^{6,33} and highlight independent association of both HLA class I and II alleles with NT1.

Antigen Presentation and T-cell involvement in NT1

We next examined whether associations with NT1 were enriched genome-wide on specific enhancers using stratified LD score regression (LDSC) on Epigenome Roadmap cell type and ENTEX tissue-specific annotations ($n = 491$ cell and tissue types)³⁴. Partitioned heritability by cell type categories was enriched in hematopoietic cell lines (observed h^2 at hematopoietic cells = 0.24[0.11], $P = 0.018$) and after partitioning the signal into specific cell subsets, ten cell types showed enrichment with $P < 0.005$. These were either helper or cytotoxic T cells or NK cells (Fig. 1C and Supplementary Data 2). As LDSC does not keep information on the HLA region due to ambiguous linkage disequilibrium, we next examined the contribution of different immune cell types using enrichment analysis with genes close to GWAS significant variants. This analysis further supported the enrichment with CD4⁺ T cells, but also implicated antigen-presenting cells such as monocytes and dendritic cells (P -enrichment < 0.01 , Table S6), reflecting, in addition to langerin and IFNAR1, expression of GWA significant, independently associated HLA-associated genes DQB1 and DPB1 (Supplementary Data 1). Together, these results indicate involvement of antigen presentation to CD4⁺ and CD8⁺ T cells in NT1.

Risk variants in T-cell receptor loci modulate $\alpha\beta$ TCR repertoire

NT1 is the only autoimmune disease with known associations in both HLA and T-cell receptor (TCR) loci (TRA and TRB) (Fig. 2A, B). TCR α and β chains heterodimerize to form biologically functional molecules that recognize peptides presented by HLA. We therefore examined the function of these leading variants by examining effects on T-cell receptor V- or J-gene chain usage using RNA sequencing in 895 individuals³⁵, as well as in 130 individuals sequenced specifically in both memory CD4⁺ and CD8⁺ T cells.

As mentioned above, rs1154155 within TRA is entirely linked with multiple SNPs across ethnic groups (Fig. 2C), one of which, rs1483979, substitutes a leucine to a phenylalanine in the CDR3 area of J24, which is predicted to interact with peptides presented by HLA (Fig. 2D; Supplementary Data 5 and 6). This substitution makes it a prime candidate for a functional effect, should an F allele J24*01 CDR3 sequence interact more favorably with an autoantigen than in the presence of




Fig. 3 | Postulated potential disease mechanisms in autoimmune narcolepsy. (1) Peripheral response: Influenza virions or vaccine protein debris are ingested by DCs facilitated by CD207; flu proteins are processed by cathepsins CTSH and CTSC for presentation by HLA molecules to specific TCR α -bearing CD4+ cells, initiating an immunological synapse and responses to influenza. Presentation by DC is modulated by IFNAR1 in the context of influenza infection and the type 1 INF response. Cross presentation of influenza antigens processed via the MHC class I pathway in concert with TNFSF4 in DCs is necessary to activate CD8+ cells that mature into cytotoxic lymphocytes (CTLs), initiating cell killing of viron infected cells. Activated Th1 CD4+ cells produce cytokines such as IFN γ and IL-2, which augment cytotoxic activity of CTLs via perforin (PRF1). SIRPG on activated T cells may also promote cell-to-cell adhesion and proliferation in this response. (2) CNS autoimmunity: Activated and primed specific CD4+ cells migrate to the CNS, where

they interact with microglia and resident DCs via DQ0602 bound to an influenza-mimic autoimmune-epitope (derived from hypocretin cells), initiating a secondary memory response. Hypocretin cell proteins are processed by cathepsins CTSH and CTSC for presentation by DQ0602 to specific TCR α -bearing CD4+ cells, initiating an immunological synapse and autoimmune response. Chain usage for TRAJ24-2, TRAJ28 and TRBV4-2 is associated with NT1 risk and may be crucial for autoantigen recognition. Further, cross presentation by resident DCs and microglial cells activates specific CD8+ cells via MHC class I binding of another HCRT neuron-derived peptide. These primed cytotoxic CD8+ cells then kill HCRT neurons after recognizing MHC class I (such as A*11:01, associated with NT1 independently of DQ0602) bound, cognate HCRT neuron-derived peptide, may be derived from *RFX4* or *LHX9*, on hypocretin neurons. SIRPG1 on DCs, microglia or activated T cells may also promote cell-to-cell adhesion and proliferation in this response.

J24*02. Making the matter more complex, however, J24 usage is also modulated by rs1154155, such that the F-associated allele is associated with decreased usage in CD4+ cells, as shown by the eQTL plot in Fig. 2F (beta = 0.33, $P < 0.001$). We also computed association in memory CD4+ T cells and in CD8+ T cells and observed a consistent effect on TRA-J28 expression specifically ($P = 2.29 \times 10^{-10}$ and $P = 4.08 \times 10^{-10}$, in CD4+ and in CD8+ T cells, respectively). Finally, we confirmed that these effects are *cis* mediated, as the ratio of J24*01 (F) over J24*02 (L) was only 0.4 in heterozygotes, indicating lower allele expression of F-alleles, with similar effects in other T-cell subpopulations (Supplementary Fig. 5).

In addition to J24-specific effects, rs1154155 is also strongly associated with TRA-J28 expression in total RNA sequencing from blood ($P = 1.36 \times 10^{-10}$, beta = -0.212, Fig. 2E) with posterior probability for shared variant $pp = 0.958$ (see Supplementary Data 3 for all rs1154155 effects) and the findings were also consistent when testing across CD4+ memory cells ($P = 2.29 \times 10^{-10}$). Interestingly, rs1154155 is entirely linked with rs3764159, a polymorphism located 14 bp upstream of TRAJ-28 within the 12-base pair recombination signal sequence spacer, possibly explaining the J28 usage effect (Fig. 2C). Controlling for the NT1 TCRA association by lead QTL SNPs for J24 and J28 usage abolishes all effects, except for a minor peak at rs72638479, itself a minor eQTL ($r^2 = 0.95$ with TRAV8-6, Supplementary Fig. 6).

Within the TRB region, rs7458379 is an eQTL for increased expression of TRBV4-2. However, based on our functional analysis, rs1108955 has the strongest evidence for increasing TRBV4-2 usage

($pp = 0.99$, Table S7, Supplementary Data 4). Furthermore, rs1008955 is in partial LD with rs7458379, but tags independent haplotypes at the TRB locus (Supplementary Fig. 7). Both variants are eQTLs for TRBV4-2 expression but reflect independent signals with NT1, such that analysis conditioning for rs7458379 shows remaining association with rs1108955 ($P = 0.00019$), whereas conditioning the association for rs1108955 removed all association at both the TRB locus and with rs7458379 ($P = 0.72$) (Supplementary Fig. 8). Taken together, this indicates that higher expression of TRBV4-2 is related to NT1 and mediated by rs7458379 and rs1108955, with the latter as the potentially causal variant at this locus.

Discussion

In this study, we explored genetic risk for NT1 and potential disease mechanisms of identified genetic risk factors. The strongest associations were seen within the HLA region. In addition, we confirmed six previously described risk loci (*TRA*, *TRB*, *CTSH*, *IFNAR1*, *ZNF365* and *TNFSF4*) and discovered seven novel associations in *CD207*, *IKZF4-ERBB3*, *NABI*, *CTSC*, *DENND1B*, *SIRPG*, and *PRF1*.

Individual associations and partitioned heritability enrichment analysis indicate a primary role of the immune system for all loci identified. Most of these loci, often to the exact same SNP, have also been involved in other autoimmune diseases (Supplementary Data 1). These findings, together with the fact that narcolepsy is associated with increased risk of other autoimmune diseases in FinnGen, suggest that NT1 is an autoimmune disease, even if it does not meet all

accepted criteria³⁶. Further, most variants identified have effects in antigen-presenting cells (*HLA*, *CTSH*, *TNFSF4*), e.g., dendritic cells (*IFNARI*, *CD207*), T cells (*TRA*, *TRB*, *SIRPG*), T helper cells (*HLA-DQ*, *HLA-DP*) or cytotoxic T cells/NK cells (*HLA-A*, *PRFI*, *NABI*), sketching a remarkably narrow potential disease pathway (Fig. 3). In addition to epidemiological data, examining genetic factors overcomes disease-associated ascertainment bias in the recruitment of constitutive cohorts. Finally, two loci were implicated at GWA significant level in 245 vaccination-associated NT1 (*TRA* and *HLA*), while five other loci replicated nominally at $P < 0.05$ (*CD207*, *NABI*, *TRB*, *IKZF4-ERBB3*, and *CTSH*), with overall strong genetic correlation between sporadic and vaccination-associated cases. This indicates that vaccination-triggered narcolepsy is essentially identical to sporadic narcolepsy.

Unlike what is reported in other autoimmune diseases, however, narcolepsy is strongly associated with TCA and TCB genetic polymorphism that modulate the TCR repertoire in very specific ways. A logical explanation for this observation could be that a TCR-mediated reactivity that involves receptors containing TRAJ24F, TRAJ28 and TRBV4-2 is an important step in the development of narcolepsy, perhaps through TCR recognition of a viral trigger or an autoantigen by CD4⁺ or CD8⁺ cells. This hypothesis is supported by recent studies suggesting usage of TRAJ24 and TRBV4-2 in the DQ0602 recognition of amidated HCRT, a likely autoantigen, as well as to specific influenza peptides with increased reactivity in narcolepsy¹⁴.

Based on these observations, we propose that NT1 is an autoimmune process where influenza A contributes to risk in the presence of HLA-DQA1*01:02-DQB1*06:02 (DQ0602). The involvement of influenza-A may explain why genetic associations found are shared across ethnic groups, as influenza is one of few viruses that act worldwide on a seasonal basis. It also relates to *IFNARI* and that both affect and respond to influenza infections, although other infectious triggers cannot be excluded. Notably, the literature suggests that langerin with Asp-288 and Ile-313 shows no binding to 6SO4-Gal-terminated glycans, increased binding to GlcNAc-terminated structures and overall decreased binding to glycans. This would make langerin more restricted in its ability to bind complex carbohydrates and more able to bind GlcNAc-terminated structures, which overall would favor influenza, but also many other organisms. Similarly, *IFNARI* has been linked to antiviral immunity more generally as well, hence specificity to flu infections cannot be concluded with complete certainty.

The universal genetic association is especially clear for HLA-DQ0602, as it is found with different nearby located HLA-DRB1 alleles: DRB1*15:01 in individuals of primary European (Europe and USA) and Asian (China, Korea, Japan and India) descent, but DRB1*15:03 or DRB1*11:01 in individuals of primary Africa descent^{5,6}. The primacy of DQ0602 over DRB1*15:01 (and thereby DRB5, as LD is complete) is also demonstrated by the fact that the DRB1*15:01-DQA1*01:02-DQB1*06:01 haplotype is not associated with narcolepsy in China and by the fact additional DQ effects are mostly mediated by DQA1 alleles that interact in trans with DQB1*06:02 (i.e., DQA1*01:01 and DQB1*01:03)³³. Consequently, the association with DRB1*15:01 was slightly less significant than association with DQ0602. In contrast to NT1, other autoimmune diseases, such as multiple sclerosis and type 1 diabetes, commonly have different HLA associations or disease presentations across countries, resulting in more complex HLA associations. Type 1 diabetes, for example, is well known to be mostly associated with HLA-DRB1*03:01 and DRB1*04 and associated DQ alleles in Europe, whereas DRB1*04:05-specific effects are evident in Japan, where the disease and these DRB1 alleles are rare^{37–39}.

In this study, a hypomorph of perforin, a gene of critical importance to NK and T-cell cytotoxicity, was protective of NT1. Supporting this, we saw enrichment through tissue-specific partitioned heritability in cytotoxic NK and T cells. Although it is conceivable that NK cells or cytotoxic CD4⁺ T cell could be involved, the most likely explanation is involvement of CD8⁺ T cell in hypocretin cell killing. Indeed, neurons

never express HLA class II, so expression of HLA class I and recognition of hypocretin neuronal antigens would be needed for hypocretin cell pathology to occur. This is also supported by the CTSC association, an enzyme of critical importance to cytotoxic CD8⁺ activation of pro-granzymes⁴⁰. Further, Bernard-Valnet et al.⁴¹ used transgenic mice with expression of a neoantigen in HCRT neurons and found that infusion of CD8⁺ T-cell targeting the neoantigen were able to cause hypocretin cell destruction, while infusion of neoantigen-specific CD4⁺ T-cell alone was insufficient, although CD4⁺ T cells migrated closely to the target neurons. Pedersen et al. (2019) also found NT1-associated CD8⁺ T-cell targeting intracellular transcription factors such as RFX4 and LXH9, known to be enriched in HCRT cells¹⁵. Finally, CD8⁺ mediation of cell killing has also been suggested by observation of a CD8⁺ T-cell infiltrate in a paraneoplastic anti-Ma2 encephalitis case with symptomatic hypocretin cell destruction, although these cases are complex and not associated with DQ0602⁴².

In summary, genetic data indicates T-cell autoimmunity in NT1 with genetic overlap of autoimmune traits. A particularity of the disease is involvement of polymorphisms such as in *IFNARI* and *CD207* that regulate antigen-presenting cell responses to infection. Another peculiarity is strong association with TCR polymorphisms, possibly reflecting oligoclonality of T-cell responses. With epidemiological studies indicating seasonality of disease onset¹⁶, and increased incidence following vaccination with Pandemrix® in Europe^{17,20}, a role of influenza or other infections is likely. CTSH implicates dendritic processing of antigens, perhaps of post-translationally modified HCRT itself^{41–45}. Presentation by DQ0602 to CD4⁺ T cells ensue, with likely involvement of very few autoantigen epitopes and a restricted number of T-cell receptors, explaining the large effect of these loci. Subsequent cell killing of hypocretin neurons by CD8⁺ cells, may be through involvement of other, intracellular autoantigens⁴³, complete the process. Altogether, this work illustrates how GWAS can identify the involvement of different cell types in a specific condition, thus ultimately providing further insight to the possible pathophysiological mechanisms underlying disease onset.

Methods

Study approval

This study has been reviewed and approved by the Stanford University Institutional Review Board (IRB) on Medical Human Subjects (Protocol # 14325, genetic and blood markers in narcolepsy and hypersomnia, Registration # IRB00005136) and by respective IRB panels in each country providing samples for the study. Informed consent was obtained from each participant in accordance with governing institutions. Patients and control subjects in FinnGen provided informed consent for biobank research, based on the Finnish Biobank Act. Alternatively, separate research cohorts, collected prior the Finnish Biobank Act came into effect (in September 2013) and start of FinnGen (August 2017), were collected based on study-specific consents and later transferred to the Finnish biobanks after approval by Fimea, the National Supervisory Authority for Welfare and Health. Recruitment protocols followed the biobank protocols approved by Fimea. The Coordinating Ethics Committee of the Hospital District of Helsinki and Uusimaa (HUS) approved the FinnGen study protocol Nr HUS/990/2017. The FinnGen study is approved by Finnish Institute for Health and Welfare (THL), approval number THL/2031/6.02.00/2017, amendments THL/1101/5.05.00/2017, THL/341/6.02.00/2018, THL/2222/6.02.00/2018, THL/283/6.02.00/2019, THL/1721/5.05.00/2019, Digital and population data service agency VRK43431/2017-3, VRK/6909/2018-3, VRK/4415/2019-3 the Social Insurance Institution (KELA) KELA 58/522/2017, KELA 131/522/2018, KELA 70/522/2019, KELA 98/522/2019, and Statistics Finland TK-53-1041-17. The Biobank Access Decisions for FinnGen samples and data utilized in FinnGen Data Freeze 5 include: THL Biobank BB2017_55, BB2017_111, BB2018_19, BB_2018_34, BB_2018_67, BB2018_71, BB2019_7, BB2019_8, BB2019_26,

Finnish Red Cross Blood Service Biobank 7.12.2017, Helsinki Biobank HUS/359/2017, Auria Biobank AB17-5154, Biobank Borealis of Northern Finland_2017_1013, Biobank of Eastern Finland 1186/2018, Finnish Clinical Biobank Tampere MH0004, Central Finland Biobank 1-2017, and Terveystalo Biobank STB 2018001. (for details, see supplementary materials 1.7 FinnGen). We confirm that all methods were carried out in accordance with relevant guidelines and regulations.

Study subjects

Six-thousand seventy-three unrelated individuals with NT1, some included in prior studies^{8,9}, and 84,856 ancestry-matched controls were included in the study. In addition, 245 individuals with vaccination-related NT1 and 18,862 controls were recruited in Finland ($N=76$ cases and 2796 controls), Sweden ($N=39$ cases and 4894 controls), Norway ($N=82$ cases and 429 controls) and United Kingdom and Ireland ($N=48$ cases and 10,743 controls)⁴⁴⁻⁴⁷. All cases had documented immunization with Pandemrix[®]. All cases had narcolepsy with clear-cut cataplexy and were *DQB1*06:02* positive or had narcolepsy with documented low hypocretin-1 in the cerebrospinal fluid. From FinnGen, we used ICD-10 code G47.4 and ICD-9 code 347, thus including individuals who have narcolepsy with or without cataplexy (for details, see Supplementary Materials 1.7 FinnGen).

Genotyping

Subjects were genotyped with Affymetrix Affy 5.0, Affy 6.0, Affymetrix Axiom CHB1, Affymetrix Axiom EUR, Axiom EAS, Axiom LAT, Axiom AFR, Axiom PMRA and Human Core Exome chip platforms (Table S1). Genotypes were called with Affypipe⁴⁸, Affymetrix genotyping console or Genome Studio. Markers with genotyping quality (call rate < 0.95) or deviation from Hardy-Weinberg equilibrium (P -value < 10^{-6}) were discarded. Samples were checked for relatedness by filtering based on proportion of identity-by-descent using cut off >0.2 in PLINK 1.9 PI_HAT score. One pair of related individuals was removed. If related individuals were a case and a control, cases were retained in the analysis. Three first principal components within each cohort were visualized and outliers removed.

Imputation

We imputed samples by pre-phasing cases and controls together using SHAPEIT v2.2 and imputed with IMPUTE2 v2.3.2^{49,50} and 1000 Genomes phase 1v3 build37 (hg19) in 5 Mb chunks across autosomes. Haplotype reference consortium data was used for the second Stanford collection. For variants having both imputed and genotyped values, the genotyped values were kept, whereas for those individuals where genotype was missing, imputed values were kept.

Genetic analyses and statistics

Analyses for all data sets were performed at Stanford University except for the Finnish and Swedish vaccination-related cases and European Narcolepsy Network samples, which were analyzed by respective study teams using the same analytical methods. GWAS was first performed in each case control group separately using SNPTEST v.2.5.2⁵¹. To adjust for cohort-specific population stratification issues, we used linear regression implemented in SNPTEST method score adjusting for the first ten principal components. Standard post imputation quality control was done. Variants with info score < 0.5 and minor allele frequency (MAF) < 0.01 were removed from the analysis. Signals specific for one genotyping platform only and variants in each locus with heterogeneity P -value < 10^{-20} were removed. We used a fixed-effects model implemented in METAV1.7 with an inverse-variance method based on a fixed-effects model for combining association results⁵². In total, 12,600,187 markers across studies were included in the final case/control meta-analysis. The significance level for statistically significant association was set to genome-wide significance (P -value < 5×10^{-8}), controlling for multiple testing. Overall, test statistics showed no genomic inflation

($\lambda < 1.05$). GCTA was used for heritability and gene-based tests⁵³. Coloc analysis was done using coloc package in R version 3.4.2 (2017-09-28)⁵⁴ and Manhattan and QQ-plots were created with QQman or FUMA. Shared and partitioned heritability was estimated using LD score regression. To compare the main characteristics of the participants, we used a multivariate logistic regression model as implemented in the R glm package. Lastly, we used FUMA and the curated list of GWAS as provided by FUMA to compute gene enrichment analysis for loci that were associated with narcolepsy at genome-wide significant level and to examine the association between narcolepsy loci and previous GWAS⁵⁵. We also computed genetic correlation between autoimmune traits and narcolepsy using LD score regression and estimated PheWAS associations per lead variant in each locus using the Open targets resource <https://genetics.opentargets.org/>.

Comparison of vaccination and non-vaccination cases

Vaccination samples were studied separately for GWAS. To compare genetic architecture of narcolepsy cases following vaccination versus other cases, we first examined association of each GWAS significant SNPs of the primary (non-vaccinated) sample in the vaccination sample. In addition, we computed polygenic risk score in non-vaccination narcolepsy cases and vaccination-related narcolepsy using PRSice⁵⁶ and looked at overlap.

Typing and imputation of HLA variants

High-resolution HLA imputation in 4-digit resolution (2-field, amino acid level) for HLA *A*, *B*, *C*, *DRB1*, *DQA1*, *DQB1*, *DPA1* and *DPB1* was performed using HLA*IMP:02 as implemented in Affymetrix HLA or the HIBAG package in R version 3.1.2 (2014-10-31). HIBAG is an HLA imputation tool that uses attribute bootstrap aggregation of several classifiers (SNPs) to select groups of SNPs that predict HLA type and allows for the use of own HLA reference panels³⁰. Reference HLA types were used from published imputation models and for individuals of primary Asian and African descent obtained with Sirona sequencing⁵⁷ in ethnic-specific populations ($N=500$ individuals of African descent, $N=2000$ individuals of European descent and $N=368$ individuals of Asian descent). Imputation accuracy was further verified by Luminex HLA typing in a subset of samples and accuracy was over 95% for all ethnic groups and common alleles with >5% frequency in population. For all alleles, the accuracies for individuals of European descent were 98% for HLA-A, 97% for HLA-B, 98% for HLA-C, 96% for HLA-DRB1, 100% for HLA-DQA1, 100% for HLA-DQB1, 100% for HLA-DPA1 and 92% for HLA-DPB1. The accuracies for individuals of Asian descent, where allele typing was also available, were 95% for HLA-DRB1, 94% for HLA-DQA1 and 98% for HLA-DQB1.

Analysis of HLA variants

HLA effects in NT1 were analyzed as described before⁶ in 23,410 individuals, including 9789 individuals of primary Asian descent and 13,621 individuals of primary European descent as ancestry-matched cases and controls. Within each ancestry group, HLA alleles were analyzed using additive models and logistic regressions after adjusting for the first 10 population-specific principal components. We identify independent associations using conditional analysis (stepwise forward regression in each cohort). Fixed-effects meta-analysis was used to combine associations using Plink 1.9⁵⁸ and R version 3.2.2. We considered alleles sustaining Bonferroni correction for correction of number of alleles with minor allele frequency over 2% ($N=110$ HLA alleles), thus significance resulting in Bonferroni cut-off $P=0.00045$.

Analysis of expression quantitative trait loci (eQTL)

We used tissue-specific summary statistics from the GTEx consortium and from⁵⁹ to examine total blood-specific effects of associating variants on gene expression^{59,60}. We used Garfield to compute

enrichments using enhancer annotation data from ENCODE provided by the Garfield software⁶¹ and stratified LD score regression to compute tissue-specific enrichment using ENCODE data as provided by the LCSC package³⁴.

Expression assessment in monocyte-derived dendritic cells

We examined how the genetic variants modulated T cell and antigen-presenting (dendritic cell and monocyte) gene expression by RNA sequencing and RNA expression. To examine environment-specific triggers for eQTLs, we challenged dendritic cells with an influenza-A infection or stimulated them with interferon. We recruited individuals free from earlier inflammatory disease, autoimmune disease, chronic metabolic disorders or chronic infectious disorders between 18 and 56 years of age (average 29.9), extracted blood mononuclear cells and differentiated into mononuclear dendritic cells, as previously described⁶². We then extracted RNA from the samples using the RNeasy 96 kit (Qiagen, CAT#74182), according to the manufacturer's protocols and sequenced the samples under baseline, influenza infected and interferon beta 1 (IFNβ1) stimulation (99 baseline, 250 influenza infected, and 227 IFNβ1 stimulated). Five hundred fifty-two pass-filter samples (94 baseline, 243 influenza, and 215 interferon) were sequenced to an average depth of 38 million 76-bp paired-end reads using the Illumina TruSeq kit. We aligned reads to hg19 genome with TopHat, assembled transcriptomes for each sample using StringTie⁶³ and computed transcript quantities using Kallisto⁶⁴. We merged transcriptomes across the same condition and then across all three conditions and removed redundant isoforms using cuffcompare⁶⁵. We performed QTL mapping using the Matrix eQTL⁶⁶ package using an empirically determined number of principal components (PCs) as covariates in each analysis. We tested 0 to 44 PCs (local eQTLs) in increments of two and the number of PCs was chosen to maximize the number of local eQTLs detected. We computed empirical *P*-values by comparing the nominal *P*-values with null *P*-values determined by permuting each gene 1000 times. False-discovery rates were calculated using the qvalue package (<https://github.com/StoreyLab/qvalue>), as previously described⁶⁷.

T-cell receptor eQTL analysis

For this analysis, we used data from 895 individuals that were originally genotyped and sequenced as part of the Depression Genes and Networks Project reported by⁶⁸, identifying short range (*cis*) SNPs and *trans* HLA alleles association with TCR V and J usage as described before³⁵. Briefly, expression/usage of each T-cell receptor alpha and beta V- and J-gene was calculated relative to total chain expression from peripheral blood RNA-sequencing. We mapped sequencing reads as in Battle et al. (Bowtie254 with Tophat55 default parameters) and counted the number of unique reads that mapped to each V/J/C-TCR/Ig gene with a modified version of HTSeq56, which allows reads to map to a sequence of more than one V/D/J/C-gene. We then removed individuals and genes with low read counts, normalized the reads using log transformation and regressed on technical and biological covariates as described in Battle et al. as well. Finally, we quantile-normalized the residuals to a normal distribution. Pearson correlations were used to test associations between genotypes and V- or J-gene expression.

Targeted TCR sequencing in NT1 cases and DQ0602-positive controls

In addition to using the data from Battle et al., (2014) we also conducted TCR sequencing in T cells in 60 individuals with NT1 and 60 healthy DQ0602 individuals using RNA from total CD4+ T cells, CD4+ T memory and CD8+ T-cell populations. We used fastqc to infer quality and trimmed low-quality reads. We then performed barcode demultiplexing, after which local blast was used to align

and extract CDR3s. Linear regression was fit for TRA usage per genotype dosage adjusting for age and gender, RNA-sequencing lane and case/control status as covariates. We also separately analyzed coding consequences for each TRAJ24 containing productive CDR3 fragment, as one of the most significantly associated SNPs was a coding SNP (rs1483979) changing an amino acid Leucine to Phenylalanine. These "LQF" and "FQF" were extracted, and their frequencies were computed. Ratio of FQF/(LQF + FQF) was further computed across all samples.

Reporting summary

Further information on research design is available in the Nature Portfolio Reporting Summary linked to this article.

Data availability

The sumstats generated in this study have been deposited in the Dryad database under <https://doi.org/10.5061/dryad.kd51c5b9b>.

References

- Hublin, C., Partinen, M., Kaprio, J., Koskenvuo, M. & Guilleminault, C. Epidemiology of narcolepsy. *Sleep* **17**, S7–S12 (1994).
- Ohayon, M. M., Priest, R. G., Zulley, J., Smirne, S. & Paiva, T. Prevalence of narcolepsy symptomatology and diagnosis in the European general population. *Neurology* **58**, 1826–1833 (2002).
- Silber, M. H., Krahn, L. E., Olson, E. J. & Pankratz, V. S. The epidemiology of narcolepsy in Olmsted County, Minnesota: a population-based study. *Sleep* **25**, 197–202 (2002).
- Peyron, C. et al. A mutation in a case of early onset narcolepsy and a generalized absence of hypocretin peptides in human narcoleptic brains. *Nat. Med.* **6**, 991–997 (2000).
- Mignot, E. et al. Complex HLA-DR and -DQ interactions confer risk of narcolepsy-cataplexy in three ethnic groups. *Am. J. Hum. Genet.* **68**, 686–699 (2001).
- Ollila, H. M. et al. HLA-DPB1 and HLA class I confer risk of and protection from narcolepsy. *Am. J. Hum. Genet.* **96**, 136–146 (2015).
- Faraco, J. et al. ImmunoChip study implicates antigen presentation to T cells in narcolepsy. *PLoS Genet.* **9**, e1003270 (2013).
- Hallmayer, J. et al. Narcolepsy is strongly associated with the T-cell receptor alpha locus. *Nat. Genet.* **41**, 708–711 (2009).
- Han, F. et al. Genome wide analysis of narcolepsy in China implicates novel immune loci and reveals changes in association prior to versus after the 2009 H1N1 influenza pandemic. *PLoS Genet.* **9**, e1003880 (2013).
- Kornum, B. R. et al. Common variants in P2RY11 are associated with narcolepsy. *Nat. Genet.* **43**, 66–71 (2011).
- Cogswell, A. C. et al. Children with Narcolepsy type 1 have increased T-cell responses to orexins. *Ann. Clin. Transl. Neurol.* **6**, 2566–2572 (2019).
- Jiang, W. et al. In vivo clonal expansion and phenotypes of hypocretin-specific CD4(+) T cells in narcolepsy patients and controls. *Nat. Commun.* **10**, 5247 (2019).
- Latorre, D. et al. T cells in patients with narcolepsy target self-antigens of hypocretin neurons. *Nature* **562**, 63–68 (2018).
- Luo, G. et al. Autoimmunity to hypocretin and molecular mimicry to flu in type 1 narcolepsy. *Proc. Natl Acad. Sci. USA* **115**, E12323–E12332 (2018).
- Pedersen, N. W. et al. CD8(+) T cells from patients with narcolepsy and healthy controls recognize hypocretin neuron-specific antigens. *Nat. Commun.* **10**, 837 (2019).
- Han, F. et al. Narcolepsy onset is seasonal and increased following the 2009 H1N1 pandemic in China. *Ann. Neurol.* **70**, 410–417 (2011).
- Weibel, D. et al. Narcolepsy and adjuvanted pandemic influenza A (H1N1) 2009 vaccines—Multi-country assessment. *Vaccine* **36**, 6202–6211 (2018).

18. Ambati, A. et al. Increased beta-haemolytic group A streptococcal M6 serotype and streptodornase B-specific cellular immune responses in Swedish narcolepsy cases. *J. Intern. Med.* **278**, 264–276 (2015).
19. Aran, A. et al. Elevated anti-streptococcal antibodies in patients with recent narcolepsy onset. *Sleep* **32**, 979–983 (2009).
20. Edwards, K. et al. Meeting report narcolepsy and pandemic influenza vaccination: What we know and what we need to know before the next pandemic? A report from the 2nd IABS meeting. *Biologicals* **60**, 1–7 (2019).
21. Partinen, M. et al. Increased incidence and clinical picture of childhood narcolepsy following the 2009 H1N1 pandemic vaccination campaign in Finland. *PLoS ONE* **7**, e33723 (2012).
22. Yang, J., Lee, S. H., Goddard, M. E. & Visscher, P. M. GCTA: a tool for genome-wide complex trait analysis. *Am. J. Hum. Genet.* **88**, 76–82 (2011).
23. Mignot, E. Genetic and familial aspects of narcolepsy. *Neurology* **50**, S16–S22 (1998).
24. Feketeova, E. et al. Narcolepsy in Slovakia—epidemiology, clinical and polysomnographic features, comorbid diagnoses: a case-control study. *Sleep. Med.* **67**, 15–22 (2020).
25. Martinez-Orozco, F. J., Vicario, J. L., De Andres, C., Fernandez-Arquero, M. & Peraita-Adrados, R. Comorbidity of narcolepsy type 1 with autoimmune diseases and other immunopathological disorders: a case-control study. *J. Clin. Med. Res.* **8**, 495–505 (2016).
26. Barateau, L. et al. Comorbidity between central disorders of hypersomnolence and immune-based disorders. *Neurology* **88**, 93–100 (2017).
27. Chen, T. Y. et al. The association between asthma and narcolepsy: a nationwide case-control study in Taiwan. *Nat. Sci. Sleep.* **13**, 1631–1640 (2021).
28. Simmons, K. M. et al. Failed genetic protection: type 1 diabetes in the presence of HLA-DQB1*06:02. *Diabetes* **69**, 1763–1769 (2020).
29. Darlay, R. et al. Amino acid residues in five separate HLA genes can explain most of the known associations between the MHC and primary biliary cholangitis. *PLoS Genet.* **14**, e1007833 (2018).
30. Zheng, X. et al. HIBAG–HLA genotype imputation with attribute bagging. *Pharmacogenomics J.* **14**, 192–200 (2014).
31. Dilthey, A. et al. Multi-population classical HLA type imputation. *PLoS Comput. Biol.* **9**, e1002877 (2013).
32. Hor, H. et al. Genome-wide association study identifies new HLA class II haplotypes strongly protective against narcolepsy. *Nat. Genet.* **42**, 786–789 (2010).
33. Ollila, H. M., Fernandez-Vina, M. & Mignot, E. HLA-DQ allele competition in narcolepsy: a comment on Tafti et al. DQB1 locus alone explains most of the risk and protection in narcolepsy with cataplexy in Europe. *Sleep* **38**, 147–151 (2015).
34. Finucane, H. K. et al. Partitioning heritability by functional annotation using genome-wide association summary statistics. *Nat. Genet.* **47**, 1228–1235 (2015).
35. Sharon, E. et al. Genetic variation in MHC proteins is associated with T cell receptor expression biases. *Nat. Genet.* **48**, 995–1002 (2016).
36. Kornum, B. R. Narcolepsy Type I as an autoimmune disorder. *Handb. Clin. Neurol.* **181**, 161–172 (2021).
37. Hu, X. et al. Additive and interaction effects at three amino acid positions in HLA-DQ and HLA-DR molecules drive type 1 diabetes risk. *Nat. Genet.* **47**, 898–905 (2015).
38. Miyadera, H. & Tokunaga, K. Associations of human leukocyte antigens with autoimmune diseases: challenges in identifying the mechanism. *J. Hum. Genet.* **60**, 697–702 (2015).
39. Tsutsumi, C. et al. Class II HLA genotype in fulminant type 1 diabetes: A nationwide survey with reference to glutamic acid decarboxylase antibodies. *J. Diabetes Investig.* **3**, 62–69 (2012).
40. Perisic Nanut, M., Sabotic, J., Jewett, A. & Kos, J. Cysteine cathepsins as regulators of the cytotoxicity of NK and T cells. *Front. Immunol.* **5**, 616 (2014).
41. Bernard-Valnet, R. CD8 T cell-mediated killing of orexinergic neurons induces a narcolepsy-like phenotype in mice. *Proc. Natl Acad. Sci. USA* **113**, 10956–10961 (2016).
42. Dauvilliers, Y. et al. Hypothalamic immunopathology in anti-Ma-associated diencephalitis with narcolepsy-cataplexy. *JAMA Neurol.* **70**, 1305–1310 (2013).
43. Wallenius, M. et al. Autoantibodies in Pandemrix(®)-induced narcolepsy: Nine candidate autoantigens fail the conformational autoantibody test. *Autoimmunity* **52**, 185–191 (2019).
44. Bomfim, I. L. et al. The immunogenetics of narcolepsy associated with A(H1N1)pdm09 vaccination (Pandemrix) supports a potent gene-environment interaction. *Genes Immun.* **18**, 75–81 (2017).
45. Heier, M. S. et al. Incidence of narcolepsy in Norwegian children and adolescents after vaccination against H1N1 influenza A. *Sleep. Med.* **14**, 867–871 (2013).
46. O’Flanagan, D. et al. Investigation of an association between onset of narcolepsy and vaccination with pandemic influenza vaccine, Ireland April 2009–December 2010. *Eur. Surveill.* **19**, 15–25 (2014).
47. Ollila, H. M. et al. Genetics of vaccination-related narcolepsy. <https://doi.org/10.1101/169623> (2017).
48. Nicolazzi, E. L., Iamartino, D. & Williams, J. L. AffyPipe: an open-source pipeline for Affymetrix Axiom genotyping workflow. *Bioinformatics* **30**, 3118–3119 (2014).
49. Howie, B., Marchini, J. & Stephens, M. Genotype imputation with thousands of genomes. *G3 (Bethesda)* **1**, 457–470 (2011).
50. Howie, B., Fuchsberger, C., Stephens, M., Marchini, J. & Abecasis, G. R. Fast and accurate genotype imputation in genome-wide association studies through pre-phasing. *Nat. Genet.* **44**, 955–959 (2012).
51. Wellcome Trust Case Control C. Genome-wide association study of 14,000 cases of seven common diseases and 3,000 shared controls. *Nature* **447**, 661–678 (2007).
52. Liu, J. Z. et al. Meta-analysis and imputation refines the association of 15q25 with smoking quantity. *Nat. Genet.* **42**, 436–440 (2010).
53. Lee, S. H., Wray, N. R., Goddard, M. E. & Visscher, P. M. Estimating missing heritability for disease from genome-wide association studies. *Am. J. Hum. Genet.* **88**, 294–305 (2011).
54. Plagnol, V., Smyth, D. J., Todd, J. A. & Clayton, D. G. Statistical independence of the colocalized association signals for type 1 diabetes and RPS26 gene expression on chromosome 12q13. *Bio-statistics* **10**, 327–334 (2009).
55. Watanabe, K., Taskesen, E., van Bochoven, A. & Posthuma, D. Functional mapping and annotation of genetic associations with FUMA. *Nat. Commun.* **8**, 1826 (2017).
56. Choi, S. W. & O’Reilly, P. F. PRSice-2: Polygenic Risk Score software for biobank-scale data. *Gigascience* **8**, giz082 (2019).
57. Wang, C. et al. High-throughput, high-fidelity HLA genotyping with deep sequencing. *Proc. Natl Acad. Sci. USA* **109**, 8676–8681 (2012).
58. Chang, C. C. et al. Second-generation PLINK: rising to the challenge of larger and richer datasets. *Gigascience* **4**, 7 (2015).
59. Westra, H. J. et al. Cell specific eQTL analysis without sorting cells. *PLoS Genet.* **11**, e1005223 (2015).
60. Consortium, G. T. Human genomics. The genotype-tissue expression (GTEx) pilot analysis: multitissue gene regulation in humans. *Science* **348**, 648–660 (2015).
61. Iotchkova, V. et al. GARFIELD classifies disease-relevant genomic features through integration of functional annotations with association signals. *Nat. Genet.* **51**, 343–353 (2019).
62. Ye, C. J. et al. Genetic analysis of isoform usage in the human antiviral response reveals influenza-specific regulation of ERAP2

- transcripts under balancing selection. *Genome Res.* **28**, 1812–1825 (2018).
63. Pertea, M. et al. StringTie enables improved reconstruction of a transcriptome from RNA-seq reads. *Nat. Biotechnol.* **33**, 290–295 (2015).
64. Bray, N. L., Pimentel, H., Melsted, P. & Pachter, L. Near-optimal probabilistic RNA-seq quantification. *Nat. Biotechnol.* **34**, 525–527 (2016).
65. Steijger, T. et al. Assessment of transcript reconstruction methods for RNA-seq. *Nat. Methods* **10**, 1177–1184 (2013).
66. Shabalina, A. A. Matrix eQTL: ultra fast eQTL analysis via large matrix operations. *Bioinformatics* **28**, 1353–1358 (2012).
67. Storey, J. D. & Tibshirani, R. Statistical significance for genomewide studies. *Proc. Natl Acad. Sci. USA* **100**, 9440–9445 (2003).
68. Battle, A. et al. Characterizing the genetic basis of transcriptome diversity through RNA-sequencing of 922 individuals. *Genome Res.* **24**, 14–24 (2014).

Acknowledgements

This study was primarily supported by grants from wake-up narcolepsy, Jazz Pharmaceuticals, donations from narcoleptic patients and, peripherally, by a previously funded NIH-23724 grant (EM). We thank GlaxoSmithKline (GSK) for a collaboration providing funds to examine effect of seasonal influenza vaccination and T-cell receptor sequencing in narcolepsy. Work was also supported by Academy of Finland (#309643), Sigrid Juselius Foundation, Finnish Cultural Foundation, Orion Farnos Research Foundation, Instrumentarium Science Foundation, Oskar Öfflund Foundation, Yrjö Jahnsson Foundation and Jalmar and Rauha Ahokas Foundation (HMO). This work was also supported by the United States Department of Defense (DoD) through the National Defense Science & Engineering Graduate Fellowship (NDSEG) program and by Stanford University through the Stanford Graduate Fellowship program. The work has been supported by the National Institute of Mental Health Grant 5RC2MH089916 for the Depression Genes and Networks Project. The Swedish post-pandemix Narcolepsy genetics, with Tomas Olsson and Ingrid Kockum as PIs, received support from the Swedish medical product agency. Han Fang was supported by the Ministry of Science and Technology (2015CB856405) and NSFC (81420108002,81670087). Selina Yogeshwar received funding from the Einstein Center for Neurosciences and the German-Academic Exchange Service (DAAD). TCR sequencing eQTL data used sample genotype and sequence as part of the depression gene and networks project; NIMH 5RC2MH089916. We want to thank Dr. Mehdi Tafti for identifying overlapping samples from EU-NN and those genotyped at Stanford University. Some of the computing for this project was performed on the Sherlock cluster. We would like to thank Stanford University and the Stanford Research Computing Center for providing computational resources and support that contributed to these research results. We want to acknowledge the participants and investigators of FinnGen study. The FinnGen project is funded by two grants from Business Finland (HUS 4685/31/2016 and UH 4386/31/2016) and the following industry partners: AbbVie Inc., Astra-Zeneca UK Ltd, Biogen MA Inc., Bristol Myers Squibb (and Celgene Corporation & Celgene International II Sàrl), Genentech Inc., Merck Sharp & Dohme Corp, Pfizer Inc., GlaxoSmithKline Intellectual Property Development Ltd., Sanofi US Services Inc., Maze Therapeutics Inc., Janssen Biotech Inc, Novartis AG, and Boehringer Ingelheim. Following biobanks are acknowledged for delivering biobank samples to FinnGen: Auriia Biobank (www.auria.fi/biobankki), THL Biobank (www.thl.fi/biobank), Helsinki Biobank (www.helsinginbiobankki.fi), Biobank Borealis of Northern Finland (<https://www.ppshp.fi/Tutkimus-ja-opetus/Biobankki/Pages/Biobank-Borealis-briefly-in-English.aspx>), Finnish Clinical Biobank Tampere (www.tays.fi/en-US/Research_and_development/Finnish_Clinical_Biobank_Tampere), Biobank of Eastern Finland (www.ita-suomenbiopankki.fi/en), Central Finland Biobank (www.ksshp.fi/fi-FI/Potilaalle/Biopankki), Finnish Red Cross Blood Service Biobank (www.veripalvelu.fi/verenluovutus/biopankkitoiminta) and Terveystalo Biobank (www.terveystalo.com/fi/Yritystietoa/Terveystalo-Biopankki/Biopankki/). All Finnish Biobanks are members of BBMRI.fi infrastructure (www.bbMRI.fi). Finnish Biobank Cooperative-FINBB (<https://finbb.fi>) is the coordinator of BBMRI-ERIC operations in Finland. The Finnish biobank data can be accessed through the Fingenious® services (<https://site.fingenious.fi/en/>) managed by FINBB.

Author contributions

H.M.O. and E.J.M. designed the study. H.M.O., E.S., L.L., R.P.H., A. Ambati, J.Y., K.T., T.M., N.R., Z.K., and N.S.A. ran experiments or performed analyses, H.M.O., E.S., N.S.A., and S.M.Y. prepared figures and tables, H.M.O., S.M.Y., and E.J.M. wrote and revised the manuscript. All authors (H.M.O., E.S., L.L., N.S.-A., A. Ambati, S.M.Y., R.P.H., O.J., J.F., M.E., G.L., J.Z., F.H., H.Y., X.S.D., J.L., J.Z., S.-C.H., T.W.K., Y.D., L.B., G.J.L., R.F., G.M., J.S., I.A., S.K., M.K.L.B., P.M.T., G.P., F.P., M.M., C.C., S.K.V.d.E., M.L., P.B., T.K., F.J.M.-O., R.P.-A., A.B., J. Montplaisir, A.D., Y.-S.H., FinnGen, P.J., S. Nevsimalova, D.K., A.I., S.O., A.W., P.G., K.S., M.H., B.H., A.S., F.M.C., V.M., E.F., M.W., N.E., H.S., P.H., P.E.H., D.R., Z.P., L.F.-S., C.L.B., J. Mathis, R.K., A. Aran, S. Nampoothiri, T.O., I.K., M. Partinen, M. Perola, B.R.K., S.R., J.W., T.M., H.T., S.S.K., M.S., K.T., M.R., J.K.P., N.R., Z.K., R.O.H., J.H., J.Y. and E.J.M) contributed data. All authors reviewed the manuscript.

Competing interests

The authors declare no competing interests.

Additional information

Supplementary information The online version contains supplementary material available at <https://doi.org/10.1038/s41467-023-36120-z>.

Correspondence and requests for materials should be addressed to Emmanuel J. Mignot.

Peer review information *Nature Communications* thanks Richard Spritz and the other, anonymous, reviewer(s) for their contribution to the peer review of this work. Peer reviewer reports are available.

Reprints and permissions information is available at <http://www.nature.com/reprints>

Publisher's note Springer Nature remains neutral with regard to jurisdictional claims in published maps and institutional affiliations.

Open Access This article is licensed under a Creative Commons Attribution 4.0 International License, which permits use, sharing, adaptation, distribution and reproduction in any medium or format, as long as you give appropriate credit to the original author(s) and the source, provide a link to the Creative Commons license, and indicate if changes were made. The images or other third party material in this article are included in the article's Creative Commons license, unless indicated otherwise in a credit line to the material. If material is not included in the article's Creative Commons license and your intended use is not permitted by statutory regulation or exceeds the permitted use, you will need to obtain permission directly from the copyright holder. To view a copy of this license, visit <http://creativecommons.org/licenses/by/4.0/>.

© The Author(s) 2023

Hanna M. Ollila^{1,2,3,4}, Eilon Sharon^{5,6}, Ling Lin¹, Nasa Sinnott-Armstrong^{5,6}, Aditya Ambati¹, Selina M. Yogeshwar^{1,7,8}, Ryan P. Hillary¹, Otto Jolanki⁵, Juliette Faraco¹, Mali Einen¹, Guo Luo¹, Jing Zhang¹, Fang Han⁹, Han Yan⁹, Xiao Song Dong⁹, Jing Li⁹, Jun Zhang¹⁰, Seung-Chul Hong¹¹, Tae Won Kim¹¹, Yves Dauvilliers¹², Lucie Barateau¹², Gert Jan Lammers^{13,14}, Rolf Fronczek^{13,14}, Geert Mayer^{15,16}, Joan Santamaria¹⁷, Isabelle Arnulf¹⁸, Stine Knudsen-Heier¹⁹, May Kristin Lyamouri Bredahl^{19,20}, Per Medbøe Thorsby²⁰, Giuseppe Plazzi^{21,22}, Fabio Pizza^{21,22}, Monica Moresco^{21,22}, Catherine Crowe²³, Stephen K. Van den Eeden²⁴, Michel Lecendreux²⁵, Patrice Bourgin²⁶, Takashi Kanbayashi^{27,28}, Francisco J. Martínez-Orozco²⁹, Rosa Peraita-Adrados³⁰, Antonio Beneto³¹, Jacques Montplaisir³², Alex Desautels³², Yu-Shu Huang³³, FinnGen³⁴, Poul Jennum³⁴, Sona Nevsimalova³⁵, David Kemlink³⁵, Alex Iranzo^{36,37,38}, Sebastiaan Overeem^{39,40}, Aleksandra Wierzbicka⁴¹, Peter Geisler⁴², Karel Sonka³⁵, Makoto Honda^{43,44}, Birgit Högl⁴⁵, Ambra Stefani⁴⁵, Fernando Morgadinho Coelho⁴⁶, Vilma Mantovani⁴⁷, Eva Feketeova⁴⁸, Mia Wadelius⁴⁹, Niclas Eriksson^{49,50}, Hans Smedje⁵¹, Pär Hallberg⁴⁹, Per Egil Hesla⁵², David Rye⁵³, Zerrin Pelin⁵⁴, Luigi Ferini-Strambi⁵⁵, Claudio L. Bassetti^{56,57}, Johannes Mathis⁵⁷, Ramin Khatami^{57,58}, Adi Aran⁵⁹, Sheela Nampoothiri⁶⁰, Tomas Olsson⁶¹, Ingrid Kockum⁶¹, Markku Partinen^{62,63}, Markus Perola⁶⁴, Birgitte R. Kornum⁶⁵, Sina Rueger⁶⁶, Juliane Winkelmann^{67,68,69}, Taku Miyagawa^{43,70}, Hiromi Toyoda⁴³, Seik-Soon Khor⁷⁰, Mihoko Shimada⁷⁰, Katsushi Tokunaga⁷⁰, Manuel Rivas⁷¹, Jonathan K. Pritchard⁵, Neil Risch⁷², Zoltan Kutalik^{66,73}, Ruth O'Hara^{74,75}, Joachim Hallmayer^{74,75}, Chun Jimmie Ye⁷⁶ & Emmanuel J. Mignot¹ ✉

¹Stanford University, Center for Sleep Sciences and Medicine, Department of Psychiatry and Behavioral Sciences, Palo Alto, CA 94304, USA. ²Institute for Molecular Medicine Finland, HiLIFE, University of Helsinki, Helsinki, Finland. ³Anesthesia, Critical Care, and Pain Medicine, Massachusetts General Hospital and Harvard Medical School, Boston, MA, USA. ⁴Center for Genomic Medicine, Massachusetts General Hospital, Boston, MA, USA. ⁵Department of Genetics, Stanford University, Stanford, CA 94305, USA. ⁶Department of Biology, Stanford University, Stanford, CA 94305, USA. ⁷Department of Neurology, Charité—Universitätsmedizin, 10117 Berlin, Germany. ⁸Charité—Universitätsmedizin Berlin, Einstein Center for Neurosciences Berlin, 10117 Berlin, Germany. ⁹Division of Sleep Medicine, The Peking University People's Hospital, Beijing, China. ¹⁰Department of Neurology, The Peking University People's Hospital, Beijing, China. ¹¹Department of Psychiatry, St. Vincent's Hospital, The Catholic University of Korea, Suwon, Korea. ¹²Sleep-Wake Disorders Center, National Reference Network for Narcolepsy, Department of Neurology, Gui-de-Chauliac Hospital, CHU Montpellier; Institute for Neurosciences of Montpellier (INM), INSERM, Université Montpellier 1, Montpellier, France. ¹³Department of Neurology, Leiden University Medical Center, Leiden, The Netherlands. ¹⁴Stichting Epilepsie Instellingen Nederland (SEIN), Sleep-Wake Centre, Heemstede, The Netherlands. ¹⁵Hephata Klinik, Schimmelpfengstr. 6, 34613 Schwalmstadt, Germany. ¹⁶Philipps Universität Marburg, Baldinger Str., 35043 Marburg, Germany. ¹⁷Neurology Service, Institut de Neurociències Hospital Clínic, University of Barcelona, Barcelona, Spain. ¹⁸Sleep Disorder Unit, Pitié-Salpêtrière Hospital, Assistance Publique—Hôpitaux de Paris, 75013 Paris, France. ¹⁹Norwegian Centre of Expertise for Neurodevelopment Disorders and Hypersomnias (NevSom), Department of Rare Disorders, Oslo University Hospital and University of Oslo, Oslo, Norway. ²⁰Hormone Laboratory, Department of Medical Biochemistry, Oslo University Hospital, Oslo, Norway. ²¹Department of Biomedical and Neuromotor Sciences (DIBINEM), University of Bologna, Via Ugo Foscolo 7, 40123 Bologna, Italy. ²²IRCCS Institute of Neurological Sciences, Bologna, Italy. ²³Mater Private Hospital, Dublin 7, Ireland. ²⁴Division of Research, Kaiser Permanente Northern California, Oakland, CA, USA. ²⁵Pediatric Sleep Center and National Reference Center for Narcolepsy and Idiopathic Hypersomnia Hospital Robert Debre, Paris, France. ²⁶Department of Sleep Medicine, Strasbourg University Hospital, Strasbourg University, Strasbourg, France. ²⁷Department of Neuropsychiatry, Akita University Graduate School of Medicine, Akita, Japan. ²⁸International Institute for Integrative Sleep Medicine (WPI-IIS), University of Tsukuba, Tsukuba, Japan. ²⁹Sleep Unit. Clinical Neurophysiology Service. San Carlos University Hospital. University Complutense of Madrid, Madrid, Spain. ³⁰Sleep and Epilepsy Unit, Clinical Neurophysiology Service, Gregorio Marañón University General Hospital and Research Institute, University Complutense of Madrid (UCM), Madrid, Spain. ³¹Sleep Unit. Medical Center Valencia, Valencia, Spain. ³²Center for Advanced Research in Sleep Medicine, Hôpital du Sacré-Coeur and Department of Neurosciences, University of Montréal, Montréal, QC, Canada. ³³Department of Child Psychiatry and Sleep Center, Chang Gung Memorial Hospital and University, Taoyuan, Taiwan. ³⁴Danish Center for Sleep Medicine, Department of Clinical Neurophysiology, University of Copenhagen, Glostrup Hospital, Glostrup, Denmark. ³⁵Department of Neurology and Centre of Clinical Neurosciences, First Faculty of Medicine, Charles University and General University Hospital, Prague, Czech Republic. ³⁶Institut d'Investigacions Biomèdiques August Pi i Sunyer (IDIBAPS), Barcelona, Spain. ³⁷Department of Neurology, Barcelona, Spain. ³⁸Multidisciplinary Sleep Disorders Unit, Barcelona, Spain. ³⁹Sleep Medicine Center Kempenhaeghe, P.O. Box 61, 5590 AB Heeze, The Netherlands. ⁴⁰Eindhoven University of Technology, Eindhoven, The Netherlands. ⁴¹Department of Clinical Neurophysiology, Institute of Psychiatry and Neurology, Warsaw, Poland. ⁴²Department of Psychiatry and Psychotherapy, University of Regensburg, Regensburg, Germany. ⁴³Department of Psychiatry and Behavioral Sciences, Tokyo Metropolitan Institute of Medical Science, Tokyo, Japan. ⁴⁴Seiwa Hospital, Neuropsychiatric Research Institute, Tokyo, Japan. ⁴⁵Department of Neurology, Medical University Innsbruck (MUI), Innsbruck, Austria. ⁴⁶Universidade Federal de São Paulo, Departamento de Psicobiologia, São Paulo, Brazil. ⁴⁷Center for Applied Biomedical Research (CRBA), St. Orsola-Malpighi University Hospital, Bologna, Italy. ⁴⁸Neurology Department, Medical Faculty of P. J. Safarik University, University Hospital of L. Pasteur Kosice, Kosice, Slovak Republic. ⁴⁹Department of Medical Sciences and Science for Life Laboratory, Uppsala University, Uppsala, Sweden. ⁵⁰Uppsala Clinical Research Center, Uppsala, Sweden. ⁵¹Division of Child and Adolescent Psychiatry, Karolinska Institutet, Stockholm, Sweden. ⁵²Coliseum on Majorstua Clinic, Oslo, Norway. ⁵³Department of Neurology, Emory University School of Medicine, Atlanta, GA, USA. ⁵⁴Faculty of Health Sciences, Hasan Kalyoncu University, Gaziantep, Turkey. ⁵⁵Sleep Disorders Center, Division of Neuroscience, Ospedale San Raffaele, Università Vita-Salute, Milan, Italy. ⁵⁶Neurology Department, EOC, Ospedale Regionale di Lugano, Lugano, Ticino, Switzerland. ⁵⁷Department of Neurology, Inselspital, Bern University Hospital, and University of Bern, Bern, Switzerland. ⁵⁸Center for Sleep Medicine and Sleep Research, Clinic Barmelweid AG, Barmelweid, Switzerland. ⁵⁹Shaare Zedek Medical Center, Jerusalem, Israel. ⁶⁰Department of Pediatric Genetics, Amrita Institute of Medical Sciences & Research Centre, Kerala, India. ⁶¹Department of Clinical Neuroscience, Karolinska Institutet, Stockholm, Sweden. ⁶²Helsinki Sleep Clinic, Vitalmed Research Centre, Helsinki, Finland. ⁶³Department of Clinical Neurosciences, University of Helsinki, Helsinki, Finland. ⁶⁴University of Helsinki, Institute for Molecular Medicine, Finland (FIMM) and Diabetes and Obesity Research Program. University of Tartu, Estonian Genome Center, Tartu, Estonia. ⁶⁵Department of Neuroscience, University of Copenhagen, Copenhagen, Denmark. ⁶⁶Swiss Institute of Bioinformatics, Lausanne, Switzerland. ⁶⁷Institute of Neurogenetics, Helmholtz Zentrum München, German Research Centre for Environmental Health,

Neuherberg, Germany. ⁶⁸Munich Cluster for Systems Neurology (SyNergy), Munich, Germany. ⁶⁹Neurologische Klinik und Poliklinik, Klinikum rechts der Isar der Technischen Universität München, Munich, Germany. ⁷⁰Department of Human Genetics, Graduate School of Medicine, The University of Tokyo, Tokyo, Japan. ⁷¹Department of Biomedical Data Science-Administration, Stanford University, Palo Alto, CA, USA. ⁷²Dept. Epidemiology and Biostatistics, UCSF, 513 Parnassus Avenue, San Francisco, CA 94117, USA. ⁷³University Center for Primary Care and Public Health, University of Lausanne, Lausanne, Switzerland, Lausanne 1010, Switzerland. ⁷⁴Department of Psychiatry and Behavioral Sciences, Stanford University, Palo Alto, CA, USA. ⁷⁵Mental Illness Research Education Clinical Centers (MIRECC), VA Palo Alto, Palo Alto, CA, USA. ⁷⁶Department of Epidemiology & Biostatistics, Institute for Human Genetics, University of California San Francisco, San Francisco, CA, USA. ✉e-mail: mignot@stanford.edu

FinnGen

Thomas Damm Als⁷⁷, Adam Ziemann⁷⁸, Ali Abbasi⁷⁸, Anne Lehtonen⁷⁸, Apinya Lertratanakul⁷⁸, Bridget Riley-Gillis⁷⁸, Fedik Rahimov⁷⁸, Howard Jacob⁷⁸, Jeffrey Waring⁷⁸, Mengzhen Liu⁷⁸, Nizar Smaoui⁷⁸, Relja Popovic⁷⁸, Adam Platt⁷⁹, Athena Matakidou⁷⁹, Benjamin Challis⁷⁹, Dirk Paul⁷⁹, Glenda Lassi⁷⁹, Ioanna Tachmazidou⁷⁹, Antti Hakanen⁸⁰, Johanna Schleutker⁸⁰, Nina Pitkänen⁸⁰, Perttu Terho⁸⁰, Petri Virolainen⁸⁰, Arto Mannermaa⁸¹, Veli-Matti Kosma⁸¹, Chia-Yen Chen⁸², Heiko Runz⁸², Sally John⁸², Sanni Lahdenperä⁸², Stephanie Loomis⁸², Susan Eaton⁸², George Okafo⁸³, Heli Salminen-Mankonen⁸³, Marc Jung⁸³, Nathan Lawless⁸³, Zhihao Ding⁸³, Joseph Maranville⁸⁴, Marla Hochfeld⁸⁴, Robert Plenge⁸³, Shameek Biswas⁸⁴, Masahiro Kanai⁸⁵, Mutaamba Maasha⁸⁵, Wei Zhou⁸⁵, Outi Tuovila⁸⁶, Raimo Pakkanen⁸⁶, Jari Laukkanen⁸⁷, Teijo Kuopio⁸⁷, Kristiina Aittomäki⁸⁸, Antti Mäkitie⁸⁹, Natalia Pujol⁹⁰, Triin Laisk⁹⁰, Katriina Aalto-Setälä⁹¹, Johanna Mäkelä⁹², Marco Hautalahti⁹², Sarah Smith⁹², Tom Southerington⁹², Eeva Kangasniemi⁹³, Henna Palin⁹³, Mika Kähönen⁹³, Sanna Siltanen⁹³, Tarja Laitinen⁹³, Felix Vaura⁹⁴, Jaana Suvisaari⁹⁴, Teemu Niiranen⁹⁴, Veikko Salomaa⁹⁴, Jukka Partanen⁹⁵, Mikko Arvas⁹⁵, Jarmo Ritari⁹⁶, Kati Hyvärinen⁹⁶, David Choy⁹⁷, Edmond Teng⁹⁷, Erich Strauss⁹⁷, Hao Chen⁹⁷, Hubert Chen⁹⁷, Jennifer Schutzman⁹⁷, Julie Hunkapiller⁹⁷, Mark McCarthy⁹⁷, Natalie Bowers⁹⁷, Rion Pendergrass⁹⁷, Tim Lu⁹⁷, Audrey Chu⁹⁸, Diptee Kulkarni⁹⁸, Fanli Xu⁹⁸, Joanna Betts⁹⁸, John Eicher⁹⁸, Jorge Esparza Gordillo⁹⁸, Laura Addis⁹⁸, Linda McCarthy⁹⁸, Rajashree Mishra⁹⁸, Janet Kumar⁹⁹, Margaret G. Ehm⁹⁹, Kirsi Auro¹⁰⁰, David Pulford¹⁰¹, Anne Pitkäranta¹⁰², Anu Loukola¹⁰², Eero Punkka¹⁰², Malla-Maria Linna¹⁰², Olli Carpén¹⁰², Taneli Raivio¹⁰², Joni A. Turunen^{103,104}, Tomi P. Mäkelä¹⁰⁵, Aino Salminen¹⁰⁶, Antti Aarnisalo¹⁰⁶, Daniel Gordin¹⁰⁶, David Rice¹⁰⁶, Erkki Isometsä¹⁰⁶, Eveliina Salminen¹⁰⁶, Heikki Joensuu¹⁰⁶, Ilkka Kalliala¹⁰⁶, Johanna Mattson¹⁰⁶, Juha Sinisalo¹⁰⁶, Jukka Koskela¹⁰⁶, Kari Eklund¹⁰⁶, Katariina Hannula-Jouppi¹⁰⁶, Lauri Aaltonen¹⁰⁶, Marja-Riitta Taskinen¹⁰⁶, Martti Färkkilä¹⁰⁶, Minna Raivio¹⁰⁶, Oskari Heikinheimo¹⁰⁶, Paula Kauppi¹⁰⁶, Pekka Nieminen¹⁰⁶, Pentti Tienari¹⁰⁶, Pirkko Pussinen¹⁰⁶, Sampsa Pikkarainen¹⁰⁶, Terhi Ollila¹⁰⁶, Tiinamaija Tuomi¹⁰⁶, Timo Hiltunen¹⁰⁶, Tuomo Meretoja¹⁰⁶, Tuula Salo¹⁰⁶, Ulla Palotie¹⁰⁶, Antti Palomäki¹⁰⁷, Jenni Aittokallio¹⁰⁷, Juha Rinne¹⁰⁷, Kaj Metsärinne¹⁰⁷, Klaus Elenius¹⁰⁷, Laura Pirilä¹⁰⁷, Leena Koulou¹⁰⁷, Markku Voutilainen¹⁰⁷, Riitta Lahesmaa¹⁰⁷, Roosa Kallionpää¹⁰⁷, Sirkku Peltonen¹⁰⁷, Tytti Willberg¹⁰⁷, Ulvi Gursoy¹⁰⁷, Varpu Jokimaa¹⁰⁷, Aarno Palotie¹⁰⁸, Anastasia Kytölä¹⁰⁸, Andrea Ganna¹⁰⁸, Anu Jalanko¹⁰⁸, Aoxing Liu¹⁰⁸, Arto Lehisto¹⁰⁸, Awaisa Ghazal¹⁰⁸, Elina Kilpeläinen¹⁰⁸, Elisabeth Widen¹⁰⁸, Elmo Saarentaus¹⁰⁸, Esa Pitkänen¹⁰⁸, Hanna Ollila¹⁰⁸, Hannele Laivuori¹⁰⁸, Henriette Heyne¹⁰⁸, Huei-Yi Shen¹⁰⁸, Jaakko Kaprio¹⁰⁸, Joel Rämö¹⁰⁸, Juha Karjalainen¹⁰⁸, Juha Mehtonen¹⁰⁸, Jyrki Pitkänen¹⁰⁸, Kalle Pärn¹⁰⁸, Kati Donner¹⁰⁸, Katja Kivinen¹⁰⁸, L. Elisa Lahtela¹⁰⁸, Mari E. Niemi¹⁰⁸, Mari Kaunisto¹⁰⁸, Mart Kals¹⁰⁸, Mary Pat Reeve¹⁰⁸, Mervi Aavikko¹⁰⁸, Nina Mars¹⁰⁸, Oluwaseun Alexander Dada¹⁰⁸, Pietro Della Briotta Parolo¹⁰⁸, Priit Palta¹⁰⁸, Rigbe Weldatsadik¹⁰⁸, Risto Kajanne¹⁰⁸, Rodos Rodosthenous¹⁰⁸, Samuli Ripatti¹⁰⁸, Sanni Ruotsalainen¹⁰⁸, Satu Strausz¹⁰⁸, Shabbeer Hassan¹⁰⁸, Shanmukha Sampath Padmanabhuni¹⁰⁸, Shuang Luo¹⁰⁸, Susanna Lemmelä¹⁰⁸, Taru Tukiainen¹⁰⁸, Timo P. Sipilä¹⁰⁸, Tuomo Kiiskinen¹⁰⁸, Vincent Llorens¹⁰⁸, Mark Daly^{108,109}, Jiwoo Lee^{85,108}, Kristin Tsuo^{85,108}, Mitja Kurki^{85,108}, Amanda Elliott^{85,108,110}, Aki Havulinna^{94,108}, Julia Partanen¹¹¹, Robert Yang¹¹², Dermot Reilly¹¹³, Alessandro Porello¹¹⁴, Amy Hart¹¹⁴, Dawn Waterworth¹¹⁴, Ekaterina Khramtsova¹¹⁴, Karen He¹¹⁴, Meijian Guan¹¹⁴, Qingqin S. Li¹¹⁵, Sauli Vuoti¹¹⁶, Eric Green¹¹⁷, Robert Graham¹¹⁷, Sahar Mozaffari¹¹⁷, Adriana Huertas-Vazquez¹¹⁸, Andrey Loboda¹¹⁸, Caroline Fox¹¹⁸, Fabiana Farias¹¹⁸, Jae-Hoon Sul¹¹⁸, Jason Miller¹¹⁸, Neha Raghavan¹¹⁸, Simonne Longrich¹¹⁸, Johannes Kettunen¹¹⁹, Raisa Serpi¹¹⁹, Reetta Hinttala¹¹⁹, Tuomo Mantere¹¹⁹, Anne Remes¹²⁰, Elisa Rahikkala¹²⁰, Johanna Huhtakangas¹²⁰, Kaisa Tasanen¹²⁰, Laura Huilaja¹²⁰, Laure Morin-Papunen¹²⁰, Maarit Niinimäki¹²⁰, Marja Väärasmäki¹²⁰, Outi Uimari¹²⁰, Peeter Karihtala¹²⁰, Terhi Pilttonen¹²⁰, Terttu Harju¹²⁰, Timo Blomster¹²⁰, Vuokko Anttonen¹²⁰, Hilikka Soinen¹²¹, Kai Kaarniranta¹²¹, Liisa Suominen¹²¹, Margit Pelkonen¹²¹, Maria Siponen¹²¹, Mikko Kiviniemi¹²¹, Oili Kaipainen-Seppänen¹²¹, Päivi Auvinen¹²¹, Päivi Mäntylä¹²¹, Reetta Kälviäinen¹²¹, Valterti Julkunen¹²¹, Chris O'Donnell¹²², Ma'en Obeidat¹²², Nicole Renaud¹²², Debby Ngo¹²³, Majd Mouded¹²³, Mike Mendelson¹²⁴, Anders Mälarstig¹²⁵, Heli Lehtonen¹²⁵, Jaakko Parkkinen¹²⁵, Kirsi Kalpala¹²⁵, Melissa Miller¹²⁵, Nan Bing¹²⁵, Stefan McDonough¹²⁵, Xinli Hu¹²⁵, Ying Wu¹²⁵, Airi Jussila¹²⁶, Annika Auranen¹²⁶, Argyro Bizaki-Vallaskangas¹²⁶, Hannu Uusitalo¹²⁶, Jukka Peltola¹²⁶, Jussi Hernesniemi¹²⁶, Katri Kaukinen¹²⁶, Laura Kotaniemi-Talonen¹²⁶, Pia Isomäki¹²⁶, Teea Salmi¹²⁶, Venla Kurra¹²⁶, Kirsi Sipilä^{127,128}, Auli Toivola⁹⁴, Elina Järvensivu⁹⁴, Essi Kaiharju⁹⁴, Hannele Mattsson⁹⁴, Kati Kristiansson⁹⁴, Lotta Männikkö⁹⁴,

Markku Laukkanen⁹⁴, Markus Perola⁹⁴, Minna Brunfeldt⁹⁴, Päivi Laiho⁹⁴, Regis Wong⁹⁴, Sami Koskelainen⁹⁴, Sini Lähteenmäki⁹⁴, Sirpa Soini⁹⁴, Teemu Paajanen⁹⁴, Terhi Kilpi⁹⁴, Tero Hiekkalinna⁹⁴, Tuuli Sistonen⁹⁴, Clément Chatelain¹²⁹, Deepak Raipal¹²⁹, Katherine Klinger¹²⁹, Samuel Lessard¹²⁹, Fredrik Åberg¹³⁰, Mikko Hiltunen¹³¹, Sami Heikkinen¹³¹, Hannu Kankaanranta^{132,133,134}, Tuula Palotie¹³⁵, Iiris Hovatta¹³⁶, Kimmo Palin¹³⁶, Niko Välimäki¹³⁶, Sanna Toppila-Salmi¹³⁶, Eija Laakkonen¹³⁷, Eeva Sliz¹³⁸, Heidi Silven¹³⁸, Katri Pylkäs¹³⁸, Minna Karjalainen¹³⁸, Riikka Arffman¹³⁸, Susanna Savukoski¹³⁸, Jaakko Tyrmi^{138,139}, Manuel Rivas¹⁴⁰, Harri Siirtola¹³⁹, Iida Vähätalo¹³⁹, Javier Garcia-Tabuenca¹³⁹, Marianna Niemi¹³⁹, Mika Helminen¹³⁹ & Tiina Luukkaala¹³⁹

⁷⁷Aarhus University, Aarhus, Denmark. ⁷⁸Abbvie, Chicago, IL, US. ⁷⁹Astra Zeneca, Cambridge, UK. ⁸⁰Auria Biobank/University of Turku/Hospital District of Southwest Finland, Turku, Finland. ⁸¹Biobank of Eastern Finland / University of Eastern Finland / Northern Savo Hospital District, Kuopio, Finland. ⁸²Biogen, Cambridge, MA, USA. ⁸³Boehringer Ingelheim, Ingelheim am Rhein, Germany. ⁸⁴Bristol Myers Squibb, New York, NY, USA. ⁸⁵Broad Institute, Cambridge, MA, USA. ⁸⁶Business Finland, Helsinki, Finland. ⁸⁷Central Finland Biobank/University of Jyväskylä/Central Finland Health Care District, Jyväskylä, Finland. ⁸⁸Department of Medical Genetics, Helsinki University Central Hospital, Helsinki, Finland. ⁸⁹Department of Otorhinolaryngology—Head and Neck Surgery, University of Helsinki and Helsinki University Hospital, Helsinki, Finland. ⁹⁰Estonian biobank, Tartu, Estonia. ⁹¹Faculty of Medicine and Health Technology, Tampere University, Tampere, Finland. ⁹²FINBB - Finnish biobank cooperative, Tampere, Finland. ⁹³Finnish Clinical Biobank Tampere/University of Tampere/Pirkanmaa Hospital District, Tampere, Finland. ⁹⁴Finnish Institute for Health and Welfare (THL), Helsinki, Finland. ⁹⁵Finnish Red Cross Blood Service/Finnish Hematology Registry and Clinical Biobank, Helsinki, Finland. ⁹⁶Finnish Red Cross Blood Service, Helsinki, Finland. ⁹⁷Genentech, San Francisco, CA, USA. ⁹⁸GlaxoSmithKline, Brentford, UK. ⁹⁹GlaxoSmithKline, Collegeville, PA, USA. ¹⁰⁰GlaxoSmithKline, Espoo, Finland. ¹⁰¹GlaxoSmithKline, Stevenage, UK. ¹⁰²Helsinki Biobank/Helsinki University and Hospital District of Helsinki and Uusimaa, Helsinki, Finland. ¹⁰³Helsinki University Hospital and University of Helsinki, Helsinki, Finland. ¹⁰⁴Eye Genetics Group, Folkhälsan Research Center, Helsinki, Finland. ¹⁰⁵HiLIFE, University of Helsinki, Finland, Finland. ¹⁰⁶Hospital District of Helsinki and Uusimaa, Helsinki, Finland. ¹⁰⁷Hospital District of Southwest Finland, Turku, Finland. ¹⁰⁸Institute for Molecular Medicine Finland (FIMM), HiLIFE, University of Helsinki, Helsinki, Finland. ¹⁰⁹Broad Institute of MIT and Harvard; Massachusetts General Hospital, Boston, MA, USA. ¹¹⁰Massachusetts General Hospital, Boston, MA, USA. ¹¹¹Institute for Molecular Medicine Finland, HiLIFE, University of Helsinki, Helsinki, Finland. ¹¹²Janssen Biotech, Beerse, Belgium. ¹¹³Janssen Research & Development, LLC, Boston, MA, USA. ¹¹⁴Janssen Research & Development, LLC, Spring House, PA, USA. ¹¹⁵Janssen Research & Development, LLC, Titusville, NJ 08560, USA. ¹¹⁶Janssen-Cilag Oy, Espoo, Finland. ¹¹⁷Maze Therapeutics, San Francisco, CA, USA. ¹¹⁸Merck, Kenilworth, NJ, USA. ¹¹⁹Northern Finland Biobank Borealis / University of Oulu / Northern Ostrobothnia Hospital District, Oulu, Finland. ¹²⁰Northern Ostrobothnia Hospital District, Oulu, Finland. ¹²¹Northern Savo Hospital District, Kuopio, Finland. ¹²²Novartis Institutes for BioMedical Research, Cambridge, MA, USA. ¹²³Novartis, Basel, Switzerland. ¹²⁴Novartis, Boston, MA, USA. ¹²⁵Pfizer, New York, NY, USA. ¹²⁶Pirkanmaa Hospital District, Tampere, Finland. ¹²⁷Research Unit of Oral Health Sciences Faculty of Medicine, University of Oulu, Oulu, Finland. ¹²⁸Medical Research Center, Oulu, Oulu University Hospital and University of Oulu, Oulu, Finland. ¹²⁹Translational Sciences, Sanofi R&D, Framingham, MA, USA. ¹³⁰Transplantation and Liver Surgery Clinic, Helsinki University Hospital, Helsinki University, Helsinki, Finland. ¹³¹University of Eastern Finland, Kuopio, Finland. ¹³²University of Gothenburg, Gothenburg, Sweden. ¹³³Seinäjäki Central Hospital, Seinäjoki, Finland. ¹³⁴Tampere University, Tampere, Finland. ¹³⁵University of Helsinki and Hospital District of Helsinki and Uusimaa, Helsinki, Finland. ¹³⁶University of Helsinki, Helsinki, Finland. ¹³⁷University of Jyväskylä, Jyväskylä, Finland. ¹³⁸University of Oulu, Oulu, Finland. ¹³⁹University of Tampere, Tampere, Finland. ¹⁴⁰University of Stanford, Stanford, CA, USA. A full list of members and their affiliations appears in the Supplementary Information.