



HAL
open science

The apical 70-pS K⁺ channel in the thick ascending limb of Henle's loop is a large-conductance Na⁺- and Cl⁻-activated, K⁺ Na⁺ 1.1-like, channel

Elise de Combiens, Nadia Frachon, Lydie Cheval, Stéphane Lourdel, Marc Paulais

► To cite this version:

Elise de Combiens, Nadia Frachon, Lydie Cheval, Stéphane Lourdel, Marc Paulais. The apical 70-pS K⁺ channel in the thick ascending limb of Henle's loop is a large-conductance Na⁺- and Cl⁻-activated, K⁺ Na⁺ 1.1-like, channel. *American Journal of Physiology. Renal Physiology*, 2024, 326 (3), pp.F485-F496. 10.1152/ajprenal.00311.2023 . hal-04556454

HAL Id: hal-04556454

<https://hal.sorbonne-universite.fr/hal-04556454>

Submitted on 23 Apr 2024

HAL is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers.

L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.

**The apical 70-pS K channel in the thick ascending limb of Henle's loop is a large-conductance Na⁺-
and Cl⁻-activated, K_{Na}1.1-like, channel**

1

2 Elise de Combiens^{a,b}, Nadia Frachon^{a,b}, Lydie Cheval^{a,b}, Stéphane Lourdel^{a,b} and Marc Paulais^{a,b}

3

4 ^aCentre de Recherche des Cordeliers, INSERM, Sorbonne Université, Université de
5 Paris, ERL8228, F-75006, Paris, France.

6 ^bCNRS-ERL8228, F-75006 Paris, France.

7

8

9

10

11 **Corresponding author**

12 Marc Paulais

13 UMR_S 1138 – ERL 8228, Centre de Recherche des Cordeliers

14 15 rue de l'école de médecine

15 75270 Paris Cedex 06, France

16 E-mail: marc.paulais@inserm.fr

17 Phone: +33688683274

18 Fax: +33144275119

19 ORCID: 0000-0002-4895-2110

20

21 **Running head:** an apical K_{Na}1.1 channel in the TAL

Abstract

Apical potassium channels are crucial for thick ascending limb (TAL) of Henle's loop transport function. The ROMK (*KCNJ1*) gene encodes a 30-pS K channel whose loss of function causes the reduced NaCl reabsorption in the TAL associated with Type 2 Bartter's syndrome. In contrast, the molecular basis of a functionally ROMK-related 70-pS K channel is still unclear. The aim of this study was to highlight new specific channel properties that may give insights on its molecular identity. Using the patch-clamp technique on the apical membrane of mouse split-open TAL tubules, we observed that 70-pS K channel activity, but not ROMK channel activity, increases with the internal Na⁺ and Cl⁻ concentrations, with relative 50 % effective concentrations (EC_{50}) and Hill coefficients (n_H) of 40.6 mM (SD 1.65) and 2.4 (SD 0.28) for Na⁺, and of 29.3 mM (SD 2.35) and 2.2 (SD 0.39) for Cl⁻. Conversely, 70-pS K channel activity was inhibited by internal K⁺ with a relative EC_{50} of 64 mM (SD 13.5) and a n_H of 3.5 (SD 2.3), and by internal NH₄⁺ and Ca²⁺. The reevaluation of channel conductive properties revealed an actual inward conductance of ~ 170 pS, with multiple subconductance levels and an inward rectification, and a substantial permeability to NH₄⁺ ($P_{NH_4}/P_K = 0.2$). We conclude that the apical 70-pS K channel in TAL cells is a large-conductance Na⁺- and Cl⁻-activated potassium channel functionally resembling a K_{Na}1.1 channel and propose that ROMK determines its functional expression possibly at the level of channel protein synthesis or trafficking.

New & Noteworthy: this study highlights new regulatory and conductive properties of the as yet poorly molecularly defined 70-pS K channel in the apical membrane of thick ascending limb of Henle's loop cells of the mouse kidney that functionally identify it as a large-conductance, Na⁺- and Cl⁻-activated K_{Na}1.1-like channel.

Key words: mouse kidney; thick ascending limb; 70-pS K channel; K_{Na}1.1 channel; ROMK channel

47

Introduction

48 The thick ascending limb (TAL) of Henle's loop of the mammalian kidney reabsorbs
49 approximately 30% of the filtered NaCl load in excess of water, hence plays a pivotal role in the
50 maintenance of the extracellular fluid volume and blood pressure, and participates to ammonia
51 cycle, acid-base metabolism and divalent cations homeostasis (1). The TAL transport function
52 involves an active transcellular process by which Na⁺ and Cl⁻ enter the cell by the apical Na⁺-K⁺-2Cl⁻
53 cotransporter NKCC2 and leave the cell by the basolateral Na⁺/K⁺ ATPase and ClCK-Barttin chloride
54 channels, and a passive paracellular reabsorption of Na⁺, Ca²⁺ and Mg²⁺ driven by a lumen-positive
55 transepithelial potential difference. It is now well established that a dominant apical membrane
56 potassium conductance is the major pathway for the exit of K⁺ into the tubular lumen necessary to
57 maintaining steady-state intracellular K⁺ concentration and to the generation of the transepithelial
58 potential difference (1).

59 It is in general agreement that of the three types of potassium channels, a low-conductance
60 (30 pS), an intermediate-conductance (70 pS) and a large-conductance (100-200 pS) Ca²⁺-activated
61 Maxi-K channels, functionally identified in the apical membrane of TAL cells (2) (3) (4) (5) (6), the 30-
62 pS and 70-pS K channels are the main contributors to the apical membrane potassium conductance
63 in native cells. During the past years, a great deal of effort has been put into their molecular
64 identification, a prerequisite for understanding their role in the physiology and pathophysiology of
65 renal salt reabsorption. There is now compelling evidence that the 30-pS K channel is formed by the
66 homomeric assembly of inwardly rectifying channel ROMK (Kir1.1; *KCNJ1*) subunits and is critical for
67 TAL function (7): the native 30-pS K channel shares many biophysical and pharmacological properties
68 with the recombinant ROMK channel, is functionally detected in ROMK-expressing cells, and is
69 absent in mice lacking *Kcnj1* and exhibiting the strongly reduced reabsorption of NaCl in the TAL
70 associated with the Type 2 Bartter's syndrome caused by loss-of-function mutations in *KCNJ1*. In
71 contrast, the molecular basis of the 70-pS K channel, a major contributor to the apical membrane
72 potassium conductance in the rat (8), is still not clear. On the one hand, that the ROMK and 70-pS K

73 channels differ by their unitary conductances, kinetics and sensitivities to inhibitors early suggested
74 two distinct molecular entities. On the other hand, studies in which the complete absence of 70-pS K
75 channel activity in *Kcnj1*-null mice clearly established the requirement of *Kcnj1* expression for its
76 functional expression (9) (10). From that emerged the paradigm of ROMK subunits forming part of
77 70-pS K channel structure, a whole apical membrane potassium conductance underlied by two
78 channels types involving the ROMK subunit thereby providing a satisfactory explanation for the
79 impaired NaCl reabsorption in the TAL of *Kcnj1*-null mice. It was then legitimately postulated that the
80 properties shared by the ROMK and 70-pS K channels are conferred by the ROMK subunit, those
81 specific to the 70-pS K channel being accounted by the contribution of a yet-to-be-identified
82 associated subunit. In either case, however, none of the currently known functional properties of the
83 70-pS K channel, largely established in the rat, is sufficiently distinctive to correlate with either one of
84 the potassium channels subunits molecularly expressed in TAL cells (11).

85 The aim of this study was therefore to identify new biophysical and regulatory properties of
86 the 70-pS K channel that may bring clues on its molecular identity. To this end, we considered the
87 briefly mentioned absence of channel opening in the rat TAL unless its inner face was exposed to
88 millimolar concentrations of Na⁺ (3), a feature not shared by the ROMK channel in the rabbit TAL (5).
89 We hypothesized that such a requirement of internal Na⁺ for 70-pS K channel activity might have
90 reflected a sensitivity to intracellular Na⁺ reminiscent that exhibited by few potassium channels,
91 namely the small-conductance G protein-coupled inwardly-rectifying (GIRK) Kir3.2 (GIRK2; *KCNJ6*)
92 and Kir3.4 (GIRK4; *KNCJ5*) channels, and the large-conductance Na⁺- and Cl⁻-activated K_{Na}1 channels
93 K_{Na}1.1 (previously listed as Slack or Slo2.2; *KCNT1*) and K_{Na}1.2 (previously listed as Slick or Slo2.1;
94 *KCNT2*) (12) (13). However, only *KCNT1* is expressed in TAL cells where it very likely encodes for a
95 potassium channel functionally resembling a K_{Na}1.1 channel in the basolateral membrane (14) (15)
96 (16) (17) (18) (19). Here, we examine whether the 70-pS K channel in the apical membrane of mouse
97 TAL cells also exhibits the ions sensitivity and the conductive properties of a K_{Na}1.1 channel with the
98 use of the patch-clamp technique on split-open tubules.

100 Animals and tissue preparation

101 Animals were handled in full compliance with the French government welfare policy and with
102 the recommendations for the care and use of animals for scientific purposes put forward by the
103 European Council Directive 2010/63/EU. The experimental procedures were approved by the Charles
104 Darwin Ethics Committee for animal experimentation (Permit 8569). Adult C57BL/6J male mice
105 (Envigo, Gannat, France) were housed in the Centre d'Explorations Fonctionnelles (UMRS1138, Paris,
106 France; Permit A1-0872) on a 12:12 light/dark cycle (Zeitgeber time, ZT) and under constant
107 temperature and humidity, and fed with pellets of a complete 0.25 % Na⁺ and 0.6 % K⁺ SAFE-A04
108 maintenance diet (Augy, France) with free access to tap water.

109 At the day of experiment, mice were euthanized at ZT 4 (11:00) by cervical dislocation. TAL
110 segments were isolated after an enzymatic treatment procedure derived from that routinely used in
111 the laboratory for the study of renal basolateral ionic channels. Briefly, the left kidney was perfused
112 with Leibovitz L-15 medium containing 150 U/ml Worthington CLS-2 collagenase (Coger SAS, Paris,
113 France), removed and cut into thin slices. Small pieces of cortex were then incubated in the same
114 medium for 25 min at 37 °C. Cortical TAL segments, identified as the tubular part upstream
115 glomerulus attachment or transition into the wider early distal convoluted tubule, were
116 microdissected in ice-cold, collagenase-free, L-15 medium under stereomicroscope, then transferred
117 into a perfusion chamber on the stage of an Axiovert™ 100 microscope (Carl Zeiss S.A.S., Rueil-
118 Malmaison, France). In order to gain access to the apical membrane, tubules were split-open with
119 the broken tip of a patch-clamp pipette filled with bath medium (see below for composition) and
120 operated by a 4MRE motorized micro-manipulator and a SM8 remote control unit (Luigs and
121 Neuman, Ratingen, Germany). Eventual debris at the luminal cell surface were removed by applying a
122 gentle suction to the pipette.

123 Single-channel recording and analysis

124 *Solutions.* Patch-clamp pipettes were filled with a solution containing (in mM) 145 KCl, 1
125 MgCl₂, 10 D-Glucose and 10 HEPES, and adjusted to pH 7.4 with KOH. Tubules were bathed in a
126 medium containing (in mM) 140 NaCl, 5 KCl, 1 CaCl₂, 1 MgCl₂, 10 D-Glucose and 10 HEPES, and
127 adjusted to pH 7.4 with NaOH. A high-Na⁺ medium typically bathed the inner face of cell-excised
128 inside-out membrane patches and was modified in order to obtain solutions with various internal
129 Na⁺, Cl⁻, K⁺, NH₄⁺ or Ca²⁺ concentrations (see Supplemental Table S1 for composition). When
130 appropriate, the osmolarity of solutions was measured by an Autocal 13DR freezing point
131 osmometer (Hermann Roebling Messtechnik, Berlin, Germany). The exchange of internal solution
132 was performed by gravity and controlled either manually or by a RSC-200 multi-channel rapid
133 solution changer (BioLogic, Seyssinet-Pariset, France) driven by the Axon Instruments® pClamp™ 10
134 Clampex software (Molecular Devices S.A.S., Villepinte, France).

135 *Current Recordings.* Currents were amplified by a L/M-EPC 7 (List Electronics, Darmstadt,
136 Germany) patch-clamp amplifier, filtered at 1 kHz by a LPBF-48DG 8-pole Bessel filter (NPI Electronics
137 GmbH, Tamm, Germany) and digitized at 2 kHz by an Axon Instruments® Digidata® 1440A A/D
138 converter (Molecular Devices) for online monitoring and recording by the Clampex software. The
139 bath reference was 0.5 M KCl in a 4% agar bridge connected to an Ag/AgCl pellet. When appropriate,
140 clamp voltage V_c (i.e. = $-V_{pipette}$) values were corrected for liquid junction potential difference as
141 measured by the Clampex software routine. All experiments were conducted at room temperature
142 (20-25 °C).

143 *Data Analysis.* Single-channel recordings were analyzed offline using the Clampex software.
144 Patch activity was quantified according to the equation $NP_o = I/i$, where I is the mean current passing
145 through N channels and i is the unitary current amplitude. Because of patch-to-patch variability in
146 channels activity, the NP_o in test condition was normalized to the NP_o in control condition measured
147 on the same membrane patch.

148 Ions sensitivity parameters were determined by the nonlinear curve fitting of a four-
149 parameter Hill equation to the normalized NP_o data using

$$NP_o = A + \{(B - A)/[1 + C^{n_H}]\} \quad (1)$$

150 where A and B are the minimum and maximum normalized NP_o , respectively, and n_H is the Hill
 151 coefficient. C is the $K/[X]$ or $[X]/K$ ratio for channel activation or inhibition, respectively, $[X]$ being
 152 the concentration of the tested ion X and K the $[X]$ value giving a response halfway between A and
 153 B (relative EC_{50}).

154 The channel inward slope conductance in cell-excised inside-out membrane patches exposed
 155 to asymmetrical transmembrane K^+ concentrations was determined by first fitting the Goldman-
 156 Hodgkin-Katz (GHK) constant field current equation for K^+ and Na^+ to i/V_c relationships data over the
 157 -100 to +40 mV V_c range using

$$i = \frac{V_c F^2}{RT} \left[P_K \left(\frac{a_o^K - a_i^K e^{\left(\frac{V_c F}{RT}\right)}}{1 - e^{\left(\frac{V_c F}{RT}\right)}} \right) - P_{Na} \left(\frac{a_i^{Na} e^{\left(\frac{V_c F}{RT}\right)}}{1 - e^{\left(\frac{V_c F}{RT}\right)}} \right) \right] \quad (2)$$

158 where F , R and T have their conventional thermodynamic meanings, P_X is the permeability to the ion
 159 X , a_o^X and a_i^X are the respective external and internal ionic activities of X , and taking activity
 160 coefficients of K^+ and Na^+ as 0.755 and 0.775, respectively (20), then by fitting the Boltzmann
 161 equation the first derivative of GHK data using

$$g = g_{-\infty}^{in} + \frac{g_{+\infty}^{out} - g_{-\infty}^{in}}{1 + e^{(V_c - V_c^0)/k}} \quad (3)$$

162 where g is the conductance at a given V_c , $g_{-\infty}^{in}$ is the inward conductance at infinite negative V_c , $g_{+\infty}^{out}$
 163 is the outward conductance at infinite positive V_c , V_c^0 is the V_c value at $g_{-\infty}^{in} - g_{+\infty}^{out}$ and k is the
 164 logarithmic sensitivity for an e-fold increase in V_c .

165 The NH_4^+ to K^+ permeability ratio (P_{NH_4}/P_K) was determined by fitting the GHK equation to
 166 i/V_c data over the -80 to +30 mV V_c range obtained in inside-out membrane patches under opposite
 167 K^+ and NH_4^+ gradients, and in the presence of internal Na^+ , using

$$i = \frac{V_c F^2}{RT} \left[\left(\frac{a_o^K - a_i^K e^{\left(\frac{V_c F}{RT}\right)}}{1 - e^{\left(\frac{V_c F}{RT}\right)}} \right) - \frac{P_{Na}}{P_K} \left(\frac{a_i^{Na} e^{\left(\frac{V_c F}{RT}\right)}}{1 - e^{\left(\frac{V_c F}{RT}\right)}} \right) - \frac{P_{NH_4}}{P_K} \left(\frac{a_i^{NH_4} e^{\left(\frac{V_c F}{RT}\right)}}{1 - e^{\left(\frac{V_c F}{RT}\right)}} \right) \right] \quad (4)$$

168 taking a NH_4^+ activity coefficient of 0.77 (21).

169 The OriginLab Corporation Origin® software (Ritme Informatique, Paris, France) was used to
170 perform linear and nonlinear regression analyses, to generate amplitude histograms and, when
171 appropriate, to fit a multiple peak Gaussian function to histogram data.

172 **Data presentation and statistics**

173 Comparisons of parameters between groups were performed with the Student *t* test for two
174 independent variables. Comparisons of the parameters of two related values were performed by
175 using the Student *t* test for paired samples. Data are reported as individual values or as mean (SD) for
176 the indicated number of experiments. *P* values < 0.05 were taken to represent statistically significant
177 differences.

178

Results

179 With a detectable activity in only 16 % of the cell-attached membrane patches (Table 1), the
180 incidence of potassium channels in the apical membrane of mouse TAL tubules was low but actually
181 comparable to the ~ 15 % previously reported by others in mice and rats (2) (3). The ROMK and 70-pS
182 K channels were the only potassium channels detected as distinguished by the well established high,
183 rapidly flickering and voltage-independent ROMK channel activity and the lower, voltage-dependent
184 and "box-like" bursting 70-pS K channel activity (Figure 1a), together with their respective unitary
185 conductance at resting membrane voltage (Figure 1b and g_o in Table 1). In the rat TAL, the 70-pS K
186 channel may account for as much as 80 % of the whole apical membrane potassium conductance (8).
187 Here, the 70-pS K channel was the most frequently observed in situ, and the ROMK and 70-pS K
188 channels exhibited similar apparent numbers of active channels (N_{app}) and activities (NP_o) (Table 1).
189 Thus, according to the mean conductance per membrane patch one may calculate for each channel
190 type (g_{patch} in Table 1), the 70-pS K channel may mediate most of the apical membrane potassium
191 conductance in the mouse TAL.

192 Cell-attached membrane patches were systematically excised into the high- Na^+ medium. In
193 this condition, the incidence of the ROMK and 70-pS K channels remained low (~ 21 % of the
194 membrane patches), with respective frequencies, N_{app} and NP_o values as in the cell-attached
195 configuration (Supplemental Table S2). Because openings of the 70-pS K channel in the rat TAL were
196 observed only in the presence of internal Na^+ (3), an absolute prerequisite for the activity of the
197 cloned $K_{Na}1.1$ channel (22), we first determined whether internal Na^+ is also required for 70-pS K
198 channel activity in the mouse TAL. Figure 2 shows a continuous recording illustrating the effect of the
199 complete removal of internal Na^+ on 70-pS K channel activity. In this recording, the membrane patch
200 was initially exposed to 140 mM Na^+ and current levels corresponding to the simultaneous opening
201 of up to six channels were observed. Then, channels activity rapidly disappeared when all internal
202 Na^+ was replaced by Li^+ , a weak substitute for Na^+ in activating the native $K_{Na}1$ and cloned
203 $K_{Na}1.1$ channels (23) (24), and was readily restored by the return back to the 140 mM Na^+ -containing

204 medium. On the average of seven results obtained according to this sequence, the complete removal
205 of internal Na^+ reduced the normalized NP_o by 97 % (SD 3.5) ($P = 0.096$ vs 100 % inhibition). From the
206 five patches where very rare and brief openings could be observed in the absence of Na^+ , the single-
207 channel current amplitude was not significantly different from that in the presence of 140 mM Na^+ (P
208 = 0.45).

209 The dependence of 70-pS K channel activity on internal Na^+ concentration is summarized in
210 Figure 3. The dose-response curve established in four separate membrane patches exposed to
211 various Na^+ concentrations shows that channel activity increased with internal Na^+ in a sigmoidal
212 fashion with a relative EC_{50} of 40.6 mM Na^+ (SD 1.65) and a n_H of 2.4 (SD 0.28).

213 Since the Na^+ -dependent $K_{\text{Na}1.1}$ channel is also activated by internal Cl^- (22) (24), the effect of
214 altering the internal Cl^- concentration on the activity of the 70-pS K channel was then investigated.
215 Figure 4 shows that its activity in a membrane patch bathed throughout with 140 mM Na^+ was
216 strongly reduced when the internal Cl^- concentration was lowered from 145 mM down to 5 mM but
217 rapidly recovered upon the reintroduction of 145 mM Cl^- . On the average of nine separate
218 measurements, the lowering of internal Cl^- concentration reduced the normalized NP_o by 79 % (SD
219 10.3) ($P = 0.0002$ vs 100 % inhibition), and did not affect the single-channel current amplitude ($P =$
220 0.4). Channel activity increased with the internal Cl^- concentration, cumulated data from five
221 membrane patches yielding a relative EC_{50} of 29.3 mM Cl^- (SD 2.35), a n_H of 2.2 (SD 0.39) and a
222 predicted normalized NP_o of 0.14 (SD 0.037) in the complete absence of Cl^- (Figure 5).

223 Figure 6 shows that, in contrast to the 70-pS K channel, the ROMK channel in the mouse TAL
224 was not sensitive to a change in internal Na^+ or Cl^- concentration. Indeed, the substitution of Li^+ for
225 all Na^+ in the bathing medium reduced ROMK channel normalized NP_o by only 0.28 % (SD 4.9) ($n = 4$;
226 $P = 0.91$ vs no inhibition). Because less than 3% of the cell-excised membrane patches exhibited
227 ROMK channel activity (see Supplemental Table S2), only two of them could be tested for an
228 alteration in internal Cl^- concentration. In these membrane patches, however, the lowering of
229 internal Cl^- concentration from 145 mM down to 5 mM did not affect ROMK normalized NP_o .

230 Unlike the apical Maxi-K channel in cultured TAL cells whose activity is increased by a rise in
231 cytosolic Ca^{2+} (6), the 70-pS K channel in the rat TAL is dramatically inhibited by millimolar internal
232 Ca^{2+} concentrations (3), an unusual feature shared with the cloned $\text{K}_{\text{Na}}1.1$ channel (25). Supplemental
233 Figure S1 shows that the switch from a Ca^{2+} -free to a 1 mM Ca^{2+} -containing internal solution also
234 reduced 70-pS K channel activity in the mouse TAL by 91 % (SD 7.5) ($P = 0.1$ vs 100% inhibition).

235 An elevation in internal K^+ concentration inhibits the Na^+ -induced activity of the native $\text{K}_{\text{Na}}1$
236 channels in guinea-pig ventricular myocytes and of the $\text{K}_{\text{Na}}1.1$ -like channel in the basolateral
237 membrane of TAL cells (Supplemental Figure S2) (26). Figure 7 shows that the 70-pS K channel is also
238 negatively regulated by internal K^+ . Here, the activity of at least five channels in a membrane patch
239 clamped at $V_c = -60$ mV and bathed throughout with 140 mM Na^+ and 145 mM Cl^- rapidly decreased
240 when the internal K^+ concentration was increased from 9 mM to 145 mM and fully recovered upon
241 the return back to the 9 mM K^+ -containing medium. On the average of eight measurements, the high
242 internal K^+ concentration decreased the normalized NP_o by 89 % (SD 10.3) ($P = 0.02$ vs 100%
243 inhibition). Of note, this maneuver also reduced the single-channel current amplitude ($P = 0.003$), but
244 most likely by the slight effect on current size of the left-shifted equilibrium potential for K^+ in this
245 condition (see Figure 10 for clarity). Since this experimental series was performed by adding 136 mM
246 K-gluconate to the high- Na^+ medium (see Supplemental Table S1), the effect of raising internal K^+
247 concentration on channel activity could have been related to an increased ionic strength or
248 osmolarity. However, Supplemental Figure S3 shows that the exposure to a high internal K^+
249 concentration under constant ionic strength and osmolarity still markedly inhibited channel activity
250 ($n = 3$). As summarized in Figure 8, the combined results from eleven separate membrane patches
251 showed that internal K^+ concentration-dependently decreased 70-pS K channel activity with a relative
252 EC_{50} of 64 mM K^+ (SD 13.5) and a n_H of 3.5 (SD 2.3).

253 Given the inhibition of the 70-pS K channel by internal K^+ , the effect of NH_4^+ , a good
254 surrogate for K^+ through ionic channels and transporters, was assessed in the presence of 140 mM
255 Na^+ . Figure 9 shows that the channel was also negatively regulated by internal NH_4^+ , the normalized

256 NP_o being reduced by 25 % (SD 12.8) ($P = 0.03$ vs no inhibition) when 10 mM NH_4^+ were added to the
257 bathing medium ($n = 4$), and by 98 % (SD 3.8) ($P = 0.35$ vs 100 % inhibition) with 145 mM NH_4^+ ($n = 4$).

258 The 70-pS K channel conductance of ~ 85 pS we determined in cell-attached membrane
259 patches at $V_c = 0$ mV (g_o in Table 1) was in reasonable agreement with the 60-75 pS range of
260 conductance reported by others at this voltage and hitherto defining the channel as an intermediate-
261 conductance channel (2) (3) (9), but did not conform to the large inward conductance of ~ 180 pS
262 characterizing the cloned $K_{Na}1.1$ channel (22). However, the steepness of the i/V_c relationship at
263 negative membrane potential differences shown in Figure 1b, as well as of previously published ones
264 (2) (3) (9), strongly suggested a 70-pS K channel of larger inward conductance. Therefore, its
265 conductive properties were reinvestigated in cell-excised inside-out membrane patches under
266 precisely controlled ionic conditions (Figure 10). In the presence of 145 mM K^+ in the pipette, and of
267 9 mM K^+ and 140 mM Na^+ in the bath, the GHK equation (equation 2) fitted well the obtained i/V_c
268 data with a P_K of $0.41 \cdot 10^{-12} \text{ cm}^3 \text{ s}^{-1}$ (SD $0.012 \cdot 10^{-12}$) ($n = 5$) close to the values obtained for the channel
269 in the rat TAL and for native $K_{Na}1$ channels (3) (26). The fit of the Boltzmann equation (equation 3) to
270 the first derivative of GHK data then yielded an inward conductance at an infinite negative clamp
271 potential difference of 172 pS (SD 5.2) ($n = 5$) (Figure 10 inset), indicating a 70-pS K channel of actual
272 large conductance.

273 By taking advantage of the remaining activity when the channel was exposed to 145 mM
274 internal K^+ or NH_4^+ in the conditions given in Figures 7 and 9, we were able to establish the first i/V_c
275 curves under a high internal K^+ or NH_4^+ concentration (Figure 10). Under symmetrical 145 mM K^+
276 condition, the currents through the channel reversed at -0.1 mV (SD 1.2) and the i/V_c relationship for
277 inward currents was linear with a slope (g_{in}) of 173 pS (SD 4.5) ($n = 4$) that further established the
278 large channel inward conductance. In addition, the outward currents were smaller than expected
279 from Ohm's law with an outward chord conductance ($g_{chord-out}$), as defined by the slope of the curve
280 between the reversal potential and $V_c + 40$ mV, of 101 pS (SD 3.4) ($n = 4$) that yielded a rectification
281 coefficient ($g_{in} / g_{chord-out}$) of 1.7. With 145 mM NH_4^+ added to the high- Na^+ medium, the channel

282 inward conductance was 164 pS (SD 8.1) (n = 4) and the fit of the GHK equation (equation 4) to the
283 mean inward current data yielded a P_{NH_4}/P_K of around 0.2.

284 Lastly, the 70-pS K channel exhibited the multiple conductance substates described for the
285 $K_{Na1.1}$ channel (22). An example is given in Supplemental Figure S4 where partial channel openings
286 and closures during a burst of activity are visible. The corresponding all-points current amplitude
287 histogram showed that the data points between the fully open and closed states exceeded the values
288 predicted from a double-Gaussian curve by the contribution of several substates.

289

Discussion

290 In this study, we used the patch-clamp technique on mouse split-open TAL tubules in order
291 to determine whether the apical 70-pS K channel exhibits the regulatory and conductive properties
292 of a $K_{Na}1.1$ channel. Basically designed and widely used for accessing apical ionic channels all along
293 the nephron, this approach also allowed the study of ionic channels in the lateral membrane of renal
294 cells (27). Since a large-conductance Na^+ - and Cl^- -activated, $K_{Na}1.1$ -like, potassium channel is already
295 functionally present in the basolateral membrane of mouse TAL cells (17) (19), it was of prime
296 importance to ensure the apical location of the channel studied here. However, by taking care to
297 precisely position the patch-clamp pipette tip onto the exposed apical membrane of cells within an
298 intact open area, none of the membrane patches exhibited the typical activity of Kir4.1/Kir5.1
299 potassium or ClC-K/Barttin chloride channels that could have revealed an accidental patch-clamping
300 of the lateral cell membrane (28) (29).

301 Our results show that the 70-pS K channel recapitulates the salient functional properties of
302 the basolateral $K_{Na}1.1$ -like and cloned $K_{Na}1.1$ channels. First, its activity increases with the internal
303 Na^+ concentration, with a relative EC_{50} of ~ 40 mM Na^+ and a n_H of ~ 2.5 that are within the ranges of
304 EC_{50} (8 to 100 mM Na^+) and n_H (2.4 to 4) values reported for the cloned $K_{Na}1.1$ channel (24) (25) (30)
305 (31) and in good agreement with the EC_{50} of 30 mM Na^+ and the n_H of ~ 4 we previously determined
306 for the basolateral channel (17). Second, the channel is also activated by internal Cl^- , with a relative
307 EC_{50} of ~ 30 mM Cl^- in the presence of an elevated internal Na^+ concentration that falls well within
308 the ranges of EC_{50} (8-35 mM Cl^-) reported for the basolateral and the cloned channels under similar
309 conditions (17) (24) (30), and a n_H of ~ 2 close to the n_H of ~ 1 reported for the basolateral channel
310 and that can be deduced from data on the cloned $K_{Na}1.1$ channel (17) (30). Furthermore, as observed
311 for the basolateral $K_{Na}1.1$ -like and cloned $K_{Na}1.1$ channels which are inactive in the absence of
312 internal Na^+ but exhibit a significant activity in a Cl^- -free medium (17) (22) (30), the presence of
313 internal Na^+ , but not that of Cl^- , is an absolute prerequisite for 70-pS K channel activity. Third, its
314 inhibition by a millimolar internal Ca^{2+} concentration, also observed for the basolateral channel (17),

315 is in accordance with that of the rat and *Drosophila* $K_{Na}1.1$ channels by a wide range of divalent
316 cations (25). And fourth, it exhibits the large conductance and multiple conductance substates that
317 biophysically characterize the cloned $K_{Na}1.1$ channels (22), and a rectification coefficient of 1.7 close
318 to the ratio of 1.8 we found for the basolateral channel (17).

319 The 70-pS K channel also exhibits much less documented properties of Na^+ -activated $K_{Na}1$
320 channels. Thus, its activity is inhibited by internal K^+ , a feature so far reported for only native $K_{Na}1$
321 channels in guinea-pig ventricular myocytes and in avian trigeminal ganglion neurons (26) (32), and
322 shared with the basolateral $K_{Na}1.1$ -like channel (this study). Here, internal K^+ concentration-
323 dependently decreased 70-pS K channel activity in the presence of a high- Na^+ medium with a relative
324 EC_{50} of around 70 mM K^+ and a n_H of 3.5 of the same order of magnitude as the EC_{50} of ~ 100 mM K^+
325 and the n_H of ~ 5 that can be deduced from published data in cardiac cells under similar conditions
326 (26). Similarly, we found that internal NH_4^+ , which is unable to replace Na^+ in activating the cloned
327 $K_{Na}1.1$ channel (24), inhibited 70-pS K channel activity. Although we did not establish a complete
328 dose-response curve for the inhibitory effect of internal NH_4^+ , our results indicate that it could be at
329 least as potent as K^+ . To our knowledge, this constitutes the first observation of a native $K_{Na}1$ channel
330 inhibited by both internal K^+ and NH_4^+ . Incidentally, the absence of current through the rat TAL 70-pS
331 K channel upon the substitution of K^+ or NH_4^+ for all internal Na^+ at that time suggested that K^+ and
332 NH_4^+ are not conducted by possibly acting as internal blockers of their own conductance (3). Our
333 results in mice show that the channel, while inhibited by a high internal K^+ or NH_4^+ concentration, is
334 actually able to conduct current in this condition.

335 Structurally, it is well established that $K_{Na}1.1$ channels are formed by the homotetrameric
336 association of $K_{Na}1.1-\alpha$ subunits. Indeed, size-exclusion chromatography experiments showed that
337 the purified chicken $K_{Na}1.1$ channel extended C-terminal domain elutes as a very stable
338 homotetrameric complex (33). In addition, the assembly of the C-terminus regulator of K^+
339 conductance (RCK) domain pairs of four $K_{Na}1.1-\alpha$ subunits is required to form the octameric ring
340 regulating the channel gate within which resides the Na^+ -binding site, and the crossing of the

341 transmembrane S6 domains of four $K_{Na}1.1-\alpha$ subunits forms part of the selectivity filter within the
342 ion conduction pathway (34). Nevertheless, as mentioned in the introduction, it is currently
343 acknowledged that the 70-pS K channel is formed by the heteromeric association of the ROMK
344 subunit with an unidentified one, each conferring specific properties (9). ROMK channel activity is not
345 sensitive to changes in internal Na^+ , Cl^- , K^+ or Ca^{2+} concentration in the rabbit TAL (5), and the present
346 study shows that the same holds true at least for Na^+ and Cl^- in the mouse TAL. Therefore, one can
347 reasonably infer that the sensitivity to internal ions of a 70-pS K channel structurally involving the
348 ROMK subunit would very likely have accounted for the contribution of the associated subunit. Na^+ -
349 activated potassium channels are endowed with a DxR/KxxH motif located in their C-terminal domain
350 that confers Na^+ -sensitivity (12). To our knowledge, TAL cells express only two types of potassium
351 channel subunits that possess such a motif, namely the Kir5.1 and $K_{Na}1.1-\alpha$ subunits (12) (28).
352 However, the observations that, among the renal cells expressing both ROMK and Kir5.1 subunits,
353 only TAL cells exhibit functional 70-pS K channels, and that these two subunits already do not
354 physically associate in vitro (35), make very unlikely the formation of a large-conductance Na^+ -
355 sensitive potassium channel by the association of ROMK and Kir5.1 subunits in native cells. We
356 obviously attempted to directly determine whether the ROMK and $K_{Na}1.1-\alpha$ subunits may
357 molecularly colocalize in the apical membrane of TAL cells and physically interact. Unfortunately,
358 none of the two commercially available anti-*Kcnt1* antibodies (clone N3/26) we tested gave a reliable
359 result (see the limitation section below). Nonetheless, indirect data from the literature argues
360 against a ROMK/ $K_{Na}1.1-\alpha$ subunits interaction. Indeed, supporting the notion that the
361 heteromerization of potassium channels subunits generally occurs between members of the same
362 family, the Kir-type subunit and the voltage-gated potassium channel Kv- α -type subunit, which
363 exhibits an overall structural organization in the transmembrane domains similar to that of the
364 $K_{Na}1.1-\alpha$ subunit but very distant from that of Kir- α subunits (36), do not coassemble with each other
365 to form heteromeric channels (37). Therefore, the $K_{Na}1.1$ -like 70-pS K channel in the apical
366 membrane of mouse TAL cells is most likely a channel structurally not involving the ROMK subunit.

367 The ROMK-independent structure of the 70-pS K channel obviously questions the mechanism
368 accounting for the requirement of *Kcnj1* for its functional expression leading to the complete
369 absence of apical ROMK and 70-pS K channels activity in *Kcnj1*-null mice. An interesting possibility
370 might be an indirect alteration in $K_{Na}1.1\text{-}\alpha$ protein synthesis or in the trafficking of the $K_{Na}1.1$ channel
371 upon *Kcnj1* deletion. Indeed, the excretion of Uromodulin (*UMOD*), or Tamm-Horsfall glycoprotein, is
372 significantly reduced in *Kcnj1*-null mice, after the pharmacological inhibition of the ROMK channel in
373 primary mouse TAL cells and in patients with Type 2 Bartter's syndrome (38). In addition, *Umod*
374 deficiency in mice decreases NKCC2 cotransporter, and ROMK and ClC/K channels mRNA levels in TAL
375 cells (39), and NKCC2 and ROMK proteins expression at the apical membrane by their defective
376 intracellular trafficking (39) (40) (41). Since a missense *Umod*^{R186S/+} mutation was recently shown to
377 cause a major retention of premature Uromodulin in the endoplasmic reticulum and to decrease the
378 amount of *Kcnt1* mRNA transcripts in mouse TAL cells (42), further experiments are needed in order
379 to determine whether the absence of functional apical 70-pS K channels in ROMK-deficient mice
380 might have been accounted for a *Kcnj1*-dependent, Uromodulin-mediated, defect in $K_{Na}1.1\text{-}\alpha$
381 channel protein expression or trafficking to the apical membrane.

382 To be functionally relevant, the sensitivity of the apical 70-pS K channel to internal ions
383 should occur in the ionic conditions that exist in TAL cells. However, channel sensitivity to internal
384 ions in cell-free membrane patches predicts a very low activity at the respective resting intracellular
385 Na^+ , K^+ and Cl^- concentrations of 15, 155 and 35 mM prevailing in TAL cells (43) (44) (45).
386 Furthermore, given the known complex interplay between internal Na^+ , Cl^- and K^+ in regulating native
387 $K_{Na}1$ channels in excitable cells or the cloned $K_{Na}1.1$ channel, it is very likely that the sensitivity of the
388 70-pS K channel to internal ions might have been underestimated. Indeed, the cloned $K_{Na}1.1$ channel is
389 cooperatively activated by intracellular Na^+ and Cl^- ions so that an increase in the internal
390 concentration of either of them increases the sensitivity to the other (30). Accordingly, the lowering
391 of internal Cl^- concentration from 145 mM to 40 mM causes a ~ 2.5-fold decrease in the sensitivity to
392 internal Na^+ of the basolateral $K_{Na}1.1$ -like channel (17). Conversely, an increase in internal K^+

393 decreases the sensitivity to internal Na^+ of native $\text{K}_{\text{Na}1}$ channels in excitable cells and vice versa (26)
394 (32). Therefore, one would expect a channel activity in situ even lower than that directly
395 extrapolated from our measurements. Yet, the 70-pS K channel exhibits the significant activity in cell-
396 attached membrane patches (NP_o of ~ 0.9 , see Table 1) previously observed by others for the channel
397 in mice and rats (2) (3) (4) (8) (9) (43). Such a paradox has been frequently reported regarding the
398 sensitivity of $\text{K}_{\text{Na}1}$ channels in excitable cells to internal Na^+ and early proposed to reflect its
399 modulation by an endogenous cytosolic factor(s) lost upon membrane patch excision (23). Very little
400 is currently known on the modulation of the sensitivity of native and cloned $\text{K}_{\text{Na}1}$ channels to internal
401 ions. Indeed, only one study reports a shift in the EC_{50} for Na^+ from 50 mM to 17 mM upon the direct
402 binding of the oxidized form of nicotinamide dinucleotide (NAD^+) to $\text{K}_{\text{Na}1}$ channels in rat dorsal root
403 ganglion neurons (46). Whether NAD^+ increases the sensitivity of the 70-pS K channel to internal Na^+
404 also in TAL cells, thereby making plausible its regulation by low cytosolic Na^+ concentrations in situ,
405 remains to be established. More generally, the identification of factors modulating the sensitivity of
406 the apical and basolateral $\text{K}_{\text{Na}1.1}$ -like channels not only to internal Na^+ but also to Cl^- , K^+ and NH_4^+ ,
407 whether they be a diffusible cytosolic factor or some other undefined process, should more precisely
408 define the conditions under which physiologically relevant ion concentrations control channels
409 activity in their native environment.

410 For now this uncertainty makes difficult to ascertain the role of intracellular ions as
411 regulators of $\text{K}_{\text{Na}1.1}$ -like channels activity in TAL cells. In neurons, the sensitivity of $\text{K}_{\text{Na}1}$ channels to
412 internal Na^+ is crucial to the control of electrical activity where the change in membrane potential
413 caused by the influx of Na^+ during action potentials is counteracted by the afterhyperpolarisation
414 generated by channels activation (47). In the TAL, apical and basolateral potassium channels are
415 essential to TAL transport function by establishing the potential differences across cell membranes
416 and generating the transepithelial potential difference (1). In addition, as emphasized by others (48),
417 the TAL is the only nephron segment where the large transcellular movements of Na^+ and Cl^-
418 associated with NaCl reabsorption cause massive influxes of K^+ via the apical NKCC2 cotransporter

419 and the basolateral Na^+/K^+ ATPase. In order to maintain steady-state intracellular K^+ concentration,
420 and hence the potential difference across apical and basolateral membranes, these K^+ influxes must
421 be coordinately counterbalanced by appropriate K^+ effluxes (*i.e.* K^+ recycling) through apical and
422 basolateral potassium channels. We previously proposed that the sensitivity of the basolateral $\text{K}_{\text{Na}1.1}$
423 channel to Na^+ and Cl^- may provide a signaling linkage between the entry and the basolateral exit of
424 K^+ (17). The identification of internal Na^+ and Cl^- as new regulators of the apical 70-pS K channel
425 suggests that a similar coupling between the entry of K^+ and its apical recycling through the channel
426 may exist. On the other hand, the inhibitory effect of internal K^+ on the activity of the apical and
427 basolateral channels in the context of NaCl reabsorption is puzzling since the associated increase in
428 intracellular K^+ concentration would decrease channels activity and hence antagonize the apical and
429 basolateral exits of K^+ . Thus, as in cardiac cells, $\text{K}_{\text{Na}1.1}$ channels activation in TAL cells might also take
430 place in specific pathological conditions, such as digitalis toxicity where the increase in internal Na^+
431 concentration is accompanied by a decrease in K^+ concentration in rat TAL cells (49), and protect
432 from damage by maintaining hyperpolarized the cell membranes. Finally, regarding the apical 70-pS
433 K channel, also intriguing is its inhibition by internal NH_4^+ . The TAL is an important site of NH_4^+
434 reabsorption and numerous studies established that apical NH_4^+ entry into TAL cells is mediated
435 predominantly by NKCC2 functioning as a $\text{Na}^+/\text{NH}_4^+/\text{2Cl}^-$ cotransporter (50). Our observations that the
436 channel is substantially permeable to NH_4^+ and able to conduct current in the presence of internal
437 NH_4^+ would make it in a position to also contribute to the luminal uptake of NH_4^+ . However, the
438 extent to which its inhibition by internal NH_4^+ , which may partly account for its reduced activity, and
439 lower apical membrane potassium conductance and voltage in intact rat TAL tubules lumenally
440 exposed to 10 mM NH_4^+ (51), may limit its contribution to the net NH_4^+ absorption by the TAL needs
441 further investigation.

442 A limitation to this study is the lack of strong and definite conclusion on the precise
443 molecular nature of the 70-pS K channel, since neither data from *Knct1*-null mice nor data using
444 molecular biology tools was provided. When combined, these data should in firm or confirm the

445 expression of the $K_{Na}1.1-\alpha$ subunit protein in TAL cells apical membrane, its involvement in forming
446 the channel and the ROMK-independent channel structure.

447 In conclusion, this study highlights new regulatory and conductive properties of the 70-pS K
448 channel in the apical membrane of mouse TAL cells that are most consistent with those of a $K_{Na}1.1$
449 channel. This strongly indicates that the ROMK subunit is not a structural component of the channel
450 and we suggest that the ROMK-dependent 70-pS K channel functional expression, which accounts for
451 the absence of apical potassium channels activity in *Kcnj1*-null mice and their severe salt-wasting
452 Type 2 Bartter's syndrome phenotype, may involve a mechanism by which *Kcnj1* expression possibly
453 affects *Kcnt1* transcription, $K_{Na}1.1-\alpha$ subunit protein synthesis or channel trafficking.

454 **Data availability:** data will be made available upon reasonable request.

455

456 **Supplemental material:** Supplemental Tables S1 and S2, and Figures S1 to S4:

457 <https://figshare.com/s/8f234b08cbf16ec42f02>

458 DOI : 10.6084/m9.figshare.24988854

459

460 **Acknowledgements:** we gratefully acknowledge the Functional Explorations Center (CEF) staff at the
461 Centre de Recherche des Cordeliers for technical assistance with animals.

462

463 **Grants:** this study was supported by grants from the French National Agency (ANR-14-CE12-0013-02
464 and ANR-17-CE14-0040). Elise de Combiens holds a Ph.D. fellowship from the French Ministère de la
465 Recherche et de l'Enseignement Supérieur.

466

467 **Disclosure:** no conflict of interest, financial or otherwise, is declared by the authors.

468

469 **Author contributions:** M.P. conceived and designed research; E. de C., N.F., L.C. and M.P. performed
470 experiments and analyzed the data; E. de C., N.F., L.C. and M.P. interpreted the results of
471 experiments; M.P. prepared figures; M.P. drafted the manuscript; E. de C., N.F., L.C., S.L. and M.P.
472 edited and revised the manuscript; E. de C., N.F., L.C., S.L. and M.P. approved the final version of the
473 manuscript.

- 475 1. **Mount DB.** Thick ascending limb of the loop of Henle. *Clin J Am Soc Nephrol* 9: 1974–1986,
476 2014. doi: 10.2215/CJN.04480413.
- 477 2. **Lu M, Wang W.** Two types of K(+) channels are present in the apical membrane of the thick
478 ascending limb of the mouse kidney. *Kidney Blood Press Res* 23: 75–82, 2000. doi:
479 10.1159/000025957.
- 480 3. **Bleich M, Schlatter E, Greger R.** The luminal K+ channel of the thick ascending limb of Henle’s
481 loop. *Pflugers Arch* 415: 449–460, 1990. doi: 10.1007/BF00373623.
- 482 4. **Wang WH.** Two types of K+ channel in thick ascending limb of rat kidney. *Am J Physiol* 267:
483 F599–605, 1994. doi: 10.1152/ajprenal.1994.267.4.F599.
- 484 5. **Wang WH, White S, Geibel J, Giebisch G.** A potassium channel in the apical membrane of
485 rabbit thick ascending limb of Henle’s loop. *Am J Physiol* 258: F244–253, 1990. doi:
486 10.1152/ajprenal.1990.258.2.F244.
- 487 6. **Guggino SE, Guggino WB, Green N, Sacktor B.** Ca²⁺-activated K+ channels in cultured
488 medullary thick ascending limb cells. *Am J Physiol* 252: C121–127, 1987. doi:
489 10.1152/ajpcell.1987.252.2.C121.
- 490 7. **Welling PA.** ROMK and Bartter Syndrome Type 2. In: *Ion Channels and Transporters of Epithelia*
491 *in Health and Disease*. Springer, New York, NY, p. 643–658.
- 492 8. **Wang W, Lu M.** Effect of arachidonic acid on activity of the apical K+ channel in the thick
493 ascending limb of the rat kidney. *J Gen Physiol* 106: 727–743, 1995. doi: 10.1085/jgp.106.4.727.
- 494 9. **Lu M, Wang T, Yan Q, Wang W, Giebisch G, Hebert SC.** ROMK is required for expression of the
495 70-pS K channel in the thick ascending limb. *Am J Physiol Renal Physiol* 286: F490–495, 2004.
496 doi: 10.1152/ajprenal.00305.2003.
- 497 10. **Lu M, Wang T, Yan Q, Yang X, Dong K, Knepper MA, Wang W, Giebisch G, Shull GE, Hebert SC.**
498 Absence of small conductance K+ channel (SK) activity in apical membranes of thick ascending
499 limb and cortical collecting duct in ROMK (Bartter’s) knockout mice. *J Biol Chem* 277: 37881–
500 37887, 2002. doi: 10.1074/jbc.M206644200.
- 501 11. **Manis AD, Hodges MR, Staruschenko A, Palygin O.** Expression, localization, and functional
502 properties of inwardly rectifying K+ channels in the kidney. *Am J Physiol Renal Physiol* 318:
503 F332–F337, 2020. doi: 10.1152/ajprenal.00523.2019.
- 504 12. **Rosenhouse-Dantsker A, Sui JL, Zhao Q, Rusinova R, Rodríguez-Menchaca AA, Zhang Z,
505 Logothetis DE.** A sodium-mediated structural switch that controls the sensitivity of Kir channels
506 to PIP₂. *Nat Chem Biol* 4: 624–631, 2008. doi: 10.1038/nchembio.112.
- 507 13. **Kaczmarek LK, Aldrich RW, Chandy KG, Grissmer S, Wei AD, Wulff H.** International Union of
508 Basic and Clinical Pharmacology. C. Nomenclature and Properties of Calcium-Activated and
509 Sodium-Activated Potassium Channels. *Pharmacol Rev* 69: 1–11, 2017. doi:
510 10.1124/pr.116.012864.
- 511 14. **Limbutara K, Chou C-L, Knepper MA.** Quantitative Proteomics of All 14 Renal Tubule Segments
512 in Rat. *J Am Soc Nephrol* 31: 1255–1266, 2020. doi: 10.1681/ASN.2020010071.

- 513 15. **Chen L, Chou C-L, Knepper MA.** A Comprehensive Map of mRNAs and Their Isoforms across All
514 14 Renal Tubule Segments of Mouse. *J Am Soc Nephrol* 32: 897–912, 2021. doi:
515 10.1681/ASN.2020101406.
- 516 16. **Onoda N, Kawabata A, Hasegawa K, Sakakura M, Urakawa I, Seki M, Zenkoh J, Suzuki A,**
517 **Suzuki Y.** Spatial and single-cell transcriptome analysis reveals changes in gene expression in
518 response to drug perturbation in rat kidney. *DNA Res* 29: dsac007, 2022. doi:
519 10.1093/dnares/dsac007.
- 520 17. **Paulais M, Lachheb S, Teulon J.** A Na⁺- and Cl⁻-activated K⁺ channel in the thick ascending limb
521 of mouse kidney. *J Gen Physiol* 127: 205–215, 2006. doi: 10.1085/jgp.200509360.
- 522 18. **Wu H, Kirita Y, Donnelly EL, Humphreys BD.** Advantages of Single-Nucleus over Single-Cell RNA
523 Sequencing of Adult Kidney: Rare Cell Types and Novel Cell States Revealed in Fibrosis. *J Am Soc*
524 *Nephrol* 30: 23–32, 2019. doi: 10.1681/ASN.2018090912.
- 525 19. **Fan L, Wang X, Zhang D, Duan X, Zhao C, Zu M, Meng X, Zhang C, Su X-T, Wang M-X, Wang W-**
526 **H, Gu R.** Vasopressin-induced stimulation of the Na⁺-activated K⁺ channels is responsible for
527 maintaining the basolateral K⁺ conductance of the thick ascending limb (TAL) in EAST/SeSAME
528 syndrome. *Biochim Biophys Acta* 1852: 2554–2562, 2015. doi: 10.1016/j.bbadis.2015.08.023.
- 529 20. **Kielland J.** Individual Activity Coefficients of Ions in Aqueous Solutions. *J Am Chem Soc* 59:
530 1675–1678, 1937. doi: 10.1021/ja01288a032.
- 531 21. **Hamer WJ, Wu Y.** Osmotic Coefficients and Mean Activity Coefficients of Uni-univalent
532 Electrolytes in Water at 25°C. *Journal of Physical and Chemical Reference Data* 1: 1047–1100,
533 1972. doi: 10.1063/1.3253108.
- 534 22. **Kaczmarek LK.** Slack, Slick and Sodium-Activated Potassium Channels. *ISRN Neurosci* 2013:
535 354262, 2013. doi: 10.1155/2013/354262.
- 536 23. **Dryer SE.** Na⁽⁺⁾-activated K⁺ channels: a new family of large-conductance ion channels. *Trends*
537 *Neurosci* 17: 155–160, 1994. doi: 10.1016/0166-2236(94)90093-0.
- 538 24. **Zhang Z, Rosenhouse-Dantsker A, Tang Q-Y, Noskov S, Logothetis DE.** The RCK2 Domain Uses a
539 Coordination Site Present in Kir Channels to Confer Sodium Sensitivity to Slo2.2 Channels.
540 *Journal of Neuroscience* 30: 7554–7562, 2010. doi: 10.1523/JNEUROSCI.0525-10.2010.
- 541 25. **Budelli G, Sun Q, Ferreira J, Butler A, Santi CM, Salkoff L.** SLO2 Channels Are Inhibited by All
542 Divalent Cations That Activate SLO1 K⁺ Channels*. *Journal of Biological Chemistry* 291: 7347–
543 7356, 2016. doi: 10.1074/jbc.M115.709436.
- 544 26. **Niu XW, Meech RW.** Potassium inhibition of sodium-activated potassium (KNa) channels in
545 guinea-pig ventricular myocytes. *J Physiol* 526: 81–90, 2000. doi: 10.1111/j.1469-
546 7793.2000.00081.x.
- 547 27. **Wang WH, McNicholas CM, Segal AS, Giebisch G.** A novel approach allows identification of K
548 channels in the lateral membrane of rat CCD. *Am J Physiol* 266: F813-822, 1994. doi:
549 10.1152/ajprenal.1994.266.5.F813.
- 550 28. **Hamilton KL, Devor DC.** Basolateral membrane K⁺ channels in renal epithelial cells. *Am J Physiol*
551 *Renal Physiol* 302: F1069-1081, 2012. doi: 10.1152/ajprenal.00646.2011.

- 552 29. **Teulon J, Planelles G, Sepúlveda FV, Andrini O, Lourdel S, Paulais M.** Renal Chloride Channels
553 in Relation to Sodium Chloride Transport. *Compr Physiol* 9: 301–342, 2018. doi:
554 10.1002/cphy.c180024.
- 555 30. **Yuan A, Santi CM, Wei A, Wang ZW, Pollak K, Nonet M, Kaczmarek L, Crowder CM, Salkoff L.**
556 The sodium-activated potassium channel is encoded by a member of the Slo gene family.
557 *Neuron* 37: 765–773, 2003. doi: 10.1016/s0896-6273(03)00096-5.
- 558 31. **Bhattacharjee A, Joiner WJ, Wu M, Yang Y, Sigworth FJ, Kaczmarek LK.** Slick (Slo2.1), a Rapidly-
559 Gating Sodium-Activated Potassium Channel Inhibited by ATP. *J Neurosci* 23: 11681–11691,
560 2003. doi: 10.1523/JNEUROSCI.23-37-11681.2003.
- 561 32. **Haimann C, Magistretti J, Pozzi B.** Sodium-activated potassium current in sensory neurons: a
562 comparison of cell-attached and cell-free single-channel activities. *Pflugers Arch* 422: 287–294,
563 1992. doi: 10.1007/BF00376215.
- 564 33. **Yuan P, Leonetti MD, Pico AR, Hsiung Y, MacKinnon R.** Structure of the Human BK Channel
565 Ca²⁺-Activation Apparatus at 3.0 Å Resolution. *Science* 329: 182–186, 2010. doi:
566 10.1126/science.1190414.
- 567 34. **Hite R, Yuan P, Li Z, Hsiung Y, Walz T, MacKinnon R.** Cryo-EM structure of the Slo2.2 Na⁺-
568 activated K⁺ channel. *Nature* 527: 198–203, 2015. doi: 10.1038/nature14958.
- 569 35. **Konstas A-A, Korbmacher C, Tucker SJ.** Identification of domains that control the heteromeric
570 assembly of Kir5.1/Kir4.0 potassium channels. *Am J Physiol, Cell Physiol* 284: C910-917, 2003.
571 doi: 10.1152/ajpcell.00479.2002.
- 572 36. **Hibino H, Inanobe A, Furutani K, Murakami S, Findlay I, Kurachi Y.** Inwardly rectifying
573 potassium channels: their structure, function, and physiological roles. *Physiol Rev* 90: 291–366,
574 2010. doi: 10.1152/physrev.00021.2009.
- 575 37. **Tytgat J, Buyse G, Eggermont J, Droogmans G, Nilius B, Daenens P.** Do voltage-gated Kv1.1 and
576 inward rectifier Kir2.1 potassium channels form heteromultimers? *FEBS Lett* 390: 280–284,
577 1996. doi: 10.1016/0014-5793(96)00674-6.
- 578 38. **Schiano G, Glaudemans B, Olinger E, Goelz N, Müller M, Loffing-Cueni D, Deschenes G, Loffing
579 J, Devuyst O.** The Urinary Excretion of Uromodulin is Regulated by the Potassium Channel
580 ROMK. *Sci Rep* 9: 19517, 2019. doi: 10.1038/s41598-019-55771-x.
- 581 39. **Bernascone I, Janas S, Ikehata M, Trudu M, Corbelli A, Schaeffer C, Rastaldi MP, Devuyst O,
582 Rampoldi L.** A transgenic mouse model for uromodulin-associated kidney diseases shows
583 specific tubulo-interstitial damage, urinary concentrating defect and renal failure. *Human
584 Molecular Genetics* 19: 2998–3010, 2010. doi: 10.1093/hmg/ddq205.
- 585 40. **Bachmann S, Mutig K, Bates J, Welker P, Geist B, Gross V, Luft FC, Alenina N, Bader M, Thiele
586 BJ, Prasad K, Raffi HS, Kumar S.** Renal effects of Tamm-Horsfall protein (uromodulin)
587 deficiency in mice. *American Journal of Physiology-Renal Physiology* 288: F559–F567, 2005. doi:
588 10.1152/ajprenal.00143.2004.
- 589 41. **Renigunta A, Renigunta V, Saritas T, Decher N, Mutig K, Waldegger S.** Tamm-Horsfall
590 glycoprotein interacts with renal outer medullary potassium channel ROMK2 and regulates its
591 function. *J Biol Chem* 286: 2224–2235, 2011. doi: 10.1074/jbc.M110.149880.

- 592 42. **Schiano G, Lake J, Mariniello M, Schaeffer C, Harvent M, Rampoldi L, Olinger E, Devuyst O.**
593 Allelic and Gene Dosage Effects Involving Uromodulin Aggregates Drive Autosomal Dominant
594 Tubulointerstitial Kidney Disease. *bioRxiv*: 2022.09.13.507770, 2022.
- 595 43. **Lu M, Zhu Y, Balazy M, Reddy KM, Falck JR, Wang W.** Effect of angiotensin II on the apical K⁺
596 channel in the thick ascending limb of the rat kidney. *J Gen Physiol* 108: 537–547, 1996. doi:
597 10.1085/jgp.108.6.537.
- 598 44. **Greger R, Weidtko C, Schlatter E, Wittner M, Gebler B.** Potassium activity in cells of isolated
599 perfused cortical thick ascending limbs of rabbit kidney. *Pflugers Arch* 401: 52–57, 1984. doi:
600 10.1007/BF00581532.
- 601 45. **Greger R, Oberleithner H, Schlatter E, Cassola AC, Weidtko C.** Chloride activity in cells of
602 isolated perfused cortical thick ascending limbs of rabbit kidney. *Pflugers Arch* 399: 29–34,
603 1983. doi: 10.1007/BF00652518.
- 604 46. **Tamsett TJ, Picchione KE, Bhattacharjee A.** NAD⁺ activates KNa channels in dorsal root
605 ganglion neurons. *J Neurosci* 29: 5127–5134, 2009. doi: 10.1523/JNEUROSCI.0859-09.2009.
- 606 47. **Cole BA, Clapcote SJ, Muench SP, Lippiat JD.** Targeting KNa1.1 channels in KCNT1-associated
607 epilepsy. *Trends in Pharmacological Sciences* 42: 700–713, 2021. doi:
608 10.1016/j.tips.2021.05.003.
- 609 48. **Sepúlveda FV, Pablo Cid L, Teulon J, Niemeyer MI.** Molecular aspects of structure, gating, and
610 physiology of pH-sensitive background K2P and Kir K⁺-transport channels. *Physiol Rev* 95: 179–
611 217, 2015. doi: 10.1152/physrev.00016.2014.
- 612 49. **Rajerison RM, Faure M, Morel F.** Effects of temperature, ouabain and diuretics on the cell
613 sodium and potassium contents of isolated rat kidney tubules. *Pflugers Arch* 406: 285–290,
614 1986. doi: 10.1007/BF00640915.
- 615 50. **Good DW.** Ammonium transport by the thick ascending limb of Henle's loop. *Annu Rev Physiol*
616 56: 623–647, 1994. doi: 10.1146/annurev.ph.56.030194.003203.
- 617 51. **Bleich M, Köttgen M, Schlatter E, Greger R.** Effect of NH₄⁺/NH₃ on cytosolic pH and the K⁺
618 channels of freshly isolated cells from the thick ascending limb of Henle's loop. *Pflugers Arch*
619 429: 345–354, 1995. doi: 10.1007/BF00374149.

620

621

Figures legends

622 **Figure 1. In situ apical potassium channels in the mouse TAL.** ROMK and 70-pS K channels properties
623 in cell-attached membrane patches with 145 mM KCl in the patch pipette, and 140 mM NaCl and 5
624 mM KCl in the bath. (a) Left : single-channel recordings in two separate membrane patches at the
625 indicated clamp voltage. C- is the current level corresponding to the closure of all channels, and O₁
626 and O₂ are the current levels corresponding to the opening of one and two channels, respectively.
627 Right : voltage-dependence of ROMK and 70-pS K channels activities. For each channel type, data
628 were normalized to the NP_o at resting membrane potential difference (V_c = 0 mV). Each point is the
629 mean of three (□) or seven (■) measurements and SD is shown as error bars. * P < 0.05 vs NP_o at V_c =
630 0 mV. (b) ROMK and 70-pS K channels *i*/V_c relationships in the conditions given in a. Each point is the
631 mean of three (□) or thirteen (■) measurements and SD is shown as error bars.

632

633 **Figure 2. Effect of internal Na⁺ removal on 70-pS K channel activity.** Continuous recording of 70-pS K
634 channel activity in a cell-excised inside-out membrane patch whose internal face is exposed to the
635 indicated Na⁺ concentrations (in mM). Bath K⁺ and Cl⁻ concentrations were kept at 9 and 145 mM,
636 respectively, and V_c was set at -40 mV. C- is the current level corresponding to the closure of all
637 channels. The lower left *a* and *b* traces are excerpts at an expanded time scale (*) taken from the
638 corresponding sections of the recording. The lower right graph summarizes NP_o data from seven
639 membrane patches, shown as individual values (□ and dashed lines) and as mean (■ and solid line)
640 with SD shown as error bars when larger than symbol. Data were normalized to the NP_o during the
641 initial exposure to 140 mM internal Na⁺. * P < 0.0001 vs NP_o = 1.

642

643 **Figure 3. Dependence of 70-pS K channel activity on internal Na⁺ concentration.** (a) Left : recordings
644 of channel activity in a cell-excised inside-out membrane patch at the indicated internal Na⁺
645 concentrations (in mM) in the conditions given in Figure 2, except V_c set at 0 mV. C- is the current
646 level corresponding to the closure of all channels and O₁ to O₆ are the current levels corresponding to

647 the opening of one to six channels. Right : relative amplitude histograms (0.2 pA bins) obtained at the
648 indicated internal Na^+ concentrations (in mM). (b) Dose-response relationship of the effect of
649 internal Na^+ concentration on channel activity in the conditions given in a. Data from four membrane
650 patches were normalized to the NP_o in the presence of 140 mM Na^+ , and are shown as individual
651 values (\square , $n = 4$ for each condition) and as mean (\blacksquare) with SD shown as error bars when larger than
652 symbol. The dashed line is a nonlinear fit of the Hill equation to the mean data (adjusted $R^2 = 0.999$).
653 * $P < 0.05$ vs $NP_o = 1$.

654
655 **Figure 4. Effect of low internal Cl^- on 70-pS K channel activity.** Continuous recording of 70-pS K
656 channel activity in a cell-excised inside-out membrane patch whose internal face is exposed to the
657 indicated Cl^- concentrations (in mM). Bath Na^+ and K^+ concentrations were kept at 140 and 9 mM,
658 respectively, and V_c was set at -40 mV. C- is the current level corresponding to the closure of all
659 channels. The lower left a and b traces are excerpts at an expanded time scale (*) taken from the
660 corresponding sections of the recording. The lower right graph summarizes NP_o data from nine
661 membrane patches, shown as individual values (\square and dashed lines) and as mean (\blacksquare and solid line)
662 with SD shown as error bars. Data were normalized to the NP_o during the initial exposure to 145 mM
663 internal Cl^- . * $P < 0.0001$ vs $NP_o = 1$.

664
665 **Figure 5. Dependence of 70-pS K channel activity on internal Cl^- concentration.** (a) Left : recordings
666 of channel activity in a cell-excised inside-out membrane patch at the indicated internal Cl^-
667 concentrations (in mM) in the conditions given in Figure 4. C- is the current level corresponding to
668 the closure of all channels, and O_1 is the current level corresponding to the opening of one channel.
669 Right : relative amplitude histograms (0.2 pA bins) obtained at the indicated internal Cl^-
670 concentrations (in mM). (b) Dose-response relationship of the effect of internal Cl^- concentration on
671 channel activity in the conditions given in a. Data from five membrane patches were normalized to
672 the NP_o in the presence of 145 mM Cl^- , and are shown as individual values (\square) and as mean (\blacksquare) with

673 SD shown as error bars. The number of observations in each condition is given in parentheses. The
674 dashed line is a nonlinear fit of the Hill equation to the mean data (adjusted $R^2 = 0.998$). * $P < 0.001$
675 vs $NP_o = 1$.

676

677 **Figure 6. Effect of internal Na^+ or Cl^- removal on ROMK channel activity.** Continuous recordings of
678 ROMK channel activity in two separate cell-excised inside-out membrane patches whose internal
679 face is exposed to the indicated Na^+ (**a**) or Cl^- (**b**) concentration (in mM). In **a**, bath K^+ and Cl^-
680 concentrations were kept at 9 and 145 mM, respectively, and V_c was set at -60 mV. In **b**, bath Na^+ and
681 K^+ concentrations were kept at 140 and 9 mM, respectively, and V_c was set at -40 mV. C- is the
682 current level corresponding to the closure of all channels. In **a** and **b**, the lower *a* and *b* traces are
683 excerpts at an expanded time scale (*) taken from the corresponding sections of the recording.

684

685 **Figure 7. Effect of elevated internal K^+ concentration on 70-pS K channel activity.** (a) Continuous
686 recording of channel activity in a cell-excised inside-out membrane patch whose internal face is
687 exposed to the indicated K^+ concentrations (in mM). Bath Na^+ and Cl^- concentrations were kept at
688 140 and 145 mM, respectively, and V_c was set at -60 mV. C- is the current level corresponding to the
689 closure of all channels. *a* and *b* are excerpts at an expanded time scale (*) taken from the indicated
690 sections of the recording. (b) Summary of NP_o data from eight membrane patches in the conditions
691 given in **a**, shown as individual values (\square and dashed lines) and as mean (\blacksquare and solid line) with SD
692 shown as error bars. Data were normalized to the NP_o during the initial exposure to 9 mM internal K^+ .
693 * $P < 0.0001$ vs $NP_o = 1$.

694

695 **Figure 8. Dependence of 70-pS K channel activity on internal K^+ concentration.** (a) Left : recordings
696 of channel activity in a cell-excised inside-out membrane patch at the indicated internal K^+
697 concentrations (in mM) in the conditions given in Figure 7. C- is the current level corresponding to
698 the closure of all channels and O_1 to O_5 are the current levels corresponding to the opening of one to

699 five channels. Right : relative amplitude histograms (0.2 pA bins) obtained at the indicated internal K^+
700 concentrations (in mM). (b) Dose-response relationship of the effect of internal K^+ concentration on
701 channel activity in the conditions given in a. Data from eight membrane patches were normalized to
702 the NP_o in the presence of 9 mM K^+ , and are shown as individual values (\square) and as mean (\blacksquare) with SD
703 shown as error bars. The number of observations in each condition is given in parentheses. The
704 dashed line is a nonlinear fit of the Hill equation to the mean data (adjusted $R^2 = 0.98$). * $P < 0.001$ vs
705 $NP_o = 1$.

706

707 **Figure 9. 70-pS K channel sensitivity to NH_4^+ .** Effect of internal NH_4^+ concentration on 70-pS K
708 channel activity. (a) Recordings of channel activity in two separate cell-excised inside-out membrane
709 patches with 145 mM K^+ in the pipette and 140 mM Na^+ in the bath and exposed to the indicated
710 NH_4^+ concentrations (in mM), at $V_c = 0$ mV. C- is the current level corresponding to the closure of all
711 channels. (b) Summary of the corresponding NP_o data, shown as individual values (\square and dashed
712 lines) and as mean (\blacksquare and solid line) with SD shown as error bars. Data were normalized to the NP_o
713 during the initial control (0 NH_4^+) condition. * $P < 0.05$ vs $NP_o = 1$, ** $P < 0.0001$ vs $NP_o = 0$.

714

715 **Figure 10. 70-pS K channel conductive properties.** 70-pS K channel i/V_c relationships in cell-excised
716 inside-out membrane patches with 145 mM K^+ in the patch pipette and 140 mM Na^+ in the bath, in
717 the presence of 9 mM internal K^+ (\blacksquare), 145 mM internal K^+ (\square) or 145 mM internal NH_4^+ (\circ). Each point
718 is the mean of five (\blacksquare) or four (\square , \circ) measurements and SD is shown as error bars when larger than
719 symbol. Under asymmetric K^+ concentration condition (\blacksquare), the solid line is the average of five non-
720 linear fits of the GHK equation to i/V_c curves (adjusted $R^2 = 0.942 - 0.98$). The inset shows the inward
721 slope conductance determined by the first derivative of the individual GHK data as function of V_c . For
722 clarity, only points at the applied clamp voltages are shown, with SD shown as error bars. The
723 continuous line is a non-linear fit of the Boltzmann equation to the mean first derivative data
724 (adjusted $R^2 = 0.998$). Under symmetric 145 mM K^+ concentration condition (\square), the dashed line is a

- 725 linear regression fit to inward currents data (adjusted $R^2 = 0.997$). With 145 mM NH_4^+ in the bath (o),
- 726 the dotted line is a fit of the GHK equation to the mean data (adjusted $R^2 = 0.998$).