



HAL
open science

Jet-cooled mid-infrared laser spectroscopy of centrosymmetric and n-bearings pahs

P Asselin, Sathapana Chawananon, Olivier Pirali, Manuel Goubet

► To cite this version:

P Asselin, Sathapana Chawananon, Olivier Pirali, Manuel Goubet. Jet-cooled mid-infrared laser spectroscopy of centrosymmetric and n-bearings pahs. The 27th International Conference on High-Resolution Molecular Spectroscopy, Université de Bologne, Italie, Sep 2024, Bologna (ITALY), Italy. hal-04793552

HAL Id: hal-04793552

<https://hal.sorbonne-universite.fr/hal-04793552v1>

Submitted on 20 Nov 2024

HAL is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers.

L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.

JET-COOLED MID-INFRARED LASER SPECTROSCOPY OF CENTROSYMMETRIC AND N-BEARINGS PAHS

P. ASSELIN, Sorbonne Université, CNRS, MONARIS, UMR 8233, ,
F-75005 Paris (France); **S. CHAWANANON**, **O. PIRALI**, Université
Paris-Saclay, CNRS, Institut des Sciences Moléculaires d'Orsay, 91405
Orsay (France); **M. GOUBET**, Université Lille, CNRS, UMR 8523 -
PhLAM - Physique des Lasers Atomes et Molecules, F-59000 Lille, (France)

The recent detection of cyano naphthalenes within TMC-1 using radioastronomy [1] provided the first unambiguous confirmation of the interstellar PAH's hypothesis and raised new questions concerning their formation and destruction pathways. In the mid-infrared (IR) domain, the launch of the James Webb Space Telescope opens exciting perspectives to collect rovibrational information about low volatile polycyclic aromatic compounds.

In this context, rotationally resolved IR studies of large aromatic species are still very scarce and mainly used synchrotron-based Fourier Transform (FT) spectroscopy coupled to room temperature long path cells [2]. However, their spectral analysis remains very challenging due to the congested rotational structures and the presence of many hot bands.

Nowadays, few set-ups combining high resolution (HR) IR spectroscopy to supersonic jets, using FT and cavity ring down spectroscopic methods were implemented to target PAH compounds [3]. Recently, a mid-IR laser spectrometer coupled to a pulsed jet (SPIRALES set-up at MONARIS) allows recording rovibrational spectra of large molecules at low temperature [4].

We report here the jet-cooled rovibrational IR study of centrosymmetric PAH molecules in both regions of ring C-H bending and C-C stretching vibrations to extract reliable spectroscopic parameters both in ground and excited vibrational states [5]. HR measurements of N-bearing PAHs with up to two N atoms substituted on the same ring evidenced abnormal Q-branch structures which make puzzling rovibrational analyses and suggests the presence of vibrational perturbations in spectral regions with high density of states.

Keywords — astrophysics, PAH, rovibrational spectroscopy, supersonic jets

References

- [1] B.A. Mc Guire, R.A. Loomis, A. M. Burkhardt, *et al. Science* **371**, 1265 (2021)
 - [2] S. Albert, K. K. Albert, P. Lerch *et al. Faraday Discuss.* **150**, 71 (2011)
 - [3] B.E. Brumfield, J. T. Stewart, B. J. McCall, *et al. J. Phys. Chem. Lett.* **3**, 1985 (2012)
O. Pirali, M. Goubet, T. Huet, *et al. Phys. Chem.. Chem. Phys.* **15**, 10141 (2013)
 - [4] P. Asselin, Y. Berger, T. Huet, *et al. Phys. Chem.. Chem. Phys.* **19**, 4576 (2017)
 - [5] S. Chawananon, O. Pirali, M. Goubet, *et al. J. Chem. Phys.* **157**, 064301 (2022)
-