



Human basophils may not undergo modulation by DC-SIGN and mannose receptor–targeting immunotherapies due to absence of receptors

Mrinmoy Das, Caroline Galeotti, Emmanuel Stephen-Victor, Anupama Karnam, Srini V. Kaveri, Jagadeesh Bayry

► To cite this version:

Mrinmoy Das, Caroline Galeotti, Emmanuel Stephen-Victor, Anupama Karnam, Srini V. Kaveri, et al.. Human basophils may not undergo modulation by DC-SIGN and mannose receptor–targeting immunotherapies due to absence of receptors. *Journal of Allergy and Clinical Immunology*, 2017, 139 (4), pp.1403-1404.e1. 10.1016/j.jaci.2016.09.062 . inserm-02455463v2

HAL Id: inserm-02455463

<https://hal.sorbonne-universite.fr/inserm-02455463v2>

Submitted on 25 Jan 2017

HAL is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers.

L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.

1 **Human basophils may not undergo modulation by DC-SIGN and mannose**
2 **receptor-targeting immunotherapies due to absence of receptors**

3

4 **Source of funding:** Supported by Institut National de la Santé et de la Recherche
5 Médicale (INSERM), Université Pierre et Marie Curie, Université Paris Descartes and
6 CSL Behring, Switzerland. C.G. is a recipient of fellowship from La Fondation pour
7 la Recherche Médicale (FDM20150633674), France and E.S-V and A.K. are
8 recipients of fellowships from Indo-French Center for Promotion of Advanced
9 Research.

10

11 **Conflict of Interest Declaration:** None

12

13 **Total word count: 503**

14

15 **Key words:** Basophils; DC-SIGN; CD206; dendritic cells; mannose receptor;
16 immunotherapy

17

18 **Abbreviations:**

19 CLEC12A: C-type lectin domain family 12 member A (CLEC12A)

20 DC: dendritic cells

21 DCIR: dendritic cell immunoreceptor

22 DC-SIGN: dendritic cell-specific intercellular adhesion molecule-3-grabbing
23 nonintegrin

24 TLR-Toll-like receptor

25

To the editor,

Sirvent et al., recently showed that novel vaccines targeting dendritic cells (DCs) by coupling glutaraldehyde-polymerized grass pollen allergoids to nonoxidized mannan enhance allergen uptake and induce functional regulatory T cells through programmed death ligand 1.¹ Mechanistically, they found that nonoxidized mannan-coupled glutaraldehyde-polymerized grass pollen allergoids are captured and internalized by two lectin receptors on DCs: mannose receptor (CD206) and DC-specific intercellular adhesion molecule-3-grabbing nonintegrin (DC-SIGN or CD209). These data thus indicated that DCs could be targeted by C-type lectin receptors for efficient allergen immunotherapy.

Basophils are one of the key players of allergic responses. They mediate allergic inflammation by secretion of Th2-polarizing cytokines IL-4, IL-13 and by the release of effector molecules like histamines and leukotrienes upon FcεRI signalling by IgE-allergen complexes.² Basophils receive activation signals not only via allergen-IgE complexes³ but also via toll-like receptors (TLRs)⁴ and possibly C-type lectin receptors. In fact, basophils express several lectin receptors like C-type lectin domain family 12 member A (CLEC12A) and dendritic cell immunoreceptor (DCIR).^{5,6} Thus, it is likely that in addition to DCs, nonoxidized mannan-coupled allergoids might also modulate basophil functions to exert immunotherapeutic benefits. Therefore, we analysed the expression of mannose receptor and DC-SIGN on steady state circulating human basophils and on stimulated basophils.

We analyzed basophils in whole blood of healthy donors without their purification in order to avoid any loss of cells and consequently misinterpretation of data (see Online Repository at www.jacionline.org). Further, erythrocyte-lysed whole blood cells were stimulated with IL-3 (100 ng/10⁶ cells) for 24 hours. IL-3-stimulated basophils were

also stimulated for degranulation with anti-IgE antibodies (100 ng/10⁶ cells) for 30 minutes. As controls for the expression of DC-SIGN and CD206, we used CD14⁺ peripheral blood monocytes (negative control), and rhIL-4 (500 IU/10⁶ cells) and rhGM-CSF (1000 IU/10⁶ cells)-differentiated monocyte-derived DCs (positive control).⁷

Circulating basophils were identified as positive for FcεRI and CD123 and negative for BDCA-4. We found that human basophils at steady state are negative for DC-SIGN and CD206 (Fig.1A-B). As basophils display enhanced expression of various receptors upon receiving activation stimuli, we explored if they express these lectin receptors upon activation. However, irrespective of stimulation (IL-3 or degranulation stimuli) basophils remained negative for DC-SIGN and CD206 (Fig.1B-C). Absence of DC-SIGN was also confirmed on isolated basophils. Further, the absence of DC-SIGN and CD206 on basophils in our report is not due to non-reactivity of antibodies used in the flow-cytometry as monocyte-derived DCs, used as positive control, uniformly expressed CD206 and DC-SIGN (Fig.1A-B). As expected, CD14⁺ circulating monocytes, used as negative control, did not stain for both the markers, thus confirming lack of non-specific binding of antibodies (Fig.1A-B).

Our results thus indicate that human basophils lack DC-SIGN and mannose receptors and hence unlike DCs, they may not directly respond and modulated by DC-SIGN- and mannose receptor-binding nonoxidized mannan-coupled allergoids. In addition, our data also suggest that basophils do not get activated by DC-SIGN- and mannose receptor-binding allergens unless they are IgE-bound. Thus, expression pattern of DC-SIGN and mannose receptor among innate cells diversifies allergic as well as tolerogenic responses.

Mrinmoy Das, MSc,^{a,b,c,*}, Caroline Galeotti, MD, ^{a,b,c,d,*}, Emmanuel Stephen-Victor, MSc, ^{a,b,c,*}, Anupama Karnam, MSc, ^{a,b,c}, Srini V Kaveri, DVM, PhD, ^{a,b,c,e} and Jagadeesh Bayry, DVM, PhD ^{a,b,c,e}

^a Institut National de la Santé et de la Recherche Médicale Unité 1138, Paris, F-75006, France

^b Sorbonne Universités, UPMC Univ Paris 06, UMR S 1138, Paris, F-75006, France

^c Centre de Recherche des Cordeliers, Equipe - Immunopathologie et immuno-intervention thérapeutique, Paris, F-75006, France

^d Department of Pediatric Rheumatology, National Referral Centre of Auto-inflammatory Diseases, CHU de Bicêtre, le Kremlin Bicêtre, F-94270, France

^e Université Paris Descartes, Sorbonne Paris Cité, UMR S 1138, Paris, F-75006, France

* These authors contributed equally to this work

Correspondence to: Jagadeesh Bayry, DVM, PhD, Institut National de la Santé et de la Recherche Médicale Unité 1138, Centre de Recherche des Cordeliers, 15 rue de l'Ecole de Médecine, Paris, F-75006, France. Tel: 00 33 1 44 27 82 03; Fax: 00 33 1 44 27 81 94




E-mail: jagadeesh.bayry@crc.jussieu.fr

REFERENCES

1. Sirvent S, Soria I, Cirauqui C, Cases B, Manzano AI, Diez-Rivero CM et al. Novel vaccines targeting dendritic cells by coupling allergoids to nonoxidized mannan enhance allergen uptake and induce functional regulatory T cells through programmed death ligand 1. *J Allergy Clin Immunol.* 2016;138:558-67 e11.
2. Voehringer D. Protective and pathological roles of mast cells and basophils. *Nat Rev Immunol.* 2013;13:362-75.
3. Oettgen HC. Fifty years later: Emerging functions of IgE antibodies in host defense, immune regulation, and allergic diseases. *J Allergy Clin Immunol.* 2016;137:1631-45.
4. Suurmond J, Stoop JN, Rivellese F, Bakker AM, Huizinga TW, Toes RE. Activation of human basophils by combined toll-like receptor- and FcepsilonRI-triggering can promote Th2 skewing of naive T helper cells. *Eur J Immunol.* 2014;44:386-96.
5. Bates EE, Fournier N, Garcia E, Valladeau J, Durand I, Pin JJ, et al. APCs express DCIR, a novel C-type lectin surface receptor containing an immunoreceptor tyrosine-based inhibitory motif. *J Immunol.* 1999;163:1973-83.
6. Divekar A, Conklin J, Yang X, Ransom J. Role of C-type lectin receptor CLEC12A in human basophil activation (48.6). *J Immunol.* 2012;188:48.6.
7. Maddur MS, Sharma M, Hegde P, Stephen-Victor E, Pulendran B, Kaveri SV, et al. Human B cells induce dendritic cell maturation and favour Th2 polarization by inducing OX-40 ligand. *Nat Commun.* 2014;5:4092.

Figure Legend

FIG.1. Human basophils are deficient for DC-SIGN and CD206. **A, B, C,** Flow cytometric analysis of DC-SIGN and CD206 on steady state basophils, stimulated basophils (IL-3 or IL-3 and anti-IgE), monocytes and monocyte-derived DCs. Representative dot-plots and percentage of cells (mean \pm SEM, n= 3) positive for DC-SIGN and CD206. ***P < .001.



142

143

Online Repository

**Human basophils are deficient for the expressions of DC-SIGN and mannose
receptor**

**Mrinmoy Das, MSc,^{a,b,c,*}, Caroline Galeotti, MD,^{a,b,c,d,*}, Emmanuel Stephen-
Victor, MSc,^{a,b,c,*}, Anupama Karnam, MSc,^{a,b,c}, Srini V Kaveri, DVM, PhD,
^{a,b,c,d} and Jagadeesh Bayry, DVM, PhD^{a,b,c,d}**

^a Institut National de la Santé et de la Recherche Médicale Unité 1138, Paris, F-75006,
France

^b Sorbonne Universités, UPMC Univ Paris 06, UMR S 1138, Paris, F-75006, France

^c Centre de Recherche des Cordeliers, Equipe - Immunopathologie et immuno-
intervention thérapeutique, Paris, F-75006, France

^d Department of Pediatric Rheumatology, National Referral Centre of Auto-
inflammatory Diseases, CHU de Bicêtre, le Kremlin Bicêtre, University of Paris Sud,
F-94270, France

^e Université Paris Descartes, Sorbonne Paris Cité, UMR S 1138, Paris, F-75006, France

* These authors contributed equally to this work

Correspondence to: Jagadeesh Bayry, DVM, PhD, Institut National de la Santé et de
la Recherche Médicale Unité 1138, Centre de Recherche des Cordeliers, 15 rue de
l'Ecole de Médecine, Paris, F-75006, France. Tel: 00 33 1 44 27 82 03; Fax: 00 33 1
44 27 81 94

E-mail: jagadeesh.bayry@crc.jussieu.fr

Total word count: 302

METHODS

Cells and stimulation

Buffy bags of healthy donors were obtained from Centre Necker-Cabanel (EFS, Paris) and INSERM-EFS ethical committee approval (N°15/EFS/012) for the use of such material was obtained. Experiments were performed in accordance with the approved guidelines of INSERM.

Red blood cells were lysed using ACK (Ammonium-Chloride-Potassium) Lysing Buffer (Lonza). Briefly, blood was spun down and resuspended and incubated in ACK lysing buffer for 30-60 seconds. Cells were washed with medium and resuspended in serum-free X-VIVO medium. Cells were stimulated with IL-3 (100 ng/million cells; ImmunoTools) for 24 hours. In addition, cells were also cultured with IL-3 (100 ng/million cells) for up to 24 hours and during last 30 minutes, cells were treated with anti-IgE antibodies (100 ng/million cells; Sigma-Aldrich). Phenotype of basophils was analysed in steady state and stimulated conditions by flow cytometry (LSR II, BD Biosciences) and the data was analyzed using FACSDiva™ software (BD Biosciences).

Peripheral blood mononuclear cells (PBMCs) were obtained from buffy bags of healthy donors by Ficoll density gradient centrifugation. Monocytes were isolated from peripheral blood mononuclear cells by using CD14 microbeads (Miltenyi Biotec) and were cultured for 5 days in rhIL-4 (500 IU/10⁶ cells) and rhGM-CSF (1000 IU/10⁶ cells) (both from Miltenyi Biotec) to obtain monocyte-derived DCs.

Antibodies for the flow cytometry

The following antibodies were used for the flow cytometry. FcεRIα-BV510 (Clone: AER37 (CRA-1)) was from BioLegend and BDCA-4 (CD304)-APC (Clone: AD5-

17F6) was obtained from Miltenyi Biotec. CD123-BV421 (Clone: 9F5), CD209-FITC (Clone: DCN46), CD206-PE (Clone: 19.2), HLA-DR-APC or PE (G46-6, BD Biosciences) and CD14-APC (Clone: M5E2) antibodies were from BD Biosciences

Statistical analysis

Levels of significance for comparison between samples were determined by One-way analysis of variance (repeated measures with Tukey's multiple comparison test). P<0.05 was considered significant. Statistical analysis was performed by Prism 5 GraphPad Software. Data are presented as mean \pm SEM.